

ANNALES DE L'INSTITUT FOURIER

BAHMAN SAFFARI

R. C. VAUGHAN

On the fractional parts of x/n and related sequences. I

Annales de l'institut Fourier, tome 26, n° 4 (1976), p. 115-131

[<http://www.numdam.org/item?id=AIF_1976__26_4_115_0>](http://www.numdam.org/item?id=AIF_1976__26_4_115_0)

© Annales de l'institut Fourier, 1976, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE FRACTIONAL PARTS OF x/n AND RELATED SEQUENCES. I

by **B. SAFFARI** and **R. C. VAUGHAN**

1. Introduction.

1. Throughout this paper $\{x\} = x - [x]$ denotes the fractional part of the real number x . We write $\|x\| = \min_{k \in \mathbf{Z}} |x - k|$ and $e(x) = e^{2\pi i x}$.

Also, the implied constants in the O symbol of Landau and the \gg and \ll symbols of Vinogradov are absolute.

Finally, by a distribution function we always mean a distribution function in the sense of probability theory, defined on the real line.

2. Let (x_n) be a sequence of real numbers. The usual study of the distribution modulo 1 of (x_n) is essentially that of the distribution of the sequence $(e(x_n))$ on the circle \mathbf{T} . The main problems are those of investigating

(i) the existence of the asymptotic (or limit) distribution measure

$$(1.1) \quad \mu = \lim_{k \rightarrow \infty} \mu_k$$

where

$$(1.2) \quad \mu_k = \frac{1}{k} \sum_{n=1}^k \delta_{e(x_n)}$$

with δ_v denoting the Dirac measure at $v \in \mathbf{T}$, and

(ii) the size of the discrepancy

$$(1.3) \quad \sup_{\omega} |\mu_k(\omega) - \mu(\omega)|$$

where ω runs through those arcs of \mathbf{T} whose end points have μ -measure zero.

It is classical that the existence of μ together with the assumption that the point $1 \in \mathbf{T}$ has μ -measure zero is equivalent to the existence of a distribution function F such that

$$(1.4) \quad F(0+) = 0, \quad F(1-) = 1$$

and

$$(1.5) \quad F(\alpha) = \lim_{k \rightarrow \infty} \frac{1}{k} A([0, \alpha), k, (x_n))$$

at every α at which F is continuous, the counting function

$$(1.6) \quad A([\alpha, \beta), k, (x_n)) \\ = \text{Card} \{n : 1 \leq n \leq k, \alpha \leq \{x_n\} < \beta\}$$

being here defined for all real numbers α and β . The conditions (1.4) mean that F is continuous at 0 and 1, and imply that F is constant on the intervals $(-\infty, 0]$ and $[1, \infty)$. In this case F is called the asymptotic (or limit) distribution function modulo 1 of the sequence (x_n) , and the discrepancy (1.3) is equal to

$$(1.7) \quad \sup_{0 \leq \alpha < \beta \leq 1} \left| \frac{1}{k} A([\alpha, \beta), k, (x_n)) - (F(\beta) - F(\alpha)) \right|$$

where α and β run through the continuity points of F .

In some situations it may be more appropriate to consider the existence of the A -asymptotic distribution function modulo 1, namely the existence (outside a countable set), and the continuity at $\alpha = 0$ and $\alpha = 1$, of

$$(1.8) \quad \lim_{k \rightarrow \infty} \sum_{n=1}^k a_{k,n} c_\alpha(x_n)$$

where

$$(1.9) \quad c_\alpha(u) = \begin{cases} 1 & 0 \leq \{u\} < \alpha \\ 0 & \text{otherwise} \end{cases}$$

is the characteristic function modulo 1 of $[0, \alpha)$, and $A = (a_{k,n})$ is a positive Toeplitz matrix. Here by a positive Toeplitz matrix we mean that

$$a_{k,n} \geq 0, \quad \sum_{n=1}^{\infty} a_{k,n} < \infty \quad \text{and} \quad \lim_{k \rightarrow \infty} \sum_{n=1}^{\infty} a_{k,n} = 1.$$

3. The sequence (x_n) is, of course, independent of k . Our object is to investigate the distribution modulo 1 of $xh(n)$ with x a large real number, $h(n)$ an arithmetical function, and the integer n belonging to $S \cap [1, k]$ where $S \subset \mathbb{N}$ and k depends on x . For our purposes it is somewhat more convenient to replace k by a real parameter y . We call $\mathcal{A} = (a_n(y) : y \in [1, \infty), n = 1, 2, \dots)$ a positive Toeplitz transformation if $a_n(y) \geq 0$ for all n and y , $\sum_{n=1}^{\infty} a_n(y) < \infty$ for every y , and $\lim_{y \rightarrow \infty} \sum_{n=1}^{\infty} a_n(y) = 1$. We are particularly interested in the special case where the $a_n(y)$ are the simple Riesz means (R, λ_n) given by

$$(1.10) \quad \lambda_n \geq 0 \quad (n = 1, 2, \dots), \quad \lambda_1 > 0$$

and

$$(1.11) \quad a_n(y) = \begin{cases} \lambda_n / \sum_{m \leq y} \lambda_m & (m \leq y) \\ 0 & (m > y) \end{cases}$$

which we assume henceforward, although several of our proofs go through in the general case (see Appendix). Let

$$(1.12) \quad \Phi_{x,y}(\alpha, h) = \sum_{n=1}^{\infty} a_n(y) c_{\alpha}(xh(n)).$$

A good deal of our attention will be taken up with $h(n) = 1/n$ and we write

$$(1.13) \quad \Phi_{x,y}(\alpha) = \sum_{n=1}^{\infty} a_n(y) c_{\alpha}(x/n).$$

The problems arising from the study of $\Phi_{x,y}(\alpha)$ as x and $y = y(x)$ tend together to infinity are closely related to the Dirichlet divisor problem.

If there exists a distribution function Φ_h such that

$$(1.14) \quad \Phi_h(0+) = 0, \quad \Phi_h(1-) = 1$$

and

$$(1.15) \quad \Phi_h(\alpha) = \lim_{x \rightarrow \infty} \Phi_{x,y(x)}(\alpha, h)$$

at every α at which Φ_h is continuous, then we call Φ_h the

\mathcal{A} -asymptotic distribution function modulo 1. This situation is equivalent to the existence on the circle \mathbf{T} of the \mathcal{A} -limit (or \mathcal{A} -asymptotic) distribution measure

$$(1.16) \quad \nu = \lim_{x \gg \infty} \sum_{n=1}^{\infty} a_n(y) \delta_{e(xh(n))}$$

together with the fact that the point $1 \in \mathbf{T}$ has ν -measure zero. However, if there exists no distribution function Φ_h satisfying both (1.14) and (1.15), then it is more appropriate to investigate the distribution modulo 1 of $xh(n)$ via (1.16).

4. Our interest in this problem arose from investigating the asymptotic behaviour of

$$\sum_{n \leq y} c_\alpha(x/n).$$

During our investigation it became obvious that there were methods which could be applied in a much more general situation. In this paper we present these methods, deferring to the sequel the study of special methods.

As an example of the application of Theorem 2, consider a subset A of \mathbf{N}^* such that the counting function

$$A(x) = \sum_{\substack{a \leq x \\ a \in A}} 1$$

satisfies

$$A(x) = x^\sigma L(x)$$

where σ is a constant with $0 < \sigma \leq 1$ and L is a slowly varying function, that is

$$\lim_{x \gg \infty} \frac{L(cx)}{L(x)} = 1$$

for any positive constant c . Then

$$(1.17) \quad \lim_{x \gg \infty} \frac{1}{A(x)} \sum_{\substack{a \leq x \\ a \in A}} c_\alpha(x/a) = \sum_{n=1}^{\infty} (n^{-\sigma} - (n+\alpha)^{-\sigma}).$$

Moreover, there exists a function $y_0(x)$ such that if $y > y_0(x)$

and $y = o(x)$ as $x \rightarrow \infty$, then

$$(1.18) \quad \lim_{x \rightarrow \infty} \frac{1}{A(y)} \sum_{\substack{a \leq y \\ a \in A}} c_a(x/a) = \alpha.$$

Relation (1.18) means that the fractional parts $\{x/a\}$, where a runs over $[0, y] \cap A$, are asymptotically uniformly distributed, whereas (1.17) means that if a runs over the whole of $[0, x] \cap A$, then the $\{x/a\}$ have the asymptotic distribution function

$$\sum_{n=1}^{\infty} (n^{-\sigma} - (n + \alpha)^{-\sigma}).$$

2. Theorems and proofs.

1. We first of all state a theorem which gives a sufficient condition for the (R, λ_n) -asymptotic distribution to be uniform. This is essentially due to Erdős and Turan [1], [2] and is a finite form of Weyl's criterion. It is also possible, of course, to give a necessary condition corresponding to Weyl's criterion, and to give results when the asymptotic distribution is non-uniform but continuous, but we have no applications in mind for these.

Theorem 1 is somewhat divorced from the following theorems. However, it clearly applies to the general situation. As an application we have in mind the case

$$(2.1) \quad h(n) = \log n.$$

THEOREM 1. — *Let the discrepancy $D_{x,y}(h)$ be defined by*

$$(2.2) \quad D_{x,y}(h) = \sup_{0 \leq \alpha < \beta \leq 1} |\Phi_{x,y}(\beta, h) - \Phi_{x,y}(\alpha, h) - (\beta - \alpha)|.$$

Then, for any positive integer m ,

$$(2.3) \quad D_{x,y}(h) < \frac{6}{m+1} + \frac{4}{\pi} \sum_{k=1}^m \left(\frac{1}{k} - \frac{1}{m+1} \right) \left| \sum_{n=1}^{\infty} a_n(y) e(kxh(n)) \right|.$$

Theorem 1 is a generalization of Theorem 2.2.5 of Kuipers and Niederreiter [3], and can be proved in exactly the same way.

2. The following theorem (together with the observations made in Lemmas 2, 3, 4) shows that the (R, λ_n) asymptotic distribution function modulo 1 of x/n can exist under very general conditions provided that y is not too small compared with x .

Whenever $\xi \geq 1$ and $\sigma \geq 0$ define

$$(2.4) \quad F(\alpha; \xi, \sigma) = \begin{cases} 0 & (\alpha \leq 0) \\ 1 & (\alpha \geq 1) \\ \theta(\alpha; \xi)(1 - \xi^\sigma([\xi] + \alpha)^{-\sigma} \\ \quad + \xi^\sigma \sum_{k > \xi} (k^{-\sigma} - (k + \alpha)^{-\sigma}) & (0 < \alpha < 1, \sigma > 0) \\ \alpha & (0 < \alpha < 1, \sigma = 0) \end{cases}$$

where

$$(2.5) \quad \theta(\alpha; \xi) = \begin{cases} 1 & \text{if } (\xi - \alpha, \xi] \cap \mathbf{N} \neq \emptyset \\ 0 & \text{otherwise.} \end{cases}$$

THEOREM 2. — Suppose that for every real number t with $0 < t < 1$ the limit

$$(2.6) \quad \lim_{y \rightarrow \infty} \sum_{n \leq ty} a_n(y)$$

exists and for at least one value of t is non-zero. Then there is a non-negative real number σ such that for every real number ε with $0 < \varepsilon < \frac{1}{2}$ there is a real number $y_0(\varepsilon, \sigma) \geq 1$ so that whenever $y_0(\varepsilon, \sigma) \leq y \leq x$ we have

$$(2.7) \quad \Phi_{x,y}(\alpha) = F(\alpha; x/y, \sigma) + O(\varepsilon^{1+\sigma}xy^{-1}) + O(2^\sigma\varepsilon^\sigma).$$

Lemma 1 below will show that the limit (2.6) is t^σ , which defines σ . We observe that when $\sigma = 0$ Theorem 2 fails to give non-trivial information. Very likely $\Phi_{x,y}(\alpha) \rightarrow \alpha$ still holds in this case, at least when $\sum_{n \leq y} \lambda_n \rightarrow \infty$, but even when $\lambda_n = 1/n$ this is a deep result.

Before proceeding with the proof of Theorem 2 we state a corollary concerning the case when the integer n is allowed only to run through a shorter interval $[y, z]$.

COROLLARY 2.1. — *With the assumptions of Theorem 2, if*

$$y_0(\varepsilon, \sigma) \leq y < z \leq x/2, (y/z)^\sigma < 1 - \varepsilon^{2+\sigma}, \varepsilon^z \leq y,$$

and $\sum_{y < n \leq z} \lambda_n > 0$, then

$$(2.8) \quad \frac{\sum_{y < n \leq z} \lambda_n c_\alpha(x/n)}{\sum_{y < n \leq z} \lambda_n} - \alpha \\ \ll (\sigma 2^\sigma z x^{-1} + \varepsilon^{1+\sigma} x y^{-1} + 2^\sigma \varepsilon^\sigma) (1 - y^\sigma z^{-\sigma} - \varepsilon^{2+\sigma})^{-1}.$$

We remark that, in this case, the asymptotic distribution is always the uniform one, at least when $\sigma > 0$.

3. The proof of Theorem 2 requires the following lemma.

LEMMA 1. — *On the hypothesis of Theorem 2 there is a non-negative real number σ such that for every real number ε with $0 < \varepsilon < 1/2$ there is a real number $y_0(\varepsilon, \sigma) \geq 1$ so that whenever $y \geq y_0(\varepsilon, \sigma)$ we have, for every t with $\varepsilon \leq t \leq 1$,*

$$(2.9) \quad \left| t^\sigma - \sum_{n \leq ty} a_n(y) \right| < \varepsilon^{2+\sigma}.$$

Proof. — The existence of (2.6) for every real number t with $0 < t < 1$ together with the assumption that for some t in this range the limit is non-zero imply that there is a non-negative real number σ such that for every t with $0 < t \leq 1$ we have

$$\lim_{y \rightarrow \infty} \sum_{n \leq ty} a_n(y) = t^\sigma.$$

Let

$$N = [2e^{\varepsilon^{-2-\sigma}} \max(1, \sigma)] + 1$$

and choose $y_0(\varepsilon, \sigma) \geq 1$ so that if $y \geq y_0(\varepsilon, \sigma)$, then for every integer r with $1 \leq r \leq N$ we have

$$(2.10) \quad \left| \left(\frac{r}{N} \right)^\sigma - \sum_{n \leq ry/N} a_n(y) \right| < \frac{1}{2} \varepsilon^{2+\sigma}.$$

Now choose an integer q such that

$$(2.11) \quad \frac{1}{N} \leq \frac{q}{N} < t \leq \frac{q+1}{N} \leq 1,$$

which is always possible if $\varepsilon \leq t \leq 1$. Note that

$$\begin{aligned} \left(\frac{q+1}{N}\right)^\sigma - \left(\frac{q}{N}\right)^\sigma &= \int_{q/N}^{(q+1)/N} \sigma u^{\sigma-1} du \\ &\leq \frac{\sigma}{N} \max \left(\left(\frac{q+1}{N}\right)^{\sigma-1}, \left(\frac{q}{N}\right)^{\sigma-1} \right) \\ &\leq \sigma \max(N^{-1}, N^{-\sigma}) \leq \max(\sigma N^{-1}, (e \log N)^{-1}) \\ &< \frac{1}{2} \varepsilon^{2+\sigma}. \end{aligned}$$

Thus, by (2.10) and (2.11),

$$\begin{aligned} \sum_{n \leq t\gamma} a_n(y) &\leq \sum_{n \leq (q+1)\gamma/N} a_n(y) < \left(\frac{q+1}{N}\right)^\sigma + \frac{1}{2} \varepsilon^{2+\sigma} \\ &< \left(\frac{q}{N}\right)^\sigma + \varepsilon^{2+\sigma} \leq t^\sigma + \varepsilon^{2+\sigma} \end{aligned}$$

and

$$\begin{aligned} \sum_{n \leq t\gamma} a_n(y) &\geq \sum_{n \leq q\gamma/N} a_n(y) > \left(\frac{q}{N}\right)^\sigma - \frac{1}{2} \varepsilon^{2+\sigma} \\ &> \left(\frac{q+1}{N}\right)^\sigma - \varepsilon^{2+\sigma} \geq t^\sigma - \varepsilon^{2+\sigma}. \end{aligned}$$

These last two inequalities give (2.9) as required.

4. Proof of Theorem 2. — Since (2.7) is trivially true when $\alpha \leq 0$ or $\alpha \geq 1$, we may assume $0 < \alpha < 1$. Let

$$K = \left\lfloor \frac{x}{\varepsilon y} - \alpha \right\rfloor.$$

Then, by (1.13), (1.11), (1.9), Lemma 1 and (2.5),

$$\begin{aligned} \Phi_{x,y}(\alpha) &= \sum_{\substack{\frac{x}{K+\alpha} < n \leq y \\ n \leq y}} a_n(y) c_\alpha(x/n) + O\left(\sum_{n \leq 2\varepsilon y} a_n(y)\right) \\ &= \sum_{k=1}^K \sum_{\substack{n \leq y \\ x/(k+\alpha) < n \leq x/k}} a_n(y) + O(2^\sigma \varepsilon^\sigma) \\ &= \theta(\alpha; x/y) \left(\sum_{n \leq y} a_n(y) - \sum_{n \leq x/([x/y] + \alpha)} a_n(y) \right) \\ &\quad + \sum_{x/y \leq k \leq K} \left(\sum_{n \leq x/k} a_n(y) - \sum_{n \leq x/(k+\alpha)} a_n(y) \right) + O(2^\sigma \varepsilon^\sigma). \end{aligned}$$

Hence, by Lemma 1 and (2.4),

$$\Phi_{x,y}(\alpha) = F(\alpha; x/y, \sigma) + O(\varepsilon^{2+\sigma}K) + O(2^\sigma \varepsilon^\sigma) \\ + O\left(\sum_{k>K} \left(\frac{x}{y}\right)^\sigma (k^{-\sigma} - (k+\alpha)^{-\sigma})\right).$$

The proof of (2.7) is completed by observing that $\varepsilon K \leq x/y$ and

$$\sum_{k>K} (k^{-\sigma} - (k+\alpha)^{-\sigma}) = \sum_{k>K} \int_k^{k+\alpha} \sigma u^{-\sigma-1} du \leq \sum_{k>K} \int_k^{k+1} \sigma u^{-\sigma-1} du \\ = (K+1)^{-\sigma} < (2\varepsilon y/x)^\sigma.$$

5. *Proof of Corollary 2.1.* — We use (2.7) and Lemma 1. The condition that $(y/z)^\sigma < 1 - \varepsilon^{2+\sigma}$ means that we can assume that $\sigma > 0$. Suppose that $\xi > 1$. Then, by (2.4),

$$F(\alpha; \xi, \sigma) \leq \xi^\sigma \int_{[\xi]}^{[\xi]+\alpha} u^{-\sigma} du + O\left(\theta(\alpha; \xi) \int_{\xi/([\xi]+\alpha)}^1 \sigma u^{-\sigma-1} du\right) \\ \leq \alpha(\xi/[\xi])^\sigma \\ + O\left(\theta(\alpha; \xi)^\sigma (1 - \xi/([\xi] + \alpha)) \max\left(1, \left(\frac{\xi}{[\xi] + \alpha}\right)^{\sigma-1}\right)\right) \\ = \alpha + O(\sigma 2^\sigma \xi^{-1}).$$

Similarly

$$F(\alpha; \xi, \sigma) \geq \xi^\sigma \int_{\xi+1}^{\xi+1+\alpha} u^{-\sigma} du \\ \geq \alpha \left(1 + \frac{1+\alpha}{\xi}\right)^{-\sigma} \geq \alpha - \frac{\sigma\alpha(1+\alpha)}{\xi}.$$

Hence, if $y_0(\varepsilon, \sigma) \leq y \leq x/2$, then by (1.11), (1.13) and (2.7),

$$\sum_{n \leq y} \lambda_n c_\alpha(x/n) = (\alpha + O(\sigma 2^\sigma y x^{-1} + x \varepsilon^{1+\sigma} y^{-1} + 2^\sigma \varepsilon^\sigma)) \sum_{n \leq y} \lambda_n.$$

Thus, if $y_0(\varepsilon, \sigma) \leq y < z \leq \frac{1}{2}$, then

$$\sum_{y < n \leq z} \lambda_n c_\alpha(x/n) = \alpha \sum_{y < n \leq z} \lambda_n \\ + O\left((\sigma 2^\sigma y x^{-1} + x \varepsilon^{1+\sigma} y^{-1} + 2^\sigma \varepsilon^\sigma) \sum_{n \leq y} \lambda_n\right).$$

We complete the proof of (2.8) by observing that by (1.11)

and Lemma 1,

$$\begin{aligned} \left(\sum_{n \leq z} \lambda_n \right) / \sum_{y < n \leq z} \lambda_n \\ = \left(1 - \left(\sum_{n \leq y} \lambda_n \right) / \sum_{n \leq z} \lambda_n \right)^{-1} < (1 - (y/z)^\sigma - \varepsilon^{2+\sigma})^{-1}. \end{aligned}$$

6. In this section we make some observations concerning the nature of $F(\alpha; \xi, 0)$.

LEMMA 2. — Suppose that $0 \leq \alpha \leq 1$ and $\xi \geq 1$. Then

$$(2.12) \quad F(\alpha; \xi, \sigma) = \alpha + O(\sigma 2^\sigma \xi^{-1}) \quad (\sigma > 0),$$

$$(2.13) \quad \lim_{\sigma \rightarrow 0^+} F(\alpha; \xi, \sigma) = \alpha = F(\alpha; \xi, 0)$$

and

$$(2.14) \quad F(\alpha; 1, \sigma) = \sum_{k=1}^{\infty} (k^{-\sigma} - (k + \alpha)^{-\sigma}) \quad (\sigma > 0).$$

By (2.14) with $\sigma = 1$, $F(\alpha; 1, 1) = \Gamma'(\alpha)/\Gamma(\alpha) + \gamma + 1/\alpha$ where Γ is the gamma function and γ is Euler's constant.

Proof. — The asymptotic formula (2.12) was established in the proof of (2.8), (2.13) then follows trivially, and (2.14) is immediate from (2.4).

LEMMA 3. — For each $\xi \geq 1$ and $\sigma > 0$ the function $F(\alpha; \xi, 0)$ is a continuous function of α and is analytic on $\mathbb{R} \setminus \{0, \{\xi\}, 1\}$ with

$$(2.15) \quad F'(\alpha) = \begin{cases} 0 & (\alpha < 0, \alpha > 1) \\ \sigma \xi^\sigma \sum_{k > \xi} (k + \alpha)^{-\sigma-1} & (0 < \alpha < \{\xi\}) \\ \sigma \xi^\sigma ([\xi] + \alpha)^{-\sigma-1} + \sigma \xi^\sigma \sum_{k > \xi} (k + \alpha)^{-\sigma-1} & (\{\xi\} < \alpha < 1). \end{cases}$$

The points $0, \{\xi\}$ and 1 are angular points of F .

LEMMA 4. — Suppose that $0 < \alpha < 1$ and $\sigma > 0$. Then considered as a function of ξ , $F(\alpha; \xi, \sigma)$ is continuous on $[1, \infty) \setminus \{2, 3, 4, \dots\}$ and for each integer $n \geq 2$,

$$(2.16) \quad \lim_{\xi \rightarrow n^-} F(\alpha; \xi, \sigma) = n^\sigma \sum_{k=n+1}^{\infty} (k^{-\sigma} - (k + \alpha)^{-\sigma})$$

and

$$(2.17) \quad \lim_{\xi \rightarrow n^+} F(\alpha; \xi, \sigma) \\ = n^\sigma \sum_{k=n}^{\infty} (k^{-\sigma} - (k + \alpha)^{-\sigma}) = F(\alpha; n, \sigma).$$

7. We now establish upper and lower bounds for the mean square of $\Phi_{x,y}(\alpha) - \alpha$ which in turn imply respectively

(i) that if y is small compared with x then the only possible (R, λ_n) asymptotic distribution modulo 1 is the uniform one, and

(ii) that the discrepancy cannot be too small.

THEOREM 3. — *Suppose that x_0 and x are non-negative real numbers, $y \geq 1$ and $0 < \alpha < 1$. Then*

$$(2.18) \quad \int_{x_0}^{x_0+x} |\Phi_{u,y}(\alpha) - \alpha|^2 du \leq \min(I_1, I_2)$$

where

$$(2.19) \quad I_1 = \frac{1}{3} (x + y^2) \sum_{n=1}^{\infty} \left(\sum_{m=1}^{\infty} \frac{1}{m} a_{mn}(y) \right)^2$$

and

$$(2.20) \quad I_2 = \sum_{n=1}^{\infty} \left(\frac{1}{3} x + \frac{1}{2} yn \right) \left(\sum_{m=1}^{\infty} \frac{1}{m} a_{mn}(y) \right)^2.$$

This theorem can be thought of in a rather loose way as a law of the iterated logarithm. This will be discussed further in a later paper. (See [5]).

THEOREM 4. — *On the hypothesis of Theorem 3,*

$$(2.21) \quad \int_{x_0}^{x_0+x} |\Phi_{u,y}(\alpha) - \alpha|^2 du \geq \max(J_1, J_2)$$

where

$$(2.22) \quad J_1 = \frac{1}{2} \pi^{-2} (x - y^2) \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} \frac{1}{m} a_{mn}(y) (1 - e(\alpha m)) \right|^2$$

and

$$(2.23) \quad J_2 = ((2\pi)^{-2} \sum_{n=1}^{\infty} (2x - 3yn) \left| \sum_{m=1}^{\infty} \frac{1}{m} a_{mn}(y) (1 - e(\alpha m)) \right|^2).$$

By taking the real part of the innermost sum in (2.22) and

(2.23) and then discarding all the terms with $m > 1$ one obtains in (2.21) the particularly simple lower bound $\max(L_1, L_2)$, where

$$L_1 = 2\pi^{-2} (\sin \pi\alpha)^4 (x - y^2) \sum_{n=1}^{\infty} a_n^2(y)$$

and

$$L_2 = \pi^{-2} (\sin \pi\alpha)^4 \sum_{n=1}^{\infty} (2x - 3yn) a_n^2(y).$$

However, in certain circumstances this loses a factor as large as $\log \log y$.

COROLLARY 4.1. — *Let the discrepancy $D_{x,y}$ be given by*

$$(2.24) \quad D_{x,y} = \sup_{0 \leq \alpha < \beta \leq 1} |\Phi_{x,y}(\beta) - \Phi_{x,y}(\alpha) - (\beta - \alpha)|.$$

Then

$$(2.25) \quad \int_{x_0}^{x+x_0} D_{u,y}^2 du \geq \sup_{\alpha \in [0,1]} \max(J_1, J_2).$$

By analogous methods it is possible to obtain corresponding inequalities for

$$\sum_{n=M+1}^{M+N} |\Phi_{n,y}(\alpha) - \alpha|^2$$

but the bounds obtained are more complicated and not so illuminating.

8. To prove Theorems 3 and 4 we require the following lemma which is Theorem 2 of Montgomery and Vaughan [4].

LEMMA 5. — *Suppose that x_1, x_2, \dots, x_R are R distinct real numbers, and that $\varphi_1, \varphi_2, \dots, \varphi_R$ are R complex numbers. Also, let*

$$(2.26) \quad \delta = \min_{\substack{r, s \\ r \neq s}} |x_r - x_s| \quad \text{and} \quad \delta_r = \min_{\substack{s \\ s \neq r}} |x_r - x_s|.$$

Then

$$(2.27) \quad \left| \sum_{\substack{r=1 \\ r \neq s}}^R \sum_{s=1}^R \frac{\varphi_r \bar{\varphi}_s}{x_r - x_s} \right| \leq \pi \min(K_1, K_2)$$

where

$$(2.28) \quad K_1 = \delta^{-1} \sum_{r=1}^R |\nu_r|^2$$

and

$$(2.29) \quad K_2 = \frac{3}{2} \sum_{r=1}^R |\nu_r|^2 \delta_r^{-1}.$$

9. *Proofs of Theorems 3 and 4.* — Let K be a positive integer. Then it is easily seen that the function $c_\alpha(u)$ given by (1.9) can be written in the form.

$$(2.30) \quad c_\alpha(u) = \alpha + \sum_{0 < |k| \leq K} \frac{1 - e(-\alpha k)}{2\pi i k} e(uk) + O\left(\min\left(1, \frac{1}{K\|u\|}\right)\right) + O\left(\min\left(1, \frac{1}{K\|u - \alpha\|}\right)\right).$$

Clearly

$$(2.31) \quad \int_{x_0}^{x_0+x} \min\left(1, \frac{1}{K\left\|\frac{u}{n} - \beta\right\|}\right) du \ll (x+n) \frac{\log K}{K} \quad (0 \leq \beta \leq 1).$$

Hence, by (1.9) and (2.30),

$$(2.32) \quad \int_{x_0}^{x_0+x} \left| \sum_{n=1}^{\infty} a_n(y) c_\alpha(u/n) - \alpha \right|^2 du = I + O\left((x+y) \frac{\log K}{K}\right)$$

where

$$(2.33) \quad I = \int_{x_0}^{x_0+x} \left| \sum_{n=1}^{\infty} \sum_{\substack{0 < |k| \leq K \\ (n, k) = 1}} \left(\sum_{m \leq K/|k|} \frac{a_{nm}(y)(1 - e(-\alpha km))}{2\pi i k m} \right) e\left(\frac{uk}{n}\right) \right|^2 du.$$

Clearly, if $n_j \leq y$, $0 < |k_j| \leq K$, $(n_j, k_j) = 1$ for $j = 1, 2$ and $k_1/n_1 \neq k_2/n_2$, then $|k_1/n_1 - k_2/n_2| \geq 1/(yn_1) \geq y^{-2}$.

Therefore, by (2.33) and Lemma 5,

$$(2.34) \quad I = \sum_{n=1}^{\infty} \sum_{\substack{0 < |k| \leq K \\ (n, k) = 1}} (x + \theta_1 y^2) \left| \sum_{m \leq K/|k|} \frac{a_{nm}(y)(1 - e(-\alpha km))}{2\pi i k m} \right|^2$$

and

(2.35)

$$I = \sum_{n=1}^{\infty} \sum_{\substack{0 < |k| \leq K \\ (n, k) = 1}} \left(x + \frac{3}{2} \theta_2 n y \right) \left| \sum_{m \leq K/|k|} \frac{a_{nm}(y)(1 - e(-\alpha km))}{2\pi i k m} \right|^2$$

where $|\theta_1| \leq 1$, $|\theta_2| \leq 1$. Theorem 3 now follows from (2.32) on letting $K \rightarrow \infty$. Theorem 4 follows in the same way on discarding all the terms with $|k| \neq 1$.

Sometimes, when the simple Riesz means (R, λ_n) are specified, it may be more appropriate to use (2.34) and (2.35) rather than appeal to Theorems 3 and 4.

10. By (2.7), (2.8) and (2.13) we see that if y is small compared with x but not too small, then under very general conditions

$$(2.36) \quad \lim_{x \gg \infty} \Phi_{x, y(x)}(\alpha) = \alpha.$$

We now show, as a consequence of Theorem 3, and again under very general conditions, that even if y is very small compared with x , then (2.36) still holds.

THEOREM 5. — Suppose that $0 < \theta < 1$, $0 < \alpha < 1$,

(2.37)

$$\lim_{y \gg \infty} \left(\left(1 + y^{\frac{3\theta-1}{2\theta}} \right) \left(\sum_{n \leq y - y^{(\theta-1)/2\theta}} \lambda_n \right)^{-2} \sum_{n \leq y} \left(\sum_{m \leq y/n} \frac{1}{m} \lambda_{mn} \right)^2 \right) = 0$$

and

$$(2.38) \quad \lim_{x \gg \infty} \Phi_{x, x^\theta}(\alpha)$$

exists, Then

$$(2.39) \quad \lim_{x \gg \infty} \Phi_{x, x^\theta}(\alpha) = \alpha.$$

We remark that (2.37) is rather a weak condition. For instance, if $\lambda_n = 1$ for every n , then it holds for every θ with $0 < \theta < 1$.

Proof. — Let y be large and define $z = y - y^{(3\theta-1)/2\theta}$. Then by Theorem 3, (1.13) and (1.11),

$$(2.40) \quad \int_{z^{1/\theta}}^{y^{1/\theta}} \left| \sum_{n \leq z} \lambda_n \left(c_\alpha \left(\frac{u}{n} \right) - \alpha \right) \right|^2 du \\ \ll (y^2 + y^{1/\theta} - z^{1/\theta}) \sum_{n \leq y} \left(\sum_{m \leq y/n} \frac{1}{m} \lambda_{mn} \right)^2.$$

Furthermore, by Cauchy's inequality (inégalité de Schwarz en français !),

$$\int_{z^{1/\theta}}^{y^{1/\theta}} \left| \sum_{z < n \leq u^\theta} \lambda_n \left(c_\alpha \left(\frac{u}{n} \right) - \alpha \right) \right|^2 du \ll (y^{1/\theta} - z^{1/\theta})(1 + y - z) \sum_{n \leq y} \lambda_n^2.$$

Hence, by (2.40),

$$(2.41) \quad \int_{z^{1/\theta}}^{y^{1/\theta}} \left| \sum_{n \leq u^\theta} \lambda_n \left(c_\alpha \left(\frac{u}{n} \right) - \alpha \right) \right|^2 du \\ \ll \left(y^2 + (y^{1/\theta} - z^{1/\theta}) \left(1 + y^{\frac{3\theta-1}{2\theta}} \right) \right) \sum_{n \leq y} \left(\sum_{m \leq y/n} \frac{1}{m} \lambda_{mn} \right)^2.$$

It is easily verified that

$$y^2 \ll (y^{1/\theta} - z^{1/\theta}) y^{(3\theta-1)/2\theta}.$$

Thus, by (2.41) and (2.37),

$$\inf_{z^{1/\theta} \leq u \leq y^{1/\theta}} |\Phi_{u, u^\theta}(\alpha) - \alpha| \rightarrow 0 \quad \text{as } y \rightarrow \infty.$$

This gives the desired result.

3. Appendix.

1. Theorem 1 does not require that the $a_n(y)$ be the simple Riesz means (R, λ_n) . It is valid provided that

$$\sum_{n=1}^{\infty} a_n(y) = 1.$$

2. Theorem 2 can be generalized in the following way. We say that the positive Toeplitz transformation $\mathcal{A} = (a_n(y))$ has asymptotic (or limit) distribution function φ with respect to the ordinary Cesaro method $(C, 1)$ if there exists a distribution function φ such that

$$(3.1) \quad \lim_{y \gg \infty} \sum_{n \leq tY} a_n(y) = \varphi(t)$$

at every t at which φ is continuous. For example, if the $a_n(y)$ are the simple Riesz means (R, λ_n) and if φ exists, then by Lemma 1 it is either a continuous function given by

$$(3.2) \quad \varphi(t) = \begin{cases} 0 & (t \leq 0) \\ t^\sigma & (0 < t < 1) \\ 1 & (t \geq 1) \end{cases} \quad (\text{with } \sigma > 0),$$

or is one of the « Heaviside » functions Y_0 and Y_1 , where $Y_a(t) = 0$ if $t < a$, $Y_a(t) = 1$ if $t \geq a$. (In the general case, necessarily $\varphi(t) = 0$ for $t < 0$). On examining the proof of Theorem 2, one sees that provided φ exists, is continuous and satisfies $\varphi(0) = 0$, $\varphi(1) = 1$, then it is possible to replace Theorem 2 by a similar but more general statement. In particular $F(\alpha; \xi, \sigma)$ is to be replaced by

$$(3.3) \quad G(\alpha; \xi, \varphi) = \begin{cases} 0 & (\alpha \leq 0) \\ 1 & (\alpha \geq 1) \\ \theta(\alpha, \xi) \left(1 - \varphi \left(\frac{\xi}{[\xi] + \alpha} \right) \right) + \sum_{k > \xi} \left(\varphi \left(\frac{\xi}{k} \right) - \varphi \left(\frac{\xi}{k + \alpha} \right) \right) & (\text{when } 0 < \alpha < 1), \end{cases}$$

but some care is needed with the error terms. Besides the above example where φ is given by (3.2), there are other interesting instances in which φ exists.

3. Theorems 3 and 4 do not require the $a_n(y)$ to be the simple Riesz means (R, λ_n) . They remain valid without modification provided that $a_n(y) = 0$ for $n > y$. Otherwise, there are extra error-terms involving $\sum_{n > y} a_n(y)$. Thus one can still obtain meaningful information in case $\lim_{y \rightarrow \infty} \sum_{n > y} a_n(y) = 0$.

BIBLIOGRAPHIE

- [1] P. ERDŐS and P. TURÁN, On a problem in the theory of uniform distributions, I, *Proc. Nederl. Acad. Wetensch.*, 51 (1948), 1146-1154, *Indagationes Math.*, 10 (1948), 370-378.
- [2] P. ERDŐS and P. TURÁN, On a problem in the theory of uniform distribution, II, *Proc. Nederl. Acad. Wetensch.*, 51 (1948), 1262-1269, *Indagationes Math.*, 10 (1948), 406-413.

- [3] L. KUIPERS and H. NIEDERREITER, Uniform distribution of sequences, Wiley, New York, 1974.
- [4] H. L. MONTGOMERY and R. C. VAUGHAN, Hilbert's inequality, *J. London Math. Soc.*, (2), 8 (1974), 73-82.
- [5] B. SAFFARI and R. C. VAUGHAN, On the fractional parts of x/n and related sequences, II (to appear).

Manuscrit reçu le 17 juillet 1975
Proposé par J.-P. Kahane.

B. SAFFARI,
Université de Paris-Sud
Orsay.

R. C. VAUGHAN,
University of Michigan
Ann Arbor.
and Imperial College
London.
