

ANNALES DE L'INSTITUT FOURIER

ROBERT A. BLUMENTHAL

Transversely homogeneous foliations

Annales de l'institut Fourier, tome 29, n° 4 (1979), p. 143-158

[<http://www.numdam.org/item?id=AIF_1979__29_4_143_0>](http://www.numdam.org/item?id=AIF_1979__29_4_143_0)

© Annales de l'institut Fourier, 1979, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

TRANSVERSELY HOMOGENEOUS FOLIATIONS

by Robert A. BLUMENTHAL

1. Introduction and statement of main results.

One way of defining a smooth codimension q foliation $\mathfrak{F}$ of a manifold M is by a smooth N^q -cocycle $\{(U_\alpha, f_\alpha, g_{\alpha\beta})\}_{\alpha, \beta \in A}$ where N^q is a smooth q -dimensional manifold and

- (i) $\{U_\alpha\}_{\alpha \in A}$ is an open cover of M .
- (ii) $f_\alpha : U_\alpha \rightarrow N^q$ is a smooth submersion whose level sets are the leaves of $\mathfrak{F}/U_\alpha$.
- (iii) $g_{\alpha\beta} : f_\beta(U_\alpha \cap U_\beta) \rightarrow f_\alpha(U_\alpha \cap U_\beta)$ is a diffeomorphism satisfying
$$f_\alpha = g_{\alpha\beta} \circ f_\beta \quad \text{on} \quad U_\alpha \cap U_\beta.$$

If N^q is a homogeneous space G/K (here G is a Lie group and $K \subset G$ is a closed subgroup) and each $g_{\alpha\beta}$ is (the restriction of) a G -translation of G/K , then $\mathfrak{F}$ is called a (transversely) homogeneous G/K -foliation.

Let us consider an important example due to Roussarie. Let $G = \mathrm{SL}(2, \mathbf{R})$, $K = \left\{ \begin{pmatrix} a & 0 \\ b & c \end{pmatrix} : ac=1, a>0 \right\}$, and let Γ be a uniform discrete subgroup of $\mathrm{SL}(2, \mathbf{R})$. The foliation of $\mathrm{SL}(2, \mathbf{R})$ whose leaves are the left cosets of K induces on $M = \Gamma \backslash \mathrm{SL}(2, \mathbf{R})$ a homogeneous $\mathrm{SL}(2, \mathbf{R})/K \cong S^1$ -foliation $\mathfrak{F}$. Moreover, $\mathfrak{F}$ is defined by a smooth nowhere zero one-form ω on M satisfying $d\omega = \omega \wedge \omega_1$, $d\omega_1 = \frac{1}{2}\omega_2 \wedge \omega$, $d\omega_2 = \omega_1 \wedge \omega_2$. Later in this paper we shall show that this set of equations completely characterizes the homogeneous $\mathrm{SL}(2, \mathbf{R})/K \cong S^1$ -foliations.

Let G be a Lie group acting effectively on the connected homogeneous space G/K and let $\mathfrak{F}$ be a homogeneous G/K -foliation of a connected manifold M .

THEOREM 1. — *To a homogeneous G/K -foliation $\mathfrak{F}$ on M is associated a homomorphism $\Phi : \pi_1(M) \rightarrow G$ well-defined up to conjugation. Let Γ be its image. The induced foliation $\tilde{\mathfrak{F}}$ on the cover $\tilde{M}$ of M associated to the kernel of Φ is given by a Γ -equivariant submersion $f : \tilde{M} \rightarrow G/K$ (Γ acting on $\tilde{M}$ by covering transformations). The holonomy group of a leaf L of $\mathfrak{F}$ is isomorphic to the isotropy subgroup $\Gamma_{\tilde{L}}$ of Γ at $\tilde{L}$, where $\tilde{L}$ is a leaf of $\tilde{\mathfrak{F}}$ projecting to L . If M is compact (whence each leaf of $\mathfrak{F}$ has a well-defined growth type), the growth of L is dominated by the growth of the orbit $\Gamma(x)$, $x = f(\tilde{L})$. Thus, if $\pi_1(M)$ has non-exponential growth (respectively, polynomial growth of degree d), then all the leaves of $\mathfrak{F}$ have non-exponential growth (respectively, polynomial growth of degree d).*

See [3] for a more general statement of the first part of the theorem.

In Section 3 we provide a differential forms characterization of a large class of homogeneous foliations. Let $\{\theta_1, \dots, \theta_{m-k}, \theta_{m-k+1}, \dots, \theta_m\}$ be a basis of the space of left-invariant one-forms on G such that $\{\theta_{m-k+1}, \dots, \theta_m\}$ is a basis of the left-invariant one-forms on K . We then have the structure equations of G relative to this basis :

$$d\theta_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \theta_j \wedge \theta_l, \quad i = 1, \dots, m \text{ where } c_{jl}^i \in \mathbb{R}.$$

THEOREM 2. — *If $H^1(M; K)$ is trivial, then the normal bundle of $\mathfrak{F}$ is trivial and there exist $m - k$ independent one-forms $\omega_1, \dots, \omega_{m-k} \in A^1(M)$ defining $\mathfrak{F}$ and one-forms $\omega_{m-k+1}, \dots, \omega_m \in A^1(M)$ satisfying*

$$d\omega_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \omega_j \wedge \omega_l, \quad i = 1, \dots, m.$$

COROLLARY 1. — *A codimension one foliation of M is a homogeneous $SL(2, \mathbb{R})/K \cong S^1$ -foliation if and only if it is defined by a smooth nowhere zero one-form $\omega \in A^1(M)$ satisfying $d\omega = \omega \wedge \omega_1$, $d\omega_1 = \frac{1}{2} \omega_2 \wedge \omega$, $d\omega_2 = \omega_1 \wedge \omega_2$ where $\omega_1, \omega_2 \in A^1(M)$.*

In Section 4 we shall establish the following :

THEOREM 3. — *If M and K are compact, then*

i) *The universal cover of M fibers over the universal cover of G/K , the fibers being the leaves of the lifted foliation.*

ii) The closure of each leaf $L \in \mathfrak{F}$ is a submanifold of M (thus $\bar{\mathfrak{F}} = \{\bar{L} : L \in \mathfrak{F}\}$ is a foliation, possibly with singularities) foliated by the leaves of $\mathfrak{F}$, this foliation being a homogeneous G'/K_L -foliation where G' is a Lie group and K_L is a compact subgroup of G' .

iii) If G/K is compact with finite fundamental group (e.g., if G is a compact semi-simple Lie group), then there exists a connected, open, dense, $\bar{\mathfrak{F}}$ -saturated submanifold of M which fibers over a connected Hausdorff manifold, the fibers being the leaves of $\bar{\mathfrak{F}}$. Moreover if $L \in \mathfrak{F}$ is a compact leaf whose holonomy group has non-exponential growth, then all the leaves of $\mathfrak{F}$ have polynomial growth.

COROLLARY 2. — If M and K are compact and if the universal cover $(\widetilde{G/K})$ of G/K is contractible, then the universal cover of M is a product $\tilde{L} \times (\widetilde{G/K})$ where $\tilde{L}$ is the (common) universal cover of the leaves of $\mathfrak{F}$ and the leaves of the lifted foliation become identified with the sets $\tilde{L} \times \{\text{pt.}\}$. Furthermore, the inclusion of a leaf $L \hookrightarrow M$ induces a monomorphism $\pi_1(L) \rightarrow \pi_1(M)$ between fundamental groups.

In Section 5 we study several particular types of homogeneous foliations and the case where the leaves are one-dimensional. For instance, it is proved that if $\mathfrak{F}$ is a one-dimensional homogeneous $SO(2q+1)/SO(2q) \cong S^{2q}$ -foliation of a compact manifold M , then $\pi_1(M)$ has polynomial growth of degree ≤ 1 and $\mathfrak{F}$ has a compact leaf. If M^3 is compact and $\pi_1(M^3)$ is not solvable, then M^3 does not support a codimension 2 Euclidean (homogeneous $SO(2) \cdot \mathbb{R}^2/SO(2)$ -) foliation.

2. Proof of Theorem 1.

Let $\{(U_\alpha, f_\alpha, \lambda_{g_{\alpha\beta}})\}_{\alpha, \beta \in A}$ be a G/K -cocycle defining $\mathfrak{F}$. Here $g_{\alpha\beta} \in G$ and $\lambda_{g_{\alpha\beta}}$ denotes the diffeomorphism of G/K sending aK to $g_{\alpha\beta}aK$. Let $P = \{[\lambda_g \circ f_\alpha]_x : x \in U_\alpha, \alpha \in A, g \in G\}$, where $[\lambda_g \circ f_\alpha]_x$ denotes the germ of $\lambda_g \circ f_\alpha$ at x . By analyticity and the connectivity of G/K , P admits a differentiable structure such that the natural projection $\pi : P \rightarrow M$ is a smooth regular covering with G as the group of covering transformations. Let $\tilde{M}$ be a connected component of P . The group of covering transformations of the covering $\pi : \tilde{M} \rightarrow M$ is a subgroup Γ of G and is the image of a homomorphism $\Phi : \pi_1(M) \rightarrow G$. The evaluation map $f : \tilde{M} \rightarrow G/K$ is a smooth Γ -equivariant submersion constant along the leaves of $\pi^{-1}(\mathfrak{F})$. Let L be a leaf of $\mathfrak{F}$ and choose a leaf $\tilde{L}$ of $\pi^{-1}(\mathfrak{F})$ which projects to L . Then

$\pi/\tilde{L} : \tilde{L} \rightarrow L$ is a regular covering whose group of covering transformations, namely $\Gamma_{\tilde{L}}$, is isomorphic to the holonomy group of L . Note that the holonomy group of L can be realized as a subgroup of K and that if σ is a loop in L which is homotopically trivial in M , then the element of holonomy determined by σ is trivial.

We now assume that M is compact. Let $\{U_i, f_i, \lambda_{\gamma_{ij}}\}_{i,j=1}^m$ be a finite G/K -cocycle defining $\mathfrak{F}$ such that $\{U_i\}_{i=1}^m$ is a regular covering of M in the sense of [6, pp. 336-337] and such that $\gamma_{ij} \in \Gamma$ for $i, j = 1, \dots, m$. By a plaque $\rho \subset U_i$ of the leaf $L \in \mathfrak{F}$ is meant a connected component of $L \cap U_i$. Fix $L \in \mathfrak{F}$ and let ρ be a plaque of L . Without loss of generality, we may assume that $\rho \subset U_1$. For each $i = 1, \dots, m$, let $v_i(n)$ be the number of distinct plaques in U_i which can be reached from the plaque ρ by a chain of plaques of length $\leq n$. If g_ρ denotes the growth function of L at ρ with respect to the regular covering $\{U_i\}_{i=1}^m$, then $g_\rho(n) = v_1(n) + \dots + v_m(n)$, $n \in \mathbb{Z}^+$ [6]. Let $\Gamma^1 \subset \Gamma$ be a finite symmetric generating set for Γ such that $\gamma_{ij} \in \Gamma^1$ for all $i, j = 1, \dots, m$. Set $z = f_1(\rho) \in G/K$. We may assume that $\Gamma(z)$ is the orbit $\Gamma(f(\tilde{L}))$ where $\tilde{L} \in \pi^{-1}(\mathfrak{F})$ is a leaf projecting to L . The growth function of Γ at z is $g_z(n) = |\Gamma^n(z)|$, $n \in \mathbb{Z}^+$ where $|\cdot|$ denotes cardinality and $\Gamma^n(z) = \{z' \in G/K : z' = \lambda_{\gamma_j} \circ \dots \circ \lambda_{\gamma_1}(z) \text{ for some } \gamma_1, \dots, \gamma_j \in \Gamma^1 \text{ where } j \leq n\}$. Let τ be a plaque of L in U_i which can be reached from the plaque ρ by a chain of plaques of length $\leq n$. Let $(\rho = \rho_1, \rho_2, \dots, \rho_l = \tau)$ be such a chain, $l \leq n$. For each $k = 1, \dots, l$ choose U_{i_k} such that $\rho_k \subset U_{i_k}$, $U_{i_1} = U_1$, $U_{i_l} = U_i$. Then $\lambda_{\gamma_{i_1 i_2}} \circ \dots \circ \lambda_{\gamma_{i_{l-1} i_l}}(z) = f_i(\tau)$. Thus $f_i(\tau) \in \Gamma^{n-1}(z)$. If $\tau' \neq \tau$ is another such plaque, then $f_i(\tau') \neq f_i(\tau)$. Hence $v_i(n) \leq g_z(n-1)$ and so $g_\rho(n) \leq m g_z(n-1)$ which completes the proof of the theorem.

2.1. COROLLARY. — *If G is nilpotent or if $\pi_1(M)$ is nilpotent then all the leaves of $\mathfrak{F}$ have polynomial growth. Moreover, the holonomy group of every leaf is finitely generated and has polynomial growth.*

Proof. — Since Γ is a finitely generated nilpotent group, it has polynomial growth [1], [11] and hence all the leaves of $\mathfrak{F}$ have polynomial growth. Let L be a leaf of $\mathfrak{F}$ and $\tilde{L} \in \pi^{-1}(\mathfrak{F})$ a leaf projecting to L . Since $\Gamma_{\tilde{L}} \subset \Gamma$ is a subgroup of a finitely generated nilpotent group, we have that $\Gamma_{\tilde{L}}$ is finitely generated [7] and hence has polynomial growth.

2.2. COROLLARY. — *If G is solvable and $\pi_1(M)$ has non-exponential growth, then all the leaves of $\mathfrak{F}$ have polynomial growth.*

Proof. — Since Γ is a finitely generated solvable group with non-exponential growth, it follows that Γ has polynomial growth [11].

2.3. COROLLARY. — *If $L \in \mathfrak{F}$ is a leaf satisfying*

i) $[\pi_1(M) : i_*\pi_1(L)] < \infty$,

ii) *The holonomy group of L has non-exponential growth (respectively, polynomial growth of degree d),*

then all the leaves of $\mathfrak{F}$ have non-exponential growth (respectively, polynomial growth of degree d).

Proof. — If $\tilde{L} \in \pi^{-1}(\mathfrak{F})$ is a leaf projecting to L , then we have $[\Gamma : \Gamma_{\tilde{L}}] < \infty$. Hence $\Gamma_{\tilde{L}}$ is finitely generated [1] and, by (ii), has non-exponential growth (respectively, polynomial growth of degree d). Hence Γ has non-exponential growth (respectively, polynomial growth of degree d) [1].

3. Structure equations and the normal bundle.

The following is established using arguments similar to those found, e.g., in Chapter 10 of [9].

3.1. PROPOSITION. — *Let $\mathfrak{F}$ be a codimension $m - k$ foliation of M defined by $m - k$ linearly independent one-forms $\omega_1, \dots, \omega_{m-k} \in A^1(M)$ and suppose that there are also one-forms $\omega_{m-k+1}, \dots, \omega_m \in A^1(M)$ such that*

$$d\omega_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \omega_j \wedge \omega_l, \quad i = 1, \dots, m.$$

Then $\mathfrak{F}$ is a homogeneous G/K -foliation.

We now prove Theorem 2. The canonical projection $p : G \rightarrow G/K$ makes G a smooth principal K -bundle over G/K . We may pull back this bundle via f to obtain a smooth principal K -bundle $\rho : f^*(G) \rightarrow \tilde{M}$ where $f^*(G) = \{(y, g) \in \tilde{M} \times G : f(y) = p(g)\}$ and $\rho(y, g) = y$. We also have a map $\bar{f} : f^*(G) \rightarrow G$, defined by $\bar{f}(y, g) = g$, such that $p \circ \bar{f} = f \circ \rho$. Define a left action of Γ on $\tilde{M} \times G$ by $\gamma(y, g) = (\gamma y, \gamma g)$ for $\gamma \in \Gamma$, $(y, g) \in \tilde{M} \times G$. This action of Γ preserves $f^*(G)$ and thus defines a smooth left action of Γ on $f^*(G)$ such that $\gamma \circ \rho = \rho \circ \gamma$ for each $\gamma \in \Gamma$. Let $\Gamma \backslash f^*(G)$ denote the space of orbits and let $\tau : f^*(G) \rightarrow \Gamma \backslash f^*(G)$ be the natural projection. The map $\rho : f^*(G) \rightarrow \tilde{M}$ induces a continuous surjection $\bar{\rho} : \Gamma \backslash f^*(G) \rightarrow \tilde{M}$ satis-

fying $\bar{\rho} \circ \tau = \pi \circ \rho$. Since the right action of K on $f^*(G)$ commutes with the left action of Γ , there is a free right action of K on $\Gamma \backslash f^*(G)$ such that $\bar{\rho} : \Gamma \backslash f^*(G) \rightarrow M$ is a smooth principal K -bundle. We remark that Γ acts freely and properly discontinuously on $f^*(G)$ and hence $\tau : f^*(G) \rightarrow \Gamma \backslash f^*(G)$ is a smooth regular covering with Γ as the group of covering transformations.

Now $p \circ \bar{f}$ is a submersion, whence $\bar{f}$ is transverse to $\mathfrak{F}_0$ (where $\mathfrak{F}_0$ is the foliation of G by the left cosets of K) and so $\bar{f}^{-1}(\mathfrak{F}_0)$ is a well-defined foliation of $f^*(G)$. The foliation $\mathfrak{F}_0$ of G is defined by $\theta_1, \dots, \theta_{m-k}$ and hence $\rho^{-1}(\pi^{-1}(\mathfrak{F})) = \bar{f}^{-1}(\mathfrak{F}_0)$ is defined by the $m-k$ linearly independent smooth one-forms $\bar{f}^*\theta_1, \dots, \bar{f}^*\theta_{m-k} \in A^1(f^*(G))$ which satisfy

$$d(\bar{f}^*\theta_i) = \sum_{1 \leq j < l \leq m} c_{jl}^i (\bar{f}^*\theta_j) \wedge (\bar{f}^*\theta_l), i = 1, \dots, m.$$

Note that $L_\gamma \circ \bar{f} \circ \gamma$ for each $\gamma \in \Gamma$ where $L_\gamma : G \rightarrow G$ denotes left translation by γ . Now $\mathfrak{F}_0$ and $\theta_1, \dots, \theta_m$ are invariant under left translation by elements of Γ and hence $\bar{f}^{-1}(\mathfrak{F}_0)$ and $\bar{f}^*\theta_1, \dots, \bar{f}^*\theta_m$ are invariant under the action of Γ on $f^*(G)$. Thus $\bar{f}^{-1}(\mathfrak{F}_0)$ projects to a well-defined foliation $\mathfrak{F}$ of $\Gamma \backslash f^*(G)$ and $\bar{f}^*\theta_1, \dots, \bar{f}^*\theta_m$ project to well-defined smooth one-forms $\alpha_1, \dots, \alpha_m$ respectively such that $\mathfrak{F}$ is defined by the $m-k$ linearly independent smooth one-forms

$$\alpha_1, \dots, \alpha_{m-k} \in A^1(\Gamma \backslash f^*(G))$$

which satisfy

$$d\alpha_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \alpha_j \wedge \alpha_l,$$

$i = 1, \dots, m$. Note that $\mathfrak{F} = \bar{\rho}^{-1}(\mathfrak{F})$. Since $H^1(M; K)$ is trivial, there exists a smooth section $s : M \rightarrow \Gamma \backslash f^*(G)$. Now s is transverse to $\bar{\rho}^{-1}(\mathfrak{F})$ and $s^{-1}(\bar{\rho}^{-1}(\mathfrak{F})) = \mathfrak{F}$; hence, setting $\omega_i = s^*\alpha_i$ for $i = 1, \dots, m$, we see that $\mathfrak{F}$ is defined by the $m-k$ independent one-forms $\omega_1, \dots, \omega_{m-k} \in A^1(M)$ which satisfy $d\omega_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \omega_j \wedge \omega_l$, $i = 1, \dots, m$. In particular, the normal bundle of $\mathfrak{F}$ is trivial.

Noting that the two-dimensional affine group

$$K = \left\{ \begin{pmatrix} a & 0 \\ b & c \end{pmatrix} : ac = 1, a > 0 \right\}$$

is contractible, we see that Corollary 1 follows from Proposition (3.1) and Theorem 2.

3.2. COROLLARY. — Suppose G/K is a simply connected Riemannian homogeneous space of constant curvature. Let $\mathfrak{F}$ be a codimension $m - k$ foliation of M with trivial normal bundle. Then $\mathfrak{F}$ is a homogeneous G/K -foliation if and only if there exist $m - k$ linearly independent one-forms $\omega_1, \dots, \omega_{m-k} \in A^1(M)$ defining $\mathfrak{F}$ and one-forms $\omega_{m-k+1}, \dots, \omega_m \in A^1(M)$ satisfying $d\omega_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \omega_j \wedge \omega_l$, $i = 1, \dots, m$.

Proof. — If $\mathfrak{F}$ is defined by such one-forms, then $\mathfrak{F}$ is a homogeneous G/K -foliation by Proposition (3.1). If $\mathfrak{F}$ is a homogeneous G/K -foliation, then the metric on G/K induces a smooth Riemannian metric on the normal bundle of $\mathfrak{F}$. The hypothesis on G/K implies that K is the full orthogonal group $O(m-k)$ and hence the principal K -bundle $\bar{\rho} : \Gamma \backslash \Gamma^*(G) \rightarrow M$ constructed in the proof of the theorem is just the bundle of orthonormal frames of the normal bundle of $\mathfrak{F}$. The conclusion now follows from the triviality of this principal K -bundle.

3.3. COROLLARY. — Let $\mathfrak{F}$ be a codimension two foliation of M with trivial normal bundle. Then

a) $\mathfrak{F}$ is transversely Euclidean (homogeneous $SO(2) \cdot \mathbf{R}^2 / SO(2)$) if and only if it is defined by independent one-forms ω_1, ω_2 satisfying

$$d\omega_1 = \frac{1}{2} \omega_2 \wedge \omega_3, \quad d\omega_2 = -\frac{1}{2} \omega_1 \wedge \omega_3, \quad d\omega_3 = 0.$$

b) $\mathfrak{F}$ is transversely hyperbolic (homogeneous $SL(2, \mathbf{R}) / SO(2)$) if and only if it is defined by independent one-forms ω_1, ω_2 satisfying $d\omega_1 = \frac{1}{2} \omega_2 \wedge \omega_3$, $d\omega_2 = \omega_1 \wedge \omega_2 - 2\omega_1 \wedge \omega_3$, $d\omega_3 = -\omega_1 \wedge \omega_3$.

c) $\mathfrak{F}$ is transversely elliptic (homogeneous $SO(3) / SO(2)$) if and only if it is defined by independent one-forms ω_1, ω_2 satisfying $d\omega_1 = \frac{1}{2} \omega_2 \wedge \omega_3$, $d\omega_2 = -\frac{1}{2} \omega_1 \wedge \omega_3$, $d\omega_3 = \frac{1}{2} \omega_1 \wedge \omega_2$.

3.4. COROLLARY. — Let $\mathfrak{F}$ be a codimension q foliation of M with trivial normal bundle. Then $\mathfrak{F}$ is transversely affine (homogeneous $GL(q, \mathbf{R}) \cdot \mathbf{R}^q / GL(q, \mathbf{R})$) if and only if there exist $q = m - k$ independent one-

forms $\omega_1, \dots, \omega_{m-k} \in A^1(M)$ defining $\mathfrak{F}$ (where $k = q^2$) and one-forms $\omega_{m-k+1}, \dots, \omega_m \in A^1(M)$ satisfying

$$d\omega_i = \sum_{1 \leq j < l \leq m} c_{jl}^i \omega_j \wedge \omega_l, \quad i = 1, \dots, m.$$

Proof. — If $\mathfrak{F}$ is transversely affine, then the principal K -bundle $\bar{\rho} : \Gamma \backslash f^*(G) \rightarrow M$ is just the bundle of frames of the normal bundle of $\mathfrak{F}$.

4. Riemannian homogeneous foliations.

Throughout this section we assume that M and K are compact. Pick and fix a G -invariant Riemannian metric on G/K .

4.1. THEOREM. — *The universal cover of M fibers over the universal cover of G/K , the fibers being the leaves of the lifted foliation.*

Proof. — Let $\pi : \tilde{M} \rightarrow M$ and $f : \tilde{M} \rightarrow G/K$ be as in Theorem 1. Let $\hat{M}$ be the universal cover of M and choose a covering map $\pi' : \hat{M} \rightarrow \tilde{M}$. Let $\hat{\mathfrak{F}}$ be the foliation of $\hat{M}$ obtained by lifting $\mathfrak{F}$ via $\pi \circ \pi'$ and let $\hat{f} = f \circ \pi'$. Then $\hat{f}$ is a submersion defining $\hat{\mathfrak{F}}$. Let $E \subset T(M)$ be the bundle tangent to the leaves of $\mathfrak{F}$. Choose a subbundle $Q \subset T(M)$ such that $T(M) = E \oplus Q$. If $\{(U_\alpha, f_\alpha, \lambda_{g_{\alpha\beta}})\}_{\alpha, \beta \in A}$ is a G/K -cocycle defining $\mathfrak{F}$, then Q inherits a smooth Riemannian metric by the requirement that $f_{\alpha, x} : Q_x \rightarrow T_{f_\alpha(x)}(G/K)$, $x \in U_\alpha$ be a vector space isometry. Choose a Riemannian metric on E and define a Riemannian metric on M by the requirement that E_x is orthogonal to Q_x for all $x \in M$. Then this metric on the foliated manifold $(M, \mathfrak{F})$ is bundle-like in the sense of [8]. Moreover, since M is compact, this bundle-like metric is complete.

The complete bundle-like metric on M lifts via $\pi \circ \pi'$ to a complete bundle-like metric on the foliated manifold $(\hat{M}, \hat{\mathfrak{F}})$. But $\hat{\mathfrak{F}}$ is regular since it is defined by a global submersion. Hence, by Corollary 3 in [8], the space of leaves $\hat{M}/\hat{\mathfrak{F}}$ of $\hat{\mathfrak{F}}$ is a complete, Riemannian, Hausdorff manifold and the natural projection $\hat{M} \rightarrow \hat{M}/\hat{\mathfrak{F}}$ is a fibration. Since $\hat{M}$ is simply connected, it follows that $\hat{M}/\hat{\mathfrak{F}}$ is simply connected. Now $\hat{f}$ induces a local isometry $\hat{M}/\hat{\mathfrak{F}} \rightarrow G/K$ which lifts to a local isometry $\hat{M}/\hat{\mathfrak{F}} \rightarrow (\widetilde{G/K})$ where $(\widetilde{G/K})$ denotes the simply connected Riemannian cover of G/K . Hence $\hat{M}/\hat{\mathfrak{F}}$ and

$(\widetilde{G/K})$ are isometric [4] since each is a connected, simply connected, complete, analytic, Riemannian manifold.

Remark. — Corollary 2 is an immediate consequence of Theorem (4.1). If we do not assume that $(\widetilde{G/K})$ is contractible, but only that $\pi_2(\widetilde{G/K}) = 0$, then the leaves of $\mathfrak{F}$ are simply connected and we still have, that $i_* : \pi_1(L) \rightarrow \pi_1(M)$ is injective for all $L \in \mathfrak{F}$.

4.2. COROLLARY. — *Suppose the universal cover of G/K is contractible.*

- i) *If $\dim \mathfrak{F} = 1$, then $\hat{M}$ is contractible.*
- ii) *If $\dim \mathfrak{F} = 2$, then either $\hat{M}$ is contractible or has the homotopy type of S^2 . In the latter case, all the leaves of $\mathfrak{F}$ are compact with universal cover S^2 .*

We adopt the following notation for the rest of this section and the next. Let N denote the connected Riemannian manifold G/K . Thus G is a transitive group of isometries of N . Let $\tilde{N}$ be the universal cover of N endowed with the Riemannian metric lifted from N and let $\tilde{G}$ be the full group of isometries of $\tilde{N}$. Then $\tilde{G}$ acts transitively on $\tilde{N}$. Theorem (4.1) tells us that we have a fibration $\rho : \hat{M} \rightarrow \hat{M}/\mathfrak{F} \cong \tilde{N}$. Let $\pi_1(M)$ denote the group of covering transformations of $\hat{M}$. Each element $\tau \in \pi_1(M)$ induces an isometry $\psi(\tau) : \hat{M}/\mathfrak{F} \rightarrow \hat{M}/\mathfrak{F}$. We regard $\psi(\tau)$ as an isometry of $\tilde{N}$. Hence $\psi(\tau) \in \tilde{G}$ and we have a homomorphism $\psi : \pi_1(M) \rightarrow \tilde{G}$. Let $\Sigma = \text{image } \psi \subset \tilde{G}$. For $x \in \tilde{N}$, let $\Sigma_x = \{\sigma \in \Sigma : \sigma(x) = x\}$ and $\Sigma(x) = \{\sigma(x) : \sigma \in \Sigma\}$. Let $L \in \mathfrak{F}$ and choose a leaf $L' \in \mathfrak{F}$ which projects to L . Then the orbit $\Sigma(x)$ of $x = \rho(L')$ under Σ depends only on L and we denote this orbit by Σ^L . The following lemma is elementary.

4.3. LEMMA. — *Let L be a leaf of $\mathfrak{F}$. Then*

- i) *L is proper (i.e., L is an imbedded submanifold of M) if and only if Σ^L is a discrete subset of $\tilde{N}$.*
- ii) *L is compact if and only if Σ^L is discrete and closed.*
- iii) *L is dense if and only if Σ^L is dense.*
- iv) *The space of leaves of $\mathfrak{F}$ is homeomorphic to the orbit space $\Sigma \backslash \tilde{N}$.*
- v) *Let $\bar{\Sigma}$ denote the closure of Σ in $\tilde{G}$. Then for $x \in \tilde{N}$, we have $\overline{\Sigma(x)} = \bar{\Sigma}(x)$; that is, the orbit of x under $\bar{\Sigma}$ is the closure of the orbit of x under Σ .*

4.4. THEOREM. — *Let $L \in \mathfrak{F}$. Then $\bar{L}$ is a submanifold of M and the foliation of $\bar{L}$ by the leaves of $\mathfrak{F}$ is a homogeneous G'/K_L -foliation where G' is a closed subgroup of $\tilde{G}$ and K_L is a compact subgroup of G' .*

Proof. — Let $\pi : \hat{M} \rightarrow M$ be the universal cover of M . We have a fibration $\rho : \hat{M} \rightarrow \hat{N}$ whose fibers are the leaves of $\hat{\mathfrak{F}} = \pi^{-1}(\mathfrak{F})$. Choose a leaf $L' \subset \pi^{-1}(L)$ and let $x = \rho(L') \in \hat{N}$. Then

$$\pi^{-1}(\bar{L}) = \overline{\pi^{-1}(L)} = \overline{\rho^{-1}(\Sigma(x))} = \rho^{-1}(\overline{\Sigma(x)}) = \rho^{-1}(\bar{\Sigma}(x))$$

and so we have a fibration $\rho/\pi^{-1}(\bar{L}) : \pi^{-1}(\bar{L}) \rightarrow \bar{\Sigma}(x)$. Now $\bar{\Sigma}(x)$ is the orbit of x under the action of the Lie group $\bar{\Sigma}$ and hence is a submanifold of $\hat{N}$ from which it follows that $\pi^{-1}(\bar{L})$ is a submanifold of $\hat{M}$. Hence $\bar{L}$ is a submanifold of M .

It remains to demonstrate that the foliation of $\bar{L}$ by the leaves of $\mathfrak{F}$ is a homogeneous foliation. Now $\pi/\pi^{-1}(\bar{L}) : \pi^{-1}(\bar{L}) \rightarrow \bar{L}$ is a regular covering and $\rho/\pi^{-1}(\bar{L}) : \pi^{-1}(\bar{L}) \rightarrow \bar{\Sigma}(x)$ is a submersion defining $\hat{\mathfrak{F}}/\pi^{-1}(\bar{L}) = \pi^{-1}(\mathfrak{F}/\bar{L})$. Moreover, for each covering transformation τ of $\pi^{-1}(\bar{L})$ we have $(\psi(\tau)/\bar{\Sigma}(x)) \circ (\rho/\pi^{-1}(\bar{L})) = (\rho/\pi^{-1}(\bar{L})) \circ (\tau/\pi^{-1}(\bar{L}))$. Hence there exists a $\bar{\Sigma}(x)$ -cocycle $\{(U_\alpha f_\alpha \sigma_{\alpha\beta})\}_{\alpha, \beta \in A}$ defining $\mathfrak{F}/\bar{L}$ such that $\sigma_{\alpha\beta} \in \Sigma \subset \bar{\Sigma}$ for all $\alpha, \beta \in A$. But $\bar{\Sigma}(x)$ inherits a Riemannian metric from $\hat{N}$ and $\psi(\tau)/\bar{\Sigma}(x) : \bar{\Sigma}(x) \rightarrow \bar{\Sigma}(x)$ is an isometry. Moreover, $\bar{\Sigma}$ acts transitively on $\bar{\Sigma}(x)$ and so $\bar{\Sigma}(x) \cong \bar{\Sigma}/K_L$ where K_L is a compact subgroup of $\bar{\Sigma}$. Taking $G' = \bar{\Sigma}$, we have that $\mathfrak{F}/\bar{L}$ is a homogeneous G'/K_L -foliation.

Remark. — Since $\{\bar{\Sigma}(x) : x \in \hat{N}\} : \{\bar{\Sigma}(x) : x \in \hat{N}\}$ partitions $\hat{N}$, we see that $\{\bar{L} : L \in \mathfrak{F}\}$ partitions M and hence $\mathfrak{F} = \{\bar{L} : L \in \mathfrak{F}\}$ is a foliation of M , possibly with singularities.

4.5. THEOREM. — *If $\hat{N}$ is compact (i.e., if $N = G/K$ is compact with finite fundamental group), there exists a connected, open, dense, $\hat{\mathfrak{F}}$ -saturated submanifold V of M which fibers over a connected Hausdorff manifold, the fibers being the leaves of $\hat{\mathfrak{F}}$. Thus, in particular, the leaves of $\hat{\mathfrak{F}}$ contained in V are mutually diffeomorphic.*

Proof. — Since $\hat{N}$ is compact, it follows that $\hat{G}$ is compact. Thus $\bar{\Sigma}$ is a compact Lie group acting smoothly on $\hat{N}$. Let $W \subset \hat{N}$ be the union of the principal orbits [2]. Then W is an open, dense, saturated subset of $\hat{N}$ and $\bar{\Sigma} \backslash W$ is a connected Hausdorff manifold [2]. Let $V = \pi(\rho^{-1}(W)) \subset M$. Then V is an open, dense, $\hat{\mathfrak{F}}$ -saturated submanifold of M . We have a smooth submersion $V \rightarrow \bar{\Sigma} \backslash W$ defining $\hat{\mathfrak{F}}/V$ and hence $\hat{\mathfrak{F}}/V$ is a regular foliation of V with all leaves compact. Thus $V \rightarrow \bar{\Sigma} \backslash W = V/\hat{\mathfrak{F}}$ is a fibration [5], the fibers being the leaves of $\hat{\mathfrak{F}}/V$. In particular, V is connected.

4.6. THEOREM. — Suppose $(\widetilde{G}/\widetilde{K})$ is compact and that $L \in \mathfrak{F}$ is a compact leaf. If the holonomy group of L has non-exponential growth (respectively, polynomial growth of degree d), then all the leaves of $\mathfrak{F}$ have polynomial growth (respectively, polynomial growth of degree d). If the fundamental group of L has non-exponential growth (respectively, polynomial growth of degree d), then $\pi_1(M)$ has non-exponential growth (respectively, polynomial growth of degree d) and all the leaves of $\mathfrak{F}$ have polynomial growth (respectively, polynomial growth of degree d).

Proof. — Since L is compact we have that Σ^L is discrete and closed, hence finite. Thus $\pi^{-1}(L)$ is a finite union of leaves. Let L' be a leaf of $\mathfrak{F}$ which projects to L and let $\pi_1(M)_{L'} = \{\tau \in \pi_1(M) : \tau(L') = L'\}$. Then the index of $\pi_1(M)_{L'}$ in $\pi_1(M)$ is finite and hence $[\pi_1(M) : i_*\pi_1(L)] < \infty$. The holonomy group of L is isomorphic to the linear holonomy group of L and hence is a finitely generated linear group with non-exponential growth, hence polynomial growth [10] (respectively, polynomial growth of degree d). Hence, by Corollary (2.3), all the leaves of $\mathfrak{F}$ have polynomial growth (respectively, polynomial growth of degree d). If $\pi_1(L)$ has non-exponential growth (respectively, polynomial growth of degree d), then so does $i_*\pi_1(L)$. Since $[\pi_1(M) : i_*\pi_1(L)] < \infty$, it follows from [1] that $\pi_1(M)$ has non-exponential growth (respectively, polynomial growth of degree d).

Combining Theorems (4.1), (4.4), (4.5) and (4.6), we obtain Theorem 3.

4.7. PROPOSITION. — If $\pi_1(\bar{L})$ is finite for some leaf $L \in \mathfrak{F}$, then all the leaves of $\mathfrak{F}$ are compact. If in addition $(\widetilde{G}/\widetilde{K})$ is compact, then $\pi_1(M)$ is finite.

Proof. — Let $\pi : \hat{M} \rightarrow M$ be the universal cover of M and let $\pi^{-1}(\bar{L})_0$ be a connected component of $\pi^{-1}(\bar{L})$. Then $\pi/\pi^{-1}(\bar{L})_0 : \pi^{-1}(\bar{L})_0 \rightarrow \bar{L}$ is a covering of $\bar{L}$. Since $\bar{L}$ is compact with finite fundamental group, it follows that $\pi^{-1}(\bar{L})_0$ is compact. Let L' be a leaf of $\mathfrak{F}$ contained in $\pi^{-1}(\bar{L})_0$. Then L' is a closed subset of the compact space $\pi^{-1}(\bar{L})_0$ and hence L' is compact. Thus all the leaves of $\mathfrak{F}$ are compact and so all the leaves of $\mathfrak{F}$ are compact. If in addition $\tilde{N}$ is compact, then $\pi : \hat{M} \rightarrow \tilde{N}$ is a fibration with compact base and compact fiber. Thus $\hat{M}$ is compact and so $\pi_1(M)$ is finite.

5. Elliptic, Euclidean, and hyperbolic foliations.

Throughout this section M denotes a closed manifold.

5.1. PROPOSITION. — Let $\mathfrak{F}$ be a one-dimensional homogeneous $SO(2q+1)/SO(2q) \cong S^{2q}$ -foliation of M . Then $\pi_1(M)$ has polynomial growth of degree $d \leq 1$ and $\mathfrak{F}$ has a compact leaf.

Proof. — Suppose that no leaf of $\mathfrak{F}$ is compact. Then all the leaves of $\mathfrak{F}$ are simply connected and hence $i_*\pi_1(L)$ is trivial for all $L \in \mathfrak{F}$. Let $\tau \in \pi_1(M)$. Then, since $\psi(\tau) : S^{2q} \rightarrow S^{2q}$ has a fixed point, there exists a leaf $L' \in \mathfrak{F}$ such that $\tau(L') = L'$. If $L = \pi(L') \in \mathfrak{F}$, we have $\tau \in \pi_1(M)_L = \{\mu \in \pi_1(M) : \mu(L') = L'\} \cong i_*\pi_1(L)$ and hence τ is the identity covering transformation and so M is simply connected. Thus M fibers over S^{2q} , the fibers being the leaves of $\mathfrak{F}$. But this implies that all the leaves of $\mathfrak{F}$ are circles which is impossible. Hence there exists a compact leaf $L \in \mathfrak{F}$. Since the fundamental group of L has polynomial growth of degree 1, it follows from Theorem (4.6) that $\pi_1(M)$ has polynomial growth of degree $d \leq 1$.

5.2. PROPOSITION. — *Let $\mathfrak{F}$ be a codimension two elliptic foliation of M . Then either*

- i) *all the leaves of $\mathfrak{F}$ are compact,*
- ii) *all the leaves of $\mathfrak{F}$ are dense, or*
- iii) *all the leaves of $\mathfrak{F}$ have polynomial growth and there exists a compact leaf.*

Proof. — In this case $N = \tilde{N} = S^2$, $\tilde{G} = SO(3)$ and so we have $\psi : \pi_1(M) \rightarrow SO(3)$ with image $\Sigma \subset SO(3)$. Thus $\bar{\Sigma}$ is a compact Lie group and hence has only finitely many connected components. Let $(\bar{\Sigma})_0$ be the connected component of the identity. Then $(\bar{\Sigma})_0$ is a compact connected Lie group and $[\bar{\Sigma} : (\bar{\Sigma})_0] < \infty$. Let $\Sigma_0 = \Sigma \cap (\bar{\Sigma})_0$. Then Σ_0 is a subgroup of Σ and $[\Sigma : \Sigma_0] < \infty$. Hence, by [1], Σ_0 is finitely generated and has the same growth type as Σ . We consider four cases :

a) $(\bar{\Sigma})_0$ is zero-dimensional : Then Σ is discrete and hence finite. Thus $\Sigma(x)$ is finite for all $x \in S^2$ and so all the leaves of $\mathfrak{F}$ are compact by Lemma (4.3).

b) $(\bar{\Sigma})_0$ is one-dimensional : Then $(\bar{\Sigma})_0$ is isomorphic to S^1 and so Σ_0 is (finitely generated) abelian. Hence Σ has polynomial growth. Thus, by arguments identical to those used to establish Theorem 1, all the leaves of $\mathfrak{F}$ have polynomial growth. Now Σ_0 is not trivial for otherwise $(\bar{\Sigma})_0$ would be zero-dimensional. Choose a non-identity element $A \in \Sigma_0$. Then A has exactly two fixed points $x, y \in S^2$. If $B \in \Sigma_0$ is nontrivial, then $ABx = BAx = Bx$ and $AB y = B A y = B y$. Hence either $Bx = x$ and $By = y$ or else $Bx = y$ and $By = x$. Either way, the orbit of x under Σ_0 is finite. Thus $\Sigma(x)$ is finite and so $\mathfrak{F}$ has a compact leaf.

c) $(\bar{\Sigma})_0$ is two-dimensional : Then $(\bar{\Sigma})_0$ is isomorphic to the two-dimensional torus which is impossible.

d) $(\bar{\Sigma})_0$ is three-dimensional : Then $(\bar{\Sigma})_0 = \text{SO}(3)$ and so Σ is dense in $\text{SO}(3)$. Hence $\Sigma(x)$ is dense in S^2 for all $x \in S^2$ and so all the leaves of $\mathfrak{F}$ are dense.

5.3. PROPOSITION. — *Let $\mathfrak{F}$ be a codimension two Euclidean, elliptic or hyperbolic foliation of M . If $L \in \mathfrak{F}$ is a compact leaf with $H^1(L) = 0$, then all the leaves of $\mathfrak{F}$ are compact. If $L \in \mathfrak{F}$ is a (not necessarily compact) leaf with $H_1(L, \mathbb{Z}) = 0$, then $i_*\pi_1(L)$ is a normal subgroup of $\pi_1(M)$.*

Proof. — Let $L \in \mathfrak{F}$ and choose a leaf $L' \in \hat{\mathfrak{F}}$ which projects to L . Let $x = \rho(L') \in \tilde{N}$. Since Σ_x is abelian, the composition

$$\pi_1(L) \xrightarrow{i_*} i_*\pi_1(L) \cong \pi_1(M)_{L'} \xrightarrow{\psi} \Sigma_x$$

induces a surjection $H_1(L, \mathbb{Z}) \rightarrow \Sigma_x$. Suppose L is compact. Then $\Sigma(x) = \Sigma^L$ is discrete and closed. If $H^1(L) = 0$, then $H_1(L, \mathbb{Z})$ is finite and so Σ_x is finite. Thus Σ is a discrete subgroup of $\tilde{G}$ and hence $\Sigma(x)$ is discrete and closed for all $x \in \tilde{N}$ [4]. Thus all the leaves of $\mathfrak{F}$ are compact. If $H_1(L, \mathbb{Z}) = 0$, then Σ_x is trivial and hence $\pi_1(M)_{L'} = \text{kernel } \psi$. Thus $i_*\pi_1(L)$ is normal in $\pi_1(M)$.

5.4. PROPOSITION. — *Let $\mathfrak{F}$ be a codimension two Euclidean or hyperbolic foliation of M . If $i_*\pi_1(L_0)$ is trivial for some leaf $L_0 \in \mathfrak{F}$, then the fundamental group of every leaf is abelian.*

Proof. — Choose a leaf $L'_0 \subset \pi^{-1}(L_0)$. Then

$$\text{kernel } \psi \subset \pi_1(M)_{L'_0} \cong i_*\pi_1(L_0)$$

and hence ψ is injective. Let $L \in \mathfrak{F}$. Choose $L' \subset \pi^{-1}(L)$ and let $z = \rho(L') \in \tilde{N}$. Since $\pi_2(\tilde{N}) = 0$ we know that $i_* : \pi_1(L) \rightarrow \pi_1(M)$ is one-one and so $\psi \circ i_*$ maps $\pi_1(L)$ isomorphically onto Σ_z . But $\Sigma_z \subset \{\tilde{g} \in \tilde{G} : \tilde{g}(z) = z\} \cong \text{SO}(2)$ and so Σ_z is abelian.

5.5. PROPOSITION. — *Suppose $\tilde{G}$ is solvable. If $\mathfrak{F}$ contains a leaf whose fundamental group is solvable, then $\pi_1(M)$ is solvable. Thus if $\tilde{G}$ is solvable and $\dim \mathfrak{F} = 1$, then $\pi_1(M)$ is solvable.*

Proof. — Suppose $L \in \mathfrak{F}$ is such that $\pi_1(L)$ is solvable. Then $i_*\pi_1(L)$ is solvable. Choose a leaf $L' \in \tilde{\mathfrak{F}}$ which projects to L . Then we have $\text{kernel } \psi \subset \pi_1(M)_{L'} \cong i_*\pi_1(L)$ and hence $\text{kernel } \psi$ is solvable. But $\pi_1(M)/\text{kernel } \psi \cong \Sigma$ is solvable since $\Sigma \subset \tilde{G}$. Thus $\pi_1(M)$ is solvable.

5.6. COROLLARY. — *If $\pi_1(M^3)$ is not solvable, then M^3 does not support a codimension two Euclidean foliation.*

5.7. PROPOSITION. — *If $H_1(M, \mathbb{Z}) = 0$, then M does not support a codimension two Euclidean foliation.*

Proof. — If $\mathfrak{F}$ is a codimension two Euclidean foliation of M , then we have $\psi : \pi_1(M) \rightarrow \tilde{G} = \text{SO}(2) \cdot \mathbb{R}^2$. Let $h : \text{SO}(2) \cdot \mathbb{R}^2 \rightarrow \text{SO}(2)$ be the projection. Since $H_1(M, \mathbb{Z}) = 0$ and $\text{SO}(2)$ is abelian, it follows that $h \circ \psi$ is the trivial homomorphism. Thus if $\tau \in \pi_1(M)$, then $\psi(\tau)$ is a translation of $\mathbb{R}^2$ and hence $\mathfrak{F}$ is a Lie $\mathbb{R}^2$ -foliation. Thus $\mathfrak{F}$ is defined by two linearly independent closed one-forms and hence, since $H^1(M) = 0$, there exists a submersion $M \rightarrow \mathbb{R}^2$ defining $\mathfrak{F}$ which is impossible.

5.8. PROPOSITION. — *Let $\mathfrak{F}$ be a codimension two Euclidean foliation of M . If all the leaves of $\mathfrak{F}$ are simply connected, then $\mathfrak{F}$ is a Lie $\mathbb{R}^2$ -foliation and $\pi_1(M)$ is abelian.*

Proof. — Let $\sigma \in \Sigma$. Suppose $\sigma(x) = x$ for some $x \in \mathbb{R}^2$. Choose $\tau \in \pi_1(M)$ such that $\psi(\tau) = \sigma$ and let $L' \in \tilde{\mathfrak{F}}$ be a leaf such that $\rho(L') = x$. Then $\tau(L') = L'$. Setting $L = \pi(L') \in \mathfrak{F}$, we have that $\tau \in \pi_1(M)_{L'} \cong i_*\pi_1(L)$. Since L is simply connected, it follows that τ , and hence σ , is the identity transformation. Thus Σ acts freely on $\mathbb{R}^2$ and so Σ is a group of translations. Hence $\mathfrak{F}$ is a Lie $\mathbb{R}^2$ -foliation. Finally, $\psi : \pi_1(M) \rightarrow \Sigma$ is an isomorphism and so $\pi_1(M)$ is abelian.

5.9. COROLLARY. — *Let $\mathfrak{F}$ be a codimension two Euclidean foliation of the 3-manifold M .*

- i) *If $\pi_1(M)$ is not abelian, then $\mathfrak{F}$ has a compact leaf.*
- ii) *If $\mathfrak{F}$ is not a Lie $\mathbb{R}^2$ -foliation, then $\mathfrak{F}$ has a compact leaf.*

5.10. PROPOSITION. — *Let $\mathfrak{F}$ be a codimension two Euclidean foliation of M and suppose that $\pi_1(M)$ is abelian. Then either*

- i) *$\mathfrak{F}$ is a Lie $\mathbb{R}^2$ -foliation, or*

ii) $\mathfrak{F}$ has a compact leaf L such that $i_* : \pi_1(L) \rightarrow \pi_1(M)$ is an isomorphism.

Proof. — Since $\pi_1(M)$ is abelian, we have that Σ is abelian. Hence all the non-identity elements of Σ have the same fixed point set Z . Either Z is empty or Z has one element. If Z is empty, then Σ is a group of translations and so $\mathfrak{F}$ is a Lie $\mathbf{R}^2$ -foliation. Suppose $Z = \{x\}$, $x \in \mathbf{R}^2$. Let $L' \in \hat{\mathfrak{F}}$ be a leaf such that $\rho(L') = x$ and let $L = \pi(L') \in \mathfrak{F}$. Then $\Sigma^L = \Sigma(x) = \{x\}$ and hence L is compact. Let $\tau \in \pi_1(M)$. Then, since $\psi(\tau)(x) = x$, we must have that $\tau(L') = L'$. Thus $\tau \in \pi_1(M)_{L'}$ and so

$$i_*\pi_1(L) \cong \pi_1(M)_{L'} = \pi_1(M).$$

5.11. COROLLARY. — Let $\mathfrak{F}$ be a codimension two Euclidean foliation of M where M is a 3-manifold with $\pi_1(M)$ abelian, $\pi_1(M) \neq \mathbf{Z}$. Then $\mathfrak{F}$ is a Lie $\mathbf{R}^2$ -foliation.

Proof. — If not, then $\mathfrak{F}$ has a compact leaf L such that $i_* : \pi_1(L) \rightarrow \pi_1(M)$ is an isomorphism. Since $\mathfrak{F}$ is one-dimensional, we have that $L \cong S^1$ and hence $\pi_1(M) = \mathbf{Z}$, a contradiction.

5.12. COROLLARY. — Let $\mathfrak{F}$ be a codimension two Euclidean foliation of M where M is an orientable 4-manifold with $\pi_1(M)$ abelian, $\pi_1(M) \neq \mathbf{Z} \times \mathbf{Z}$. Then $\mathfrak{F}$ is a Lie $\mathbf{R}^2$ -foliation.

Proof. — If not, then $\mathfrak{F}$ has a compact leaf L such that $i_* : \pi_1(L) \rightarrow \pi_1(M)$ is an isomorphism. Since M is orientable and $\mathfrak{F}$ is transversely orientable, we have that the leaves of $\mathfrak{F}$ are orientable. Thus L is a compact orientable surface. If the genus of L is one, then $\pi_1(M) = \mathbf{Z} \times \mathbf{Z}$, contrary to assumption. If the genus of L is greater than one, then $\pi_1(M)$ is not abelian, contrary to assumption. If the genus of L is zero, then M is simply connected and hence doesn't support a codimension two Euclidean foliation.

5.13. PROPOSITION. — If $\pi_1(M)$ has non-exponential growth, then M does not support a codimension two hyperbolic foliation.

Proof. — In this case $\tilde{G} = \mathrm{SL}(2, \mathbf{R})$ and we have $\psi : \pi_1(M) \rightarrow \mathrm{SL}(2, \mathbf{R})$ with image $\psi = \Sigma \subset \mathrm{SL}(2, \mathbf{R})$. By Lemma (4.3), $\Sigma \backslash \tilde{N}$ is compact and hence $\Sigma \backslash \tilde{G}$ is compact. Thus $\Sigma \backslash \mathrm{SL}(2, \mathbf{R})$ is compact which is impossible since Σ has non-exponential growth.

BIBLIOGRAPHY

- [1] H. BASS, The degree of polynomial growth of finitely generated nilpotent groups, *Proc. London Math. Soc.*, (3), 25 (1972), 603-614.
- [2] G. BREDON, Introduction to compact transformation groups, Academic Press, New York, 1972.
- [3] A. HAEFLIGER, Structures feuilletées et cohomologie à valeur dans un faisceau de groupoides, *Comm. Math. Helv.*, 32 (1958), 248-329.
- [4] S. KOBAYASHI and K. NOMIZU, Foundations of differential geometry, vol. I, *Interscience Tracts in Pure and Appl. Math.*, 15, Interscience, New York, 1963.
- [5] R. PALAIS, A global formulation of the Lie theory of transformation groups, *Memoirs of the Amer. Math. Soc.*, 22 (1957).
- [6] J. F. PLANTE, Foliations with measure preserving holonomy, *Ann. of Math.*, 102 (1975), 327-361.
- [7] M. S. RAGHUNATHAN, Discrete subgroups of Lie groups, *Ergebnisse der Mathematik und ihrer Grenzgebiete* (68), Springer-Verlag, Berlin, 1972.
- [8] B. REINHART, Foliated manifolds with bundle-like metrics, *Ann. of Math.*, (1), 69 (1959), 119-132.
- [9] M. SPIVAK, A comprehensive introduction to differential geometry, vol. I, Publish or Perish, Boston, 1970.
- [10] J. TITS, Free subgroups in linear groups, *J. of Alg.*, 20 (1972), 250-270.
- [11] J. WOLF, Growth of finitely generated solvable groups and curvature of Riemannian manifolds, *J. of Diff. Geom.*, 2 (1968), 421-446.

Manuscrit reçu le 20 décembre 1978
révisé le 9 juillet 1979.

Robert A. BLUMENTHAL,
Department of Mathematics
St. Louis University
St. Louis, Missouri 63103 (USA).
and
U.E.R. de Mathématiques,
Université de Lille I
59650 Villeneuve d'Ascq (France).
