

ANNALES DE L'INSTITUT FOURIER

ROBERT KAUFMAN

On the weak L^1 space and singular measures

Annales de l'institut Fourier, tome 32, n° 1 (1982), p. 119-128

[<http://www.numdam.org/item?id=AIF_1982__32_1_119_0>](http://www.numdam.org/item?id=AIF_1982__32_1_119_0)

© Annales de l'institut Fourier, 1982, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE WEAK L^1 SPACE AND SINGULAR MEASURES

by Robert KAUFMAN

Introduction.

The class R of finite, complex measures μ on $(-\infty, \infty)$ such that $\hat{\mu}(\infty) = 0$, has been intensively investigated (since 1916). For this class $o(1)$ is trivial and for absolutely continuous measures, we have the Riemann-Lebesgue Lemma. We investigate the corresponding $o(1)$ condition for the partial-sum operators

$$S_T(x, \mu) \equiv \int D_T(x-t) \mu(dt),$$

$$D_T(t) \equiv (\pi t)^{-1} \sin Tt, T > 0.$$

The $o(1)$ condition for S_T depends on the weak L^1 norm, defined by

$$\|u\|_1^* \equiv \sup_Y m\{|u| > Y\};$$

$$\|S_T(\mu)\|_1^* \leq C \|\mu\|, 0 < T < +\infty.$$

The weak estimate is an easy consequence of Kolmogorov's estimate for the Hilbert transform [2, Chapter II]. Elementary approximations show that when $\mu = f(x)dx$, then $\lim \|S_T(\mu) - f\|_1^* = 0$. When μ is singular and $\lim \|S_T(\mu) - g\|_1^* = 0$ for a certain measurable g , two conclusions can be obtained without great difficulty (see below):

- a) $\|S_k(\mu) - S_{k+1}(\mu)\|_1^* \rightarrow 0$ whence $\hat{\mu}(\infty) = 0$;
- b) $S_T(\mu) \rightarrow 0$ in measure as $T \rightarrow +\infty$

whence $g = 0$ a.e. This leads us to define:

(*) Presented at the Italian-American Conference on harmonic analysis, Minnesota, 1981.

W_0 is the class of measures μ for which $\|S_T(\mu)\|_1^* \rightarrow 0$ as $T \rightarrow +\infty$.

We present an elementary structural property of W_0 , and then show by example that

(A) There exist M_0 -sets F carrying no measure $\mu \neq 0$ in W_0 .

The sets F are defined by a purely metrical property, and they need not be especially small. Their construction is based on an idea from the theory of divergent Fourier series [3I, Chapter VIII].

(B) The set F_θ of all sums $\sum_{n=0}^{\infty} \pm \theta^n$ ($0 < \theta < 1/2$) carries a measure $\lambda \neq 0$ in W_0 , provided F_θ is an M_0 -set.

To elucidate example (B) and the next one we recall that F_θ fails to be an M_0 -set (or even an M -set) unless $\mu_\theta \in R$, where μ_θ is the Bernoulli convolution carried by F_θ and that $\mu_\theta \in R$ except for certain algebraic numbers θ [3II, p. 147-156]. Therefore the next example is somewhat unexpected.

(C) When $0 < \theta < 1/2$, then $\mu_\theta \notin W_0$, in fact

$$\|S_T(\mu_\theta)\|_1^* \geq c(\theta) > 0$$

for large $T > 0$. We observe in passing that μ is not known to be singular for $1/2 < \theta < 1$ except when $\mu_\theta \notin R$, e.g., for $\theta^{-1} = (1 + \sqrt{5})/2$.

1.

From the weak estimate for S_T it is clear that W_0 is norm-closed in the space of all measures. We shall prove that when $\mu \in W_0$ and $\psi \in C^1 \cap L^\infty$, then $\psi\mu \in W_0$; consequently the same is true if only $\psi \in L^1(\mu)$. We need two lemmas; the first was already used implicitly.

LEMMA 1. — *Let μ be a measure such that $S_k(\mu) - S_{k+1}(\mu) \rightarrow 0$ in measure (over finite intervals). Then $\hat{\mu}(\infty) = 0$, i.e., $\mu \in R$.*

Proof. — $|D_k(t) - D_{k+1}(t)| \leq \min(1, |t|^{-1}) \equiv K(t)$, say, and $K \in L^2(-\infty, \infty)$. Thus the functions $|S_k(\mu) - S_{k+1}(\mu)|$ have a common majorant $\int K(x-t) |\mu|(dt)$ in L^2 . The hypothesis on

$S_k - S_{k+1}$ then yields $\|S_k - S_{k+1}\|_2 \rightarrow 0$. This means that $\int_k^{k+1} (|\hat{\mu}(t)|^2 + |\hat{\mu}(-t)|^2) dt \rightarrow 0$ so $\hat{\mu}(\infty) = 0$, because $\hat{\mu}$ is uniformly continuous.

LEMMA 2. — Let $\mu \in R$ and $\psi \in C^1 \cap L^\infty$. Then as $T \rightarrow +\infty$ $\|S_T(x, \psi \cdot \mu) - \psi(x) S_T(x, \mu)\|_1^* \rightarrow 0$.

Proof. — Since μ can be approximated in norm by measures $\mu_n \in R$, each of compact support, we can suppose that μ itself has compact support, say $|t| \leq a$. Now $S_T(\psi \cdot \mu) - \psi S_T(\mu)$ converges to 0 uniformly on $[-a-1, a+1]$, being equal to

$$\pi^{-1} \int \sin T(t-x) \cdot \varphi(x, t) \mu(dt),$$

with $\varphi(x, t) = (t-x)^{-1} [\psi(t) - \psi(x)]$; $\varphi(x, t)$ is jointly continuous. This is sufficient to obtain the uniform convergence claimed.

For $|x| > a+1$ we write

$$x S_T(x, \mu) = \pi^{-1} \int \sin T(t-x) \cdot \sigma(x, t) \mu(dt)$$

with $\sigma(x, t) = x(t-x)^{-1}$; now $|\sigma| \leq a+1$ and

$$\left| \frac{\partial}{\partial t} \sigma(x, t) \right| \leq a+1,$$

for $|t| \leq a$. Therefore $x S_T(\mu, x) \rightarrow 0$ as $T \rightarrow +\infty$, uniformly for $|x| \geq a+1$. The same applies to $x S_T(x, \psi \cdot \mu)$, because $\psi \cdot \mu \in R$, and these inequalities show that $\psi S_T(\mu) - S_T(\psi \cdot \mu) \rightarrow 0$.

2. Examples.

I. Let F be a compact set in $(-\infty, \infty)$, $0 < \alpha < 1$, (ϵ_j) a sequence decreasing to 0; for each j , let $F = \cup F_k^j$, where

$$\text{diam}(F_k^j) \leq \epsilon_j, d(F_k^j, F_\ell^j) \geq \epsilon_j^\alpha, k \neq \ell.$$

Then F carries no probability measure μ in W_0 (and hence no signed measure $\mu \neq 0$ in W_0).

We define the following property of a number β in $[0, 1)$, relative to μ and the sequence of partitions $F = \cup F_k^j$:

(**) The total μ -measure of the sets F_k^j , such that $\mu(F_k^j) > \epsilon_j^\beta$, tends to 0, as $j \rightarrow +\infty$.

Plainly $\beta = 0$ has property (**), because μ , being an element of R , can have no discontinuities. We shall prove that if β has property (**), and $0 \leq \beta < \alpha$, then $\gamma = \beta + (1 - \alpha)/2$ has property (**). This leads to a contradiction as soon as $\gamma > \alpha$, since the number of sets $F_k^j \neq \emptyset$ is $O(\epsilon_j^{-\alpha})$.

Assuming that β has property (**), we form $\lambda = \lambda_j$, by omitting from F_k the intervals F_k^j of μ -measure $> \epsilon_j^\beta$. By Kolmogorov's estimate, $\|S_T(\lambda_j)\|_1^* \rightarrow 0$, as $j \rightarrow +\infty$ and $T \rightarrow +\infty$, independently. Let now \int^* denote an integral over the domain $|x - t| > \epsilon_j^\alpha/2$. Then

$$\begin{aligned} \int^* |x - t|^{-1} \lambda_j(dt) &= O(\epsilon_j^{-\alpha}), \text{ if } \beta = 0, \\ \int^* |x - t|^{-1} \lambda_j(dt) &= O(\epsilon_j^{\beta-\alpha} (\log \epsilon_j)), \quad 0 < \beta < \alpha. \end{aligned}$$

The first of these is obvious; the second is obtained by packing the subsets F_k^j as close to x as is consistent with the condition $d(F_k, F_\emptyset) \geq \epsilon_j^\alpha$.

For each k such that $\lambda_j(F_k^j) > \epsilon_j^\gamma$, we let ξ_k belong to F_k^j and consider the set defined by

$$\begin{aligned} (S_k^j): \quad \frac{1}{2} \lambda(F_k^j) \epsilon_j^\sigma &< |x - \xi_k| < \lambda(F_k^j) \epsilon_j^\sigma, \\ |\sin \epsilon_j^{-\tau} (x - \xi_k)| &> \frac{1}{2} \end{aligned}$$

where $\sigma = -\beta + 3\alpha/4 + 1/4$, $\tau = (1 + \gamma + \sigma)/2$.

The number $\lambda(F_k^j) \epsilon_j^\sigma$ lies between $\epsilon_j^{\beta+\sigma}$ and $\epsilon_j^{\gamma+\sigma}$; we note that $\beta + \sigma > \alpha$, and $\gamma + \sigma = 3/4 + \alpha/4 < 1$. Moreover $\epsilon_j^{-\tau} \epsilon_j = o(1)$, while $\epsilon_j^{-\tau} \lambda(F_k^j) \epsilon_j^\sigma \rightarrow +\infty$.

For each k in question, the Lebesgue measure of S_k^j is asymptotically $c\lambda(F_k^j) \epsilon_j^\sigma$, and the different sets are disjoint, because $\lambda(F_k^j) \epsilon_j^\sigma = o(\epsilon_j^\alpha)$. We shall prove that $|S_T(\lambda_j)| > c' \epsilon_j^{-\sigma}$ for a certain $c' > 0$, with $T = \epsilon_j^{-\tau} \rightarrow +\infty$. This will prove that the total μ -measure of the subsets F_k^j , such that $\epsilon_j^\gamma < \epsilon_j \leq \epsilon_j^\beta$, is $o(1)$.

When $x \in S_k^j$,

$$|S_T(x) - \int_{F_k^j} D_T(x - t) \lambda(dt)| < \int^* |x - t|^{-1} \lambda(dt),$$

and the error term on the right is $o(\epsilon_j^{-\sigma})$, because $\sigma > \alpha - \beta$.

When $t \in F_k^j$, $t - \xi_k = o(x - \xi_k)$ because $\gamma + \sigma < 1$, and $\sin T(t - x) = \sin T(\xi_k - x) + o(1)$ because $\tau < 1$. This easily leads to the lower bound on $|S_T(x)|$.

Our construction is adapted from Kolmogorov's divergent Fourier series [31, Chapter VIII].

To complete our example, we must present a set F that is also an M_0 -set. This is known for various M_0 -sets, but seems to occur explicitly in [1]: there exists a closed set $E \subseteq [0, 1]$ and a sequence of integers $N_k \rightarrow +\infty$ such that

(1) $|N_k x| < N_k^{-1}$ (modulo 1) for $x \in E$, $k \geq 1$,

(2) The mapping $y = e^x$ transforms E onto an M_0 -set. Then $y(E)$ is covered by intervals of length $\leq 2eN_k^{-2}$, whose distances are at least $(N_k^{-1} - 2N_k^{-2})$.

In the remaining examples it is occasionally convenient to write $S_T(y)$ in place of $S_T(y, \mu)$, when $\mu = \mu_\theta$.

II. We present example (C) first, because (B) is based on an improvement in one of the inequalities used in (C). For each $n = 0, 1, 2, 3, \dots$, F_θ is a union of 2^{n+1} sets E_k of diameter $2\theta^{n+1}(1 - \theta)^{-1}$, and mutual distances at least

$$2\theta^{n+1}(1 - 2\theta)(1 - \theta)^{-1} \equiv c_1 \theta^{n+1}; \mu(E_k) = 2^{-n-1}.$$

The lower bound on the mutual distances gives a Hölder condition on $\mu: \mu(B) \leq c_2 (\text{diam } B)^\alpha$, where $\alpha = -\log 2/\log \theta < 1$. If ξ_k is the center of E_k , we have an identity

$$\int_{E_k} f(t) \mu(dt) = 2^{-n-1} \int f(\xi_k + \theta^{n+1}t) \mu(dt).$$

For each set E_k , we define the set E_k^\sim by the inequality $d(x, E_k) < c_1 \theta^{n+1}/3$, so the sets E_k^\sim have distances at least $2c_1 \theta^{n+1}/3$. If $x \in E_k^\sim$, then

$$|S_T(x, \mu) - \int_{E_k} D_T(x - t) \mu(dt)| < \int_{R-E_k} |x - t|^{-1} \mu(dt),$$

and in the last integral, $|x - t| \geq 2c_1 \theta^{n+1}/3$. Hence, by the Hölder condition, the integral is $\leq c_3 (\theta^n)^{\alpha-1} = c_3 2^{-n} \theta^{-n}$. The principal term can be evaluated by the identity above, and simplified to the form $2^{-n} \theta^{-n-1} S_{T\theta^{n+1}}(\theta^{-n-1}x - \theta^{-n-1}\xi_k)$.

We observe that

$$\lim \int S_T(x, \mu) f(x) dx = \int f(x) \mu(dx),$$

for suitable test functions f ; for example, this is true if f and \dot{f} are integrable. Since μ is singular, we can find a test function f , such that $\|f\|_1 < 1$ and $|\int f(x) \mu(dx)| > 2c_3 + 2c_1^{-1}$. Hence $\max |D_T(\mu)| > 2c_3 + 2c_1^{-1}$ for large T , say for $T > T_0$.

Let $T > \theta^{-1} T_0$, and let $n \geq 0$ be chosen so that $T^* = \theta^{n+1} T$ satisfies the inequalities $T_0 \leq T^* \leq \theta^{-1} T_0$. Suppose that

$$|D_{T^*}(\theta^{-n-1}x - \theta^{-n-1}\xi_k)| > c_3 + c_1^{-1}.$$

Then $d(\theta^{-n-1}x - \theta^{-n-1}\xi_k, F_\theta) < c_1/3$, since $\pi > 3$, or $d(x, \xi_k + \theta^{n+1}F_\theta) < c_1\theta^{n+1}/3$, so $x \in E_k^\sim$. Hence

$$|D_T(x, \mu)| > c_3 \cdot 2^{-n-1}\theta^{-n-1} - c_3 2^{-n}\theta^{-n} = c_4 2^{-n}\theta^{-n}.$$

But it is easy to see that the set of x 's in question has measure at least $c_5 2^n \theta^n$, because $T_0 \leq T^* \leq \theta^{-1} T_0$, and the functions D_{T^*} have derivatives bounded by $\theta^{-2} T_0^2$. Hence $\|D_T(\mu)\|_1^* \geq c_4 c_5$.

III. The example (B) requires a complicated construction, but relies in essence on small improvements on estimates already used. To estimate $S_T(\mu, x)$ we divide the range of integration into the subsets $\{|x - t| < T^{-1}\}$ and $\{|x - t| > T^{-1}\}$. The second yields an integral $O(T^{1-\alpha})$, by the Hölder condition, and the first yields $T \cdot O(T^{-\alpha}) = O(T^{1-\alpha})$ for the same reason (and the inequality $|D_T| < T$).

We give another estimate on $S_T(x, \mu)$ for large T , supposing that $\mu \in R$.

LEMMA 3. — To each $\epsilon > 0$ there is a T_0 such that

$$|S_T(x, \mu)| < \epsilon d(x, F_\theta)^{-1}$$

whenever $T \geq T_0$ and $d \equiv d(x, F_\theta) \geq \epsilon$.

Proof. — Let $\delta = d(x, F)$ and observe that

$$\delta S_T(x, \mu) = \pi^{-1} \int \sin T(x - t) \cdot \delta \cdot (x - t)^{-1} \mu(dt).$$

The function $g(t) = \delta \cdot (x - t)^{-1}$ is bounded by 1 on F , and

$|g(t_1) - g(t_2)| \leq \delta^{-1} |t_1 - t_2|$ for numbers t_1, t_2 in F_θ . Hence the conclusion follows from our assumption that $\mu \in R$ and the Tietze extension theorem.

The inequality of the Lemma can be written in a more useful way. When $t \in F_\theta$, then $|x - t| \leq d + 2 \leq d(1 + 2\epsilon^{-1})$. Hence $d(x, F_\theta)^{-1} \leq (1 + 2\epsilon^{-1}) \int |x - t|^{-1} \mu(dt)$. Suppose now that $x \notin E_k^\sim$ so that $d(\theta^{-n-1}x - \theta^{-n-1}\xi_k, F_0) \geq c_1 \theta^{n+1}/3$. Using the identity for integrals over E_k , we find the following estimate:

If $x \notin E_k^\sim$ and $T\theta^{n+1} > T_{00}$, then

$$\left| \int_{E_k} D_T(x - t) \mu(dt) \right| < \epsilon \int_{E_k} |x - t|^{-1} \mu(dt).$$

Consequently, when $x \in E_\ell^\sim$ and $T\theta^{n+1}$ is sufficiently large (depending on $\epsilon > 0$)

$$|S_T(x, \mu) - 2^{-n-1} \theta^{-n-1} S_{T\theta^{n+1}}(\theta^{-n-1}x - \theta^{-n-1}\xi_\ell)| < \epsilon \theta^{n(\alpha-1)}.$$

LEMMA 4. — To each $\epsilon > 0$ there is a $\delta > 0$ so that, when $\theta^{-1} < Y < \delta T^{1-\alpha}$ then $Ym\{|S_T(x, \mu)| > Y\} < \epsilon$.

Proof. — We choose $n \geq 0$ so that $1 < \theta^{n+1} Y^{1/1-\alpha} < \theta^{-1}$; this leads to the inequalities $\theta^{n(\alpha-1)} > Y$, and $T\theta^{n+1} > \delta^{-1}$. For fixed ℓ , we must estimate the Lebesgue measure of the set defined by

$$|S_{T\theta^{n+1}}(\mu, \theta^{-n-1}x - \theta^{-n-1}\xi_\ell)| > \frac{1}{2} \cdot 2^{n+1} \theta^{n+1} Y.$$

The right hand side exceeds $\frac{1}{2} \theta^{-1}$; when $T\theta^{n+1}$ is large, the measure of the set is at most $\epsilon \theta^{n+1}$; the total for all ℓ is at most $\epsilon 2^{n+1} \theta^{n+1} < \epsilon Y^{-1}$. Hence $Ym\{|S_T(x, \mu)| > Y\} < \epsilon$.

In view of the inequality $|S_T(\mu, x)| = O(T^{1-\alpha})$, the conclusion of the last lemma holds when $Y > \delta^{-1} T^{1-\alpha}$, $T > 1$, for a certain $\delta > 0$.

In preparation for the next lemma, we recall the identity ($n = 1, 2, 3, \dots$)

$$\int f(t) \mu(dt) \equiv 2^{-n} \sum_{k=1}^{2^n} \int f(\xi_k + \theta^n t) \mu(dt).$$

We define $\int f(t) \sigma_n(dt) \equiv 2^{-n} \sum_k \int f(\xi_k + \theta^{n+k} t) \mu(dt)$. Then

$\sigma_n = g_n \cdot \mu$, where $g_n \geq 0$, g_n is continuous on F_θ and takes the values 0 and 2^k ($1 \leq k \leq 2^n$). Using the formula for σ_n we get an identity

$$S_T(x, \sigma_n) = 2^{-n} \theta^{-n} \sum_k \theta^{-k} S_{T\theta^{n+k}}(\theta^{-n-k}x - \theta^{-n-k}\xi_k).$$

LEMMA 5. — To each $\epsilon > 0$, there is an $N > 1$ such that $\limsup_{T \rightarrow +\infty} \|S_T(\sigma_n)\|_1^* < \epsilon$, if $n \geq N$.

Proof. — In calculating $\limsup_{T \rightarrow +\infty} \|S_T(\sigma_n)\|_1^*$ we can omit x 's outside $(-3, 3)$, because $\sigma_n \in \mathbb{R}$. In an obvious notation we write $\sigma_n = \sum_k \sigma_{n,k}$, and observe that, for $T > T_{n,\epsilon}$

$$|S_T(\sigma_n)| < \max_k |S_T(\sigma_{n,k})| + \epsilon/12.$$

When $Y > \epsilon/6$ (the others are trivial, since we suppose that $|x| < 6$),

$$\begin{aligned} m\{|S_T(\sigma_n)| > 2Y\} &\leq \sum_k m\{|S_T(\sigma_{n,k})| > Y\} \\ &= \sum_k \theta^{n+k} m\{|S_{T\theta^{n+k}}(x, \mu)| > 2^n \theta^{n+k} Y\}. \end{aligned}$$

Each summand is $O(2^{-n} Y^{-1})$ by Kolmogorov's inequality; if $T\theta^{n+k} > 1$, then the k -th term exceeds $\epsilon 2^{-n} Y$ only if

$$\delta(T\theta^{n+k})^{1-\alpha} < Y < \delta^{-1}(T\theta^{n+k})^{1-\alpha},$$

by Lemma 4 and the remark after it, and this inequality occurs for at most $2(1-\alpha)^{-1} \cdot \log \delta / \log \theta$ indices $k = 1, \dots, 2^n$. (We assume that $Y > \theta^{-1}$, since $S_T(\sigma_n) \rightarrow 0$ almost everywhere as $T \rightarrow +\infty$.) This proves our lemma.

A further property of σ_n , obtained simply by increasing n , is the inequality $|\sigma_n(I) - \mu(I)| < \epsilon$ for all intervals I .

The next lemma establishes a property of the functional $\|\cdot\|_1^*$ to simplify the remaining calculations.

LEMMA 6. — Let $a_i = \|f_i\|_1^*$ $1 \leq i \leq N$. Then $\|\sum f_i\|_1^* \leq (\sum a_i^{1/2})^2$.

Proof. — Let $0 \leq t_i \leq 1$, and $\sum t_i = 1$. Then

$$m\{|\sum f_i| \geq Y\} \leq \sum m\{|f_i| \geq t_i Y\} \leq \sum t_i^{-1} Y^{-1} a_i.$$

The minimum of the sum is $Y^{-1}(\sum a_i^{1/2})^2$. With a little more effort, we can obtain the bound $c(1-p)^{-1}(\sum a_i^p)^{1/p}$, $0 < p < 1$.

We are now in a position to construct the measure λ . We shall find probability measures $\lambda_k = f_k \mu$, with $f_k \geq 0$, $\int f_k d\mu = 1$, such that $\|S_T(\lambda_k)\|_1^* < k^{-1}$ for $T > T_k > T_{k-1} \dots$ and $|\hat{\lambda}_k(u)| < k^{-2}$ for $u > T_k$. Lemma 5 provides λ_1 ; let us suppose that λ_k and T_k are known. We find σ_k so that $|\sigma_k(I) - \lambda_k(I)| < k^{-1}(1 + T_k)^{-2}$ and $\|S_T(\sigma_k)\|_1^* < k^{-4}/25$, and $|\hat{\sigma}_k(u)| < k^{-1}$, for $u > T_{k+1}^0 > T_k$. (The construction of $f_{k+1}\mu$ from $f_k\mu$ follows Lemma 5). We now set $\lambda_{k+1} = (1 - k^{-1/2})\lambda_k + k^{-1/2}\sigma_k$; by Lemma 6, we have for $T > T_{k+1}^0$

$$\|S_T(\lambda_{k+1})\|_1^{*1/2} \leq (1 - k^{-1/2})^{1/2} k^{-1/2} + k^{-2}/5.$$

When $k = 1$, the last bound is $1/5$, while $(k+1)^{-1} = \frac{1}{2}$. For $k \geq 2$, we need the inequality

$$(1 - k^{-1/2})^{1/2} k^{-1/2} + k^{-2}/5 < (k+1)^{-1/2},$$

which can be verified with the aid of calculus. Clearly, we have $|\hat{\lambda}_{k+1}(u)| < (k+1)^{-2}$ for $T > T_{k+1}^{00}$; we take $T_{k+1} = T_{k+1}^0 + T_{k+1}^{00}$.

By the construction, and integration by parts,

$$|\hat{\lambda}_k(u) - \hat{\lambda}_{k+1}(u)| \leq k^{-3/2}(1 + T_k)^{-2}|u|;$$

consequently $|\hat{\lambda}_k(u) - \hat{\lambda}_{k+1}(u)| \leq k^{-3/2}$ unless $|u| > 1 + T_k$. However, if $|u| > T_{k+1} > T_k$, then $|\hat{\lambda}_k(u) - \hat{\lambda}_{k+1}(u)| < 2k^{-2}$. Since $|\hat{\lambda}_k - \hat{\lambda}_{k+1}| \leq 2k^{-1/2}$, we have a limit $\varphi(u)$, with

$$|\varphi - \hat{\lambda}_k| = O(k^{-1/2}).$$

Hence $\varphi = \hat{\lambda}$, with λ carried by F_θ and $\lambda \in R$.

In verifying that $\lim \|S_T(\lambda)\|_1^* = 0$ we can calculate the weak norms over $(-3, 3)$. Suppose that $T_{k-1} \leq T \leq T_k$; then

$$|S_T(\lambda_k) - S_T(\lambda)| = O(k^{-1/2}).$$

Since $T \geq T_{k-1}$, $\|S_T(\lambda_{k-1})\|_1^* < (k-1)^{-1}$; and finally

$$\|S_T(\lambda_k) - S_T(\lambda_{k-1})\|_1^* = O(k^{-1/2}).$$

Hence $\|S_T(\lambda)\|_1^* = O(k^{-1/2})$ over $(-3, 3)$.

BIBLIOGRAPHY

- [1] R. KAUFMAN, On transformations of exceptional sets, *Bull. Greek Math. Soc.*, 18 (1977), 176-185.
- [2] E.M. STEIN, *Singular Integrals and Differentiability Properties of Functions*, Princeton, 1970.
- [3] A. ZYGMUND, *Trigonometric Series, I, II*, Cambridge, 1959 and 1968.

Manuscrit reçu le 23 février 1981.

Robert KAUFMAN,
University of Illinois at
Urbana-Champaign
Department of Mathematics
Urbana, Ill. 61801 (U.S.A.).