

ALGEBRAIC COMBINATORICS

Sridhar P. Narayanan, Digjoy Paul, Amritanshu Prasad & Shraddha Srivastava

Character Polynomials and the Restriction Problem

Volume 4, issue 4 (2021), p. 703-722.

[<http://alco.centre-mersenne.org/item/ALCO_2021__4_4_703_0>](http://alco.centre-mersenne.org/item/ALCO_2021__4_4_703_0)

© The journal and the authors, 2021.
Some rights reserved.




This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>

Access to articles published by the journal *Algebraic Combinatorics* on
the website <http://alco.centre-mersenne.org/> implies agreement with the
Terms of Use (<http://alco.centre-mersenne.org/legal/>).



Algebraic Combinatorics is member of the
Centre Mersenne for Open Scientific Publishing
www.centre-mersenne.org



Character Polynomials and the Restriction Problem

Sridhar P. Narayanan, Digjoy Paul, Amritanshu Prasad
 & Shraddha Srivastava

ABSTRACT Character polynomials are used to study the restriction of a polynomial representation of a general linear group to its subgroup of permutation matrices. A simple formula is obtained for computing inner products of class functions given by character polynomials. Character polynomials for symmetric and alternating tensors are computed using generating functions with Eulerian factorizations. These are used to compute character polynomials for Weyl modules, which exhibit a duality. By taking inner products of character polynomials for Weyl modules and character polynomials for Specht modules, stable restriction coefficients are easily computed. Generating functions of dimensions of symmetric group invariants in Weyl modules are obtained. Partitions with two rows, two columns, and hook partitions whose Weyl modules have non-zero vectors invariant under the symmetric group are characterized. A reformulation of the restriction problem in terms of a restriction functor from the category of strict polynomial functors to the category of finitely generated FI-modules is obtained.

1. INTRODUCTION

Let K be a field of characteristic 0. Let $P = K[X_1, X_2, \dots]$, a ring of polynomials in infinitely many variables. Regard P as a graded algebra where the variable X_i has degree i . In this grading, the monomial $X_1^{a_1} X_2^{a_2} \dots$ has degree $\sum_i i a_i$.

DEFINITION 1.1. For each $n \geq 1$, let V_n be a representation of the symmetric group S_n . The collection $\{V_n\}_{n=1}^\infty$ is said to have eventually polynomial character, if there exists $q \in P$ and a positive integer N such that, for each $n \geq N$ and each $w \in S_n$,

$$\text{trace}(w; V_n) = q(X_1(w), X_2(w), \dots),$$

where $X_i(w)$ is the number of i -cycles in w . The collection $\{V_n\}$ is said to have polynomial character if $N = 1$. The polynomial q is called the character polynomial of $\{V_n\}$.

Character polynomials have been used to study characters of families of representations of symmetric groups that occur naturally in combinatorics, topology and other areas. A survey of their history can be found in the article of Garsia and Goupil [6]. More recently, Church, Ellenberg and Farb [3] developed the theory of FI-modules. They showed that each finitely generated FI-module gives rise to a family of representations with eventually polynomial character.

Manuscript received 7th February 2020, revised 9th April 2021, accepted 14th April 2021.

KEYWORDS. character polynomial, restriction problem.

ACKNOWLEDGEMENTS. AP was partially supported by Swarnajayanti Fellowship grant DST/SJF/MSA-02/2014-15.

Any polynomial $q \in P$ gives rise to a class function on S_n for every positive integer n . The value of this function at $w \in S_n$ is obtained by substituting for X_i the number of i -cycles in w . For each n , we define the moment of q as the average value of the associated class function on S_n . The ring P has a basis indexed by integer partitions, which we call the binomial basis (Definition 2.2). We give an explicit formula for the moment of a binomial basis element (Theorem 2.3). This formula can be used to compute inner products of class functions coming from character polynomials. It implies that such an inner product achieves a constant value for large n (Corollary 2.4). This is a character-theoretic analogue of [3, Theorem 1.13], which establishes representation stability for finitely generated FI-modules.

The homogeneous polynomial representations of degree d general linear groups $GL_n(K)$ and the representations of the symmetric group S_d are linked via Schur–Weyl duality (see [7]). The restriction problem explores a different relationship between polynomial representation of $GL_n(K)$ and S_n . It asks how an irreducible polynomial representation of $GL_n(K)$ decomposes when it is restricted to S_n sitting inside $GL_n(K)$ as the subgroup of permutation matrices.

For each partition λ , let W_λ denote the Weyl functor (see [1, Definition II.1.3]) associated to λ . Let $P(n, d)$ denote the set of all partitions of d with at most n parts. Then $W_\lambda(K^n)$, as λ runs over $P(n, d)$, are the irreducible polynomial representations of the general linear group $GL_n(K)$ of degree d .

Following the notation and terminology of [3, Definition 2.2.5], for a partition $\mu = (\mu_1, \dots, \mu_m)$ of size $|\mu|$ and an integer $n \geq \mu_1 + |\mu|$, let $\mu[n]$ denote the padded partition $(n - |\mu|, \mu_1, \mu_2, \dots, \mu_m)$. Let $V_{\mu[n]}$ denote the Specht module of S_n corresponding to $\mu[n]$.

Consider the decomposition of the restriction of $W_\lambda(K^n)$ to S_n into Specht modules:

$$\text{Res}_{S_n}^{GL_n(K)} W_\lambda(K^n) = \bigoplus_{\mu} V_{\mu[n]}^{\oplus r_{\lambda\mu}(n)},$$

where the sum is over partitions μ such that $n - |\mu| \geq \mu_1$. It is well-known that the coefficients $r_{\lambda\mu}(n)$ are eventually constant for large n (this result is attributed to D. E. Littlewood by Assaf and Speyer [2]). Let $r_{\lambda\mu}$ be their eventually constant value, which is called the *stable restriction coefficient*. Finding a combinatorial interpretation of $r_{\lambda\mu}$ is known as the *restriction problem*.

In this article we show that the family $\{\text{Res}_{S_n}^{GL_n(K)} W_\lambda(K^n)\}$ has polynomial character. We determine its character polynomial S_λ (Theorem 2.7) by applying the Jacobi–Trudi identities to the character polynomials of symmetric and exterior powers of K^n (Theorem 2.6). The character polynomials of symmetric and exterior tensor powers of K^n have generating functions with Eulerian factorization (Theorem 2.6). Multiplying S_λ by the character polynomial q_μ of Specht modules $\{V_{\mu[n]}\}$ (which was computed by Macdonald [16, Example I.7.14(b)] and Garsia–Goupil [6]), and then taking moments (Theorem 2.3) gives an algorithm to compute stable restriction coefficients (Theorem 3.3). Assaf and Speyer [2] and independently, Orellana and Zabrocki [19] introduced *Specht symmetric functions* to study the restriction problem. In Section 3.3 we explain the relationship between these two approaches.

Notwithstanding several interesting recent developments [2, 8, 19, 18], a solution to the restriction problem remains elusive. Even interpreting $r_{\lambda\emptyset}$ combinatorially (here $\emptyset$ denotes the empty partition of 0, so $r_{\lambda\emptyset}$ is the dimension of the space of S_n -invariant vectors in $W_\lambda(K^n)$ for large n) appears to be a non-trivial and interesting problem. We provide generating functions in λ for the dimension of the space of S_n -invariant vectors in $W_\lambda(K^n)$ (Corollary 4.3). Using our main generating function (Theorem 4.1) for the dimension of S_n -invariants in mixed tensors, we are able to

characterize partitions with two rows, two columns and hook partitions which have non-zero S_n -invariant vectors. To the best of our knowledge, the specific problem of determining the S_n -invariant vectors of a Weyl-module has not been studied before.

We conclude this paper by placing the restriction problem in the context of strict polynomial functors and FI-modules. Friedlander and Suslin [5] introduced strict polynomial functors of degree d . The polynomial representations of degree d of $GL_n(K)$ are obtained by evaluating strict polynomial functors of degree d at K^n . Similarly, families of representations of S_n with stability properties can be obtained by evaluating finitely generated FI-modules at $\{1, \dots, n\}$ (see [3]). We define a functor from the category of strict polynomial functors of degree d to the category of finitely generated FI-modules for every d (Section 5.3). This functor corresponds to restriction of representations from $GL_n(K)$ to S_n under evaluation functors (Theorem 5.1).

2. CHARACTER POLYNOMIALS AND THEIR MOMENTS

2.1. MOMENTS AND STABILITY.

DEFINITION 2.1 (Moment). *The moment of $q \in P$ at n is defined as:*

$$\langle q \rangle_n = \frac{1}{n!} \sum_{w \in S_n} q(X_1(w), X_2(w), \dots).$$

We shall express integer partitions in *exponential notation*: given a partition α with largest part r , we write:

$$\alpha = 1^{a_1} 2^{a_2} \dots r^{a_r},$$

where a_i is the number of parts of α of size i for each $1 \leq i \leq r$. Thus α is a partition of the integer $|\alpha| := a_1 + 2a_2 + \dots + ra_r$. For every integer partition $\alpha = 1^{a_1} \dots r^{a_r}$ define $\binom{X}{\alpha} \in P$ by:

$$\binom{X}{\alpha} = \binom{X_1}{a_1} \binom{X_2}{a_2} \dots \binom{X_r}{a_r}.$$

DEFINITION 2.2 (Binomial basis). *The basis of P consisting of elements*

$$\left\{ \binom{X}{\alpha} \mid \alpha \text{ is an integer partition} \right\}$$

is called the binomial basis of P .

For an integer partition $\alpha = 1^{a_1} 2^{a_2} \dots r^{a_r}$, define $z_\alpha = \prod_{i=1}^r i^{a_i} a_i!$. This is the order of the centralizer in S_n of a permutation with cycle type α .

THEOREM 2.3. *For every integer partition $\alpha = 1^{a_1} 2^{a_2} \dots$, we have:*

$$\left\langle \binom{X}{\alpha} \right\rangle_n = \begin{cases} 0 & \text{if } n < |\alpha|, \\ 1/z_\alpha & \text{otherwise.} \end{cases}$$

Proof. We have:

$$\sum_{n \geq 0} \left\langle \binom{X}{\alpha} \right\rangle_n v^n = \sum_{n \geq 0} \frac{1}{n!} \sum_{w \in S_n} \prod_{i \geq 1} \binom{X_i(w)}{a_i} v^{iX_i(w)}.$$

Replace the sum $w \in S_n$ by a sum over conjugacy classes in S_n . If $\beta = 1^{b_1} 2^{b_2} \dots$ is a partition of n , then the number of elements in S_n with cycle type β is $\frac{n!}{\prod_i i^{b_i} b_i!}$. We

get:

$$\begin{aligned}
 \sum_{n \geq 0} \left\langle \binom{X}{\alpha} \right\rangle_n v^n &= \sum_{n \geq 0} \sum_{\beta \vdash n} \prod_{i \geq 1} \frac{v^{ib_i}}{i^{b_i} b_i!} \binom{b_i}{a_i} \\
 &= \prod_{i \geq 1} \sum_{b_i \geq a_i} \frac{v^{ib_i}}{i^{b_i} b_i!} \binom{b_i}{a_i} \\
 &= \prod_{i \geq 1} \sum_{b_i \geq a_i} \frac{v^{ib_i}}{i^{b_i} a_i! (b_i - a_i)!} \\
 &= \prod_{i \geq 1} \sum_{b_i \geq a_i} \frac{v^{ia_i}}{i^{a_i} a_i!} \frac{v^{i(b_i - a_i)}}{i^{b_i - a_i} (b_i - a_i)!}.
 \end{aligned}$$

Setting $c_i = b_i - a_i$ gives:

$$\begin{aligned}
 \sum_{n \geq 0} \left\langle \binom{X}{\alpha} \right\rangle_n v^n &= \frac{v^{|\alpha|}}{z_\alpha} \prod_{i \geq 1} \sum_{c_i \geq 0} \frac{v^{ic_i}}{i^{c_i} c_i!} \\
 &= \frac{v^{|\alpha|}}{z_\alpha} \sum_{n \geq 0} v^n \sum_{\gamma \vdash n} \frac{1}{z_\gamma}.
 \end{aligned}$$

Since $\sum_{\gamma \vdash n} 1/z_\gamma = 1$ for every n , we get:

$$(1) \quad \sum_{n \geq 0} \left\langle \binom{X}{\alpha} \right\rangle_n v^n = \frac{v^{|\alpha|}}{z_\alpha} \frac{1}{1-v},$$

from which Theorem 2.3 follows. $\square$

For two representations V and W of S_n , let:

$$\langle V, W \rangle_n = \dim \operatorname{Hom}_{S_n}(V, W),$$

which is the same as the Schur inner product of their characters:

$$\langle V, W \rangle_n = \frac{1}{n!} \sum_{w \in S_n} \operatorname{trace}(w; V) \operatorname{trace}(w, W).$$

COROLLARY 2.4. *For any $q \in P$ of degree d , $\langle q \rangle_n = \langle q \rangle_d$ for all $n \geq d$. In particular, if $\{V_n\}$ and $\{W_n\}$ are families of representations with polynomial characters of degree d_1 and d_2 , then $\langle V_n, W_n \rangle_n$ stabilizes for $n \geq d_1 + d_2$.*

Proof. This follows from the fact that the polynomials $\binom{X}{\alpha}$, as α runs over the set of integer partitions, form a basis of P . $\square$

DEFINITION 2.5 (Stable moment). *For a polynomial $q \in P$ we define the stable moment $\langle q \rangle$ of q to be the eventually constant value of $\langle q \rangle_n$:*

$$\langle q \rangle = \lim_{n \rightarrow \infty} \langle q \rangle_n.$$

Let $V_n = V_{\lambda[n]}$, the Specht module of S_n corresponding to the padded partition $\lambda[n]$. It is well-known that $\{V_n\}$ is a family of representations with eventually polynomial character [6, Proposition I.1]. In other words, for every partition λ , there exists a polynomial $q_\lambda \in P$ such that

$$(2) \quad \chi_{\lambda[n]}(w) = q_\lambda(X_1(w), X_2(w), \dots) \text{ for } n \geq |\lambda| + \lambda_1,$$

where $\chi_{\lambda[n]}$ denotes the character of the Specht module $V_{\lambda[n]}$. Given three partitions λ , μ , and ν of the same integer k , let $g_{\lambda\mu\nu}(n)$ denote the multiplicity of $V_{\lambda[n]}$ in $V_{\mu[n]} \otimes V_{\nu[n]}$. Then

$$g_{\lambda\mu\nu}(n) = \langle q_{\lambda} q_{\mu} q_{\nu} \rangle_n.$$

By Corollary 2.4, $g_{\lambda\mu\nu}(n)$ is eventually constant, recovering a well-known theorem of Murnaghan (see [15]). Church, Ellenberg, and Farb [3, Section 3.4] point out that this result can also be obtained by showing that the families $V_{\mu[n]} \otimes V_{\nu[n]}$ and $V_{\lambda[n]}$ come from finitely generated FI-modules.

2.2. SYMMETRIC AND ALTERNATING TENSORS. Let Sym^d and $\wedge^d$ denote the symmetric and alternating tensor functors respectively. For every $n \geq 0$, $\text{Sym}^d(K^n)$ and $\wedge^d(K^n)$ can be regarded as representations of S_n . In this section, we will prove that they have polynomial character by direct computation.

We shall work with generating functions that live in the ring $P[[t]]$ of formal power series in the variable t and coefficients in P . Let $\left(\begin{smallmatrix} X_i \\ j \end{smallmatrix}\right) = \binom{X_i+j-1}{j}$. The ring $P[[t]]$ admits expressions of the kind

$$(3) \quad (1-t^i)^{-X_i} = \sum_{j \geq 0} \left(\begin{smallmatrix} X_i \\ j \end{smallmatrix}\right) t^{ij},$$

$$(4) \quad (1+(-t)^i)^{X_i} = \sum_{j \geq 0} (-1)^{ij} \left(\begin{smallmatrix} X_i \\ j \end{smallmatrix}\right) t^{ij},$$

which will be needed later. Also, given $w \in S_n$ for any n and $q \in P$, we write $q(w) = q(X_1(w), X_2(w), \dots)$.

THEOREM 2.6. *Let $\{H_d\}_{d=0}^{\infty}$ be the sequence of polynomials in P defined by:*

$$(5) \quad \sum_{d=0}^{\infty} H_d t^d = \prod_{i=1}^{\infty} (1-t^i)^{-X_i},$$

an identity in the formal power series ring $P[[t]]$.

Then for every $n \geq 1$ and every $w \in S_n$,

$$H_d(w) = \text{trace}(w; \text{Sym}^d(K^n)).$$

Let $\{E_d\}_{d=0}^{\infty}$ be the sequence of polynomials in P defined by:

$$(6) \quad \sum_{d=0}^{\infty} E_d t^d = \prod_{i=1}^{\infty} (1-(-t)^i)^{X_i}.$$

Then for every $n \geq 1$ and every $w \in S_n$,

$$E_d(w) = \text{trace}(w; \wedge^d(K^n)).$$

For every positive integer d , we have:

$$(7) \quad H_d = \sum_{\alpha \vdash d} \prod_{i=1}^d \left(\begin{smallmatrix} X_i \\ a_i \end{smallmatrix}\right),$$

$$(8) \quad E_d = \sum_{\alpha \vdash d} (-1)^{a_2+a_4+\dots} \prod_{i=1}^d \left(\begin{smallmatrix} X_i \\ a_i \end{smallmatrix}\right).$$

Proof. The standard basis of K^n is indexed by the set $[n] = \{1, \dots, n\}$. The space $\text{Sym}^d K^n$ has an induced basis indexed by multisets of size d with elements drawn from $[n]$. The trace of $w \in S_n$ on $\text{Sym}^d K^n$ is the number of such multisets that are fixed by w . In a multiset that is fixed by w , the elements in each cycle of w

appear with the same multiplicity. Hence a multiset fixed by w may be regarded as a multiset of cycles of w . Assign weight $\text{wt}(C) = t^i$ to each i -cycle C of w . To a multiset $M = \{C_1^{m_1} \cdots C_r^{m_r}\}$ of cycles of w , assign weight $\text{wt}(M) = \prod_{1 \leq j \leq r} \text{wt}(C_j)^{m_j}$. Then

$$\sum_{d \geq 0} \text{trace}(w; \text{Sym}^d K^n) t^d = \sum_M \text{wt}(M),$$

where the sum runs over all multisets M of cycles of w . Each of the $X_i(w)$ i -cycles of w contributes a factor $(1 - t^i)^{-1}$ to this generating function, so

$$\sum_{d \geq 0} \text{trace}(w; \text{Sym}^d K^n) t^d = \prod_{i \geq 1} (1 - t^i)^{-X_i(w)}.$$

Similarly, $\wedge^d K^n$ has a basis indexed by subsets of $[n]$ with d elements. Only subsets fixed by w contribute to the trace, and these are unions of the cycles of w . A cycle of length i changes the sign of the corresponding basis vector by a factor of $-(-1)^i$. Assign weight $\text{wt}(C) = -(-t)^i$ to each i -cycle C of w . To a subset $N = \{C_1, \dots, C_r\}$ of cycles of w , assign weight $\text{wt}(N) = \prod_{1 \leq i \leq r} \text{wt}(C_i)$. Then

$$\sum_{d \geq 0} \text{trace}(w; \wedge^d K^n) t^d = \sum_N \text{wt}(N),$$

where the sum runs over all subsets N of the set of cycles of w . Each of the $X_i(w)$ i -cycles of w contributes a factor $(1 - (-t)^i)$ to this generating function, so

$$\sum_{d \geq 0} \text{trace}(w; \wedge^d K^n) t^d = \prod_{i \geq 1} (1 - (-t)^i)^{X_i(w)}.$$

Expansion of the products in (5) and (6) using (3) and (4) gives (7) and (8) respectively. $\square$

2.3. CHARACTER POLYNOMIALS OF WEYL MODULES. Applying the Jacobi–Trudi identities [16, Section I.3] to the character polynomials of Sym^d and $\wedge^d$ gives character polynomials for Weyl functors. For a partition λ , let λ' denote its conjugate partition.

THEOREM 2.7. *For every partition λ , the element of P defined by*

$$(9) \quad S_\lambda = \det(H_{\lambda_i + j - i}) = \det(E_{\lambda'_i + j - i})$$

is such that for every positive integer n and every $w \in S_n$,

$$S_\lambda(w) = \text{trace}(w; W_\lambda(K^n)).$$

The polynomials S_λ for partitions λ of integers at most 5 are given in Table 1. The highest degree coefficients in these expansions are character values of symmetric groups. More precisely, we have:

THEOREM 2.8. *Let λ be a partition of a positive integer d . For every partition $\alpha = 1^{a_1} 2^{a_2} \cdots$ of d , the coefficient of $\binom{X}{\alpha}$ in the expansion of S_λ in the binomial basis (Definition 2.2) is $\chi_\lambda(w_\alpha)$, where w_α is a permutation with cycle type α .*

The theorem will be a consequence of the following lemma:

LEMMA 2.9. *Let λ be a partition of a positive integer d . For every partition α of d , the coefficient of $\binom{X}{\alpha}$ in the expansion of H_λ in the binomial basis is $\sigma_\lambda(w_\alpha)$, the value of the character σ_λ of the permutation representation of S_d induced from the trivial representation of the Young subgroup $S_{\lambda_1} \times \cdots \times S_{\lambda_l}$ (see [9, Section 2.2] or [20, Section 2.3]) at a permutation w_α with cycle type α .*

λ	S_λ
$n = 1$	
(1)	X_1
$n = 2$	
(2)	$\frac{1}{2}X_1^2 + \frac{1}{2}X_1 + X_2$
(1 ²)	$\frac{1}{2}X_1^2 - \frac{1}{2}X_1 - X_2$
$n = 3$	
(3)	$\frac{1}{6}X_1^3 + \frac{1}{2}X_1^2 + X_1X_2 + \frac{1}{3}X_1 + X_3$
(2, 1)	$\frac{1}{3}X_1^3 - \frac{1}{3}X_1 - X_3$
(1 ³)	$\frac{1}{6}X_1^3 - \frac{1}{2}X_1^2 - X_1X_2 + \frac{1}{3}X_1 + X_3$
$n = 4$	
(4)	$\frac{1}{24}X_1^4 + \frac{1}{4}X_1^3 + \frac{1}{2}X_1^2X_2 + \frac{11}{24}X_1^2 + \frac{1}{2}X_1X_2 + \frac{1}{2}X_2^2 + X_1X_3 + \frac{1}{4}X_1 + \frac{1}{2}X_2 + X_4$
(3, 1)	$\frac{1}{8}X_1^4 + \frac{1}{4}X_1^3 + \frac{1}{2}X_1^2X_2 - \frac{1}{8}X_1^2 - \frac{1}{2}X_1X_2 - \frac{1}{2}X_2^2 - \frac{1}{4}X_1 - \frac{1}{2}X_2 - X_4$
(2, 2)	$\frac{1}{12}X_1^4 - \frac{1}{12}X_1^2 + X_1X_2 + X_2^2 - X_1X_3$
(2, 1 ²)	$\frac{1}{8}X_1^4 - \frac{1}{4}X_1^3 - \frac{1}{2}X_1^2X_2 - \frac{1}{8}X_1^2 - \frac{1}{2}X_1X_2 - \frac{1}{2}X_2^2 + \frac{1}{4}X_1 + \frac{1}{2}X_2 + X_4$
(1 ⁴)	$\frac{1}{24}X_1^4 - \frac{1}{4}X_1^3 - \frac{1}{2}X_1^2X_2 + \frac{11}{24}X_1^2 + \frac{1}{2}X_1X_2 + \frac{1}{2}X_2^2 + X_1X_3 - \frac{1}{4}X_1 - \frac{1}{2}X_2 - X_4$
$n = 5$	
(5)	$\frac{1}{120}X_1^5 + \frac{1}{12}X_1^4 + \frac{1}{6}X_1^3X_2 + \frac{7}{24}X_1^3 + \frac{1}{2}X_1^2X_2 + \frac{1}{2}X_1X_2^2 + \frac{1}{2}X_1^2X_3 + \frac{5}{12}X_1^2 + \frac{5}{6}X_1X_2 + \frac{1}{2}X_1X_3 + X_2X_3 + X_1X_4 + \frac{1}{5}X_1 + X_5$
(4, 1)	$\frac{1}{30}X_1^5 + \frac{1}{6}X_1^4 + \frac{1}{3}X_1^3X_2 + \frac{1}{6}X_1^3 + \frac{1}{2}X_1^2X_3 - \frac{1}{6}X_1^2 - \frac{1}{3}X_1X_2 - \frac{1}{2}X_1X_3 - X_2X_3 - \frac{1}{5}X_1 - X_5$
(3, 2)	$\frac{1}{24}X_1^5 + \frac{1}{12}X_1^4 + \frac{1}{6}X_1^3X_2 - \frac{1}{24}X_1^3 + \frac{1}{2}X_1^2X_2 + \frac{1}{2}X_1X_2^2 - \frac{1}{2}X_1^2X_3 - \frac{1}{12}X_1^2 - \frac{1}{6}X_1X_2 + \frac{1}{2}X_1X_3 + X_2X_3 - X_1X_4$
(3, 1 ²)	$\frac{1}{20}X_1^5 - \frac{1}{4}X_1^3 - X_1^2X_2 - X_1X_2^2 + \frac{1}{5}X_1 + X_5$
(2 ² , 1)	$\frac{1}{24}X_1^5 - \frac{1}{12}X_1^4 - \frac{1}{6}X_1^3X_2 - \frac{1}{24}X_1^3 + \frac{1}{2}X_1^2X_2 + \frac{1}{2}X_1X_2^2 - \frac{1}{2}X_1^2X_3 + \frac{1}{12}X_1^2 + \frac{1}{6}X_1X_2 - \frac{1}{2}X_1X_3 - X_2X_3 + X_1X_4$
(2, 1 ³)	$\frac{1}{30}X_1^5 - \frac{1}{6}X_1^4 - \frac{1}{3}X_1^3X_2 + \frac{1}{6}X_1^3 + \frac{1}{2}X_1^2X_3 + \frac{1}{6}X_1^2 + \frac{1}{3}X_1X_2 + \frac{1}{2}X_1X_3 + X_2X_3 - \frac{1}{5}X_1 - X_5$
(1 ⁵)	$\frac{1}{120}X_1^5 - \frac{1}{12}X_1^4 - \frac{1}{6}X_1^3X_2 + \frac{7}{24}X_1^3 + \frac{1}{2}X_1^2X_2 + \frac{1}{2}X_1X_2^2 + \frac{1}{2}X_1^2X_3 - \frac{5}{12}X_1^2 - \frac{5}{6}X_1X_2 - \frac{1}{2}X_1X_3 - X_2X_3 - X_1X_4 + \frac{1}{5}X_1 + X_5$

TABLE 1. Character polynomials of Weyl modules

Proof. Consider the set of ordered set partitions of $[d]$ with parts of sizes given by λ :

$$X_\lambda = \{(T_1, \dots, T_l) \mid [d] = T_1 \cup \dots \cup T_l \text{ is a set partition, with } |T_i| = \lambda_i\}.$$

Let $K[X_\lambda]$ be the permutation representation associated to the action of S_d on X_λ . This representation is isomorphic to the representation of S_d induced from the trivial

representation of its Young subgroup $S_{\lambda_1} \times \cdots \times S_{\lambda_l}$. Therefore σ_λ is the character of $K[X_\lambda]$, and $\sigma_\lambda(w_\alpha) = |X_\lambda^{w_\alpha}|$, the number of fixed points of a permutation w_α in X_λ . Take $(T_1, \dots, T_l) \in X_\lambda^{w_\alpha}$. Then each T_i is formed by taking a union of cycles of w_α . Suppose that b_{ij} is the number of j cycles of w_α in T_i . Then the array (b_{ij}) satisfies the constraints:

$$\begin{aligned} (10) \quad & b_{i1} + 2b_{i2} + \cdots = \lambda_i \text{ for each } i, \\ (11) \quad & b_{1j} + b_{2j} + \cdots = a_j \text{ for each } j, \end{aligned}$$

where a_j is the number of j -cycles in α . Let $B(\lambda; \alpha)$ denote the set of such arrays. We have:

$$(12) \quad \text{trace}(w_\alpha, K[X_\lambda]) = \sum_{(b_{ij}) \in B(\lambda; \alpha)} \prod_j \binom{a_j}{b_{1j} \ b_{2j} \ \cdots}.$$

On the other hand, by (7),

$$H_\lambda = \sum_{b_{i1} + 2b_{i2} + \cdots = \lambda_i} \prod_{i \geq 1} \prod_{j \geq 1} \binom{X_j}{b_{ij}}.$$

The terms of homogeneous degree d in this product come from the top degree terms in each factor. When $|\alpha| = |\lambda|$, such a term has leading coefficient $\prod_j X_j^{a_j}$ if and only if $(b_{ij}) \in B(\lambda; \alpha)$. Hence the coefficient of $\binom{X}{\alpha}$ is the expression on the right hand side of (12) and the lemma follows. $\square$

Proof of Theorem 2.8. For partitions λ and μ of d , let $K_{\mu\lambda}$ denote the number of semistandard Young tableaux of shape μ and weight λ . Then $K = (K_{\mu\lambda})$ is a unitriangular integer matrix with rows and columns indexed by partitions of d . We have:

$$(13) \quad H_\lambda = \sum_\mu K_{\mu\lambda} S_\mu,$$

$$(14) \quad \sigma_\lambda = \sum_\mu K_{\mu\lambda} \chi_\mu.$$

Let $K_{\mu\lambda}^{-1}$ be the entries of the inverse matrix K^{-1} . Then

$$\begin{aligned} S_\lambda &= \sum_\mu K_{\mu\lambda}^{-1} H_\mu && \text{by (13)} \\ &\equiv \sum_\mu K_{\mu\lambda}^{-1} \sum_\alpha \sigma_\mu(\alpha) \binom{X}{\alpha} && \text{ignoring lower deg. terms (Lemma 2.9)} \\ &= \sum_{\alpha \vdash d} \sum_\mu K_{\mu\lambda}^{-1} \sigma_\mu(\alpha) \binom{X}{\alpha} \\ &= \sum_\alpha \chi_\lambda(\alpha) \binom{X}{\alpha} && \text{by (14),} \end{aligned}$$

thereby completing the proof of Theorem 2.8. $\square$

THEOREM 2.10. *For every partition $\lambda = (\lambda_1, \dots, \lambda_l)$, S_λ is the coefficient of $t_1^{\lambda_1} \cdots t_l^{\lambda_l}$ in*

$$\prod_{i < j} (1 - t_j/t_i) \prod_{r=1}^l \prod_{i \geq 1} (1 - t_r^i)^{-X_i}.$$

Proof. For every vector $\lambda = (\lambda_1, \dots, \lambda_l)$ with non-negative integer coefficients, define:

$$S_\lambda = \det(H_{\lambda_i - i + j}).$$

When λ is a partition this coincides with the character polynomial of the Weyl module W_λ . Then, for every partition λ , S_λ is the coefficient of t^λ in $\sum_{\lambda \geq 0} S_\lambda t^\lambda$. Here $\lambda \geq 0$ indicates that the sum is over all vectors in $\mathbf{Z}_{\geq 0}^l$, and $t^\lambda = t_1^{\lambda_1} \cdots t_l^{\lambda_l}$. Now

$$\begin{aligned} \sum_{\lambda \geq 0} S_\lambda t^\lambda &= \sum_{\lambda \geq 0} \sum_{w \in S_l} \operatorname{sgn}(w) \prod_{r=1}^l H_{\lambda_r - r + w(r)} t_r^{\lambda_r} \\ &= \sum_{w \in S_l} \operatorname{sgn}(w) \prod_{r=1}^l t_r^{r-w(r)} \sum_{\lambda_r \geq 0} H_{\lambda_r - r + w(r)} t_r^{\lambda_r - r + w(r)} \\ &= \sum_{w \in S_l} \operatorname{sgn}(w) \prod_{r=1}^l t_r^{r-w(r)} \sum_{\lambda_r \geq 0} H_{\lambda_r} t_r^{\lambda_r} \\ &= \prod_{i < j} (1 - t_j/t_i) \prod_{r=1}^l \prod_{i \geq 1} (1 - t_r^i)^{-X_i}. \end{aligned}$$

Here we have used the convention that $H_d = 0$ when $d < 0$. □

2.4. DUALITY. Going through the entries of Table 1, the reader may have noticed that the coefficients in the expansion of S_λ agree up to sign with those of $S_{\lambda'}$ for every partition λ . For each vector $\mu = (\mu_1, \dots, \mu_m)$ with non-negative integer entries, let $X^\mu = X_1^{\mu_1} \cdots X_m^{\mu_m}$, and $|\mu| = \mu_1 + \cdots + \mu_m$.

THEOREM 2.11. *For every partition λ , if $S_\lambda = \sum_{\mu} a_\mu^\lambda X^\mu$, then*

$$S_{\lambda'} = \sum_{\mu} (-1)^{|\lambda| - |\mu|} a_\mu^\lambda X^\mu.$$

Proof. Let $\tau_d : P \rightarrow P$ denote the linear involution defined by $X^\mu \mapsto (-1)^{d - |\mu|} X^\mu$. Comparing equations (5) and (6) shows that:

$$\tau_d(H_d) = E_d.$$

It follows that, if $|\mu| = d$, then

$$\tau_d(H_{\mu_1} \cdots H_{\mu_m}) = E_{\mu_1} \cdots E_{\mu_m}.$$

When λ is a partition of d , then every term in the expansion of the Jacobi-Trudi determinants $\det(H_{\lambda_i + j - i})$ is of the form H_μ or E_μ for integer vector μ with $|\mu| = d$. Therefore $\tau_d(\det(H_{\lambda_i + j - i})) = \det(E_{\lambda_i + j - i})$. By the Jacobi-Trudi identities (9),

$$\tau_d(S_\lambda) = \tau_d(\det(H_{\lambda_i + j - i})) = \det(E_{\lambda_i + j - i}) = S_{\lambda'},$$

as claimed. □

3. THE RESTRICTION PROBLEM

3.1. CHARACTER POLYNOMIALS OF SPECHT MODULES. Recall that for partitions λ and μ , we say that $\lambda - \mu$ is a vertical strip if the Young diagram of μ is contained inside the Young diagram of λ , and each row of the Young diagram of λ contains at most one box that is not in the Young diagram of μ [16, Section I.1].

For any partition λ , Macdonald [16, Example I.7.14(b)] gave the character polynomials $q_\lambda \in P$ of (2) as follows:

$$(15) \quad q_\lambda = \sum_{\{\mu \mid \lambda - \mu \text{ is a vertical strip}\}} (-1)^{|\lambda| - |\mu|} \sum_{\alpha \vdash |\mu|} \chi_\mu(\alpha) \binom{X}{\alpha}.$$

It immediately follows that the leading coefficients of q_λ in the binomial basis are the same as those of S_λ (see Theorem 2.8):

THEOREM 3.1. *Let λ be a partition of a positive integer d . For every partition α of d , the coefficient of $\binom{X}{\alpha}$ in the expansion of q_λ in the binomial basis of P is $\chi_\lambda(\alpha)$.*

COROLLARY 3.2. *The sets:*

$$\begin{aligned} \mathbf{S} &= \{S_\lambda \mid \lambda \text{ is an integer partition}\}, \\ \mathbf{q} &= \{q_\lambda \mid \lambda \text{ is an integer partition}\} \end{aligned}$$

are bases of P .

Proof. Regard P as a graded algebra where the degree of X_i is i for each $i \geq 1$. Let P_d denote the homogeneous elements of degree d in P . The degree d homogeneous parts of $\binom{X}{\alpha}$, as α runs over all partitions of d , form a basis of P_d . Theorem 2.8 and the identity (15) imply that the degree d homogeneous parts of S_λ and q_λ also form such a basis as λ runs over all partitions of d , since the character table of S_d forms a non-singular matrix. Therefore $\mathbf{S}$ and $\mathbf{q}$ are bases of P . $\square$

3.2. STABLE RESTRICTION COEFFICIENTS. The coefficients in the expansion of the elements of the basis $\mathbf{S}$ in terms of the basis $\mathbf{q}$:

$$S_\lambda = \sum_{\mu} r_{\lambda\mu} q_\mu$$

are called the stable restriction coefficients. They determine the decomposition of a Weyl module $W_\lambda(K^n)$ into irreducible representations of S_n :

$$\text{Res}_{S_n}^{GL_n(K)} W_\lambda(K^n) = \bigoplus_{\mu} V_{\mu[n]}^{\oplus r_{\lambda\mu}}.$$

The following result, which is now immediate, is an algorithm for computing the stable restriction coefficients:

THEOREM 3.3. *For any partitions λ and μ ,*

$$r_{\lambda\mu} = \langle S_\lambda q_\mu \rangle.$$

The polynomial S_λ can be computed using Theorem 2.7, q_μ using (15). After expanding the product in the binomial basis, the moment can be computed using Theorem 2.3. The matrix of the stable restriction coefficients $r_{\lambda\mu}$, as λ and μ run over

partitions of $0 \leq n \leq 5$ is given by:

$$\left(\begin{array}{cccc|cccc|cccc|cccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 3 & 4 & 2 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 2 & 2 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 5 & 7 & 5 & 2 & 2 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 7 & 5 & 6 & 2 & 3 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 3 & 4 & 1 & 1 & 2 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 3 & 0 & 2 & 2 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 7 & 12 & 9 & 5 & 5 & 3 & 0 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 5 & 14 & 13 & 12 & 6 & 9 & 3 & 2 & 3 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 4 & 10 & 11 & 8 & 6 & 8 & 2 & 1 & 3 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 3 & 4 & 8 & 1 & 7 & 6 & 0 & 2 & 1 & 3 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 3 & 4 & 3 & 2 & 5 & 1 & 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 3 & 0 & 0 & 0 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

The blocks demarcate the partitions of each integer n , and within each block, the partitions of n are enumerated in reverse lexicographic order.

3.3. RELATION TO SYMMETRIC FUNCTIONS. Let Λ denote the ring of symmetric functions (as in [16, Section I.2]). Macdonald [16, Example I.7.13] constructed an isomorphism $\phi : \Lambda \rightarrow P$ taking the Schur function s_λ to the character polynomial q_λ . In this section we study a different isomorphism $\Phi : \Lambda \rightarrow P$, due to Orellana and Zabrocki [19], which takes s_λ to S_λ . Under this isomorphism q_λ is the image of $\tilde{s}_\lambda$ the *Specht symmetric functions* of [2, 19].

Following [19, Proposition 12], define an algebra homomorphism $\Phi : \Lambda \rightarrow P$ by:

$$(16) \quad \Phi : p_k \mapsto \sum_{d|k} dX_d.$$

For each $k > 0$, define

$$\Xi_k = 1, e^{2\pi i/k}, e^{4\pi i/k}, \dots, e^{2(k-1)\pi i/k},$$

and for an integer partition $\mu = (\mu_1, \dots, \mu_m)$,

$$\mathbb{I}_\mu = \mathbb{I}_{\mu_1}, \dots, \mathbb{I}_{\mu_m}.$$

Let R_n denote the space of K -valued class functions on S_n . For every $n \geq 0$ there is a map $\text{ev}_\Lambda^n: \Lambda \rightarrow R_n$ defined by:

$$\mathrm{ev}_\Lambda^n f(w) = f(\Xi_\mu),$$

where μ is the cycle type of w . In other words, the symmetric function is evaluated on $|\mu|$ variables, whose values are given by the list Ξ_μ , the remaining variables being set to 0. With this definition, $\text{ev}_\lambda^n(s_\lambda)$ is the character of $\text{Res}_{S_n}^{GL_n(K)} W_\lambda(K^n)$.

For each $q \in P$ consider the function $\text{ev}_P^n(q) \in R_n$ given by:

$$\mathrm{ev}_P^n(q)(w) = q(X_1(w), X_2(w), \dots).$$

This defines a ring homomorphism $\text{ev}_P^n : P \rightarrow R_n$.

Observe that $\oplus_{n=1}^{\infty} \text{ev}_P^n : P \rightarrow \oplus_{n=1}^{\infty} R_n$ is injective, for if $\text{ev}_P^n(q) \equiv 0$ for all n , then q vanishes whenever $X_1, X_2, \dots$ take non-negative integer values, and hence q must be identically 0.

THEOREM 3.4. *The algebra homomorphism Φ is the unique K -linear map $\Lambda \rightarrow P$ such that the diagram*

$$(17) \quad \begin{array}{ccc} \Lambda & \xrightarrow{\Phi} & P \\ & \searrow \text{ev}_\Lambda^n & \swarrow \text{ev}_P^n \\ & R_n & \end{array}$$

commutes for every $n \geq 1$.

Proof. From the definition of ev_Λ^n ,

$$\text{ev}_\Lambda^n(p_k)(w) = \sum_d X_d(w) \sum_{j=0}^{d-1} (e^{2\pi i/d})^{jk}.$$

Now observe that

$$\sum_{j=0}^{d-1} (e^{2\pi i/d})^{jk} = \begin{cases} d & \text{if } d \mid k, \\ 0 & \text{otherwise.} \end{cases}$$

It follows that

$$\text{ev}_\Lambda^n(p_k)(w) = \sum_{d \mid k} d X_d(w) = \text{ev}_P^n(\Phi(p_k)).$$

Since $\oplus_n \text{ev}_P^n : P \rightarrow \oplus R_n$ is injective, $\Phi(p_k)$ is completely determined by the commutativity of (17). Since the polynomials $\{p_k\}_{k \geq 1}$ generate Λ , Φ is completely determined by its values on p_k . $\square$

LEMMA 3.5. *The homomorphism $\Phi : \Lambda \rightarrow P$ is an isomorphism of rings.*

Proof. The inverse of Φ is obtained using the Möbius inversion formula:

$$X_k \mapsto \frac{1}{d} \sum_{d \mid k} \mu(k/d) p_d,$$

where μ denotes the number-theoretic Möbius function (see, e.g. [13, Section 3.1.1]). $\square$

THEOREM 3.6. *For every partition λ , we have:*

$$(18) \quad \Phi(s_\lambda) = S_\lambda,$$

$$(19) \quad \Phi(\tilde{s}_\lambda) = q_\lambda.$$

Proof. This follows immediately from Theorem 3.4. $\square$

REMARK 3.7. The second identity (19) is [19, Prop. 12].

4. MOMENT GENERATING FUNCTIONS

4.1. THE MAIN GENERATING FUNCTION. In this and the following subsections, we shall frequently use the following elementary identities:

$$(A) \quad \exp(t/i) = \sum_{b \geq 0} \frac{1}{i^b b!} t^b,$$

$$(B) \quad \log \frac{1}{1-t} = \sum_{i=1}^{\infty} t^i / i.$$

We shall use $\alpha = 1^{a_1} 2^{a_2} \dots$, $\beta = 1^{b_1} 2^{b_2} \dots$, $\gamma = 1^{c_1} 2^{c_2} \dots$. Let Par denote the set of all integer partitions. We use the notation $\lambda = (\lambda_1, \dots, \lambda_l)$, $\mu = (\mu_1, \dots, \mu_m)$. Also, $t^\lambda = t_1^{\lambda_1} \dots t_l^{\lambda_l}$ and $u^\mu = u_1^{\mu_1} \dots u_m^{\mu_m}$. We shall interpret $\lambda \geq 0$ as $\lambda_i \geq 0$ for $i = 1, \dots, l$ and $\mu \geq 0$ as $\mu_i \geq 0$ for $i = 1, \dots, m$.

We use the notation $R \sqsubset [l]$ to signify that R is a multiset with elements drawn from $[l]$. We write t^R for the monomial where t_i is raised to the multiplicity of i in R . Similarly, for any $S \subset [m]$, we write $u^S = \prod_{i \in S} u_i$.

THEOREM 4.1. *We have:*

$$\sum_{n \geq 0, \lambda \geq 0, \mu \geq 0} \langle H_\lambda E_\mu \rangle_n t^\lambda u^\mu v^n = \prod_{R \sqsubset [l]} \frac{\prod_{S \subset [m], |S| \text{ odd}} (1 + u^S t^R v)}{\prod_{S \subset [m], |S| \text{ even}} (1 - u^S t^R v)}.$$

Proof. Using (5) and (6), we have:

$$\sum_{\lambda \geq 0, \mu \geq 0} H_\lambda E_\mu t^\lambda u^\mu = \prod_{r=1}^l \prod_{s=1}^m \prod_{i \geq 1} \left(\frac{1 - (-u_s)^i}{1 - t_r^i} \right)^{X_i}.$$

Now proceeding as in the proof of Theorem 2.3,

$$\begin{aligned} \sum_{n \geq 0, \lambda \geq 0, \mu \geq 0} \langle H_\lambda E_\mu \rangle_n t^\lambda u^\mu v^n &= \prod_{i \geq 1} \sum_{b_i \geq 0} \frac{v^{ib_i}}{i^{b_i} b_i!} \prod_{r=1}^l \prod_{s=1}^m \left(\frac{1 - (-u_s)^i}{1 - t_r^i} \right)^{b_i} \\ &\stackrel{(A)}{=} \prod_{i \geq 1} \exp \left(\frac{v^i}{i} \prod_{r=1}^l \prod_{s=1}^m \left[\frac{1 - (-u_s)^i}{1 - t_r^i} \right] \right) \\ &= \exp \left(\sum_{i \geq 1} \sum_{R \sqsubset [l]} \sum_{S \subset [m]} (-1)^{|S|} \frac{(t^R (-1)^{|S|} u^S v)^i}{i} \right) \\ &\stackrel{(B)}{=} \prod_{R \sqsubset [l]} \prod_{S \subset [m]} \left(1 - (-1)^{|S|} t^R u^S v \right)^{(-1)^{|S|+1}}, \end{aligned}$$

which is equivalent to the desired expression. $\square$

COROLLARY 4.2. *We have:*

$$\sum_{\lambda \geq 0, \mu \geq 0} \langle H_\lambda E_\mu \rangle t^\lambda u^\mu = \prod_{R, S} (1 - (-1)^{|S|} u^S t^R)^{(-1)^{|S|+1}},$$

where the product is over $R \sqsubset [l]$, $S \subset [m]$, with at least one of R and S non-empty.

COROLLARY 4.3. *For every partition λ , $\langle W_\lambda \rangle_n$ is the coefficient of $t^\lambda v^n$ in*

$$\prod_{i < j} (1 - t_j/t_i) \prod_{R \sqsubset [l]} (1 - t^R v)^{-1}.$$

Proof. From Theorem 4.1 we get:

$$(20) \quad \sum_{\lambda \geq 0, n \geq 0} \langle H_\lambda \rangle_n t^\lambda v^n = \prod_{R \sqsubset [l]} (1 - t^R v)^{-1}.$$

Using this, the corollary can be deduced from Theorem 2.10 by taking moments. $\square$

4.2. S_n -INVARIANT VECTORS. For a representation V_n of S_n , let $V_n^{S_n}$ denote the subspace of S_n -invariant vectors. If a family $\{V_n\}$ of representations has polynomial character $q \in P$, then

$$\langle q \rangle_n = \dim(V_n^{S_n}) \text{ for all } n \geq 0.$$

Therefore, for any partition λ , $\dim W_\lambda(K^n)^{S_n} = \langle S_\lambda \rangle_n$. In particular, $W_\lambda(K^n)$ has a non-zero S_n -invariant vector if and only if $\langle S_\lambda \rangle_n \neq 0$.

THEOREM 4.4. *For every positive integer n and every partition λ with at most n parts,*

$$\dim W_\lambda(K^n)^{S_n} \leq \dim W_\lambda(K^{n+1})^{S_{n+1}}.$$

Proof. We use Littlewood's plethystic formula [15, Theorem XI] (see also, [17, Theorem 2.6]) for restriction coefficients. This formula asserts that, for every partition λ with at most n parts, and every partition μ of n , the multiplicity of the Specht module V_μ in $\text{Res}_{S_n}^{GL_n(K)} W_\lambda(K^n)$ is given by $(s_\lambda, s_\mu[H])$. Here $(-, -)$ denotes the Hall inner product on symmetric functions, and $s_\mu[H]$ denotes the plethystic substitution of H into s_μ (for definitions, see [6, Section 1]). Taking $\mu = (n)$ in Littlewood's formula gives

$$\dim W_\lambda(K^n)^{S_n} = (s_\lambda, h_n[H]).$$

Recall [6, Eq. 1.8] that

$$h_{n+1}[H] - h_n[H] = h_{n+1}[H - 1],$$

so that

$$\dim W_\lambda(K^{n+1})^{S_{n+1}} - \dim W_\lambda(K^n)^{S_n} = (s_\lambda, h_{n+1}[H - 1]) \geq 0.$$

The inequality above holds because the plethystic substitution of a Schur-positive symmetric function into another is Schur-positive. $\square$

DEFINITION 4.5 (Vector Partitions). *Let $\mathbf{v} \in \mathbf{Z}_{\geq 0}^l$. A vector partition of $\mathbf{v}$ is an unordered collection $\mathbf{v}_1, \dots, \mathbf{v}_n$ of non-zero vectors in $\mathbf{Z}_{\geq 0}^l$ such that*

$$\mathbf{v} = \mathbf{v}_1 + \dots + \mathbf{v}_n.$$

Let $p_n(\mathbf{v})$ (resp. $p_{\leq n}(\mathbf{v})$) denote the number of vector partitions of $\mathbf{v}$ with exactly (resp. at most) n parts.

THEOREM 4.6. *For every partition $\lambda = (\lambda_1, \dots, \lambda_l)$,*

$$\dim W_\lambda(K^n)^{S_n} = \sum_{w \in S_l} \text{sgn}(w) p_{\leq n}(\lambda_1 - 1 + w(1), \dots, \lambda_l - l + w(l)).$$

Proof. The coefficient of $t^\lambda v^n$ in the right hand side of (20) is $p_{\leq n}(\lambda)$. Therefore,

$$(21) \quad \langle H_\lambda \rangle_n = p_{\leq n}(\lambda) \text{ for every } \lambda \in \mathbf{Z}_{\geq 0}^l.$$

By the Jacobi-Trudi identity (9),

$$S_\lambda = \sum_{w \in S_l} \text{sgn}(w) H_{\lambda_1 - 1 + w(1), \dots, \lambda_l - l + w(l)},$$

so by (21),

$$\langle S_\lambda \rangle_n = \sum_{w \in S_l} \text{sgn}(w) p_{\leq n}(\lambda_1 - 1 + w(1), \dots, \lambda_l - l + w(l)),$$

as claimed. $\square$

REMARK 4.7. In general, we do not know of a combinatorial proof of the non-negativity of $\sum_{w \in S_l} \text{sgn}(w) p_{\leq n}(\lambda_1 - 1 + w(1), \dots, \lambda_l - l + w(l))$, which follows from Theorem 4.6. When $l = 2$, this is the main result of Kim and Hahn [10], who refer to it as a conjecture of Landman, Brown and Portier [14].

The problem of characterizing those partitions λ for which $W_\lambda(K^n)$ has a non-zero S_n -invariant vector for large n appears to be quite hard. The following result solves this problem for partition with two rows, two columns, and for hook-partitions.

THEOREM 4.8. *Let λ be a partition.*

- (4.8.1) *If $\lambda = (\lambda_1, \lambda_2)$, then $\langle S_\lambda \rangle > 0$ unless $\lambda = (1, 1)$.*
 (4.8.2) *If $\lambda' = (\lambda_1, \lambda_2)$, then $\langle S_\lambda \rangle > 0$ if and only if $\lambda_1 = \lambda_2$ (in which case $\langle S_\lambda \rangle = 2$) or $\lambda_1 = \lambda_2 + 1$ (in which case $\langle S_\lambda \rangle = 1$).*
 (4.8.3) *If $\lambda = (a + 1, 1^b)$, then $\langle S_\lambda \rangle > 0$ if and only if $a \geq \binom{b+1}{2}$.*

Proof of (4.8.1). By Theorem 4.6 we need to show that, for every $\lambda_1 \geq \lambda_2 \geq 1$,

$$p_{\leq n}(\lambda_1, \lambda_2) > p_{\leq n}(\lambda_1 + 1, \lambda_2 - 1)$$

for sufficiently large n , unless $\lambda_1 = \lambda_2 = 1$. From the main result of Kim and Hahn [10] (the result on the last line of the first page), it follows that

$$p_n(\lambda_1, \lambda_2) \geq p_n(\lambda_1 + 1, \lambda_2 - 1) \text{ for all } n \geq 1.$$

Therefore, it suffices to prove that $p_n(\lambda_1, \lambda_2) > p_n(\lambda_1 + 1, \lambda_2 - 1)$ for at least one value of n . When $k \geq l \geq 1$ are such that at least one of k and l is even, $p_2(k, l) > p_2(k + 1, l - 1)$. When both k and l are odd and $(k, l) \neq (1, 1)$, $p_3(k, l) > p_3(k + 1, l - 1)$. These inequalities will be proved in Lemmas 4.10 and 4.11 below. $\square$

LEMMA 4.9. *For all $k, l \geq 0$,*

$$(22) \quad p_2(k, l) = \begin{cases} \frac{(k+1)(l+1)-1}{2} & \text{if both } k \text{ and } l \text{ are even,} \\ \frac{(k+1)(l+1)}{2} - 1 & \text{otherwise,} \end{cases}$$

$$(23) \quad p_3(k, l) = \frac{1}{6}(A + 3B + 2C),$$

where

$$\begin{aligned} A &= \binom{k+2}{2} \binom{l+2}{2} - 3(k+1)(l+1) + 3, \\ B &= \begin{cases} (k/2 + 1)(l/2 + 1) - 2 & \text{if } k \text{ and } l \text{ are even,} \\ (k+1)(l+2)/4 - 1 & \text{if } k \text{ is odd and } l \text{ is even,} \\ (k+2)(l+1)/4 - 1 & \text{if } k \text{ is even and } l \text{ is odd,} \\ (k+1)(l+1)/4 - 1 & \text{otherwise,} \end{cases} \\ C &= \begin{cases} 1 & \text{if } k \text{ and } l \text{ are divisible by 3,} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof. Consider the set of all ordered triples $((k_1, l_1), (k_2, l_2), (k_3, l_3))$ such that $\sum_i (k_i, l_i) = (k, l)$ and no $(k_i, l_i) = (0, 0)$. The group S_3 acts by permutation on the set of all such triples, and the number of orbits is $p_3(k, l)$. The quantities A , B , and C in Lemma 4.9 are the number of such triples that are fixed by permutations in S_3 of cycle types $(1, 1, 1)$, $(2, 1)$, and (3) , respectively. The formula for $p_3(k, l)$ then follows from Burnside's lemma. The formula for $p_2(k, l)$ is obtained in a similar fashion. $\square$

LEMMA 4.10. *For all integers $k \geq l \geq 1$ such that at least one of k and l is even,*

$$p_2(k, l) > p_2(k + 1, l - 1).$$

Proof. By Lemma 4.9, we also have:

$$p_2(k + 1, l - 1) = \begin{cases} \frac{(k+2)l-1}{2} & \text{if } k \text{ and } l \text{ are odd,} \\ \frac{(k+2)l}{2} - 1 & \text{otherwise.} \end{cases}$$

Thus, if k and l are both even and $k \geq l$, then

$$\begin{aligned} p_2(k, l) - p_2(k+1, l-1) &= \frac{(k+1)(l+1)-1}{2} - \left(\frac{(k+2)l}{2} - 1 \right) \\ &= \frac{k-l}{2} + 1 > 0. \end{aligned}$$

If one of k and l is even and the other is odd, then

$$\begin{aligned} p_2(k, l) - p_2(k+1, l-1) &= \frac{(k+1)(l+1)}{2} - 1 - \left(\frac{(k+2)l}{2} - 1 \right) \\ &= \frac{k-l+1}{2} > 0, \end{aligned}$$

thereby completing the proof of Lemma 4.10. $\square$

LEMMA 4.11. *If $k \geq l \geq 1$, both k and l are odd, and $(k, l) \neq (1, 1)$, then $p_3(k, l) > p_3(k+1, l-1)$.*

Proof. When k and l are odd, Lemma 4.9 gives:

$$12(p_3(k, l) - p_3(k+1, l-1)) = \begin{cases} (kl+2l)(k-l) + k(k-3) + 3l+4 & \text{if } 3 \mid k \text{ and } 3 \mid l, \\ (kl+2l)(k-l) + k(k-3) + 3l & \text{otherwise.} \end{cases}$$

This is clearly positive for all $k \geq l$ such that $k \geq 3$ and $l \geq 1$. $\square$

Proof of (4.8.2). By the second Jacobi–Trudi identity,

$$(24) \quad W_{\lambda'} = E_{\lambda_1} E_{\lambda_2} - E_{\lambda_1+1} E_{\lambda_2-1}.$$

Taking $l = 0$ and $m = 2$ Corollary 4.2 gives:

$$\sum_{\lambda_1, \lambda_2 \geq 0} \langle E_{\lambda_1} E_{\lambda_2} \rangle u_1^{\lambda_1} u_2^{\lambda_2} = \frac{(1+u_1)(1+u_2)}{(1-u_1 u_2)}.$$

Therefore the coefficient of $u_1^{\lambda_1} u_2^{\lambda_2}$ is the number $p'(\lambda_1, \lambda_2)$ of ways of writing (λ_1, λ_2) as a sum of vectors of the form $(1, 1)$, $(1, 0)$ and $(0, 1)$, where the vectors $(0, 1)$ and $(1, 0)$ are used at most once. Clearly

$$p'(\lambda_1, \lambda_2) = \begin{cases} 2 & \text{if } \lambda_1 = \lambda_2 \geq 1, \\ 1 & \text{if } |\lambda_1 - \lambda_2| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

By (24), for any partition $\lambda = (\lambda_1, \lambda_2)$ with two parts,

$$\langle W_{\lambda'} \rangle = p'(\lambda_1, \lambda_2) - p'(\lambda_1+1, \lambda_2-1) = \begin{cases} 2 & \text{if } \lambda_1 \geq \lambda_2 \geq 1, \\ 1 & \text{if } \lambda_1 - \lambda_2 = 1, \\ 0 & \text{otherwise,} \end{cases}$$

as claimed. $\square$

Proof of (4.8.3). Using Pieri’s rule, we have:

$$h_k e_l = s_{(k-1|l)} + s_{(k|l-1)},$$

whence

$$s_{(a|b)} = h_{a+1} e_b - h_{a+2} e_{b-1} + \cdots + (-1)^b h_{a+b+1} e_0.$$

It follows that

$$(25) \quad \langle W_{(a|b)} \rangle = \sum_{j=0}^b (-1)^j \langle H_{a+j+1} E_{b-j} \rangle.$$

Taking $l = m = 1$ in Corollary 4.2 gives:

$$(26) \quad \sum_{a,b \geq 0} \langle H_i E_j \rangle t^i u^j = \frac{\prod_{k=0}^{\infty} (1 + t^k u)}{\prod_{k=1}^{\infty} (1 - t^k)}.$$

The coefficient of $t^i u^j$ in the above expression is the number $\tilde{p}(i, j)$ of ways of writing the vector (i, j) as a sum of vectors of the form $(a, 0)$ where $a > 0$, and $(a, 1)$ where $a \geq 0$, and vectors of the form $(a, 1)$ are used at most once. If $\tilde{p}(i, j) > 0$, then j distinct vectors of the form $(a, 1)$ are used, so that $i \geq \binom{j}{2}$. Therefore

$$(27) \quad \tilde{p}(i, j) = 0 \text{ for all } i < \binom{j}{2}.$$

If $a < \binom{b+1}{2}$, then $a - j < \binom{b+j+1}{2}$ for all $j \geq 0$. By (27) $\langle H_{a-j} E_{b+j+1} \rangle = 0$ for all $j \geq 0$. This allows us to extend the index of summation in the right hand side of (25) without changing the sum:

$$\langle W_{(a|b)} \rangle = \sum_{j=-a-1}^b (-1)^j \langle H_{a+j+1} E_{b-j} \rangle = \sum_{k+l=a+b+1} \langle H_k E_l \rangle = \langle W_{(0|a+b+1)} \rangle.$$

Taking $l = 0$ and $m = 1$ in Corollary 4.2 can be used to show that $\langle E_k \rangle = 0$ for all $k > 1$, so $\langle W_{(0|a+b+1)} \rangle = \langle E_{a+b+2} \rangle = 0$ for all $a, b \geq 0$. Therefore, $\langle W_{(a|b)} \rangle = 0$ for $a < \binom{b+1}{2}$.

Conversely, suppose $a \geq \binom{b+1}{2}$. By Theorem 4.4, it suffices to show that $W_{\lambda}(K^n)$ contains a non-zero S_n -invariant vector for some positive integer n . We shall show that $W_{\lambda}(K^{b+1})$ contains a non-zero S_{b+1} -invariant vector. The hook partition $\lambda = (a+1, 1^b)$ dominates the partition $\mu = (a - \binom{b}{2} + 1, b, b-1, \dots, 2, 1)$, which has $b+1$ distinct parts. Therefore $W_{(a|b)}(K^{a+b+1})$ contains a non-zero vector v with weight μ . For each $w \in S_n$ let $v_w = \rho_{(a|b)}(w)v$. Then v_w lies in the weight space of $w \cdot \mu$. Hence the vectors $\{v_w \mid w \in S_n\}$ are linearly independent, and generate a representation that is isomorphic to the regular representation of S_n . In particular, the trivial representation is contained in $W_{(a|b)}(K^{a+b+1})$. $\square$

5. STRICT POLYNOMIAL FUNCTORS AND FI-MODULES

In this section we may take K to be any field (not necessarily of characteristic zero). Friedlander and Suslin [5] introduced strict polynomial functors to unify homogeneous polynomial representations of $GL_n(K)$ of degree d across all n . Later, Church, Ellenberg and Farb [3] introduced FI-modules to unify representations of S_n across all n . In this section, we lift the restriction functor $\text{Res}_{S_n}^{GL_n(K)}$ to a functor from the category of strict polynomial functors to the category of FI-modules.

5.1. STRICT POLYNOMIAL FUNCTORS. The Schur category (also known as the divided power category, see [22]) $\mathbf{\Gamma}^d$ is the category whose objects are finite dimensional vector spaces over K . Given objects V and W ,

$$\text{Hom}_{\mathbf{\Gamma}^d}(V, W) = \text{Hom}_{S_d}(V^{\otimes d}, W^{\otimes d}).$$

The category of strict polynomial functors is the category $\text{Rep } \mathbf{\Gamma}^d$ whose objects are K -linear covariant functors from $\mathbf{\Gamma}^d$ to the category of K -vector spaces, and morphisms are natural transformations between functors (see [5, Section 2]). When K has characteristic zero, $\text{Rep } \mathbf{\Gamma}^d$ is semisimple, and its simple objects are functors known as *Weyl functors* (see [11]). For each partition λ of d let W_{λ} denote the Weyl functor corresponding to λ .

Let $\text{Rep}^d GL_n(K)$ denote the category of homogeneous polynomial representations of $GL_n(K)$ of degree d . Define a functor $\text{ev}_n : \text{Rep } \Gamma^d \rightarrow \text{Rep}^d GL_n(K)$ as follows: for each strict polynomial functor $F : \Gamma^d \rightarrow \text{Vec}$ define $\text{ev}_n(F) = F(K^n)$. Let $T \in GL_n(K)$ act on $F(K^n)$ by $F(T^{\otimes d})$. This makes $F(K^n)$ a representation of $GL_n(K)$ which turns out to be a homogeneous polynomial representation of degree d . For each $n \geq 1$ and $\lambda \in P(n, d)$, $\text{ev}_n(W_\lambda) = W_\lambda(K^n)$ is the irreducible polynomial representation of $GL_n(K)$ corresponding to λ (consistent with the notation of Section 1).

5.2. FI-MODULES. The category FI is the one that has finite sets as objects, and injective functions as morphisms. The category of FI-modules is the category FI-mod whose objects are covariant functors from FI to the category of K -vector spaces, and morphisms are natural transformations of functors. Let $\text{Rep } S_n$ denote the category of representations of S_n over K . The evaluation functor $\text{ev}_n : \text{FI-mod} \rightarrow \text{Rep } S_n$ is defined by setting $\text{ev}_n(V) = V([n])$, where $[n] = \{1, \dots, n\}$.

For each partition $\lambda = (\lambda_1, \dots, \lambda_l)$ there exists an FI-module $V(\lambda)$ (see [3, Proposition 3.4.1]) such that, for every $n \geq |\lambda| + \lambda_1$, we have:

$$\text{ev}_n(V(\lambda)) = V_{\lambda[n]}.$$

5.3. THE RESTRICTION FUNCTOR. For each object A of FI let $\mathcal{F}[A]$ be the vector space of all functions $A \rightarrow K$. Given an injective function $i : A \rightarrow B$, define $\mathcal{F}(i) : \mathcal{F}[A] \rightarrow \mathcal{F}[B]$ by:

$$\mathcal{F}(i)(f)(b) = \begin{cases} f(a) & \text{if there exists } a \in A \text{ such that } i(a) = b, \\ 0 & \text{otherwise,} \end{cases}$$

for all $f \in \mathcal{F}[A]$. Then $\mathcal{F} : \text{FI} \rightarrow \Gamma^1$ is a functor. For every positive integer d , define $\mathcal{F}^d : \text{FI} \rightarrow \Gamma^d$ by

$$\mathcal{F}^d(A) = \mathcal{F}[A], \quad \mathcal{F}^d(i) = \mathcal{F}(i)^{\otimes d}.$$

The restriction functor

$$\text{Res}^d : \text{Rep } \Gamma^d \rightarrow \text{FI-mod}$$

is defined by:

$$\text{Res}^d F = F \circ \mathcal{F}^d \text{ for every object } F \text{ of } \text{Rep } \Gamma^d.$$

THEOREM 5.1. *The diagram of functors*

$$\begin{array}{ccc} \text{Rep } \Gamma^d & \xrightarrow{\text{Res}^d} & \text{FI-mod} \\ \text{ev}_n \downarrow & & \downarrow \text{ev}_n \\ \text{Rep}^d GL_n(K) & \xrightarrow[\text{Res}_{S_n}^{GL_n(K)}]{} & \text{Rep } S_n \end{array}$$

commutes, in the sense that $\text{Res}_{S_n}^{GL_n(K)} \circ \text{ev}_n$ is naturally isomorphic to $\text{ev}_n \circ \text{Res}^d$.

Proof. Given $F \in \text{Rep } \Gamma^d$, $\text{ev}_n(K)$ is $F(K^n)$. Given $w \in S_n$, let $T_w \in GL_n(K)$ denote the linear map that takes the i th coordinate vector e_i of K^n to $e_{w(i)}$. An element $w \in S_n$ acts on $F(K^n)$ via $F(T_w^{\otimes d})$. On the other hand,

$$\text{ev}_n \circ \text{Res}^d(F) = F \circ \mathcal{F}^d([n]).$$

An element $w \in S_n$ acts on $F(\mathcal{F}^d([n]))$ by $F(\mathcal{F}^d(w)) = F(\mathcal{F}(w)^{\otimes d})$. These two actions of S_n coincide under the isomorphism $K^n \rightarrow \mathcal{F}^d([n])$ given by $e_i \mapsto \delta_i$, where δ_i is the Kronecker delta function on $[n]$ supported at i . Thus we get an isomorphism

$\text{Res}_{S_n}^{GL_n(K)} \circ \text{ev}_n \rightarrow \text{ev}_n \circ \text{Res}^d(F)$ of representations of S_n . The naturality of this isomorphism follows tautologically from unwinding the definitions. $\square$

Church, Ellenberg and Farb introduced the notion of finite generation of FI-modules [3, Definition 1.2]. In characteristic zero, finitely generated FI-modules have eventually polynomial character [3, Theorem 1.5], and in general characteristic, have eventually polynomial dimension [4, Theorem B]. Therefore, the stability of restriction coefficients (discussed in Section 3.3) is also a consequence of the following theorem.

THEOREM 5.2. *For every finitely generated strict polynomial functor F of degree d , the FI-module $\text{Res}^d F$ is finitely generated in degree d .*

Proof. Let Γ^d denote the strict polynomial functor $\Gamma^d(V) = (V^{\otimes d})^{S_d}$, the subspace of S_d -invariant tensors in $V^{\otimes d}$. For $\lambda = (\lambda_1, \dots, \lambda_l)$, let $\Gamma^\lambda(V) = \Gamma^{\lambda_1}(V) \otimes \dots \otimes \Gamma^{\lambda_l}(V)$. The functors Γ^λ , as λ runs over all partitions of d , generate $\text{Rep } \mathbf{FI}^d$ [12, Proposition 2.9]. Therefore, it suffices to prove the theorem for the functors $F = \Gamma^\lambda$. Since Γ^λ is a subobject of $\otimes^d$ in the $\text{Rep } \mathbf{FI}^d$, $\text{Res}^d \Gamma^d$ is a subobject of $\text{Res}^d \otimes^d$ in FI-mod . By the Noetherian property of FI-modules [4, Theorem A], it suffices to show that $\text{Res}^d \otimes^d$ is finitely generated. But $\text{Res}^d \otimes^d(A) = \mathcal{F}^1(A)^{\otimes d} \cong \mathcal{F}^1(A^d)$, which is finitely generated in degree d by [3, Proposition 2.3.6]. $\square$

Acknowledgements. We thank Aprameyo Pal, K. N. Raghavan, Anne Schilling, and S. Viswanath for helpful discussions and encouragement. We thank Brian Hopkins for his answer on mathoverflow <https://mathoverflow.net/q/340231> which helped us prove part (1) of Theorem 4.8. We thank Dipendra Prasad for an argument in the proof of the sufficiency of the condition in part (3) of Theorem 4.8. We thank the referees for their comments and corrections to the original manuscript.

REFERENCES

- [1] Kaan Akin, David A. Buchsbaum, and Jerzy Weyman, *Schur functors and Schur complexes*, Adv. in Math. **44** (1982), no. 3, 207–278.
- [2] Sami H. Assaf and David E. Speyer, *Specht modules decompose as alternating sums of restrictions of Schur modules*, Proc. Amer. Math. Soc. **148** (2020), no. 3, 1015–1029.
- [3] Thomas Church, Jordan S. Ellenberg, and Benson Farb, *FI-modules and stability for representations of symmetric groups*, Duke Math. J. **164** (2015), no. 9, 1833–1910.
- [4] Thomas Church, Jordan S. Ellenberg, Benson Farb, and Rohit Nagpal, *FI-modules over Noetherian rings*, Geom. Topol. **18** (2014), no. 5, 2951–2984.
- [5] Eric M. Friedlander and Andrei Suslin, *Cohomology of finite group schemes over a field*, Invent. Math. **127** (1997), no. 2, 209–270.
- [6] Adriano M. Garsia and Alain Goupil, *Character polynomials, their q -analogs and the Kronecker product*, Electron. J. Combin. **16** (2009), no. 2, Special volume in honor of Anders Björner, Research Paper 19.
- [7] James A. Green, *Polynomial representations of GL_n* , 2nd ed., Lecture Notes in Mathematics, vol. 830, Springer, Berlin Heidelberg, 2007.
- [8] N. Harman, *Representations of monomial matrices and restriction from GL_n to S_n* , <https://arxiv.org/abs/1804.04702>, 2018.
- [9] Gordon James and Adalbert Kerber, *The representation theory of the symmetric group*, Encyclopedia of Mathematics and its Applications, vol. 16, Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [10] Jun Kyo Kim and Sang Geun Hahn, *Partitions of bipartite numbers*, Graphs Combin. **13** (1997), no. 1, 73–78.
- [11] Frank M. Kouwenhoven, *Schur and Weyl functors*, Adv. Math. **90** (1991), no. 1, 77–113.
- [12] Henning Krause, *Koszul, Ringel and Serre duality for strict polynomial functors*, Compos. Math. **149** (2013), no. 6, 996–1018.
- [13] Joseph P. S. Kung, Gian-Carlo Rota, and Catherine H. Yan, *Combinatorics: the Rota way*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2009.

- [14] Bruce M. Landman, Ezra A. Brown, and Frederick J. Portier, *Partitions of bi-partite numbers into at most j parts*, Graphs Combin. **8** (1992), no. 1, 65–73.
- [15] Dudley E. Littlewood, *Products and plethysms of characters with orthogonal, symplectic and symmetric groups*, Canadian J. Math. **10** (1958), 17–32.
- [16] Ian G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Classic Texts in the Physical Sciences, The Clarendon Press, Oxford University Press, New York, 2015.
- [17] Sridhar P. Narayanan, Digjoy Paul, Amritanshu Prasad, and Shraddha Srivastava, *Polynomial induction and the restriction problem*, To appear in the proceedings of *Group Algebras, Representations and Computation*, held at ICTS Bangalore in 2019 in a special issue of *Indian J. Pure Appl. Math.*, <https://arxiv.org/abs/2004.03928>, 2020.
- [18] Rosa Orellana and Mike Zabrocki, *The Hopf structure of symmetric group characters as symmetric functions*, <https://arxiv.org/abs/1901.00378>, 2018.
- [19] ———, *Symmetric group characters as symmetric functions*, <https://arxiv.org/abs/1605.06672v4>, 2018.
- [20] Amritanshu Prasad, *Representation theory: A combinatorial viewpoint*, Cambridge Studies in Advanced Mathematics, vol. 147, Cambridge University Press, Delhi, 2015.
- [21] Thomas Scharf and Jean-Yves Thibon, *A Hopf-algebra approach to inner plethysm*, Adv. Math. **104** (1994), no. 1, 30–58.
- [22] Wilberd van der Kallen, *Lectures on bifunctors and finite generation of rational cohomology algebras*, in Lectures on functor homology, Progr. Math., vol. 311, Birkhäuser/Springer, Cham, 2015, pp. 41–65.

SRIDHAR P. NARAYANAN, The Institute of Mathematical Sciences (HBNI), CIT campus, Taramani,
Chennai 600041, India
E-mail : sridharn@imsc.res.in

DIGJOY PAUL, Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India
E-mail : digjoygpaul@gmail.com

AMRITANSHU PRASAD, The Institute of Mathematical Sciences (HBNI), CIT campus, Taramani,
Chennai 600041, India
E-mail : amri@imsc.res.in
Url : <https://www.imsc.res.in/~amri>

SHRADDHA SRIVASTAVA, Department of Mathematics, Uppsala University, Box 480, SE75106,
Uppsala, Sweden
E-mail : maths.shraddha@gmail.com