


ANNALES MATHÉMATIQUES



BLAISE PASCAL

ANNE-LAURE THIEL

Categorification of the virtual braid groups

Volume 18, n° 2 (2011), p. 231-243.

<http://ambp.cedram.org/item?id=AMBP_2011__18_2_231_0>

© Annales mathématiques Blaise Pascal, 2011, tous droits réservés.

L'accès aux articles de la revue « Annales mathématiques Blaise Pascal » (<http://ambp.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://ambp.cedram.org/legal/>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

*Publication éditée par le laboratoire de mathématiques
de l'université Blaise-Pascal, UMR 6620 du CNRS
Clermont-Ferrand — France*

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

Categorification of the virtual braid groups

ANNE-LAURE THIEL

Abstract

We extend Rouquier's categorification of the braid groups by complexes of Soergel bimodules to the virtual braid groups.

Catégorification des groupes de tresses virtuelles

Résumé

Nous généralisons la catégorification des groupes de tresses par complexes de bimodules de Soergel due à Rouquier aux groupes de tresses virtuelles.

Introduction

Virtual links have been introduced by Kauffman in [5] as a geometric counterpart of Gauss codes. A virtual knot diagram is a generic oriented immersion of circles into the plane, with the usual positive and negative crossings plus a new kind of crossings called virtual. Such crossings appear for instance when one projects a generic link in a thickened surface onto a plane (see [2] or [7]). Many invariants for classical links can be extended to virtual links. Classical oriented links can be represented by closed braids; likewise virtual links can be represented by the closures of virtual braids. There is also an analogue of Markov theorem [3]. Now virtual braids with n strands form a group, denoted \mathcal{VB}_n , which can be described by generators and relations, generalizing the generators and relations of the usual braid group with n strands \mathcal{B}_n . The aim of this note is, using this presentation, to categorify \mathcal{VB}_n in the weak sense of Rouquier. Rouquier actually proves in [10] a stronger version of the result that we want to extend and in particular he states the faithfulness of his categorification. More precisely, to any word ω in the generators of \mathcal{VB}_n we associate a bounded cochain complex $F(\omega)$ of bimodules such that if two words ω and ω' represent the

Keywords: braid group, virtual braid, categorification.

Math. classification: 20F36, 05E10, 05E18, 13D99, 18G35.

same element of \mathcal{VB}_n , then the corresponding cochain complexes $F(\omega)$ and $F(\omega')$ are homotopy equivalent. So we obtain a morphism from the group \mathcal{VB}_n to the group of isomorphism classes of invertible complexes up to homotopy. This morphism, contrary to the case of \mathcal{B}_n studied by Rouquier, is not injective. We describe a part of its kernel in Remarks 3.5.

Note that, using Mazorchuk and Stroppel's study of Arkhipov's twisting functor in [9], one can give a representation theoretic approach to the categorification of virtual braids.

1. Virtual braids

Following [13] and [8], we recall the definition of the virtual braid group \mathcal{VB}_n with n strands.

Definition 1.1. The virtual braid group \mathcal{VB}_n is the group generated by $2(n-1)$ generators $\sigma_1, \dots, \sigma_{n-1}$ and $\zeta_1, \dots, \zeta_{n-1}$ satisfying the braid group relations

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \quad \text{if } |i - j| > 1, \quad (1.1)$$

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad \text{if } 1 \leq i \leq n - 2, \quad (1.2)$$

the permutation group relations

$$\zeta_i \zeta_j = \zeta_j \zeta_i, \quad \text{if } |i - j| > 1, \quad (1.3)$$

$$\zeta_i \zeta_{i+1} \zeta_i = \zeta_{i+1} \zeta_i \zeta_{i+1}, \quad \text{if } 1 \leq i \leq n - 2, \quad (1.4)$$

$$\zeta_i^2 = 1, \quad \text{if } 1 \leq i \leq n - 1, \quad (1.5)$$

and the mixed relations

$$\sigma_i \zeta_j = \zeta_j \sigma_i, \quad \text{if } |i - j| > 1, \quad (1.6)$$

$$\sigma_i \zeta_{i+1} \zeta_i = \zeta_{i+1} \zeta_i \sigma_{i+1}, \quad \text{if } 1 \leq i \leq n - 2. \quad (1.7)$$

The classical braid group \mathcal{B}_n (see [4] for a definition) naturally embeds in \mathcal{VB}_n as a subgroup generated by $\sigma_1, \dots, \sigma_{n-1}$.

The braid group \mathcal{VB}_n can be depicted diagrammatically. To each generator σ_i (resp. σ_i^{-1}) we associate the elementary braid diagram consisting of a single positive (resp. negative) crossing between the i th and $i + 1$ st strand as shown in Figure 1.1 (resp. Figure 1.2). To each generator ζ_i we associate the elementary virtual braid diagram with a single virtual crossing between the i th and $i + 1$ st strand of Figure 1.3.

CATEGORIFICATION OF THE VIRTUAL BRAID GROUPS




FIGURE 1.1. The positive braid σ_i




FIGURE 1.2. The negative braid σ_i^{-1}




FIGURE 1.3. The virtual braid ζ_i

The multiplication law of the group \mathcal{VB}_n consists in concatenating these elementary braids. We use the convention that braids multiply from bottom to top: if D (resp. D') is a virtual braid diagram representing an element β (resp. β') of \mathcal{VB}_n , then the product $\beta\beta'$ is represented by the diagram obtained by putting D' on top of D and gluing the lower endpoints of D' to the upper endpoints of D .

The braid group relations, the permutation group relations and the mixed relations have a diagrammatical interpretation. They correspond to planar isotopies and the generalized Reidemeister moves depicted in Figures 1.4, 1.5 and 1.6.



FIGURE 1.4. Classical Reidemeister II–III moves




FIGURE 1.5. Virtual Reidemeister moves




FIGURE 1.6. The mixed Reidemeister move

2. Rouquier's categorification of the braid groups

In this section we recall how Rouquier [10] categorified the braid group \mathcal{B}_n .

2.1. Soergel bimodules

We first discuss some bimodules introduced by Soergel [11], [12] in his work on representation theory.

Let R be the subalgebra of $\mathbb{Q}[x_1, \dots, x_n]$ defined by

$$R = \mathbb{Q}[x_1 - x_2, x_2 - x_3, \dots, x_{n-1} - x_n] = \mathbb{Q}[x_1 - x_2, x_1 - x_3, \dots, x_1 - x_n].$$

The symmetric group S_n acts on $\mathbb{Q}[x_1, \dots, x_n]$ by $\omega(x_i) = x_{\omega(i)}$ for all $x_i \in R$ and $\omega \in S_n$. This action preserves R . Let R^ω be the subalgebra of elements of R fixed by ω . In particular R^{τ_i} is the subalgebra of R of elements fixed by the transposition $\tau_i = (i, i + 1)$. As an algebra,

$$R^{\tau_i} = \mathbb{Q}[x_1 - x_2, \dots, (x_1 - x_i) + (x_1 - x_{i+1}), \\ (x_1 - x_i)(x_1 - x_{i+1}), x_1 - x_{i+2}, \dots, x_1 - x_n].$$

Let us also consider the R -bimodules $B_\omega = R \otimes_{R^\omega} R$ for any $\omega \in S_n$. The R -bimodules B_{τ_i} will be denoted by B_i for simplicity of notation. We introduce a grading on R , R^{τ_i} and B_i by setting $\deg(x_k) = 2$ for all $k = 1, \dots, n$.

Two R -bimodule morphisms between these objects will be relevant to us, namely $\text{br}_i : B_i \rightarrow R$ and $\text{rb}_i : R\{2\} \rightarrow B_i$ defined by

$$\text{br}_i(1 \otimes 1) = 1 \quad \text{and} \quad \text{rb}_i(1) = (x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1}).$$

The curly brackets indicate a shift of the grading: if $M = \bigoplus_{i \in \mathbb{Z}} M_i$ is a \mathbb{Z} -graded bimodule and p an integer, then the \mathbb{Z} -graded bimodule $M\{p\}$ is defined by $M\{p\}_i = M_{i-p}$ for all $i \in \mathbb{Z}$. The maps br_i and rb_i are degree-preserving morphisms of graded R -bimodules.

2.2. Categorification of the braid groups

Following [6] and [10], to each braid generator $\sigma_i \in \mathcal{B}_n$ we assign the cochain complex $F(\sigma_i)$ of graded R -bimodules

$$F(\sigma_i) : 0 \longrightarrow R\{2\} \xrightarrow{\text{rb}_i} B_i \longrightarrow 0, \tag{2.1}$$

where B_i sits in cohomological degree 0. To σ_i^{-1} we assign the cochain complex $F(\sigma_i^{-1})$ of graded R -bimodules

$$F(\sigma_i^{-1}) : 0 \longrightarrow B_i\{-2\} \xrightarrow{\text{br}_i} R\{-2\} \longrightarrow 0, \tag{2.2}$$

where $B_i\{-2\}$ sits in cohomological degree 0. To the unit element $1 \in \mathcal{B}_n$ we assign the complex of graded R -bimodules

$$F(1) : 0 \longrightarrow R \longrightarrow 0, \tag{2.3}$$

where R sits in cohomological degree 0; the complex $F(1)$ is a unit for the tensor product of complexes so tensoring any complex of graded R -bimodules with $F(1)$ leaves the complex unchanged. Finally to any word $\sigma = \sigma_{i_1}^{\varepsilon_1} \dots \sigma_{i_k}^{\varepsilon_k}$ where $\varepsilon_1, \dots, \varepsilon_k = \pm 1$, we assign the complex of graded R -bimodules $F(\sigma) = F(\sigma_{i_1}^{\varepsilon_1}) \otimes_R \dots \otimes_R F(\sigma_{i_k}^{\varepsilon_k})$.

Rouquier proved the following result, which can be called a categorification of the braid group \mathcal{B}_n .

Theorem 2.1. [10] *If ω and ω' are words representing the same element of \mathcal{B}_n , then $F(\omega)$ and $F(\omega')$ are homotopy equivalent complexes of R -bimodules.*

3. Categorification of the virtual braid groups

Our aim is to extend Rouquier’s categorification to the virtual braid group \mathcal{VB}_n . The cochain complexes associated to the generators σ_i of \mathcal{VB}_n coming from \mathcal{B}_n will be the same as Rouquier’s complexes above. We have to assign complexes to the generators ζ_i of \mathcal{VB}_n corresponding to virtual crossings such that all these complexes satisfy the same relations as the generators of \mathcal{VB}_n up to homotopy equivalence.

3.1. Twisted bimodules

In order to achieve this categorification, we consider the R -bimodule R_ω for each permutation $\omega \in S_n$. As a left R -module, R_ω is equal to R while the right action of $a \in R$ is the multiplication by $\omega(a)$. Note that $R_{\text{id}} = R$ as an R -bimodule. The following lemma is obvious.

Lemma 3.1. *For all $\omega, \omega' \in S_n$ the map $\psi : R_\omega \otimes_R R_{\omega'} \rightarrow R_{\omega\omega'}$ defined for all $a, b \in R$ by $\psi(a \otimes b) = a\omega(b)$ is an isomorphism of R -bimodules.*

The bimodules we will mostly use are the bimodules $R_i = R_{\tau_i}$. The reason why we consider these bimodules is that, by Lemma 3.1, they possess the following interesting property:

$$R_i \otimes_R R_i \cong R$$

for all $i = 1, \dots, n - 1$.

Lemma 3.2. *For all permutations $\omega, \omega' \in S_n$ the R -bimodules $R_\omega \otimes_R B_{\omega'}$ and $B_{\omega\omega'^{-1}} \otimes_R R_\omega$ are isomorphic.*

Proof. First note that there are natural isomorphisms of R -bimodules:

$$R_\omega \otimes_R B_{\omega'} \cong R_\omega \otimes_{R^{\omega'}} R \quad \text{and} \quad B_{\omega\omega'^{-1}} \otimes_R R_\omega \cong R \otimes_{R^{\omega\omega'^{-1}}} R_\omega.$$

Now consider the map $\psi : R_\omega \otimes_{R^{\omega'}} R \rightarrow R \otimes_{R^{\omega\omega'^{-1}}} R_\omega$ defined for all $a, b \in R$ by

$$\psi(a \otimes b) = a \otimes \omega(b).$$

This map is well defined: for any $c \in R^{\omega'}$, we just have to check that $a \otimes cb$ and $a\omega(c) \otimes b$ have the same image under ψ . This is true because $c \in R^{\omega'}$ implies that $\omega(c) \in R^{\omega\omega'^{-1}}$. Moreover the map ψ is obviously a morphism of R -bimodules.

Similarly, the map $\varphi : R \otimes_{R^{\omega\omega'^{-1}}} R_\omega \rightarrow R_\omega \otimes_{R^{\omega'}} R$ defined for all $a, b \in R$ by

$$\varphi(a \otimes b) = a \otimes \omega^{-1}(b)$$

is a well defined morphism of R -bimodules as well.

Finally, ψ and φ are easily seen to be inverse of each other. \square

Now let us assign a complex of graded R -bimodules to each generator of the virtual braid group. To the elements σ_i and σ_i^{-1} we assign the complexes $F(\sigma_i)$ and $F(\sigma_i^{-1})$ defined by (2.1) and (2.2). To the element ζ_i we assign the complex concentrated in degree 0:

$$F(\zeta_i) : 0 \longrightarrow R_i \longrightarrow 0. \tag{3.1}$$

Just as in Section 2.2 we assign to the unit element 1 of \mathcal{VB}_n the complex $F(1)$ of (2.3), and to a virtual braid word we assign the tensor product over R of the complexes associated to the generators involved in the expression of the word.

Remark 3.3. Consider $\omega = \zeta_{i_1} \dots \zeta_{i_k}$ a word in $\{\zeta_1, \dots, \zeta_{n-1}\}$ and let $\tilde{\omega} = \tau_{i_1} \dots \tau_{i_k}$ be the corresponding element of S_n . It follows from Lemma 3.1 that the complex $F(\omega)$ is isomorphic to $0 \longrightarrow R_{\tilde{\omega}} \longrightarrow 0$.

3.2. Categorification of \mathcal{VB}_n

We now state our main result.

Theorem 3.4. *If ω and ω' are words representing the same element of \mathcal{VB}_n , then $F(\omega)$ and $F(\omega')$ are homotopy equivalent complexes of R -bimodules.*

Proof. By definition of \mathcal{VB}_n and in view of Theorem 2.1, it is enough to check that there are homotopy equivalences between the complexes associated to the braid words appearing in both sides of Relations (1.3)-(1.7). Actually we will prove a stronger result: these complexes are isomorphic.

Permutation group relations. Let $\omega = \omega'$ be a permutation group relation. Since the bimodules $R_{\tilde{\omega}}$ and $R_{\tilde{\omega}'}$ are equal, the complexes $F(\omega)$ and $F(\omega')$ are isomorphic in view of Remark 3.3.

Mixed relations. Let us first deal with Relation (1.6). We have to prove that for $|i - j| > 1$ the complexes $F(\zeta_j \sigma_i)$ and $F(\sigma_i \zeta_j)$ are isomorphic. We have

$$F(\zeta_j \sigma_i) : 0 \longrightarrow R_j \otimes_R R\{2\} \xrightarrow{\text{id} \otimes \text{rb}_i} R_j \otimes_R B_i \longrightarrow 0$$

and

$$F(\sigma_i \zeta_j) : 0 \longrightarrow R \otimes_R R_j\{2\} \xrightarrow{\text{rb}_i \otimes \text{id}} B_i \otimes_R R_j \longrightarrow 0.$$

First observe that $F(\zeta_j \sigma_i)$ is naturally isomorphic to the following complex of bimodules

$$0 \longrightarrow R_j\{2\} \xrightarrow{d} R_j \otimes_{R^{\tau_i}} R \longrightarrow 0$$

whose differential d sends each $a \in R_j\{2\}$ to

$$a(\tau_j(x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1})) = a((x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1})).$$

Similarly, $F(\sigma_i \zeta_j)$ is isomorphic to

$$0 \longrightarrow R_j\{2\} \xrightarrow{d'} R \otimes_{R^{\tau_i}} R_j \longrightarrow 0$$

whose differential d' sends each $a \in R_j\{2\}$ to

$$a((x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1})).$$

Since the transpositions τ_i and τ_j commute, the proof of Lemma 3.2 provides us the isomorphism of R -bimodules $\psi : R_j \otimes_{R^{\tau_i}} R \rightarrow R \otimes_{R^{\tau_i}} R_j$. Using

the invariance of $(x_i - x_{i+1})$ under the action of τ_j , we easily check that the following vertical maps and their inverse commute with the differentials.

$$\begin{array}{ccccc} 0 & \longrightarrow & R_j\{2\} & \xrightarrow{d} & R_j \otimes_{R^{\tau_i}} R \longrightarrow 0 \\ & & \downarrow \text{id} & & \downarrow \psi \\ 0 & \longrightarrow & R_j\{2\} & \xrightarrow{d'} & R \otimes_{R^{\tau_i}} R_j \longrightarrow 0 \end{array}$$

Thus the complexes $F(\zeta_j\sigma_i)$ and $F(\sigma_i\zeta_j)$ are isomorphic for all $|i - j| > 1$.

We finally deal with Relation (1.7). We have to show that the complexes $F(\zeta_{i+1}\zeta_i\sigma_{i+1})$ and $F(\sigma_i\zeta_{i+1}\zeta_i)$ are isomorphic for $i = 1, \dots, n - 2$. The complex $F(\zeta_{i+1}\zeta_i\sigma_{i+1})$ is equal to

$$0 \longrightarrow R_{i+1} \otimes_R R_i \otimes_R R\{2\} \xrightarrow{\text{id} \otimes \text{id} \otimes \text{rb}_{i+1}} R_{i+1} \otimes_R R_i \otimes_R B_{i+1} \longrightarrow 0$$

and the complex $F(\sigma_i\zeta_{i+1}\zeta_i)$ is equal to

$$0 \longrightarrow R \otimes_R R_{i+1} \otimes_R R_i\{2\} \xrightarrow{\text{rb}_i \otimes \text{id} \otimes \text{id}} B_i \otimes_R R_{i+1} \otimes_R R_i \longrightarrow 0.$$

There exist a natural isomorphism between $F(\zeta_{i+1}\zeta_i\sigma_{i+1})$ and the complex

$$0 \longrightarrow R_{\tau_{i+1}\tau_i}\{2\} \xrightarrow{d} R_{\tau_{i+1}\tau_i} \otimes_{R^{\tau_{i+1}}} R \longrightarrow 0$$

whose differential d sends each $a \in R_{\tau_{i+1}\tau_i}\{2\}$ to

$$\begin{aligned} a(\tau_{i+1}\tau_i(x_{i+1} - x_{i+2}) \otimes 1 + 1 \otimes (x_{i+1} - x_{i+2})) = \\ a((x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_{i+1} - x_{i+2})). \end{aligned}$$

Similarly, $F(\sigma_i\zeta_{i+1}\zeta_i)$ is isomorphic to

$$0 \longrightarrow R_{\tau_{i+1}\tau_i}\{2\} \xrightarrow{d'} R \otimes_{R^{\tau_i}} R_{\tau_{i+1}\tau_i} \longrightarrow 0$$

whose differential d' sends each $a \in R_{\tau_{i+1}\tau_i}\{2\}$ to

$$a((x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1})).$$

Applying Lemma 3.2 and using the relation $\tau_i\tau_{i+1}\tau_i = \tau_{i+1}\tau_i\tau_{i+1}$ and the involutivity of τ_{i+1} yields the isomorphism

$$\psi : R_{\tau_{i+1}\tau_i} \otimes_{R^{\tau_{i+1}}} R \rightarrow R \otimes_{R^{\tau_i}} R_{\tau_{i+1}\tau_i}$$

given for all $a \in R_{\tau_{i+1}\tau_i}$, $b \in R$ by

$$\psi(a \otimes b) = a \otimes \tau_{i+1}\tau_i(b).$$

Let us complete the proof by checking that the following vertical maps commute with the differentials (and similarly for their inverse).

$$\begin{array}{ccccccc}
 0 & \longrightarrow & R_{\tau_{i+1}\tau_i}\{2\} & \xrightarrow{d} & R_{\tau_{i+1}\tau_i} \otimes_{R^{\tau_{i+1}}} R & \longrightarrow & 0 \\
 & & \downarrow \text{id} & & \downarrow \psi & & \\
 0 & \longrightarrow & R_{\tau_{i+1}\tau_i}\{2\} & \xrightarrow{d'} & R \otimes_{R^{\tau_i}} R_{\tau_{i+1}\tau_i} & \longrightarrow & 0
 \end{array}$$

For any element a in $R_{\tau_{i+1}\tau_i}\{2\}$ we compute both its image under $d' \circ \text{id}$ and $\psi \circ d$. We obtain

$$d' \circ \text{id}(a) = a((x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1}))$$

and

$$\begin{aligned}
 \psi \circ d(a) &= \psi(a(x_i - x_{i+1}) \otimes 1 + a \otimes (x_{i+1} - x_{i+2})) \\
 &= a(x_i - x_{i+1}) \otimes 1 + a \otimes \tau_{i+1}\tau_i(x_{i+1} - x_{i+2}) \\
 &= a((x_i - x_{i+1}) \otimes 1 + 1 \otimes (x_i - x_{i+1}))
 \end{aligned}$$

This shows that the complexes $F(\zeta_{i+1}\zeta_i\sigma_{i+1})$ and $F(\sigma_i\zeta_{i+1}\zeta_i)$ are isomorphic for all $i = 1, \dots, n - 2$. \square

Remarks 3.5.

- Let us call *virtualisation moves* the moves consisting in squeezing a classical crossing between two virtual crossings, as shown in Figure 3.1. We observe that the complexes $F(\zeta_i\sigma_i^\varepsilon\zeta_i)$ and $F(\sigma_i^\varepsilon)$ are isomorphic for all $i = 1, \dots, n - 1$ and $\varepsilon \in \{-1, 1\}$. This is essentially due to the involutivity of τ_i , which implies (*cf.* Lemma 3.1 and Lemma 3.2) that the bimodules $R_i \otimes_R B_i \otimes_R R_i$ and B_i are isomorphic. Thus our categorification of \mathcal{VB}_n does not detect the virtualisation moves.

- Adding the relation $\zeta_i\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\zeta_{i+1}$ to the presentation of \mathcal{VB}_n , one obtains a presentation of the group of welded braids with n strands defined in [1]. Noting that the only morphism between R_ω and $R_{\omega'}$ is the

CATEGORIFICATION OF THE VIRTUAL BRAID GROUPS




FIGURE 3.1. Virtualisation moves

trivial one if $\omega \neq \omega'$, one can check that the complexes $F(\zeta_i \sigma_{i+1} \sigma_i)$

$$\begin{array}{ccc}
 & R_i \otimes_R B_{i+1}\{2\} & \\
 \text{id} \otimes \text{rb}_{i+1} \nearrow & & \searrow \text{id} \otimes \text{id} \otimes \text{rb}_i \\
 R_i \otimes_R R\{4\} & & R_i \otimes_R B_{i+1} \otimes_R B_i \\
 -\text{id} \otimes \text{rb}_i \searrow & & \nearrow \text{id} \otimes \text{rb}_{i+1} \otimes \text{id} \\
 & R_i \otimes_R B_i\{2\} &
 \end{array}$$

and $F(\sigma_{i+1} \sigma_i \zeta_{i+1})$

$$\begin{array}{ccc}
 & B_{i+1} \otimes_R R_{i+1}\{2\} & \\
 \text{rb}_{i+1} \otimes \text{id} \nearrow & & \searrow \text{id} \otimes \text{rb}_i \otimes \text{id} \\
 R \otimes_R R_{i+1}\{4\} & & B_{i+1} \otimes_R B_i \otimes_R R_{i+1} \\
 -\text{rb}_i \otimes \text{id} \searrow & & \nearrow \text{rb}_{i+1} \otimes \text{id} \otimes \text{id} \\
 & B_i \otimes_R R_{i+1}\{2\} &
 \end{array}$$

are not equivalent up to homotopy.

References

- [1] R. FENN, R. RIMÁNYI & C. ROURKE – “The braid-permutation group”, *Topology* **36** (1997), no. 1, p. 123–135.
- [2] N. KAMADA & S. KAMADA – “Abstract link diagrams and virtual knots”, *J. Knot Theory Ramifications* **9** (2000), no. 1, p. 93–106.
- [3] S. KAMADA – “Braid presentation of virtual knots and welded knots”, *Osaka J. Math.* **44** (2007), no. 2, p. 441–458.
- [4] C. KASSEL & V. TURAEV – *Braid groups*, Graduate Texts in Mathematics, vol. 247, Springer, New York, 2008.
- [5] L. H. KAUFFMAN – “Virtual knot theory”, *European J. Combin.* **20** (1999), no. 7, p. 663–690.
- [6] M. KHOVANOV – “Triply-graded link homology and Hochschild homology of Soergel bimodules”, *Internat. J. Math.* **18** (2007), no. 8, p. 869–885.
- [7] G. KUPERBERG – “What is a virtual link?”, *Algebr. Geom. Topol.* **3** (2003), p. 587–591 (electronic).
- [8] V. O. MANTUROV – *Knot theory*, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [9] V. MAZORCHUK & C. STROPPEL – “On functors associated to a simple root”, *J. Algebra* **314** (2007), no. 1, p. 97–128.
- [10] R. ROUQUIER – “Categorification of \mathfrak{sl}_2 and braid groups”, in *Trends in representation theory of algebras and related topics*, Contemp. Math., vol. 406, Amer. Math. Soc., Providence, RI, 2006, p. 137–167.
- [11] W. SOERGEL – “The combinatorics of Harish-Chandra bimodules”, *J. Reine Angew. Math.* **429** (1992), p. 49–74.
- [12] ———, “Gradings on representation categories”, in *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)* (Basel), Birkhäuser, 1995, p. 800–806.
- [13] V. V. VERSHININ – “On homology of virtual braids and Burau representation”, *J. Knot Theory Ramifications* **10** (2001), no. 5, p. 795–812, Knots in Hellas ’98, Vol. 3 (Delphi).

CATEGORIFICATION OF THE VIRTUAL BRAID GROUPS

ANNE-LAURE THIEL
Institut de Recherche Mathématique
Avancée, Université de Strasbourg et
CNRS
7 rue René Descartes, F-67084
Strasbourg Cedex, France
thiel@math.unistra.fr