



*Actions hamiltoniennes :  
invariants et classification*

RENCONTRE ORGANISÉE PAR :  
Michel Brion and Thomas Delzant

2010

Chris Woodward

**Moment maps and geometric invariant theory—Corrected version (October 2011)**

Vol. 1, n° 1 (2010), p. 121-166.

[http://ccirm.cedram.org/item?id=CCIRM\\_2010\\_\\_1\\_1\\_121\\_0](http://ccirm.cedram.org/item?id=CCIRM_2010__1_1_121_0)

Centre international de rencontres mathématiques  
U.M.S. 822 C.N.R.S./S.M.F.  
Luminy (Marseille) FRANCE

**cedram**

*Texte mis en ligne dans le cadre du  
Centre de diffusion des revues académiques de mathématiques  
<http://www.cedram.org/>*

# Moment maps and geometric invariant theory

## Corrected version (October 2011)

Chris WOODWARD

### CONTENTS

1. Introduction	121
2. Actions of Lie groups	122
3. Hamiltonian group actions	123
4. Geometric invariant theory	131
5. The Kempf-Ness theorem	134
6. Schur-Horn convexity and its generalizations	142
7. The stratifications of Hesselink, Kirwan, and Ness	147
8. Moment polytopes	151
9. Multiplicity-free actions	153
10. Localization via sheaf cohomology	157
References	163

### 1. INTRODUCTION

These are expanded notes from a set of lectures given at the school “Actions Hamiltoniennes: leurs invariants et classification” at Luminy in April 2009. The topics center around the theorem of Kempf and Ness [60], which describes the equivalence between the notion of quotient in geometric invariant theory introduced by Mumford in the 1960’s [82], and the notion of symplectic quotient introduced by Meyer [81] and Marsden-Weinstein [79] in the 1970’s. Infinite-dimensional generalizations of this equivalence have played an increasingly important role in geometry, starting with the theorem of Narasimhan and Seshadri [83] connecting unitary structures on a bundle with holomorphic stability, which by historical accident preceded the finite-dimensional theorem.

The proof of the Kempf-Ness theorem depends on the convexity of certain *Kempf-Ness functions* whose minima are zeros of the moment map. The convexity also plays an important role in the relation to geometric quantization discovered by Guillemin and Sternberg [40]: it corresponds to the fact that “invariant quantum states concentrate near zeros of the moment map”. Roughly speaking these notes were written as an exercise in “just how far” one can carry the convexity of the Kempf-Ness function. For example, using convexity I give alternative proofs of some of the results in Kirwan’s book [61] as well as finite-dimensional versions of Harder-Narasimhan and Jordan-Hölder filtrations; the former appears in the algebraic literature under the name of Hesselink one-parameter subgroups [52] but the latter seems to have been undeveloped.

The text is interspersed with applications to existence of invariants in representation theory, such as the problem of determining the existence of invariants in tensor products of irreducible representations, and various techniques for computing moment polytopes. For example, the last section describes Teleman’s improved version of quantization commutes with reduction [104] which

---

Course taught during the meeting “Hamiltonian Actions: invariants et classification” organized by Michel Brion and Thomas Delzant. 6-10 April 2009, C.I.R.M. (Luminy).  
Partially supported by NSF grants DMS060509 and DMS0904358.

also covers the behavior of the higher cohomology groups, and the non-abelian localization formula which computes the difference between the sheaf cohomology of the quotient and the invariant cohomology of the action. Some of the topics not treated are notably: Duistermaat-Heckman theory, symplectic normal forms, localization theorems in equivariant cohomology, and connections to classical invariant theory, to name a few.

The author is grateful for comments and corrections by Michel Brion, Gert Heckman, and Reyer Sjamaar, and apologizes for any omissions of work in what has become a vast literature.

## 2. ACTIONS OF LIE GROUPS

To establish notation we review the basics of Lie group actions.

**2.1. Lie groups.** A *Lie group* is a smooth manifold  $K$  equipped with a group structure so that group multiplication  $K \times K \rightarrow K$  is a smooth map. The *Lie algebra*  $\mathfrak{k}$  is the space of left-invariant vector fields on  $K$ , and may be identified with the tangent space of  $K$  at the identity  $e \in K$ . The *exponential map*  $\exp : \mathfrak{k} \rightarrow K$  is defined by evaluating the time-one flow at the identity.

Suppose that  $K$  is compact and connected. Let  $T \subset K$  be a maximal torus with Lie algebra  $\mathfrak{t}$ . We denote by  $\Lambda := \exp^{-1}(e) \cap \mathfrak{t}$  the integral lattice and by  $\Lambda^\vee \subset \mathfrak{t}^\vee$  its dual, the weight lattice. Any element  $\mu \in \Lambda^\vee$  defines a character  $T \rightarrow U(1), t \mapsto t^\mu$  given for  $\xi \in \mathfrak{t}$  by  $\exp(\xi)^\mu := \exp(2\pi i \mu(\xi))$ . The *Weyl group* of  $T$  is denoted  $W = N(T)/T$ . The Lie algebra  $\mathfrak{k}$  splits under the action of  $T$  into the direct sum of the Lie algebra  $\mathfrak{t}$  and a finite sum of *root spaces*  $\mathfrak{k}_\alpha, \alpha \in \mathcal{R}(\mathfrak{k})$  where  $\mathcal{R}(\mathfrak{k}) \subset \Lambda^\vee / \{\pm 1\}$  is the set of roots and each  $\mathfrak{k}_\alpha$  is identified with a one complex-dimensional representation on which  $T$  acts by  $\exp(\xi)^\alpha := \exp(2\pi i \alpha(\xi))$ . The kernels  $\ker(\alpha)$  of the roots  $\alpha \in \mathcal{R}(\mathfrak{k})$  divide  $\mathfrak{t}$  into a set of (open) *Weyl chambers*; given a generic linear function on  $\mathfrak{t}$  there is a unique open *positive Weyl chamber* on which the function is positive; we denote by  $\mathfrak{t}_+$  its closure and by  $\mathfrak{t}_+^\vee \subset \mathfrak{t}^\vee$  the image of  $\mathfrak{t}_+$  under an identification  $\mathfrak{t} \rightarrow \mathfrak{t}^\vee$  induced by an invariant metric on  $\mathfrak{k}$ . Simple  $K$ -modules are classified by the set of *dominant weights*  $\Lambda_+ := \Lambda \cap \mathfrak{t}_+^\vee$ .

**2.2. Smooth actions and quotients.** Let  $X$  be a smooth manifold. A (*left*) *action* of  $K$  on  $X$  is a smooth map  $K \times X \rightarrow X, (k, x) \mapsto kx$  with the properties that  $k_0(k_1x) = (k_0k_1)x$  and  $ex = x$  for all  $k_0, k_1 \in K$  and  $x \in X$ . A *K-manifold* is a smooth manifold equipped with a smooth  $K$ -action. Let  $X_0, X_1$  be  $K$ -manifolds. A smooth map  $\varphi : X_0 \rightarrow X_1$  is *K-equivariant* if  $\varphi(kx) = k\varphi(x)$  for all  $k \in K, x \in X_0$ .

Both the Lie algebra and its dual are naturally  $K$ -manifolds: The *adjoint* action of an element  $k \in K$  on the Lie algebra  $\mathfrak{k}$  is denoted  $\text{Ad}(k) \in \text{End}(\mathfrak{k})$ . The coadjoint action of  $k$  on the dual  $\mathfrak{k}^\vee$  is  $\text{Ad}^\vee(k) := (\text{Ad}(k^{-1}))^\vee$ . The group  $K$  itself is a  $K$ -manifold in three different ways: the left action, the (inverted) right action, and the *adjoint* action by conjugation  $\text{Ad}(k_0)k_1 := k_0k_1k_0^{-1}$ . The exponential map  $\exp : \mathfrak{k} \rightarrow K$  is equivariant with respect to the adjoint action on  $\mathfrak{k}$  and  $K$ . If  $K$  is compact, then the dual  $\mathfrak{t}^\vee$  of the Lie algebra  $\mathfrak{t}$  of the maximal torus  $T$  admits a canonical embedding in  $\mathfrak{k}^\vee$ , whose image is the  $T$ -fixed point set for the coadjoint action of  $T$  on  $\mathfrak{k}^\vee$ , and so  $\mathfrak{k}^\vee$  admits a canonical projection onto  $\mathfrak{t}^\vee$ .

Let  $X$  be  $K$ -manifold. Let  $\text{Diff}(X)$  denote the infinite-dimensional group of diffeomorphisms of  $X$  and  $\text{Vect}(X)$  the Lie algebra of vector fields on  $X$ . The  $K$ -action induces a canonical group homomorphism

$$K \rightarrow \text{Diff}(X), \quad k \mapsto k_X, \quad k_X(x) = kx$$

and a Lie algebra homomorphism

$$\mathfrak{k} \rightarrow \text{Vect}(X), \quad \xi \mapsto \xi_X, \quad \xi_X(x) = \frac{d}{dt} \Big|_{t=0} \exp(-t\xi)x.$$

The sign here arises because the Lie bracket is defined using left-invariant vector fields which are the generating vector fields for the *right* action of the group on itself, whereas our actions are by default from the *left*. The *orbit* of a point  $x \in X$  is the set  $Kx := \{kx | k \in K\} \subset X$ . The *stabilizer* of a point  $x \in X$  is  $K_x := \{k \in K | kx = x\}$ ; its Lie algebra is the set  $\mathfrak{k}_x := \{\xi \in \mathfrak{k} | \xi_X(x) = 0\}$ . A (co)adjoint orbit is an orbit of the (co)adjoint action of  $K$  on  $\mathfrak{k}$  resp.  $\mathfrak{k}^\vee$ .

Let  $\psi : K_0 \rightarrow K_1$  be a homomorphism of Lie groups and let  $X$  be a  $K_1$ -manifold. The action of  $K_1$  and the homomorphism  $\psi$  induce a  $K_0$ -action on  $X$  by  $k_0x := \psi(k_0)x$ . The orbits of the  $K_0$  action are those of the  $K_1$ -action, while the stabilizers  $(K_0)_x = \psi^{-1}((K_1)_x)$  are inverse images under  $\psi$ .

Let  $X$  be a  $K$ -manifold. A *slice* at  $x$  is a  $K_x$ -invariant submanifold  $V \subset X$  containing  $x$  such that  $KV$  is open in  $X$  and the natural smooth  $K$ -equivariant map  $K \times_{K_x} V \rightarrow KV$  is a diffeomorphism onto its image. It follows from the existence of geodesic flows etc. that actions of *compact* groups have slices. A *quotient* of a  $K$ -space is a pair  $(Y, \pi)$  consisting of a space  $Y$  and a  $K$ -invariant morphism  $\pi : X \rightarrow Y$  such that any other  $K$ -invariant morphism factors through  $\pi$ . The existence of slices implies that any free action of a compact group  $K$  on a manifold  $X$  has a manifold quotient  $X/K$ ; more generally if the action is not free then the quotient exists in the category of Hausdorff topological spaces. (Strictly speaking one should write the quotient on the left, since our actions are by convention left actions. However, I find this rather cumbersome since in English  $X/K$  reads “the quotient of  $X$  by  $K$ ”).

**2.3. Equivariant differential forms.** Recall that a *graded derivation* of a graded algebra  $A$  of degree  $d$  is an operator  $D \in \text{End}(A)_d$  such that  $D(a_0 a_1) = D(a_0) a_1 + (-1)^{|a_0|} a_0 D(a_1)$  for homogeneous elements  $a_0, a_1 \in A$ . The space of graded derivations  $\text{Der}(A)$  (direct sum over degrees) forms a *graded Lie algebra* with bracket given by the *graded commutator*: given graded derivations  $D_0, D_1$  of degrees  $|D_0|, |D_1|$ , define  $\{D_0, D_1\} = D_0 D_1 - (-1)^{|D_0||D_1|} D_1 D_0$ .

Let  $X$  be a smooth manifold of dimension  $n$ . We denote by  $\text{Vect}(X)$  the Lie algebra of smooth vector fields on  $X$ , and by  $\Omega(X) = \bigoplus_{j=0}^n \Omega^j(X)$  the graded algebra of smooth forms on  $X$ . For any  $v \in \text{Vect}(X)$  we have the derivations defined by *contraction*  $\iota_v : \Omega^j(X) \rightarrow \Omega^{j-1}(X)$  and *Lie derivative*  $L_v : \Omega^j(X) \rightarrow \Omega^{j+1}(X)$ . Let  $d$  denote the de Rham operator, the graded derivation  $d : \Omega^j(X) \rightarrow \Omega^{j+1}(X)$  such that  $df(v) = L_v f$ ,  $ddf = 0$  for  $f \in \Omega^0(X), v \in \text{Vect}(X)$ . The operators  $\iota_v, L_v, d$  generate a finite dimensional graded Lie algebra of  $\text{Der}(\Omega(X))$  with graded commutation relations for  $v, w \in \text{Vect}(X)$  given by

$$\begin{array}{c|ccc} \{ , \} & \iota_v & L_v & d \\ \hline \iota_w & 0 & \iota_{[v,w]} & L_w \\ L_w & \iota_{[w,v]} & L_{[w,v]} & 0 \\ d & L_v & 0 & 0 \end{array} .$$

It suffices to check the commutation relations by verifying them on generators  $f \in \Omega^0(X), dg \in \Omega^1(X)$  of  $\Omega(X)$ . We denote by  $Z^j(X)$  the space of *closed forms*  $Z^j(X) = \{\alpha \in \Omega^j(X) | d\alpha = 0\}$  by  $B^j(X) = \{\alpha \in \Omega^j(X) | \exists \beta \in \Omega^{j-1}(X), d\beta = \alpha\}$  the space of *exact forms* and by  $H^j(X)$  the *de Rham cohomology*  $H^j(X) = Z^j(X)/B^j(X)$ .

Suppose that  $X$  admits a smooth action of a Lie group  $K$ . Cartan (see [42]) introduced a space  $\Omega_K(X)$  of  *$K$ -equivariant forms*

$$\Omega_K^j(X) = \bigoplus_{2a+b=j} \text{Hom}^a(\mathfrak{k}, \Omega^b(X))^K, \quad \Omega_K(X) = \bigoplus_{j=0}^{\infty} \Omega_K^j(X)$$

where  $\text{Hom}^a(\cdot)^K$  denotes equivariant polynomial maps of homogeneous degree  $a$ . The *equivariant de Rham operator* is defined by

$$d_K : \Omega_K^j(X) \rightarrow \Omega_K^{j+1}(X), \quad (d_K(\alpha))(\xi) = (d + \iota_{\xi_x})(\alpha(\xi)).$$

Let  $Z_K^j(X)$  resp.  $B_K^j(X)$  denote the equivariant closed resp. exact forms. The *equivariant de Rham cohomology* is

$$H_K^j(X) = Z_K^j(X)/B_K^j(X), \quad H_K(X) = \bigoplus_{j=0}^{\infty} H_K^j(X).$$

If  $K$  action is free,  $H_K(X)$  is isomorphic to the cohomology  $H(X/K)$  of the quotient, see for example [42].

### 3. HAMILTONIAN GROUP ACTIONS

This section contains a quick review of equivariant symplectic geometry. More detailed treatments can be found in Cannas [23], Guillemin-Sternberg [41], Abraham-Marsden [1], or Delzant’s lectures in this volume.

**3.1. Symplectic manifolds.** Let  $X$  be a smooth manifold. A *symplectic form* on  $X$  is a closed non-degenerate two-form  $\omega \in \Omega^2(X)$ . A *symplectic manifold* is a manifold equipped with a symplectic two-form. A *symplectomorphism* of symplectic manifolds  $(X_0, \omega_0), (X_1, \omega_1)$  is a diffeomorphism  $\varphi : X_0 \rightarrow X_1$  with  $\varphi^*\omega_1 = \omega_0$ . The term *symplectic* is the Greek translation of the Latin word *complex*, and was used by Weyl to distinguish the classical groups of linear symplectomorphisms resp. complex linear transformations.

The simplest example of a symplectic manifold is  $\mathbb{R}^{2n}$  equipped with the standard two-form  $\sum_{j=1}^n dq_j \wedge dp_j$ ; Darboux's theorem says that any symplectic manifold is locally symplectomorphic to  $\mathbb{R}^{2n}$  equipped with the standard form. There are simple cohomological restrictions on which manifolds admit symplectic forms: Suppose that  $X$  has dimension  $2n$ . Non-degeneracy of a two-form  $\omega \in \Omega^2(X)$  is equivalent to the non-vanishing of the highest wedge power  $\omega^n \in \Omega^{2n}(X)$ ; if  $X$  is compact and  $\omega$  is symplectic then the cohomology class  $[\omega^n] = [\omega]^n$  must be non-zero, since its integral is non-vanishing, which implies that the classes  $[\omega], [\omega]^2, \dots, [\omega]^{n-1}$  are also non-vanishing. For example this argument rules out the existence of symplectic structures on spheres except for the two-sphere, where any area form gives a symplectic structure.

Symplectic manifolds provide a natural framework for Hamiltonian dynamics as follows. For any symplectic manifold  $(X, \omega)$  let  $\text{Symp}(X, \omega) \subset \text{Diff}(X)$  denote the group of symplectomorphisms and  $\text{Vect}^s(X) \subset \text{Vect}(X)$  the Lie subalgebra of *symplectic vector fields*  $v \in \text{Vect}(X), L_v\omega = 0$ . Any smooth function  $H \in C^\infty(X)$  defines a symplectic vector field  $H^\# \in \text{Vect}^s(X)$  by  $\iota_{H^\#}\omega = dH$ . In local Darboux coordinates,  $H^\#$  is given by

$$H^\# = \sum_{j=1}^n \frac{\partial H}{\partial p_j} \frac{\partial}{\partial q_j} - \frac{\partial H}{\partial q_j} \frac{\partial}{\partial p_j}.$$

The image of  $C^\infty(X)$  in  $\text{Vect}^s(X)$  is the space  $\text{Vect}^h(X)$  of *Hamiltonian* vector fields. Thus a vector field  $v \in \text{Vect}(X)$  is symplectic resp. Hamiltonian iff the associated closed one-form  $\iota_v\omega$  is closed resp. exact. The *Poisson bracket* is the Lie bracket on  $C^\infty(X)$  defined by the formula

$$(3.1) \quad \{H_0, H_1\} = \omega(H_0^\#, H_1^\#).$$

The map  $H \mapsto -H^\#$  extends to an exact sequence of Lie algebras

$$0 \rightarrow H^0(X, \mathbb{R}) \rightarrow C^\infty(X) \rightarrow \text{Vect}^s(X) \rightarrow H^1(X, \mathbb{R}) \rightarrow 0$$

where the Lie bracket on the de Rham cohomology groups  $H^0, H^1(X, \mathbb{R})$  is taken to be trivial. A *Hamiltonian dynamical system* is a pair  $(X, H)$  consisting of a symplectic manifold  $X$  and an *energy function*  $H \in C^\infty(X)$ . Time evolution is given by the flow of  $H^\# \in \text{Vect}(X)$ . If  $K \in C^\infty(X)$  is another function, such as a component of angular momentum, then  $\{K, H\} = -L_{K^\#}H = L_{H^\#}K$ , so  $H$  is invariant under the flow generated by  $K^\#$  iff  $K$  is conserved in time. This equivalence is often called *Noether's theorem*: for every symmetry of a Hamiltonian system there is a conserved quantity.

The cotangent bundle  $T^\vee Q$  of a smooth manifold  $Q$  possesses a canonical symplectic structure: Let  $\pi : T^\vee Q \rightarrow Q, (q, p) \rightarrow q$  be the canonical projection. The *canonical one-form* on  $T^\vee Q$  is

$$\alpha \in \Omega^1(T^\vee Q), \quad \alpha_{(q,p)}(v) = p(D\pi_{q,p}(v)).$$

Local coordinates  $q_1, \dots, q_n$  on  $Q$  induce dual coordinates  $p_1, \dots, p_n$  in which  $\alpha = \sum_{j=1}^n p_j dq_j$ . It follows that the *canonical two-form*  $\omega$  on  $T^\vee Q$  given by  $\omega = -d\alpha$  is symplectic. These forms are canonical in the sense that any diffeomorphism  $Q_0 \rightarrow Q_1$  induces an isomorphism  $T^\vee Q_0 \rightarrow T^\vee Q_1$  preserving the canonical one-forms, which is therefore a symplectomorphism. Physically  $T^\vee Q$  represents the space of states of a classical particle moving on a manifold  $Q$ . However, many Hamiltonian dynamical systems have symplectic manifolds that are not cotangent bundles. For example, the two-sphere is the natural symplectic manifold for the study of the evolution of the angular momentum vector of a rigid body.

**Proposition 3.1.1.** *The following are natural operations on symplectic manifolds:*

- (a) (*Sums*) Let  $(X_0, \omega_0), (X_1, \omega_1)$  be symplectic manifolds. Then the disjoint union  $(X_0 \sqcup X_1, \omega_0 \sqcup \omega_1)$  is a symplectic manifold.

- (b) (*Products*) Let  $(X_j, \omega_j)$  be symplectic manifolds,  $j = 0, 1$ . Then the product  $X_0 \times X_1$  equipped with two-form  $\pi_0^* \omega_0 + \pi_1^* \omega_1$  is a symplectic manifold, where  $\pi_j : X_0 \times X_1 \rightarrow X_j, j = 0, 1$  is the projection onto  $X_j$ .
- (c) (*Duals*) Let  $(X, \omega)$  be a symplectic manifold. Then the dual  $(X, -\omega)$  (or more generally,  $(X, \lambda\omega)$  for any non-zero  $\lambda \in \mathbb{R}$ ) is a symplectic manifold.

Symplectomorphism is a very restrictive notion of morphism, since in particular the symplectic manifolds must be the same dimension. A more flexible notion of morphism in the symplectic category is given by the notion of *Lagrangian correspondence* [108]. (The discussion of correspondences is only used to formulate the universal property for symplectic quotients; readers not interested in this can skip all discussion of correspondences and the symplectic category.) Let  $(X, \omega)$  be a symplectic manifold. A *Lagrangian submanifold* of  $X$  is a submanifold  $i : L \rightarrow X$  with  $i^* \omega = 0$  and  $\dim(L) = \dim(X)/2$ . Let  $(X_j, \omega_j), j = 0, 1$  be symplectic manifolds. A *Lagrangian correspondence* from  $X_0$  to  $X_1$  is a Lagrangian submanifold of  $X_0^- \times X_1$ . Let  $L_{01} \subset X_0^- \times X_1$  and  $L_{12} \subset X_1^- \times X_2$  be Lagrangian correspondences. Let  $\pi_{02}$  denote the projection from  $X_0^- \times X_1 \times X_1^- \times X_2$ . Then

$$L_{01} \circ L_{12} := \pi_{02}(L_{01} \times_{X_1} L_{12})$$

is, if smooth and embedded, a Lagrangian correspondence in  $X_0^- \times X_2$  called the *composition* of  $L_{01}$  and  $L_{12}$ . The graph  $\text{graph}(\psi_{01})$  of any symplectomorphism  $\psi_{01}$  from  $X_0$  to  $X_1$  is automatically a Lagrangian correspondence, and if  $\psi_{01}, \psi_{12}$  are two such symplectomorphisms then  $\text{graph}(\psi_{01} \circ \psi_{12}) = \text{graph}(\psi_{01}) \circ \text{graph}(\psi_{12})$ . With this notion of composition, the pair (symplectic manifolds, Lagrangian correspondences) becomes a partially defined category, with identity given by the diagonal correspondence. The partially defined composition leads to an honest category, obtained by allowing sequences of morphisms and identifying sequences if they are related by geometric composition [107].

Symplectic geometry can be considered a special case of Poisson geometry: A *Poisson bracket* on a manifold  $X$  is a Lie bracket  $\{ , \} : C^\infty(X) \times C^\infty(X) \rightarrow C^\infty(X)$  that is a derivation with respect to multiplication of functions, that is,  $\{f, gh\} = \{f, g\}h + g\{f, h\}$ . A *Poisson manifold* is a manifold equipped with a Poisson bracket. A *morphism* of Poisson manifolds is a smooth map  $\psi : X_0 \rightarrow X_1$  such that  $\{\psi^* f, \psi^* g\} = \psi^* \{f, g\}$ . Given any Poisson bracket on a manifold  $X$ , for each  $H \in C^\infty(X)$  the derivation  $\{H, \}$  is equal to  $L_{H^\#}$  for some vector field  $H^\#$ . The span of the vector fields  $H^\#$  defines a decomposition of  $X$  into *symplectic leaves*, each of which is equipped with a symplectic structure so that (3.1) holds. On the other hand, the notion of symplectic geometry as a special case of Poisson geometry is not particularly compatible with the idea that Lagrangian correspondences should serve as morphisms.

**3.2. Hamiltonian group actions.** Let  $K$  be a Lie group acting smoothly on a manifold  $X$ . The action is *symplectic* if it preserves the symplectic form, that is,  $k_X \in \text{Symp}(X, \omega)$  for all  $k \in K$ , *infinitesimally symplectic* if  $\xi_X \in \text{Vect}^s(X)$  for all  $\xi \in \mathfrak{k}$ , and *weakly Hamiltonian* if  $\xi_X \in \text{Vect}^h(X)$  for all  $\xi \in \mathfrak{k}$ . A *symplectic  $K$ -manifold* is a symplectic manifold equipped with a symplectic action of  $K$ .

Let  $(X, \omega)$  be a symplectic  $K$ -manifold. The action is *Hamiltonian* if the map  $\mathfrak{k} \rightarrow \text{Vect}(X), \xi \mapsto \xi_X$  lifts to an equivariant map of Lie algebras  $\mathfrak{k} \rightarrow C^\infty(X)$ . Such a map is called a *comoment map*. A *moment map* is an equivariant map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ , satisfying

$$(3.2) \quad \iota_{\xi_X} \omega = -d\langle \Phi, \xi \rangle, \quad \forall \xi \in \mathfrak{k}$$

Any comoment map  $\phi : \mathfrak{k} \rightarrow C^\infty(X)$  defines a moment map by  $\langle \Phi(x), \xi \rangle = (\phi(\xi))(x)$ .

*Example 3.2.1.* Let  $K = V$  be a vector space acting on  $X = T^\vee V$  by translation. After identifying  $\mathfrak{k} \rightarrow V$  and so  $\mathfrak{k}^\vee \rightarrow V^\vee$ , a moment map for the action is given by the projection  $X \cong V \times V^\vee \rightarrow V^\vee, (q, p) = p$ , that is, by the ordinary momentum, hence the terminology *moment map*.

The notion of moment map was introduced in independent work of Kirillov, Kostant, and Souriau, in connection with geometric quantization and representation theory. See [16] for a discussion of the history of the moment map and the relationship of the work between these authors. Unfortunately there is no standard sign convention for (3.2); our convention agrees with that of Kirwan [61]. More generally, if  $X$  is a smooth manifold equipped with a closed two-form  $\omega$  and an action of  $K$  leaving  $\omega$  invariant, then we say that  $\Phi$  is a moment map if (3.2) holds.

A *Hamiltonian resp. degenerate Hamiltonian  $K$ -manifold* is a datum  $(X, \omega, \Phi)$  consisting of a symplectic  $K$ -manifold  $(X, \omega)$  resp. smooth  $K$ -manifold  $X$  equipped with an invariant closed two-form  $\omega$ , and a moment map  $\Phi$  for the action. Let  $(X_0, \omega_0, \Phi_0)$  and  $(X_1, \omega_1, \Phi_1)$  be Hamiltonian  $K$ -manifolds. An *isomorphism* of Hamiltonian  $K$ -manifolds is a  $K$ -equivariant symplectomorphism  $\varphi : (X_0, \omega_0) \rightarrow (X_1, \omega_1)$  such that  $\varphi^*\Phi_1 = \Phi_0$ .

*Example 3.2.2.* Archimedes' computation of the area of the two-sphere is essentially a moment map calculation. Let  $S^2 = \{x^2 + y^2 + z^2 = 1\}$  be the unit sphere in  $\mathbb{R}^3$ . Let  $v = x\frac{\partial}{\partial x} + y\frac{\partial}{\partial y} + z\frac{\partial}{\partial z} \in \text{Vect}(\mathbb{R}^3)$ . The two-form  $\omega = \iota_v(dx \wedge dy \wedge dz) = xdy \wedge dz - ydx \wedge dz + zdx \wedge dy$  restricts to a symplectic form on  $S^2$ , invariant under rotation on  $\mathbb{R}^3$ . A moment map for the action of  $S^1$  on  $S^2$  by rotation clockwise around the  $z$ -axis is given by  $(x, y, z) \mapsto z$ , under the identification of the Lie algebra of  $S^1$  and its dual with  $\mathbb{R}$ . Indeed, the generating vector field for  $\xi = 1$  is  $\xi_X = -x\frac{\partial}{\partial y} + y\frac{\partial}{\partial x}$ . A computation shows that  $\iota_{\xi_X}\omega = -dz$ .

To relate this to Archimedes' area formula, note that if  $r, \theta, z$  are cylindrical coordinates on  $\mathbb{R}^3$ , then  $\iota_{\frac{\partial}{\partial \theta}}\omega = dz$  and so  $\omega = dz \wedge d\theta$ . Thus the area of the unit two-sphere between any two values  $z_1, z_2 \in (-1, 1)$  of  $z$  is the same as the area of the cylinder  $S^1 \times [-1, 1]$  between those two values,  $2\pi(z_2 - z_1)$ . In particular (and this is the result reported by Cicero to be inscribed on Archimedes' tombstone) the area of the unit two-sphere  $S^2$  is equal to the area of the cylinder  $S^1 \times [-1, 1]$ , namely  $4\pi$ . We can deduce from the moment map for the circle action the moment map for the full




Figure 3.1:  $S^1 \times [-1, 1]$  has the same area as  $S^2$

rotation group  $SO(3)$  as follows. We identify  $\mathfrak{so}(3) \rightarrow \mathbb{R}^3$  so that the infinitesimal rotation around the  $j$ -th basis vector  $e_j$  maps to  $e_j$ , and  $\mathfrak{so}(3)^\vee \rightarrow \mathbb{R}^3$  using the standard metric on  $\mathbb{R}^3$ . The action of  $SO(3)$  on  $S^2$  has moment map the inclusion  $S^2 \rightarrow \mathbb{R}^3$ . Indeed, by symmetry, moment maps for the rotation around the other two axes are given by  $(x, y, z) \mapsto x$  resp.  $y$ . Hence the inclusion satisfies the equation (3.2). In addition  $\Phi$  is equivariant and so defines a moment map. This ends the example.

The following are natural operations on Hamiltonian  $K$ -manifolds:

- Proposition 3.2.3.**
- (a) (*Sums*) Let  $(X_0, \omega_0, \Phi_0), (X_1, \omega_1, \Phi_1)$  be Hamiltonian  $K$ -manifolds. Then the disjoint union  $X_0 \sqcup X_1$  is a Hamiltonian  $K$ -manifold, equipped with moment map  $\Phi_0 \sqcup \Phi_1$ .
  - (b) (*Exterior Products*) Let  $(X_j, \omega_j, \Phi_j)$  be Hamiltonian  $K_j$ -manifolds,  $j = 0, 1$ . Then the product  $X_0 \times X_1$  is a Hamiltonian  $K_0 \times K_1$ -manifold, equipped with moment map  $\pi_0^*\Phi_0 \times \pi_1^*\Phi_1$ , where  $\pi_j : X_0 \times X_1 \rightarrow X_j, j = 0, 1$  is the projection onto  $X_j$ .
  - (c) (*Duals*) Let  $(X, \omega, \Phi)$  be a Hamiltonian  $K$ -manifold. Then the dual  $(X, -\omega, -\Phi)$  (or more generally, any rescaling by a non-zero constant) is a Hamiltonian  $K$ -manifold.
  - (d) (*Pull-backs*) Let  $\varphi : K_0 \rightarrow K_1$  be a homomorphism of Lie groups, and  $(X, \omega, \Phi)$  a Hamiltonian  $K_1$ -manifold. The Lie algebra homomorphism  $D\varphi : \mathfrak{k}_0 \rightarrow \mathfrak{k}_1$  induces a dual map  $D\varphi^\vee : \mathfrak{k}_1^\vee \rightarrow \mathfrak{k}_0^\vee$ . The action of  $K_0$  induced by  $\phi$  has moment map  $D\varphi^\vee \circ \Phi$ .
  - (e) (*Interior products*) Let  $(X_j, \omega_j, \Phi_j)$  be Hamiltonian  $K$ -manifolds,  $j = 0, 1$ . Then the product  $X_0 \times X_1$  is a Hamiltonian  $K$ -manifold, equipped with moment map  $\pi_0^*\Phi_0 + \pi_1^*\Phi_1$ . This is a combination of the previous two items, using the diagonal embedding  $\mathfrak{k} \rightarrow \mathfrak{k} \times \mathfrak{k}$  whose adjoint is  $\mathfrak{k}^\vee \times \mathfrak{k}^\vee \rightarrow \mathfrak{k}^\vee, (\xi_0, \xi_1) \mapsto \xi_0 + \xi_1$ .

More generally one can speak of Hamiltonian actions on Poisson manifolds. The dual  $\mathfrak{k}^\vee$  of the Lie algebra  $\mathfrak{k}$  has a canonical *Lie-Poisson* bracket,  $C^\infty(\mathfrak{k}^\vee) \times C^\infty(\mathfrak{k}^\vee) \rightarrow C^\infty(\mathfrak{k}^\vee)$  with the property that  $\{\xi, \eta\} = [\xi, \eta]$  for  $\xi, \eta \in \mathfrak{k}$ . A *Poisson moment map* for a  $K$ -action on a Poisson manifold  $X$  is

a Poisson map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ . A *Hamiltonian-Poisson  $K$ -manifold* is a Poisson  $K$ -manifold equipped with a Poisson moment map.

**Proposition 3.2.4.** *Any Hamiltonian  $K$ -manifold  $(X, \omega, \Phi)$  is a Hamiltonian-Poisson  $K$ -manifold.*

*Proof.* For  $\lambda, \xi \in \mathfrak{k}$  we have  $\Phi^*\{\lambda, \xi\} = \Phi^*[\lambda, \xi] = L_{\lambda_X} \Phi^* \xi = \{\Phi^* \lambda, \Phi^* \xi\}$ . The case of non-linear functions is similar.  $\square$

Conversely, any Poisson moment map induces an ordinary moment map on its symplectic leaves. In particular the coadjoint action is Poisson-Hamiltonian with moment map the identity, and the symplectic leaves are the coadjoint orbits. Thus as observed by Kirillov, Kostant, and Souriau,

**Proposition 3.2.5.** *Any coadjoint orbit  $K\lambda, \lambda \in \mathfrak{k}^\vee$  of  $K$  has the canonical structure of a Hamiltonian  $K$ -manifold with moment map given by the inclusion  $K\lambda \rightarrow \mathfrak{k}^\vee$ .*

*Example 3.2.6.* Identify  $\mathbb{R}^3 \cong \mathfrak{so}(3) \cong \mathfrak{so}(3)^\vee$ . The Proposition gives Hamiltonian  $SO(3)$ -structures on the orbits of  $SO(3)$  on  $\mathbb{R}^3$ , which are either spheres (for non-zero radii  $\lambda$ ) or a point (if  $\lambda = 0$ .) This reproduces the moment map in Example 3.2.2.

For any *transitive* Hamiltonian action, the moment map is a local diffeomorphism and so gives a covering of the coadjoint orbit that is its image, see Kostant [70].

The Darboux theorem has various equivariant generalizations that we will not discuss here; we only mention that as a consequence:

**Proposition 3.2.7.** *(see [61]) Let  $X$  be a Hamiltonian  $K$ -manifold,  $K$  compact. For any  $\xi \in \mathfrak{k}$ , the function  $\langle \Phi, \xi \rangle$  is a Morse function with even index.*

In the remainder of the section we explain two other ways in which moment maps can be naturally interpreted. The first is closely related to the notion of *equivariantly closed differential form* introduced in Section 2.3, see Atiyah and Bott [9]:

**Proposition 3.2.8.** *Let  $(X, \omega)$  be a symplectic  $K$ -manifold. There exists a one-to-one correspondence between moment maps for the action of  $K$ , and equivariantly closed extensions of  $\omega \in \Omega^2(X)$  to  $\Omega_K^2(X)$ .*

*Proof.* Since  $\Omega_K^2(X) \cong \Omega^2(X)^K \oplus \text{Hom}(\mathfrak{k}, \Omega^0(X))^K$  any extension in  $\Omega_K^2(X)$  is equal to  $\omega + \Phi$  for some  $\Phi \in \text{Map}_K(X, \mathfrak{k}^\vee) \cong \text{Hom}(\mathfrak{k}, \Omega^0(X))^K$ . The extension is equivariantly closed iff  $0 = d_K(\omega + \Phi) = (d\omega, \iota_{\xi_X} \omega + d\langle \Phi, \xi \rangle)$ . Since  $\omega$  is by assumption closed,  $d_K(\omega + \Phi) = 0$  iff  $\Phi$  is a moment map.  $\square$

The second interpretation of a moment map depends on the notion of *linearization* of an action, as we now explain. Suppose that  $L \rightarrow X$  is a Hermitian line bundle with unit circle bundle  $L_1$  with generating vector fields  $\xi_L \in \text{Vect}(L_1), \xi \in \mathbb{R}$ . The circle group  $U(1)$  acts on  $L_1$  by scalar multiplication. Let  $\alpha \in \Omega^1(L_1)^{U(1)}, \alpha(\xi_L) = \xi$  be a connection one-form with curvature  $(2\pi/i)\omega \in \Omega^2(X)$ . (That is, to fix conventions,  $d\alpha = \pi^* \omega$  where  $\pi : L_1 \rightarrow X$  is the projection.) The group  $\text{Aut}(L_1, \alpha)$  of unitary automorphisms of  $L$  preserving  $\alpha$  naturally maps to the symplectomorphism group  $\text{Symp}(X, \omega)$  of  $X$ , defining an exact sequence  $1 \rightarrow U(1) \rightarrow \text{Aut}(L_1, \alpha) \rightarrow \text{Symp}(X, \omega)$ . A *linearization* of the action of  $K$  on  $X$  is a lift  $K \rightarrow \text{Aut}(L_1, \alpha)$ . An *infinitesimal linearization* is a lift  $\mathfrak{k} \rightarrow \text{Vect}(L_1)^{U(1)}$ .

**Proposition 3.2.9.** *Let  $X$  be a  $K$ -manifold,  $\omega \in \Omega^2(X)^K$  a closed invariant two-form, and  $\pi : L \rightarrow X$  a Hermitian line-bundle with connection one-form  $\alpha \in \Omega^1(L_1)^{K \times U(1)}$  whose curvature is equal to  $(2\pi/i)\omega$ . The set of moment maps  $\Phi$  for the  $K$ -action is in one-to-one correspondence with the set of infinitesimal linearizations of the action of  $K$ .*

*Proof.* Let  $\pi_1 : L_1 \rightarrow X$  denote the projection. Given a lift  $\mathfrak{k} \rightarrow \text{Vect}(L_1)^{U(1)}$ , define a moment map  $\Phi : X \rightarrow \mathfrak{k}^\vee$  by  $\langle \Phi(x), \xi \rangle = (\alpha(\xi_L))(l)$ , for any  $l \in \pi^{-1}(x)$ , independent of the choice of  $l$ . Then

$$\begin{aligned} \pi_1^* d\langle \Phi, \xi \rangle &= d(\alpha(\xi_L)) = d\iota_{\xi_L} \alpha(l) = (L_{\xi_L} - \iota_{\xi_L} d)\alpha \\ &= L_{\xi_L} \alpha - \iota_{\xi_L} \pi_1^* \omega = -\pi_1^* \iota_{\xi_X} \omega. \end{aligned}$$

Since  $\alpha$  is invariant,  $\Phi$  is equivariant, and so defines a moment map. Conversely, given a moment map define  $\xi_L \in \text{Vect}(L_1)^{S^1}$  by  $\langle \Phi(x), \xi \rangle = (\alpha(\xi_L))(l)$ . Then the same computation shows that



$L_{\xi_L}\alpha = 0$ . To see that  $\xi \mapsto \xi_L$  defines a lift of  $\mathfrak{k} \rightarrow \text{Vect}^s(X, \omega)$  to  $\text{Vect}(L_1)^{U(1)}$ , note that given  $\xi, \eta \in \mathfrak{k}$ , the vectors  $[\xi, \eta]_L$  and  $[\xi_L, \eta_L]$  agree up to a vertical vector field. To see that they are equal, note  $\alpha([\xi_L, \eta_L]) = [L_{\xi_L}, L_{\eta_L}]\alpha = \pi^*L_{\xi_L}\langle \Phi, \eta \rangle = \pi^*\langle \Phi, [\xi, \eta] \rangle = \alpha([\xi, \eta]_L)$ .  $\square$

The following is immediate from the definitions:

**Proposition 3.2.10.** *Suppose that  $\Phi$  is the moment map induced by a lift of the action to a Hermitian line bundle with connection  $L$ . Then  $\exp(\xi), \xi \in \mathfrak{k}_x$  acts on the fiber  $L_x$  via  $l \mapsto \exp(i\langle \Phi(x), \xi \rangle)l$ .*

In other words, the value of the moment map at a fixed point determines the action of the identity component of the group on the fiber over that point.

The notion of Lagrangian correspondence generalizes to Hamiltonian actions as follows. (again, readers not interested in universal properties of quotients may skip this discussion):

**Definition 3.2.11.** Let  $X$  be a Hamiltonian  $K$ -manifold with moment map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ . A  $K$ -Lagrangian submanifold is a  $K$ -invariant Lagrangian submanifold on which  $\Phi$  vanishes. Let  $(X_j, \omega_j, \Phi_j)$  be Hamiltonian  $K$ -manifolds for  $j = 0, 1$ . A  $K$ -Lagrangian correspondence is a  $K$ -Lagrangian submanifold of  $X_0^- \times X_1$ .

Allowing sequences of  $K$ -Lagrangian correspondences and identifying sequences related by a geometric composition gives an honest category as in non-equivariant case.

**3.3. Symplectic quotients.** Naturally one would like a notion of quotient of a Hamiltonian  $K$ -manifold, which should be an object in the symplectic category and satisfy a universal property for morphisms in the equivariant symplectic category. It is easy to see that the most naive definition, of the actual quotient, is unsatisfactory for several reasons. For example, even if the action is free, then the quotient will not necessarily have even dimension, and so may not admit a symplectic structure. Also the action will not in general be free, and so the quotient will not even have the structure of a manifold.

The construction of Meyer [81] and Marsden-Weinstein [79] is free of these problems, at least under suitable hypotheses: Let  $(X, \omega, \Phi)$  be a Hamiltonian  $K$ -manifold with moment map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ . Define the *symplectic quotient*

$$X//K := \Phi^{-1}(0)/K.$$

**Theorem 3.3.1** (Meyer [81], Marsden-Weinstein [79]). *Let  $X$  be a Hamiltonian  $K$ -manifold. If  $K$  acts freely and properly on  $\Phi^{-1}(0)$ , then  $X//K$  has the structure of a smooth manifold of dimension  $\dim(X) - 2\dim(K)$  with a unique symplectic form  $\omega_0$  satisfying  $i^*\omega = p^*\omega_0$ , where  $i : \Phi^{-1}(0) \rightarrow X$  and  $p : \Phi^{-1}(0) \rightarrow X//K$  are the inclusion and projection respectively.*

The double slash in the notation  $X//K$  is meant to reflect that the dimension drops by  $2\dim(K)$ , in contrast to the ordinary quotient  $X/K$  for which dimension drops by  $\dim(K)$ , if the action is free. The proof depends on the following. Let  $\text{ann}(\mathfrak{k}_x) \subset \mathfrak{k}^\vee$  be the annihilator of  $\mathfrak{k}_x$ .

**Lemma 3.3.2.** *Let  $X$  be a Hamiltonian  $K$ -manifold. For any  $x \in X$ ,*

- (a)  $\text{Im } D_x\Phi = \text{ann}(\mathfrak{k}_x)$ .
- (b)  $\text{Ker } D_x\Phi = \{\xi_X(x), \xi \in \mathfrak{k}\}^{\omega_x}$ .

*Proof.* (a) We have  $\langle D_x\Phi(v), \eta \rangle = \omega_x(v, \eta_X(x))$  for  $v \in T_xX$  which vanishes for all  $v \in T_xX$  iff  $\eta_X(x) = 0$ . (b) The same identity shows  $\omega_x(\xi_X(x), v) = 0$  for  $v \in \text{Ker } D_x\Phi$ , so the left-hand-side of (b) is contained in the right. Equality now follows by a dimension count, using (a).  $\square$

*Proof of Theorem.* By part (a) of the Lemma, the pull-back  $i^*w$  vanishes on the orbits of  $K$  and is  $K$ -invariant, and so descends to a form  $\omega_0$  on  $X//K$ . Part (b) shows that  $\omega_0$  is non-degenerate. Since  $p^*d\omega_0 = di^*\omega = i^*d\omega = 0$ ,  $\omega_0$  is closed, hence symplectic.  $\square$

The following is a fundamental example:

*Example 3.3.3.* (Products of spheres) Let  $\lambda_1, \dots, \lambda_n$  be positive real numbers and  $X = S_{\lambda_1}^2 \times \dots \times S_{\lambda_n}^2$ , where  $S_\lambda^2$  denotes the unit two-sphere with invariant area form re-scaled by  $\lambda$ . The group  $K = SO(3)$  acts diagonally on  $X = (S^2)^n$  with moment map

$$\Phi : X \rightarrow \mathfrak{k}^\vee \cong \mathbb{R}^3, \quad (x_1, \dots, x_n) \mapsto x_1 + \dots + x_n$$

by 3.2.2 and 3.2.3 (e). The symplectic quotient is the *moduli space of closed  $n$ -gons with lengths  $\lambda_1, \dots, \lambda_n$*

$$X//SO(3) = \{(x_1, \dots, x_n) \in (\mathbb{R}^3)^n \mid \|x_j\| = \lambda_j, x_1 + \dots + x_n = 0\}/SO(3).$$

Its topology depends on the choice of  $\lambda_1, \dots, \lambda_n$ , see for example Hausmann-Knutson [47]. In general there are a finite number of “chambers” in which the topology of  $X//SO(3)$  is constant. The chambers in which  $X//SO(3)$  is non-empty are described by the following:

*Proposition 3.3.4.*  $X//SO(3) \neq \emptyset$  iff  $\lambda_j \leq \sum_{i \neq j} \lambda_i$  for all  $j = 1, \dots, n$ .

*Proof.* For  $n = 3$ , these are the triangle inequalities. For  $n > 3$ , we assume without loss of generality that  $\lambda_1 \geq \dots \geq \lambda_n$ . Then the inequalities above are equivalent to the single inequality  $\lambda_1 \leq \lambda_2 + \dots + \lambda_n$ . One checks that there exists  $j$  so that  $|\lambda_2 + \dots + \lambda_j - \lambda_{j+1} - \dots - \lambda_n| < \lambda_1$ . The general case follows from that for  $n = 3$ , which implies that there exists a triangle with side lengths  $\lambda_1, \lambda_2 + \dots + \lambda_j, \lambda_{j+1} + \dots + \lambda_n$ .  $\square$

This ends the example.

We end this section with two remarks on the definition of symplectic quotient. First, the symplectic quotient of a Hamiltonian action can be viewed as a symplectic leaf of the quotient of the corresponding Hamiltonian-Poisson action in the following sense. Suppose that  $X$  is a Hamiltonian-Poisson  $K$ -manifold such that  $K$  acts freely. The restriction of the Poisson bracket to  $C^\infty(X)^K$  defines a canonical Poisson structure on  $X/K$ . Then  $X//K$  is a symplectic leaf on the smooth locus in  $X/K$  [6]; the other leaves are symplectic quotients at other coadjoint orbits, discussed in Section 8.

Second, the symplectic quotient satisfies the following universal property for quotients. Suppose that  $(X, \omega, \Phi)$  is a Hamiltonian  $K$ -manifold and  $K$  acts freely on  $\Phi^{-1}(0)$ . We denote by  $L_\Phi \subset X^- \times (X//K)$  the image of  $\Phi^{-1}(0)$  under  $i \times p$ . Then  $L_\Phi$  is a  $K$ -Lagrangian correspondence.

**Theorem 3.3.5.** *Suppose that  $X$  is a Hamiltonian  $K$ -manifold. If  $Y$  is a symplectic manifold with trivial  $K$ -action, then any  $K$ -Lagrangian correspondence from  $X$  to  $Y$  factors through  $L_\Phi$ .*

*Proof.* Suppose for simplicity that the morphism consists of a single correspondence  $L \subset X^- \times Y$ . By definition of  $K$ -Lagrangian correspondence,  $L \subset \Phi^{-1}(0) \times Y$ . Since  $K$  acts freely on  $\Phi^{-1}(0)$ ,  $L/K$  is a submanifold of  $X^-//K \times Y$  and is easily checked to be Lagrangian. Then  $L = L/K \circ L_\Phi$ .  $\square$

Unfortunately the generalization of this universal property to arbitrary morphisms in the symplectic category requires rather complicated freeness assumptions.

**3.4. Fubini-Study actions.** Kähler manifolds are complex manifolds with symplectic structures that are compatible, in a certain sense, with the complex structure. An *almost complex structure* on a manifold  $X$  is an endomorphism  $J \in \text{End}(TX)$  with  $J^2 = -I$ , where  $I \in \text{End}(TX)$  is the identity. An almost complex structure  $J$  is *compatible* with a symplectic structure  $\omega$  if  $\omega(\cdot, J\cdot)$  is a Riemannian metric. Any symplectic manifold admits a compatible almost complex structure; a Kähler manifold is a symplectic manifold equipped with an integrable compatible almost complex structure.

Affine and projective space have natural *Fubini-Study* Kähler structures as follows. Any Hermitian structure  $(\cdot, \cdot) : V \times V \rightarrow \mathbb{C}$  defines a symplectic structure on  $V$  via its imaginary part,

$$\omega_{V,v}(v_1, v_2) = \text{Im}(v_1, v_2).$$

while its real part gives a Riemannian metric on  $V$ . Let  $K$  be a Lie group acting on  $V$ . If  $K$  preserves the Hermitian structure then the action is symplectic and a canonical moment map is given by

$$\langle \Phi_V(v), \xi \rangle = \text{Im}(v, \xi v)/2.$$

*Example 3.4.1.* Let  $K = \text{Sp}(V, \omega)$  be the group of linear symplectomorphisms of  $V$  then the map  $\xi \mapsto \langle \Phi_V, \xi \rangle$  defines an isomorphism of the Lie algebra  $\mathfrak{sp}(V, \omega)$  with  $\text{Sym}^2(V^\vee)$ , analogous to the isomorphism of the *orthogonal* Lie algebras  $\mathfrak{o}(V, g)$  with  $\Lambda^2(V)$ . The Lie algebra structure induced on  $\text{Sym}^2(V^\vee)$  is that induced from the Poisson bracket by the inclusion  $\text{Sym}^2(V^\vee) \subset C^\infty(V)$ .

*Example 3.4.2.* Let  $K = S^1$  act on  $V = \mathbb{C}^n$  with weights  $a_1, \dots, a_n$ . If the Hermitian structure on  $V$  is the standard one then the moment map on  $V$  is Hamiltonian with moment map

$$\Phi(z_1, \dots, z_n) = \sum_{j=1}^n -a_j |z_j|^2 / 2$$

In particular, if  $K$  acts by scalar multiplication then the moment map is

$$\Phi(z_1, \dots, z_n) = - \sum_{j=1}^n |z_j|^2 / 2.$$

The canonical symplectic quotient  $V // S^1$  is a point. If we shift the moment map by a scalar,  $\Phi_c = \Phi + c$ , then the symplectic quotient is

$$V // S^1 = \left\{ \sum_{j=1}^n |z_j|^2 / 2 = c \right\} / S^1$$

which identifies with the projective space  $\mathbb{P}(V)$  of complex lines in  $V$  via  $V // S^1 \rightarrow \mathbb{P}(V), [v] \mapsto \text{span}(v)$ .

It follows that projective space  $\mathbb{P}(V)$  naturally has a symplectic structure, called the *Fubini-Study* symplectic form  $\omega_{\mathbb{P}(V)}$ . Explicitly this is given as follows: The tangent space to  $\mathbb{P}(V)$  at  $[v], v \in V - \{0\}$  naturally identifies with the Hermitian orthogonal to  $[v]$ . Then

$$\omega_{\mathbb{P}(V), [v]}(v_1, v_2) = \frac{\text{Im}(v_1, v_2)}{(v, v)}.$$

If  $z_1, \dots, z_n$  are coordinates corresponding to a unitary basis then

$$\omega_{\mathbb{P}(V), [z]} = \frac{i \sum_{j=1}^n dz_j \wedge d\bar{z}_j}{2 \sum_{j=1}^n z_j \bar{z}_j}.$$

If  $K$  acts on  $V$  preserving the Hermitian structure, then it commutes with the action of  $S^1$ . The induced action on  $\mathbb{P}(V)$  is also symplectic, and has canonical moment map

$$\langle \Phi_{\mathbb{P}(V)}([v]), \xi \rangle = \text{Im}(v, \xi v) / (v, v).$$

Suppose that  $K = S^1$ , and acts on  $V$  with weights  $a_1, \dots, a_n \in \mathbb{Z}$ . The action of  $K$  on  $\mathbb{P}(V)$  is Hamiltonian with moment map

$$(3.3) \quad \Phi_{\mathbb{P}(V)}([z_1, \dots, z_n]) = \frac{\sum_{j=1}^n -a_j |z_j|^2 / 2}{\sum_{j=1}^n |z_j|^2 / 2}.$$

**Proposition 3.4.3.** *Let  $K$  act on  $V$  preserving the Hermitian structure. Any smooth invariant subvariety  $X \subset \mathbb{P}(V)$  inherits the structure of a Hamiltonian  $K$ -manifold from the Fubini-Study Hamiltonian  $K$ -manifold structure on  $\mathbb{P}(V)$ .*

*Proof.* It suffices to check that the restriction of  $\omega_{\mathbb{P}(V)}$  to  $X$  is non-degenerate, which holds since  $\omega_{\mathbb{P}(V)}(v, Jv) > 0$  for  $v \in T_x X, Jv \in T_x X$  since  $T_x X$  is  $J$ -invariant.  $\square$

**3.5. Geometric quantization.** The philosophy of geometric quantization played an important role in the development of equivariant symplectic geometry. Unfortunately good quantization schemes exist only for certain classes of Hamiltonian actions.

Suppose that  $Q$  is a manifold and  $T^\vee Q$  its cotangent bundle. One thinks of  $T^\vee Q$  as the space of classical states for a particle moving on  $Q$ , with a vector in  $T_q^\vee Q$  representing the momentum. In quantum mechanics the state of the system is given by a *quantum wave-function*  $\psi \in L^2(Q)$ , whose norm-square  $|\psi(q)|^2$  represents the probability of finding the particle at position  $q$ , if its position is measured. The construction of  $L^2(Q)$  from  $T^\vee Q$  can be done in two steps: first cut down the number of directions by half, then pass to functions.

One can try to extend this procedure to arbitrary symplectic manifolds  $(X, \omega)$  by axiomatizing this two-step process. A *Lagrangian distribution* resp. *complex Lagrangian distribution* is a subbundle  $P \subset TX$  resp  $TX \otimes_{\mathbb{R}} \mathbb{C}$  such that each fiber  $P_x$  is a Lagrangian subspace of  $T_x X$  resp. complex Lagrangian subspace of  $T_x X \otimes_{\mathbb{R}} \mathbb{C}$ . A *polarization* is a Hermitian line bundle  $L$

with connection  $\nabla$  such that the curvature of  $\nabla$  is  $\text{curv}(\nabla) = (2\pi/i)\omega$ . A *quantization datum* resp. *complex quantization datum* consists of a Lagrangian distribution resp. complex Lagrangian distribution together with a polarization. The original literature on geometric quantization uses polarization to refer to the Lagrangian distribution. This conflicts with the use of polarization in the geometric invariant theory literature, which we have adopted. The *geometric quantization* of  $(X, \omega)$  (depending on the choice of  $(P, L, \nabla)$ ) is the vector space of smooth sections of  $L$  which are covariant constant with respect to  $\nabla$  along  $P$ :

$$\mathcal{H}(X, \omega) := \{\sigma \in \Gamma(L), \nabla_v \sigma = 0 \ \forall v \in P\}.$$

We ignore the problem of defining a Hilbert space structure on  $\mathcal{H}(X, \omega)$ , see [40] for more details.

A case for which a good quantization procedure exists is the case that  $X$  is a compact Kähler Hamiltonian  $K$ -manifold equipped with polarization  $\mathcal{O}_X(1) \rightarrow X$ . A Lagrangian distribution is provided by the antiholomorphic directions on  $X$ , that is,  $P = T^{0,1}X \subset TX \otimes_{\mathbb{R}} \mathbb{C}$ . Then  $\mathcal{H}(X, \omega) = H^0(X, \mathcal{O}_X(1))$ . In other words, in the language of geometric quantization *holomorphic sections of the polarizing line bundle are quantum states*.

One can now compare the various operations on symplectic manifolds with those on vector spaces:

- Proposition 3.5.1.** (a) (*Duals*) If  $J$  is the complex structure for  $X$  then  $-J$  is a compatible complex structure for  $X^-$ . If  $P = T^{0,1}X$  then  $\bar{P} = T^{0,1}X^-$ . Furthermore,  $\bar{L}$  with connection  $-\alpha$  is naturally a polarization for  $X^-$ . Thus  $\mathcal{H}(X^-)$  is the space of complex-conjugates of sections of  $L$ , which is naturally identified with the dual  $\mathcal{H}(X)^\vee$  of  $\mathcal{H}(X)$ .
- (b) (*Sums*) If  $X_0, X_1$  are Kähler Hamiltonian  $K$ -manifolds with polarizations, then  $\mathcal{H}(X_0 \cup X_1) = \mathcal{H}(X_0) \oplus \mathcal{H}(X_1)$ .
- (c) (*Products*) With the same assumptions as in (b),  $\mathcal{H}(X_0 \times X_1) = \mathcal{H}(X_0) \otimes \mathcal{H}(X_1)$ .

*Example 3.5.2.* Let  $X = S^2 \cong \mathbb{P}^1$  and  $\omega$  the standard symplectic form. The moment map for the action of  $S^1$  on  $(X, d\omega)$  has image  $[-d, d]$ . The  $d$ -th tensor product  $\mathcal{O}_X(d)$  of the hyperplane bundle  $\mathcal{O}_X(1)$  is a polarization of  $(X, d\omega)$ , so that  $\mathcal{H}(X, \omega) = H^0(X, \mathcal{O}_X(d))$  is the space of homogeneous polynomials in two variables of degree  $d$ . Note that the weights of  $\mathcal{H}(X, \omega)$  are  $\{d, d-2, d-4, \dots, -d\}$ , which are the intersections points of the image  $\Phi(X)$  with the lattice  $d + 2\mathbb{Z} \subset \mathbb{Z}$ . The  $SU(2)$ -action on  $X$  induces on  $\mathcal{H}(X)$  the structure of an  $SU(2)$ -module with highest weight  $d$ . The product of spheres  $S_{\lambda_1}^2 \times \dots \times S_{\lambda_n}^2$  has quantization the tensor product of simple  $SU(2)$ -modules  $V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n}$ .

Unfortunately (i) quantizing arbitrary morphisms (i.e. Lagrangian correspondences) is quite difficult, even in this case (ii) there is no good geometric quantization scheme for arbitrary symplectic manifolds. The problem of finding good schemes for say, coadjoint orbits of real Lie groups or moduli spaces of flat connections have vast literatures attached to them.

The reader may notice that we have not said anything yet about the behavior of the quantum state spaces under the symplectic quotient construction. We take this up in Section 5.

#### 4. GEOMETRIC INVARIANT THEORY

In this section we review Mumford's geometric invariant theory [82], see also Brion's review in this volume or the reviews by Newstead [86] or Schmitt [93]. For connections to moduli problems see Newstead [85].

**4.1. Algebraic group actions and quotients.** Let  $G$  be a complex linear algebraic group.  $G$  is called *reductive* iff every  $G$ -module splits into simple  $G$ -modules, or equivalently, if  $G$  is the complexification of a compact Lie group  $K$ . A *Borel subgroup* of a reductive group  $G$  is a maximal closed connected solvable subgroup  $B \subset G$ . The set of Borel subgroups is in bijection with set of right cosets  $G/B$ , called the *generalized flag variety* for  $G$ , via the map  $gB \mapsto gBg^{-1}$ . A subgroup  $P \subset G$  is *parabolic* iff  $G/P$  is complete iff  $P$  contains a Borel subgroup. The quotient  $G/P$  is called a *generalized partial flag variety*. Let  $T$  be a maximal torus of  $G$  (for example, the complexification of a maximal torus of the maximal compact subgroup, which was also somewhat confusingly called  $T$ .) We denote by  $W = N(T)/T$  the Weyl group of  $T$ . The action of  $T$  on the Lie algebra  $\mathfrak{g}$  induces

a root space decomposition

$$\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \mathcal{R}(\mathfrak{g})} \mathfrak{g}_\alpha$$

where  $T$  acts trivially on  $\mathfrak{t}$  and on  $\mathfrak{g}_\alpha$  by  $t\xi = t^\alpha\xi$ , and  $\mathcal{R}(\mathfrak{g}) \subset \Lambda^\vee$  is the set of roots of  $\mathfrak{g}$ . Given a choice of positive Weyl chamber let  $B^\pm$  be the Borel subgroups whose Lie algebras contain the positive resp. negative root spaces of  $\mathfrak{g}$ . Each  $\lambda \in \mathfrak{t}^\vee$  determines *standard parabolic subgroups*  $P_\lambda^\pm$  with Lie algebra  $\mathfrak{p}_\lambda^\pm = \mathfrak{b}^\pm + \bigoplus_{\langle h_\alpha, \lambda \rangle = 0} \mathfrak{g}_\alpha$ , where  $h_\alpha \in \mathfrak{t}$  is the coroot corresponding to  $\alpha \in \mathfrak{t}^\vee$ . Any parabolic subgroup (in particular, any Borel) is conjugate to a standard parabolic subgroup.

An *action* of  $G$  on a variety  $X$  is a morphism  $G \times X \rightarrow X$  such that  $g_1(g_2x) = (g_1g_2)x$  and  $ex = x$ , for all  $g_1, g_2 \in G, x \in X$ . A variety  $X$  equipped with a  $G$ -action is called a  $G$ -variety. An (*étale*) *slice* for the action of  $G$  at  $x \in X$  is an affine subvariety  $V \subset X$  and a  $G$ -morphism  $G \times_{G_x} V \rightarrow X$  that is an isomorphism (*étale morphism*) onto a neighborhood of  $X$ . In contrast with the case of compact group actions, reductive group actions do not in general have slices. Luna's slice theorem [76] asserts that any *closed* orbit of an action of a reductive group on an affine variety has an étale slice. A *categorical quotient* of  $X$  by  $G$  is a pair  $(Y, \pi)$  where  $Y$  is a variety and  $\pi : X \rightarrow Y$  is a  $G$ -invariant morphism that satisfies the universal property for quotients: if  $f : X \rightarrow Z$  is a  $G$ -invariant morphism then  $f$  factors uniquely through  $Y$ . A *good quotient* of  $X$  is a pair  $(Y, \pi)$  where

- (a)  $\pi : X \rightarrow Y$  is  $G$ -invariant, affine, surjective,
- (b) if  $U \subset Y$  is open then  $\mathcal{O}_Y(U) \rightarrow \mathcal{O}_X(\pi^{-1}(U))^G$  is an isomorphism
- (c) If  $W_1, W_2$  are disjoint closed  $G$ -invariant subsets of  $X$  then  $\pi(W_1), \pi(W_2)$  are disjoint closed subsets of  $Y$ .

A good quotient is automatically a categorical quotient. A *geometric quotient* is a good quotient that separates orbits.

*Example 4.1.1.* The generalized flag variety  $X = G/B^-$  has is a  $G$ -variety for the left action. If  $G$  is connected reductive then  $X$  has a canonical decomposition into *Bruhat cells*

$$(4.1) \quad X = \bigcup_{w \in W} X_w, \quad X_w := BwB^-/B^-$$

and *opposite Bruhat cells*

$$(4.2) \quad X = \bigcup_{w \in W} Y_w, \quad Y_w := B^-wB^-/B^-.$$

The codimension resp. dimensions are given by

$$\text{codim}(X_w) = l(w), \quad \dim(Y_w) = l(w)$$

where  $l(w)$  is the minimal number of simple reflections in a decomposition of  $w$ . We denote by  $x_w = wB^-/B^- = X_w \cap Y_w$  the unique  $T$ -fixed point in  $X_w$  resp.  $Y_w$ . There is a similar decomposition of any generalized flag variety  $X = G/P_\lambda^-$  into cells  $X_{[w]}$  indexed by  $[w] \in W/W_\lambda$ . In the special case  $G = GL(r)$ , the Weyl group  $W$  is naturally identified with the symmetric group and  $B^\pm$  are the groups of invertible upper resp. lower triangular matrices. We identify  $\mathfrak{k} \rightarrow \mathfrak{k}^\vee$ ; if  $\lambda = \text{diag } i(1, \dots, 1, 0, \dots, 0)$  has rank  $s$  then  $P_\lambda$  is the group of matrices preserving the subspace  $\mathbb{C}^s \oplus 0 \subset \mathbb{C}^r$ . The quotient  $X = G/P_\lambda$  is isomorphic to the Grassmannian  $G(s, r)$  of  $s$ -dimensional subspaces of  $\mathbb{C}^r$ . The quotient  $W/W_\lambda$  is natural identified with the set of subsets  $I \subset \{1, \dots, r\}$  of size  $s$  via the map  $w \mapsto w\{1, \dots, s\}$ . Let  $F_1 \subset F_2 \subset \dots \subset F_r = \mathbb{C}^r$  be the standard flag in  $\mathbb{C}^r$ . Then the opposite Bruhat cell  $Y_I$  has closure the *Schubert variety*

$$(4.3) \quad \overline{Y}_I = \{E \in G(s, r), \dim(E \cap F_{i_j}) \geq j, j = 1, \dots, s\}.$$

This ends the example.

**4.2. Stability conditions.** Let  $G$  be a complex reductive group and  $X$  a  $G$ -variety. A *polarization* of  $X$  is an ample  $G$ -line bundle  $\mathcal{O}_X(1) \rightarrow X$ . Its  $d$ -th tensor power is denoted  $\mathcal{O}_X(d)$ . Let

$$R(X) = \bigoplus_{d \geq 0} H^0(X, \mathcal{O}_X(d)).$$

The action of  $X$  induces an action on  $R(X)$  by pull-back. We denote by  $R(X)^G \subset R(X)$  the subring of invariants, and by  $R(X)_{>0}^G$  the part of  $R(X)^G$  of positive degree.

**Definition 4.2.1.** A point  $x \in X$  is

- (a) *semistable* if  $s(x) \neq 0$  for some  $s \in R(X)_{>0}^G$ ;
- (b) *polystable* if  $x$  is semistable and  $Gx \subset X^{\text{ss}}$  is closed;
- (c) *stable* if  $x$  is polystable and has finite stabilizer;
- (d) *unstable* if  $x$  is not semistable.

*Example 4.2.2.* Suppose that  $G = \mathbb{C}^*$  acts on  $\mathbb{P}^2$  by  $g[z_0, z_1, z_2] = [g^{-1}z_0, z_1, gz_2]$ . Then  $R(X)_d$  is spanned by  $z_0^{d_0} z_1^{d_1} z_2^{d_2}$  with  $d_0 + d_1 + d_2 = d$ , which has weight  $d_0 - d_2$  under  $\mathbb{C}^*$ . Thus the invariant sections have  $d_0 = d_2$ . One sees easily that  $x$  is

- (a) semistable iff  $x \neq [1, 0, 0], [0, 0, 1]$
- (b) polystable iff  $x \in \{[0, 1, 0]\} \cup \{[z_0, z_1, z_2] \mid z_0 z_2 \neq 0\}$
- (c) stable iff  $x \in \{[z_0, z_1, z_2] \mid z_0 z_2 \neq 0\}$

Let  $X^{\text{ss}}$  resp.  $X^{\text{ps}}$  resp.  $X^{\text{s}}$  resp.  $X^{\text{us}}$  denote the semistable resp. polystable resp. stable resp. unstable locus. We will need the following alternative characterizations of poly resp. semistability, see Mumford [82] or Brion's lectures in this volume:

**Lemma 4.2.3.** *Let  $X \subset \mathbb{P}(V)$  be a  $G$ -variety. A point  $x \in X$  is polystable (resp. semistable) iff the orbit of any lift  $v$  in  $V$  is closed (resp. 0 does not lie in the closure of  $Gv$ ).*

Define an equivalence relation on orbits as follows:

**Definition 4.2.4.** *Orbit-equivalence* is the equivalence relation on  $X^{\text{ss}}$  defined by  $x_0 \sim x_1$  iff  $\overline{Gx_0} \cap \overline{Gx_1} \cap X^{\text{ss}} \neq \emptyset$ .

Transitivity of this relation follows from:

**Proposition 4.2.5.** *(see [82]) The closure  $\overline{Gx}$  of any semistable  $x$  contains a unique polystable orbit. Hence two orbits  $Gx_0, Gx_1$  are orbit-equivalent iff their closures contain the same polystable orbit.*

See Theorem 5.5.9 for an analytic proof. The following can be considered the main result of geometric invariant theory [82]:

**Theorem 4.2.6** (Mumford). *Let  $X$  be a projective  $G$ -variety equipped with polarization  $\mathcal{O}_X(1)$ .*

- (a) *There exists a categorical quotient  $\pi : X^{\text{ss}} \rightarrow X//G$ .*
- (b)  *$\pi(X^{\text{s}}) \subset X//G$  is open and  $\pi|_{X^{\text{s}}} : X^{\text{s}} \rightarrow \pi(X^{\text{s}})$  is a geometric quotient.*
- (c) *The topological space underlying  $X//G$  is the space of orbits modulo the orbit-closure relation  $X^{\text{ss}}/\sim$ .*
- (d)  *$X//G$  is isomorphic to the projective variety with coordinate ring  $R(X)^G$ .*

Some authors prefer to write  $X^{\text{ss}}//G$  for the geometric invariant theory quotient, while we drop the superscript from the notation.

**4.3. The Hilbert-Mumford criterion.** Mumford [82], based on previous work of Hilbert for the case of the special linear group acting on projective space, gave a method for explicitly identifying the semistable loci:

**Theorem 4.3.1.** *(Hilbert-Mumford criterion) Let  $X$  be a polarized projective  $G$ -variety.  $x \in X$  is semistable iff  $x$  is semistable for all one-parameter subgroups  $\mathbb{C}^* \rightarrow G$ .*

One direction of the Hilbert-Mumford criterion is trivial: Let  $X$  be a polarized  $G$ -variety. Suppose that  $x$  is  $G$ -semistable, so that there exists  $s \in R(X)_{>0}^G$  with  $s(x) \neq 0$ . Then  $s$  is also invariant for any one-parameter subgroup, hence  $x$  is semistable for any one-parameter subgroup. The other direction is somewhat harder; the proof given in Mumford [82] uses an algebraic theorem of Iwahori. We will give an alternative analytic proof using the Kempf-Ness function in Section 7.2. The following is a fundamental example:

*Example 4.3.2.* Let  $X = (\mathbb{P}^1)^n$  and  $\mathcal{O}_X(1) = \mathcal{O}_{\mathbb{P}^1}(1)^{\boxtimes n}$  the  $n$ -fold exterior tensor product. The group  $G = SL(2, \mathbb{C})$  acts diagonally on  $X$ . We wish to show

- (a)  $X^{\text{ss}} = \{(x_1, \dots, x_n) \in (\mathbb{P}^1)^n, \text{ at most } n/2 \text{ points equal}\}.$
- (b)  $X^{\text{s}} = \{(x_1, \dots, x_n) \in (\mathbb{P}^1)^n, \text{ less than } n/2 \text{ points equal}\}.$
- (c)  $X^{\text{ps}} - X^{\text{s}} = \{(x_1, \dots, x_n) \in X^{\text{ss}}, \#\{x_1, \dots, x_n\} = 2\}.$  In other words,  $n/2$  are equal and the other  $n/2$  are also equal.

Indeed, if  $z_j, w_j$  are the coordinates on the  $j$ -factor then  $H^0(\mathcal{O}_X(d))$  is spanned by  $z_1^{d_1} w_1^{d-d_1} \dots z_n^{d_n} w_n^{d-d_n}$  where  $d_j \in [0, d], j = 1, \dots, n$ . If  $\mathbb{C}^* \subset G$  is the standard maximal torus given by  $g \mapsto \text{diag}(g, g^{-1})$  then  $H^0(\mathcal{O}_X(d))^{\mathbb{C}^*}$  is spanned by the polynomials  $z_1^{d_1} w_1^{d-d_1} \dots z_n^{d_n} w_n^{d-d_n}$  with  $\sum_{j=1}^n d_j = \sum_{j=1}^n d - d_j$ , that is,  $\sum(d_j/d) = n/2$ . Since  $d_j/d \in [0, 1]$ , this means that at least  $n/2$  of the  $d_j$ 's are non-zero. Thus  $([z_1, w_1], \dots, [z_n, w_n])$  is  $\mathbb{C}^*$ -semistable iff at most  $n/2$   $z_j$ 's and at most  $n/2$   $w_j$ 's equal zero. Repeating the same for an arbitrary one-parameter subgroup (or equivalently, basis for  $\mathbb{C}^2$ ) proves the claim.

*Example 4.3.3.* More generally, suppose that  $X = (\mathbb{P}^1)^n$  is equipped with the polarization  $\mathcal{O}_X(1) := \boxtimes_{i=1}^n \mathcal{O}_{\mathbb{P}^1}(\lambda_i)$  for some positive integers  $\lambda_1, \dots, \lambda_n$ . Then  $x = (x_1, \dots, x_n)$  is semistable iff for all  $x \in \mathbb{P}^1$ ,

$$\sum_{x_j=x} \lambda_j \leq \sum_{x_j \neq x} \lambda_j.$$

For future use we mention the following equivalent form of the Hilbert-Mumford criterion and Lemma 4.2.3:

**Corollary 4.3.4.** *Let  $G$  be a reductive group acting linearly on a finite dimensional vector space  $V$ . For any  $v \in V$ ,  $\overline{Gv}$  contains 0, if and only if  $\overline{\mathbb{C}^*v}$  contains 0 for some one-parameter subgroup  $\mathbb{C}^* \subset G$ .*

*Remark 4.3.5.* The statement of the corollary does not hold for arbitrary (that is, not linear) actions resp. arbitrary points. An example I learned from Brion: Let  $X = \mathbb{P}(S^3(\mathbb{C}^2) \oplus \mathbb{C})$  with the action induced from the action of  $SL(2, \mathbb{C})$  on  $\mathbb{C}^2$  and the trivial action on  $\mathbb{C}$ . Identifying  $S^3(\mathbb{C}^2)$  with homogeneous polynomials in two variables  $u, v$ , one sees that the orbit of  $[u^2v, 1]$  contains the orbit of  $[u^3, 1]$  in its closure. The stabilizer of  $[u^3, 1]$  is a maximal unipotent subgroup of  $SL(2, \mathbb{C})$  and so does not contain a copy of  $\mathbb{C}^*$ . Thus  $[u^3, 1]$  cannot be contained in the closure of an orbit of a one-parameter subgroup. On the other hand, the lemma is true for arbitrary actions of abelian groups, as follows from, for example, Atiyah Theorem' 8.2.1 below.

## 5. THE KEMPF-NESS THEOREM

The material in this section is contained in the original paper of Kempf-Ness [60], the book of Mumford-Fogarty-Kirwan [82], and the paper of Guillemin-Sternberg [38]. The notes of Thomas [105] and the thesis of Székelyhidi [102] also describe the Kempf-Ness theorem with many examples and generalizations.

**5.1. Complexification of Lie groups and their actions.** We begin with some basic remarks on the relation between complex and compact group actions. Any compact Lie group  $K$  admits a *complexification*  $G$ , a complex reductive Lie group  $G$  containing  $K$  as a maximal compact real subgroup, and whose Lie algebra  $\mathfrak{g}$  is equal to  $\mathfrak{k} \oplus i\mathfrak{k}$ . The complexification  $G$  satisfies the universal property that any Lie group homomorphism from  $K$  to a complex Lie group  $H$  extends to a complex Lie group homomorphism from  $G$ . The complexification  $G$  admits a *Cartan decomposition*: a diffeomorphism (see Helgason [51, VI.1.1])

$$(5.1) \quad K \times \mathfrak{k} \rightarrow G, \quad (k, \xi) \mapsto k \exp(i\xi).$$

We denote by  $K \backslash G$  the quotient by the left action, which is a symmetric space of non-compact type with non-positive curvature [51], [4]. For any point  $[g] \in K \backslash G$  the geodesics through  $[g]$  are of the form  $[\exp(i\xi)g]$  for  $\xi \in \mathfrak{k}$  (see [51, Exercise 1]) and  $\xi \mapsto [\exp(i\xi)]$  defines a diffeomorphism of  $\mathfrak{k}$  onto  $K \backslash G$ .

If  $X$  is a compact complex manifold then the group  $\text{Aut}(X)$  of automorphisms is a complex Lie group, with Lie algebra given by the space  $H^0(X, TX)$  of holomorphic vector fields on  $X$ , see for example Akhiezer [3]. Any action of a compact group  $K$  therefore extends to the complexification  $G$ .

By a *Kähler Hamiltonian  $K$ -manifold* we mean a compact Hamiltonian  $K$ -manifold equipped with an integrable  $K$ -invariant complex structure. If  $X$  is compact then the  $K$ -action automatically extends to a  $G$ -action preserving the complex structure but not the symplectic structure. By the Kodaira embedding theorem, if the symplectic form is rational then a compact Kähler Hamiltonian  $K$ -manifold is isomorphic as a complex  $G$ -manifold to a smooth complex algebraic  $G$ -variety. However, the symplectic form may not be the pull-back of the Fubini-Study form under any holomorphic embedding of  $X$ , see for example Tian [106]. The generating vector fields for  $\xi \in \mathfrak{k}$  are the Hamiltonian flows corresponding to the moment map components  $\langle \Phi, \xi \rangle$ , while the generating vectors fields for  $i\xi, \xi \in \mathfrak{k}$  are the *gradient flows* corresponding to  $\langle \Phi, \xi \rangle$ . In particular, for any  $x \in X, \xi \in \mathfrak{k}$ , the trajectory  $\exp(it\xi)x$  converges to a point  $x_\infty \in X$  with  $\xi_X(x_\infty) = 0$ . Furthermore, since  $\langle \Phi, \xi \rangle$  is a Morse function by Lemma 3.2.7, this convergence is *exponentially fast* in  $t$ ; the exponential nature of convergence will be used later.

*Example 5.1.1.* The example of flag varieties will be particularly important later and we briefly describe these actions from the algebraic and symplectic points of view. Let  $V$  be a finite dimensional vector space. A *partial flag* in  $V$  is a filtration  $F = (F_1 \subset F_2 \subset \dots \subset F_m \subset V)$ . The *type* of  $F$  is the sequence of dimensions  $\dim(F_1) < \dim(F_2) < \dots < \dim(F_m)$ . Given a sequence  $t = (0 < t_1 < \dots < t_m < \dim(V)) \in \mathbb{Z}^m$  we let  $\text{Fl}(t, V)$  denote the set of partial flags of type  $t$ . The general linear group  $GL(V)$  acts transitively on  $\text{Fl}(t, V)$  with stabilizer the parabolic subgroup of transformations preserving the filtration. A  $GL(V)$ -equivariant canonical projective embedding of  $\text{Fl}(t, V)$  is given by choosing a basis  $v_1, \dots, v_n$  so that  $v_1, \dots, v_{t_j}$  is a basis for  $F_j$  for each  $j = 1, \dots, m$ , and mapping

$$\text{Fl}(t, V) \rightarrow \prod_{j=1}^m \mathbb{P}(\Lambda^{t_j} V), \quad F \mapsto \prod_{j=1}^m \Lambda_{k=1}^{t_j} v_k.$$

Given a Hermitian metric on  $V$ , any partial flag induces a Hermitian splitting

$$V = F_1 \oplus (F_2 \cap F_1^\perp) \oplus (F_3 \cap F_2^\perp) \dots \cap (F_m \cap F_{m-1}^\perp)$$

and such splittings are in one-to-one correspondence with flags. Given real numbers  $\lambda_1 > \dots > \lambda_m$  the flag defines a skew-Hermitian operator acting by  $i\lambda_j$  on  $F_j \cap F_{j-1}^\perp$ . Conversely, any such Hermitian operator determines a splitting via its eigenspace decomposition. The unitary group  $K = U(V)$  acts transitively on the space of such matrices, which form an orbit of the action of  $K$  on the Lie algebra  $\mathfrak{k}$ . Now  $\mathfrak{k}$  may be identified with its dual via any invariant inner product, so one sees that  $\text{Fl}(t, V)$  is naturally identified with the coadjoint orbit  $K\lambda$  of  $\lambda$ , identified with an element of  $\mathfrak{k}^\vee$  via the inclusion  $\mathfrak{t} \rightarrow \mathfrak{k}$  and an identification  $\mathfrak{k} \rightarrow \mathfrak{k}^\vee$ . Given a generic  $\xi \in \mathfrak{t}_+$ , the stable resp. unstable manifolds of the Morse function  $\langle \Phi, \xi \rangle$  are the Bruhat resp. opposite Bruhat cells of (4.1) resp. (4.2).

**5.2. Statement and proof.** The Kempf-Ness theorem states the equivalence of the symplectic and geometric invariant theory quotients; the affine case is treated in [60] and the projective case is similar (Theorem 8.3 in [82]).

**Theorem 5.2.1.** *Let  $K$  be a compact group and  $G$  its complexification. Let  $V$  be a  $G$ -module equipped with a  $K$ -invariant Hermitian structure. Let  $X \subset \mathbb{P}(V)$  be a smooth projective  $G$ -variety, and  $\Phi : X \rightarrow \mathfrak{k}^\vee$  the Fubini-Study moment map. Then  $\Phi^{-1}(0) \subseteq X^{\text{ps}}$  and the inclusion induces a homeomorphism  $X//K \rightarrow X//G$ .*

The proof uses the properties of a *Kempf-Ness function* for each  $v \in V - \{0\}$ :

$$\psi_v : K \backslash G \rightarrow \mathbb{R}, \quad [g] \mapsto \log \|gv\|^2/2.$$

We denote by  $\partial_\lambda \psi_v([g])$  the derivative of  $\psi_v$  along the geodesic  $[\exp(i\lambda)g]$  determined by  $\lambda$ ; note that this depends on a choice of representative  $g$  of  $[g]$ . The Kempf-Ness function can be viewed as the integral of the moment map in the following sense:

**Lemma 5.2.2.** *For all  $v \in V$  and  $\lambda \in \mathfrak{k}$  we have  $\partial_\lambda \psi_v([g]) = -\langle \Phi([gv]), \lambda \rangle$ .*



*Proof.* The proof uses the explicit formula for the Fubini-Study moment map

$$\begin{aligned}\partial_\lambda \psi_v([g]) &= \frac{d}{dt} \Big|_{t=0} \log \|\exp(it\lambda)gv\|^2/2 \\ &= \frac{(i\lambda gv, gv)}{(gv, gv)} = -\langle \Phi([gv]), \lambda \rangle.\end{aligned}$$

□

**Corollary 5.2.3.** *The gradient of  $\psi$  is equal to  $\Phi$ , that is,*

$$\text{grad } \psi([g]) = \frac{d}{dt} \Big|_{t=0} [\exp(-it\Phi(gx)g)].$$

*Proof.* By Lemma 5.2.2,

$$(\text{grad } \psi([g]), \lambda) = \frac{d}{dt} \Big|_{t=0} \psi([\exp(it\lambda)g]) = -\langle \Phi(gx), \lambda \rangle.$$

□

The basic property of the Kempf-Ness function is its *convexity*: its restriction to any geodesic in  $K \backslash G$  is a convex function, or equivalently, its second derivatives along geodesics are non-negative [30, Section 6], [59, Section 2].

**Corollary 5.2.4.** (a) *For any  $v \in V$ ,  $\psi_v$  is a convex function with critical points given by points  $[g] \in K \backslash G$  such that  $\Phi([gv]) = 0$ .*  
 (b) *The second derivative  $\partial_\lambda^2 \psi_v([e])$  along the geodesic determined by  $\lambda \in \mathfrak{k}$  is positive iff  $\lambda$  lies in  $\mathfrak{k} - \mathfrak{k}_x$ .*  
 (c) *For  $\xi \in \mathfrak{k}_x$  we have  $\psi_v([\exp(i\xi)]) = \psi_v([e]) + 2\langle \Phi(x), \xi \rangle$ .*

*Proof.* The claim on the critical points follows from Lemma 5.2.2:

$$\begin{aligned}D_{[g]}\psi_v = 0 &\iff \partial_\lambda \psi_v([g]) = 0, \forall \lambda \in \mathfrak{k} \\ &\iff \langle \Phi([gv]), \lambda \rangle = 0, \forall \lambda \in \mathfrak{k} \\ &\iff \Phi([gv]) = 0.\end{aligned}$$

To prove convexity we compute the second derivatives

$$\begin{aligned}(\partial_\lambda)^2 \psi_v([g]) &= -\frac{d}{dt} \Big|_{t=0} \langle \Phi([\exp(i\lambda t)gv]), \lambda \rangle \\ &= -2L_{J\lambda_X} \langle \Phi, \lambda \rangle([gv]) \\ &= 2\omega(\lambda_X, J\lambda_X)([gv]) \geq 0\end{aligned}$$

since  $\omega(\cdot, J\cdot)$  is a Riemannian metric. The claim on strict convexity and the formula for  $\psi_v([\exp(i\xi)])$ ,  $\xi \in \mathfrak{k}_x$  are immediate from the previous lemma. □

Note that if  $\psi_v$  is strictly convex (that is, has trivial infinitesimal stabilizer) and has a critical point, then the critical point is the unique global minimum. The following lemma characterizes for which  $v$  minima of  $\psi_v$  exist:

**Lemma 5.2.5.** *Let  $v \in V - \{0\}$  and  $x = [v] \in \mathbb{P}(V)$ .*

- (a)  *$\psi_v$  attains a minimum iff  $x$  is polystable.*
- (b)  *$\psi_v$  is bounded from below iff  $x$  is semistable.*

*Proof.* (a) Recall from 4.2.3 that  $x$  is polystable iff  $Gv$  is closed. Suppose  $Gv$  is closed. Let  $Kg_j \in K \backslash G$  be a minimizing sequence for  $\psi_v$ . Then after passing to a subsequence  $g_j v$  converges to  $gv$  for some  $g \in G$ , since  $Gv$  is closed, and  $Kg$  must be a global minimum of  $\psi_v$ , since  $\psi_v$  is convex.

Conversely, suppose that  $\psi_v$  attains a minimum at  $Kg \in K \backslash G$ . After replacing  $\psi_v$  with  $\psi_{gv}$ , we may assume that  $g = e$ . Clearly  $\psi_v$  is invariant under the stabilizer  $G_v$  of  $v$ . We claim that the induced map

$$(5.2) \quad \psi_v/G_v : K \backslash G/G_v \rightarrow \mathbb{R}$$

is proper. Suppose that  $Kg_j G_v$  is a sequence of points in  $K \backslash G$  such that  $\psi(Kg_j)$  is bounded. Let  $O$  be the image of  $G_v$  under  $G \rightarrow K \backslash G$ ; since this is proper and  $O$  is an orbit of  $G_v$ ,  $O$  is a closed

submanifold. It follows that there exists a minimal length path connecting  $Kg_j$  to  $O$ . Since any such path is a geodesic, we have

$$\gamma_j(t) = K \exp(it\xi_j)h_j, t \in [0, 1]$$

for some  $\xi_j \in \mathfrak{k}$  and  $h_j \in G_v$ . The direction of  $\gamma(t)$  at  $t = 0$ ,

$$\frac{d}{dt}\Big|_{t=0}\gamma_j(t) = \frac{d}{dt}\Big|_{t=0}K \exp(it\xi_j)h_j$$

is perpendicular to the tangent space  $T_{Kh_j}O$  to  $O$  at  $Kh_j$ ,

$$T_{Kh_j}O = \left\{ \frac{d}{dt}\Big|_{t=0}K \exp(t\mu)h_j, \mu \in \mathfrak{g}_v \right\}$$

since otherwise one could find a shorter path. Thus

$$\frac{d}{dt}\Big|_{t=0}K \exp(it\xi_j)h_j \perp \frac{d}{dt}\Big|_{t=0}K \exp(i\mu)h_j$$

for all  $\mu \in \mathfrak{g}_v$ . Since the metric is invariant under the right action of  $h_j$ ,  $i\xi_j$  is perpendicular to the projection of  $\mathfrak{g}_v$  on  $i\mathfrak{k}$ . Note that since  $\psi_v$  is bounded,  $\Phi([v])$  vanishes on  $\mathfrak{k}_x$  by the last sentence in Corollary 5.2.4. Hence  $i\mathfrak{k}_x$  is contained in  $\mathfrak{g}_v$  by Proposition 3.2.10, thus  $i\xi_j$  is perpendicular to  $i\mathfrak{k}_x$ . Strict convexity of  $\psi_v$  along any geodesic of the form  $K \exp(it\xi)$  with  $\xi \in \mathfrak{k}_x^\perp$  implies that

$$\psi_v(K \exp(it\xi)) < C_0 + C_1\|\xi\|, \quad \forall \xi \in \mathfrak{k}_x^\perp.$$

where

$$C_0 = \sup_{\|\xi\| \leq 1} \psi_v(K \exp(i\xi)), \quad C_1 = \sup_{\|\xi\|=1} \frac{d}{dt}\Big|_{t=0}\psi_v(K \exp(it\xi)).$$

Since  $\psi(K \exp(it\xi_j)h_j) = \psi(K \exp(it\xi_j))$  is bounded, so is  $\xi_j$ . Hence (5.2) is proper, which completes the proof of the claim.

To show that  $Gv$  is closed, suppose that  $g_jv$  is a sequence converging to some  $v_\infty$ . Then  $\psi_v(Kg_j)$  is bounded, so by the claim on properness of (5.2), the sequence  $g_j$  converges, after passing to a subsequence, to some  $g_\infty$ . Continuity of the action implies that  $g_jv$  converges to  $g_\infty v = v_\infty$ , so  $v_\infty$  lies in  $Gv$ .

(b) If  $\psi_v$  is bounded from below, then any minimizing sequence  $\xi_j$  has  $\exp(i\xi_j)x$  converging to a critical point of  $\psi_v$ , which is necessarily a zero of  $\Phi$ . Hence  $Gx$  contains a polystable orbit in its closure and is therefore semistable. If  $\psi_v$  is not bounded from below, then  $\overline{Gv}$  contains 0 and so  $x$  is unstable, see Lemma 4.2.3.  $\square$

**Corollary 5.2.6.**  $X^{\text{ps}} = G\Phi^{-1}(0)$ .

*Proof.* By Lemmas 5.2.5, 5.2.4, 4.2.3.  $\square$

*Proof of the Kempf-Ness theorem 5.2.1.* Consider the inclusion

$$i/K : \Phi^{-1}(0)/K \rightarrow X^{\text{ps}}/G \cong X//G.$$

First note that  $i/K$  is injective: Suppose that  $x_0, x_1 \in \Phi^{-1}(0)$  are such that  $x_0 = gx_1$  for some  $g \in G$ . Choose a lift  $v$  of  $x_0$ . Then both  $[e], [g]$  are global minimum points of  $\psi_v$ , and since  $\psi_v$  is convex this implies that the geodesic  $[\exp(it\xi)], t \in [0, 1]$  connecting  $[e], [g]$  also consists of global minima. But then  $\xi \in \mathfrak{k}_{x_0}$  and so  $Kx_0 = Kx_1$ . Next note that  $i/K$  is surjective by Corollary 5.2.6. Finally  $i/K$  is a homeomorphism: Any bijection from a Hausdorff space to a compact space is a homeomorphism. (Alternative, the gradient flow of the norm-square of the moment map discussed in Section 7 defines a continuous inverse to  $i/K$ .)  $\square$

*Remark 5.2.7.* Let  $X$  be a compact Kähler Hamiltonian  $K$ -manifold. An analog of the Kempf-Ness function may be obtained by integrating the one-form given by the moment map: Define  $\alpha \in \Omega^1(K \setminus G)$  by

$$\alpha_{[g]}([\frac{d}{dt}\Big|_{t=0}\exp(it\lambda)g]) = \langle \Phi(gx), \lambda \rangle.$$

Then anti-symmetry of  $\omega$  implies that  $\alpha$  is closed, hence exact by the Poincaré lemma, hence  $\alpha_x = d\psi_x$  for some  $\psi_x : K \setminus G \rightarrow \mathbb{R}$ . Say that a point  $x \in X$  is *polystable* iff  $\psi_x$  attains a minimum, *semistable* iff  $\psi_x$  is bounded from below. With these definitions the following Kähler analog of the Kempf-Ness theorem holds, c.f. Mundet [56], Heinzner-Loose [49], Heinzner-Huckleberry [50],

Bruasse-Teleman [22], Teleman [103]: Let  $X//G$  be the quotient of the semistable locus by the orbit closure equivalence relation. Then the same arguments show that  $\Phi^{-1}(0)$  is contained in the semistable locus and the inclusion induces a homeomorphism  $X//K \rightarrow X//G$ .

We discuss the geometry of the Kempf-Ness function further in Theorem 5.4.2.

*Example 5.2.8.* We illustrate the theorem with the *Clebsch-Gordan* theory of existence of invariants in tensor products of representations of  $G = SL(2, \mathbb{C})$ . The weight lattice  $\Lambda^\vee$  for  $G$  is naturally identified with the set  $\mathbb{Z}/2$  of non-negative half-integers and for any  $\lambda \in \Lambda^\vee, \lambda \geq 0$  we denote by  $V_\lambda$  the corresponding simple  $G$ -module. Given  $\lambda_1, \dots, \lambda_n$  we ask whether  $V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n}$  contains an invariant vector. Now  $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d)) \cong V_{d/2}$  and so  $R(\mathbb{P}^1) = \bigoplus_\lambda V_\lambda$ . If we equip  $X = (\mathbb{P}^1)^n$  with the ample line bundle  $\mathcal{O}_X(1) := \bigotimes_{j=1}^n \mathcal{O}_{\mathbb{P}^1}(\lambda_j)$  then

$$R(X) = \bigoplus_{d \geq 0} \bigotimes_{j=1}^n H^0(\mathcal{O}_{\mathbb{P}^1}(d\lambda_j)) = \bigoplus_{d \geq 0} \bigotimes_{j=1}^n V_{d\lambda_j}.$$

So

$$R(X//G) = R(X)^G = \left( \bigoplus_{d \geq 0} \bigotimes_{j=1}^n V_{d\lambda_j} \right)^G.$$

This is non-zero if and only if  $X//G$  is empty. The Kempf-Ness Theorem 5.2.1 gives  $X//G \cong X//K \cong (S_{\lambda_1}^2 \times \dots \times S_{\lambda_n}^2)//SU(2)$  where  $S_\lambda^2$  denotes the two-sphere equipped with re-scaled symplectic form  $\lambda$  and  $SU(2)$  acts via the double cover  $SU(2) \rightarrow SO(3)$ . By Proposition 3.3.4,

*Corollary 5.2.9.*  $(\bigotimes_{j=1}^n V_{d\lambda_j})^G$  is non-trivial for some  $d$  iff

$$(5.3) \quad \lambda_j \leq \sum_{i \neq j} \lambda_i, j = 1, \dots, n.$$

This gives a geometric proof of the well-known Clebsch-Gordan rules. A basis for the space of invariants is induced from a choice of parenthesization of the tensor product above, see for example [25]. The relation between the different invariants is also connected to symplectic geometry [92].

**5.3. Quantization commutes with reduction.** The proof of the Kempf-Ness Theorem 5.2.1, which seems otherwise somewhat miraculous, has a conceptual interpretation given by Guillemin-Sternberg [38] in terms of geometric quantization (Section 3.5) as follows. Namely, rather than choosing a lift of  $x \in X$  to  $V - \{0\}$ , which is the total space of  $\mathcal{O}_X(-1)$ , it is more natural from the viewpoint of geometric quantization to choose a lift  $l$  in the positive line bundle  $\mathcal{O}_X(1) \rightarrow X$ . Define the *Guillemin-Sternberg stability function*

$$\psi_l^\vee : K \backslash G \rightarrow \mathbb{R}, \quad g \mapsto \log \|gl\|^2/2.$$

The same computation as in the Kempf-Ness case, except for a change of sign, implies that the gradient of  $\psi_l^\vee$  is minus the moment map, and  $\psi_l$  is concave. In particular, suppose that  $s \in H^0(X, \mathcal{O}_X(1))^G$  is an invariant section. Then

$$\psi_{s(x)}^\vee([g]) = \log \|gs(x)\|^2/2 = \log \|s(gx)\|^2/2.$$

Now concavity of  $\psi_{s(x)}^\vee$  implies that any critical point of  $\|s\|^2$  occurs at  $\Phi^{-1}(0)$  and is a local maximum, and  $s$  is approximately Gaussian. This type of behavior is quite standard for “typical quantum states”, which introductory physics lectures often show as concentrating near some submanifold of the corresponding classical state space in Gaussian fashion.

Suppose that  $K$  acts freely on the zero level set  $\Phi^{-1}(0)$ . The complex structure  $J$  on  $X$  induces an almost complex structure  $J//K$  on  $X//K$  by identifying  $\pi^*T(X//K)$  with the subbundle of  $TX|_{\Phi^{-1}(0)}$  perpendicular to the generating vector fields  $\xi_X, \xi \in \mathfrak{k}$ . This complex structure is integrable since the Nijenhuis tensor vanishes. Similarly the polarization  $\mathcal{O}_X(1) \rightarrow X$  naturally descends to a polarization  $\mathcal{O}_{X//K}(1) \rightarrow X//K$ , defined by restricting to  $\Phi^{-1}(0)$  and quotienting by the action of  $K$ .

**Theorem 5.3.1** (Quantization commutes with reduction). *Let  $X$  be a compact Hamiltonian  $K$ -manifold equipped with moment map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ , polarization  $\mathcal{O}_X(1) \rightarrow X$  and a compatible  $K$ -invariant Kähler structure  $J$ , such that  $K$  acts freely on the zero level set  $\Phi^{-1}(0)$ , and let*

$R(X)_d$  denote the space of sections of  $\mathcal{O}_X(d)$  as above. For each  $d \geq 0$  there is a canonical isomorphism  $\rho : R(X)_d^K \rightarrow R(X//K)_d$

*Proof.* For smooth projective varieties  $X \subset \mathbb{P}(V)$  this is a combination of Mumford’s Theorem 4.2.6 and the Kempf-Ness Theorem 5.2.1. More generally let  $X$  be a compact polarized Kähler Hamiltonian  $K$ -manifold. Any section  $s \in H^0(X, \mathcal{O}_X(1))^K$  naturally defines a section  $\rho(s) \in H^0(X//K, \mathcal{O}_{X//K}(1))$  by restriction to  $\Phi^{-1}(0)$  and descent to the quotient. Then  $\rho$  is an injection, since any invariant section has maximum norm on  $\Phi^{-1}(0)$ . Proving surjectivity required a somewhat complicated argument in the approach of Guillemin-Sternberg, and the following alternative algebraic argument is substantially easier: By Kodaira embedding  $X$  is biholomorphic to smooth subvariety of  $\mathbb{P}(V)$ , and the polarization  $\mathcal{O}_X(1)$  is isomorphic as a holomorphic line bundle to the pull-back of the hyperplane bundle on  $\mathbb{P}(V)$ , although the symplectic structure and moment map may not be pull-backs. By the extension of Kempf-Ness to Kähler varieties discussed in 5.2.7, the semistable locus corresponding to the polarization  $\mathcal{O}_X(1)$  has quotient by  $G$  diffeomorphic to  $X//K$ . Given a section  $s \in H^0(X//K, \mathcal{O}_{X//K}(1))$ ,  $s$  naturally lifts to an invariant section on the semistable locus  $X^{\text{ss}}$  with maximum on  $\Phi^{-1}(0)$ . Since the norm of this section is bounded, it extends over all of  $X$ .  $\square$

Guillemin-Sternberg also proved “quantization commutes with reduction” for another of class of Hamiltonian actions for which there exists a good quantization scheme, namely cotangent bundles [39]. Quantization commutes with reduction was generalized to arbitrary compact Hamiltonian manifolds using “Spin-c” quantization by Meinrenken [80], and further generalized to “non-abelian localization” by Teleman and Paradan, see the last section of these notes.

**5.4. Convex functions on  $K \backslash G$ .** In this section we further investigate the geometry of the Kempf-Ness functional, mostly following Kapovich-Leeb-Millson [59, Section 3.1]. The discussion uses some geometry of  $K \backslash G$  for which the reader may refer to [4]. We already mentioned that the quotient  $K \backslash G$  is a *Hadamard manifold*, that is, a space of non-positive sectional curvature. Such a manifold has a natural compactification by adding a *boundary at infinity*  $\partial_\infty(K \backslash G)$ , given by equivalence classes of unit-speed geodesics where two unit-speed geodesics are equivalent if they have bounded distance. This boundary is naturally equipped with a *Tits metric* defined as the supremum of angles of formed by a pair of geodesics approaching the given points at infinity. The topology defined by the Tits metric is discrete iff  $K \backslash G$  is higher rank, and defines the standard topology on  $(T \cap K) \backslash T \subset K \backslash G$  for any complex maximal torus  $T$ .

Let  $\psi : K \backslash G \rightarrow \mathbb{R}$  be a Lipschitz continuous convex function. The *slope at infinity* of  $\psi : K \backslash G \rightarrow \mathbb{R}$  is the function

$$\mu : \partial_\infty(K \backslash G) \rightarrow \mathbb{R}, \xi \mapsto \lim_{t \rightarrow \infty} \frac{\psi(\rho(t))}{t}$$

where  $\rho$  is any geodesic ray asymptotic to  $\xi$ . By [59, Lemma 3.2],  $\mu$  is Lipschitz continuous on  $\partial_\infty(K \backslash G)$  with respect to the Tits metric with the same Lipschitz constant. The boundary  $\partial_\infty(K \backslash G)$  has curvature bounded by 1; one says that a function on  $\partial_\infty(K \backslash G)$  is convex if it is convex along any geodesic of length at most  $\pi$ .

**Definition 5.4.1.** Let  $C_{<0}(x)$  resp.  $C_{\leq 0}(x)$  resp.  $C_0(x)$  denote the subset of  $\partial_\infty(K \backslash G)$  with negative resp. non-positive resp. zero slope.

**Theorem 5.4.2.** (a)  $C_{\leq 0}(x)$  is convex, and the function  $\mu$  is convex on  $C_{\leq 0}(x)$  and strictly convex on  $C_{<0}(x)$ .

(b)  $\psi$  is proper and bounded below iff  $\mu > 0$  everywhere on  $\partial_\infty(K \backslash G)$ .

(c) If  $C_{<0}(x) \neq \emptyset$ , then

(i)  $\mu$  has a unique minimum  $\mu(\xi_{\min})$ ,

(ii)  $C_{\leq 0}(x)$  is the closure of  $C_{<0}(x)$ , and

(iii) any gradient trajectory of  $\psi$  has asymptotic direction  $\xi_{\min}$  and asymptotic slope  $\mu(\xi_{\min})$ .

(d) If  $C_0(x)$  is open, then  $\mu \geq 0$  everywhere.

*Proof.* Except for the assertion about gradient trajectories, this is Eberlein [32, 4.1.1’] and Kapovich-Leeb-Millson [59, Section 3.1], and is essentially a consequence of convexity of  $\psi$ . Indeed convexity of  $\psi$  implies that if  $\xi_1 \in \partial_\infty(K \backslash G)$  is the midpoint of a geodesic segment connecting

$\xi_0, \xi_2 \in \partial_\infty(K \setminus G)$  with angles less than  $\pi$  then

$$\mu(\xi_1) \leq \frac{\mu(\xi_0) + \mu(\xi_2)}{2 \cos(d(\xi_0, \xi_2)/2)}$$

and most of the claims follow from this inequality. Angles of  $\pi$  must be dealt with separately; in particular, for example, in the case  $G = SL(2, \mathbb{C})$  the Tits metric on  $G/K$  assigns distance  $\pi$  to every pair of distinct points and so the above argument is not particularly helpful.

Suppose that  $\psi$  has a direction of negative slope. By Lemma 5.4.3 below,  $\|\text{grad}(\psi)\|$  is bounded below by a positive constant. By Caprace-Lytchak [24, Proposition 4.2] all gradient trajectories converge to the same point at infinity and at the same rate of escape. (For the special case of a Kempf-Ness function the existence of a limiting direction follows from Duistermaat's result Lemma 7.1.5 and Corollary 5.2.3.) Hence there exists  $\xi_\infty \in \mathfrak{k}$  such that any gradient trajectory  $[g_t]$  of  $-\psi$  has limiting direction  $\xi_\infty$ , that is,

$$\left(\frac{d}{dt}[g_t]\right)g_t^{-1} \rightarrow \frac{d}{dt}\Big|_{t=0}[\exp(-it\xi_\infty)]$$

for some  $\xi_\infty \in \mathfrak{k}$ , then

$$(5.4) \quad \lim_{t \rightarrow \infty} [g_t] = \lim_{t \rightarrow \infty} [\exp(i\xi_\infty t)] \in \partial_\infty(K \setminus G)$$

see Kaimanovich [58, Theorem 2.1]. (In fact Chen-Sun [26] show that, in the Kempf-Ness setting, any gradient trajectory is asymptotic to a geodesic ray.) It follows from Lipschitz continuity of  $\psi$  that the rate of decay of  $\psi$  along  $[g_t]$  is the same as that along  $[\exp(ti\xi_\infty)]$ , so that  $\mu(\xi_{\min}) \leq -\|\xi_\infty\|$ . If  $\xi_\infty \neq \xi_{\min}\|\xi_\infty\|$  then one obtains a contradiction by connecting  $[g_t]$  to  $[\exp(-it\xi_{\min}\|\xi_\infty\|)]$  by a geodesic  $[\exp(is\zeta_t)g_t]$ ,  $s \in [0, 1]$  and using convexity of  $\psi$ : Since  $\psi([\exp(is\zeta_t)g_t])$  goes to  $-\infty$  at least as fast for  $s = 1$  as for  $s = 0$  as  $t \rightarrow \infty$ ,

$$\frac{d}{ds}\Big|_{s=0}\psi([\exp(is\zeta_t)g_t]) < \epsilon, \forall \epsilon > 0, t \gg 0.$$

Now  $\text{grad}(\psi) \rightarrow \xi_\infty$  implies that  $\frac{d}{ds}\Big|_{s=0}\psi([\exp(is\zeta_t)g_t])$  is approximately  $(\xi_\infty, \zeta_t)$  for  $t \gg 0$ . On the other hand, by angle comparison the angle formed by  $\frac{d}{dt}\Big|_{t=0}[g_t]$  and  $\frac{d}{ds}\Big|_{s=0}\psi([\exp(is\zeta_t)g_t])$  is bounded from below by  $\pi/2 + d(\xi_{\min}, \xi_\infty/\|\xi_\infty\|)/2$  for  $t \gg 0$ . This implies that  $(\xi_\infty, \zeta_t) > 0$ , which is a contradiction.  $\square$

It remains to show

**Lemma 5.4.3.** *Let  $\psi, \mu, \xi_{\min}$  be as in Theorem 5.4.2. Then  $\mu(\xi_{\min}) \geq \inf_{[g] \in K \setminus G} \|\text{grad}(\psi)([g])\|$ .*

*Proof.* Convexity of  $\psi$  along  $[\exp(it\xi)g]$  implies that for  $t \geq 0$ ,

$$\begin{aligned} \frac{d}{dt}\psi([\exp(it\xi)g]) &\geq \frac{d}{dt}\Big|_{t=0}\psi([\exp(it\xi)g]) \\ &= (\xi, \text{grad}(\psi)([g])) \geq -\|\text{grad}(\psi)([g])\|. \end{aligned}$$

Taking the infimum over  $[g] \in K \setminus G$  and  $\xi \in \mathfrak{k}$  of unit norm gives the result.  $\square$

*Remark 5.4.4.* The direction of maximal descent in Theorem 5.4.2 (c) is not necessarily rational. However, if  $\mu$  is negative somewhere then it is negative on some rational vector, since the Tits metric is the standard one on  $(T \cap K) \setminus T$  for any maximal torus  $T$  and rational directions are dense in  $\mathfrak{k}$ . That is, non-negativity of  $\mu$  is equivalent to non-negativity of  $\mu$  on the rational vectors, i.e., those generating one-parameter subgroups.

**5.5. Polystable points.** By Lemma 5.2.6, the polystable orbits are the orbits of points  $x \in \Phi^{-1}(0)$ . In this section we investigate these and the orbit-closure equivalence relation in more detail. The following was observed by Kempf-Ness [60] in the linear case and by Slodowy [101] in general, see also Sjamaar [99].

**Proposition 5.5.1.** *Let  $X$  be a Kähler Hamiltonian  $K$ -manifold, and  $x \in \Phi^{-1}(0)$ . Then  $G_x$  is the complexification of  $K_x$ ; in particular,  $G_x$  is reductive.*

*Proof.* Suppose that  $x \in \Phi^{-1}(0)$  and  $gx = x$ . Write  $g = k \exp(i\xi)$  for some  $\xi \in \mathfrak{k}, k \in K$ . Let  $\psi_x$  be a Kempf-Ness function for  $x$ . Since  $x, gx \in \Phi^{-1}(0)$ , we have

$$\text{grad } \psi_x([k \exp(i\xi)]) = \text{grad } \psi_x([\exp(i\xi)]) = \text{grad } \psi_x([e]) = 0.$$

By convexity,  $\psi_x$  is constant along the geodesic  $[\exp(it\xi)]$ , so  $\xi \in i\mathfrak{k}_x$  by Corollary 5.2.4. Hence  $x = kx$  so  $k \in K_x$ , which implies  $g \in (K_x)_{\mathbb{C}}$ . The reverse inclusion  $(K_x)_{\mathbb{C}} \subset G_x$  is obvious.  $\square$

*Remark 5.5.2.* Stabilizer groups are not in general reductive. For example let  $X = SL(2, \mathbb{C}) \times_B \mathbb{P}^1$ . Then every stabilizer is either solvable or unipotent, and so no projective embedding of  $X$  has semistable points.

Second we show that polystable points are “seen by one-parameter subgroups.” For this we need to review some results on existence of holomorphic slices. Let  $X$  be a complex manifold with a holomorphic action of a group  $G$ . Let  $x \in X$ . Recall that a *slice* at  $x$  is a  $G_x$ -invariant submanifold  $S$  of  $X$  containing  $x$  such that  $GS$  is open in  $X$  and the natural  $G$ -equivariant map from  $G \times_{G_x} S \rightarrow X$  is an isomorphism onto  $GS$ . Sjamaar [99] has proved the following analog of slice theorems of Luna [76] and Snow:

**Theorem 5.5.3** (Sjamaar). *Let  $G$  be a connected complex reductive group with maximal compact  $K$ . Let  $X$  be a Kähler Hamiltonian  $K$ -manifold such that the action of  $K$  extends to a holomorphic action of  $G$ . Suppose that  $x \in \Phi^{-1}(0)$ . Then there exists a slice at  $x$ .*

**Corollary 5.5.4.** *An orbit  $Gx$  contains a polystable point  $y$  in its closure, iff there exists a one-parameter subgroup  $\mathbb{C}^* \subset G$  and a point  $z \in Gx$  such that  $\mathbb{C}^*z$  contains a polystable point in its closure.*

*Proof.* Let  $y$  be a polystable point. We may assume that  $\Phi(y) = 0$ . By Theorem 5.5.3, there exists a slice  $S$  at  $y$ . Now  $S$  is biholomorphic to its tangent space  $T_y S$ , equivariantly for the action of  $K_x$ , in a neighborhood  $U$  of  $y$ . Furthermore, since this map is holomorphic, the map is equivariant for the *infinitesimal*  $G$ -action. By Lemma 4.3.4, there exists a one-parameter subgroup  $\mathbb{C}^* \rightarrow G$  and a point  $v \in T_y S$  such that the closure of  $\mathbb{C}^*v$  contains  $0 \in T_y S$ . By choosing  $v$  sufficiently small, we ensure that  $\{zv, |z| \leq 1\}$  is in the image of  $U$ . Let  $s \in S$  be the pre-image of  $v$ . Then  $\{zs, |z| \leq 1\}$  contains  $y$  in its closure, as required.  $\square$

Using this corollary we prove a finite-dimensional analog of the *Jordan-Hölder* theory for semistable vector bundles, see for example Seshadri [96].

**Definition 5.5.5.** For any  $\lambda \in \mathfrak{k}$ , let  $x_\lambda = \lim_{t \rightarrow \infty} \exp(-ti\lambda)x$  the *associated graded point* of  $x$  with respect to  $\lambda$ .

*Remark 5.5.6.* The fact that  $\exp(-ti\lambda)x$  is the gradient flow of a Morse function (see 3.2.7) implies that the gradient trajectory converges *exponentially fast* to  $x_\lambda$ , that is,  $\text{dist}(\exp(-ti\lambda)x, x_\lambda) \leq C_0 e^{-C_1 t}$  for some constants  $C_0, C_1$ .

**Definition 5.5.7.**  $\lambda \in \mathfrak{k}$  is *Jordan-Hölder* for  $x \in X^{\text{ss}}$  iff  $x_\lambda$  is polystable.

*Example 5.5.8.* Let  $X = \mathbb{C}^2$  and  $G = (\mathbb{C}^*)^2$  acting by  $(g_1, g_2)(z_1, z_2) = (g_1 z_1, g_2 z_2)$ . Then any  $(\lambda_1, \lambda_2)$  with  $\lambda_1, \lambda_2 > 0$  is Jordan-Hölder.

**Theorem 5.5.9.** *Let  $X$  be a compact Kähler Hamiltonian  $K$ -manifold and  $x \in X$  a semistable point.*

- (a) *If  $x$  is semistable but not polystable then the set of Jordan-Hölder vectors for  $x$  is a non-empty  $K_x$ -invariant cone in  $\mathfrak{k}$ .*
- (b) *The orbit  $Gx_\lambda$  of the associated graded  $x_\lambda$  of a Jordan-Hölder  $\lambda$  is the unique polystable orbit in  $Gx$ .*

*Proof.* (a) Suppose  $x$  is semistable but not polystable. Since  $x$  is semistable,  $Gx$  contains a polystable  $y$  in its closure. By Corollary 5.5.4, any polystable  $y$  is in the closure  $\mathbb{C}^*z$  for some one-parameter subgroup  $\mathbb{C}^* \subset G$  and  $z \in Gx$ . Suppose that  $z = g^{-1}x$  for some  $g \in G$ . Then  $(\text{Ad}(g)\mathbb{C}^*)x = g\mathbb{C}^*z$  contains  $gy$  in its closure, and  $gy$  is polystable as well. Convexity of the set of Jordan-Hölder vectors follows from Theorem 5.4.2 applied to a Kempf-Ness function  $\psi : K \backslash G \rightarrow \mathbb{R}$ . Indeed, by compactness of  $X$  the norm  $\|\Phi\|$  is bounded, so by Corollary 5.2.3  $\psi$  is Lipschitz

continuous. Furthermore, by Lemma 5.2.4,  $\psi$  is convex. Hence Theorem 5.4.2 applies. (b) Suppose that  $y_0, y_1$  are polystable points in the closure of  $Gx$ . By Corollary 5.5.4, there exist vectors  $\lambda_j, j = 0, 1$  and points  $x_0, x_1 \in Gx$  such that  $y_j = (x_j)_{\lambda_j}$ . The distance between  $\exp(-it\lambda_j)x_j$  can be estimated as follows: Suppose that  $x_j = g_jx$  and let  $h_{s,t} = [\exp(i\delta_{s,t}s) \exp(i\lambda_0t)g_0]$  so that  $[h_{s,t}]$  is the geodesic path in  $K \setminus G$  between  $[\exp(-it\lambda_j)g_j], j = 0, 1$ . Let  $x_{s,t} = h_{s,t}x$ . The square of the distance from  $x_{0,t}$  to  $x_{1,t}$  is given by

$$\begin{aligned} \left( \int_0^1 \|\partial_s x_{s,t}\| ds \right)^2 &\leq \int_0^1 \|\partial_s x_{s,t}\|^2 ds = \int_0^1 g(\partial_s x_{s,t}, \partial_s x_{s,t}) ds \\ &= \int_0^1 \partial_s^2 \psi([h_{s,t}]) ds = \partial_s \psi([h_{s,t}])|_{s=0}^{s=1}. \end{aligned}$$

Now  $\text{grad } \psi$  converges exponentially to zero along  $[\exp(-it\lambda_j)]$  as  $t \rightarrow \infty$  for  $j = 0, 1$ , since  $\exp(-it\lambda_j)x_j$  converges exponentially fast to  $(x_j)_{\lambda_j}$ , see Remark 5.5.6. On the other hand, since  $h_{s,t}, \exp(-it\lambda_0)g_0, \exp(-it\lambda_1)g_1$  are the sides of a geodesic quadrangle with one side of fixed length, there exist constants  $C_0, C_1$  such that

$$\|\delta_t\| < C_0 + tC_1\|\lambda_0 + \lambda_1\|$$

for all  $t$ . Hence

$$\text{dist}(x_{\lambda_0}, x_{\lambda_1}) = \lim_{t \rightarrow \infty} \text{dist}(x_{0,t}, x_{1,t}) = 0$$

and the claim follows.  $\square$

*Remark 5.5.10.* We have included (b) to emphasize a somewhat confusing point: distant points in  $\mathfrak{k}$  may map to near points in  $X$  if the gradient of  $\psi$  on the path between them is sufficiently small.

*Remark 5.5.11.* In fact, the full strength of Sjamaar's (or Luna's) slice theorem is not needed here; it suffices to find a slice for the *infinitesimal* action of  $G$  which is substantially easier. Some terminology: If a Lie group with Lie algebra  $\mathfrak{g}$  acts on a manifold we say that a submanifold  $U$  is  $\mathfrak{g}$ -invariant if the generating vector fields are tangent to  $U$ . A *slice* for the infinitesimal action of  $\mathfrak{g}$  at  $x$  is a  $\mathfrak{g}_x$ -invariant holomorphic submanifold  $S$  containing  $x$ , such that the natural map  $\mathfrak{g} \times_{\mathfrak{g}_x} TS \rightarrow TX|_S$  is an isomorphism. Using the implicit function theorem, one sees that any sequence of points converging to  $x$  may be translated by the action of  $G$  (which is now only defined in a neighborhood of the identity) into a sequence of points in  $S$ . Thus if an orbit  $Gy$  in  $X$  contains  $x \in S$  in its closure, then  $Gy \cap S$  also contains  $x$  in its closure, and by Lemma 4.3.4  $\mathbb{C}^*y \cap S$  contains  $x$  in its closure for some one-parameter subgroup  $\mathbb{C}^* \subset G$ .

## 6. SCHUR-HORN CONVEXITY AND ITS GENERALIZATIONS

In this section we discuss the generalization of Clebsch-Gordan theory to arbitrary groups, in particular, the theory of existence of invariants in tensor products of representations of  $GL(r)$ , the connections (via the Kempf-Ness theorem) with eigenvalue problems, and a combinatorial answer by Knutson, Tao, and the author [68].

**6.1. The Borel-Weil theorem.** Let  $G$  be a connected complex reductive group. Let  $\lambda$  be any dominant weight for  $G$  and  $V_\lambda$  a simple  $G$ -module with highest weight  $\lambda$ . Let  $P_\lambda^-$  be the opposite standard parabolic corresponding to  $\lambda$ , and  $G/P_\lambda^-$  the generalized flag variety corresponding to  $\lambda$ . We denote by  $\mathbb{C}_\lambda^\vee$  the one-dimensional representation of  $P_\lambda^-$  corresponding to  $-\lambda$ , and by  $\mathcal{O}_X(\lambda) = G \times_{P_\lambda^-} \mathbb{C}_\lambda^\vee$ .

**Theorem 6.1.1** (Borel-Weil [95]). *Let  $X = G/P_\lambda^-$  with  $\lambda$  a weight. Then  $H^0(X, \mathcal{O}_X(\lambda)) \cong V_\lambda$  if  $\lambda$  is dominant and vanishes otherwise.*

*Proof.* First consider the case  $G = SL(2, \mathbb{C})$ . We identify  $\Lambda^\vee$  with  $\mathbb{Z}/2$ . Then  $H^0(\mathcal{O}_X(\lambda))$  is the set of homogeneous polynomials in two variables of degree  $2\lambda$ , if  $\lambda$  is non-negative, and zero otherwise. In the first case one checks easily that  $H^0(\mathcal{O}_X(\lambda))$  is simple with highest weight  $\lambda$ .

Next let  $G$  be an arbitrary connected complex reductive group. Let  $X = G/B^-$  and  $X_1 = BB^-/B^- \cong B/T \cong U$  the open Bruhat cell, (here  $U$  is a maximal unipotent) so that

$$H^0(X_1, \mathcal{O}_X(\lambda)|_{X_1})^U = H^0(U, \mathbb{C})^U \cong \mathbb{C}.$$

Thus  $H^0(X_1, \mathcal{O}_X(\lambda)|_{X_1})$  contains a unique highest weight vector, which we denote by  $s_\lambda$ . We wish to determine whether  $s_\lambda$  extends over the complement of  $X_1$  in  $X$ . It suffices to check the order of vanishing of  $s_\lambda$  on the divisors  $X_{s_\alpha}$ , as  $\alpha$  ranges over simple roots. For each root  $\alpha$ , we let  $h_\alpha \in \mathfrak{t}$  denote the corresponding coroot, so that  $\mathfrak{sl}(2, \mathbb{C})_\alpha := \mathbb{C}h_\alpha \oplus \mathfrak{g}_\alpha$  is the three-parameter Lie algebra corresponding to  $\alpha$ . Let  $SL(2, \mathbb{C})_\alpha \rightarrow G$  denote the homomorphism induced by the inclusion  $\mathfrak{sl}(2, \mathbb{C})_\alpha \rightarrow \mathfrak{g}$ . The orbit  $C_\alpha = SL(2, \mathbb{C})_\alpha B^- / B^-$  of  $SL(2, \mathbb{C})_\alpha$  on  $X$  is isomorphic to  $SL(2, \mathbb{C})_\alpha / SL(2, \mathbb{C})_\alpha \cap B^- \cong \mathbb{P}^1$ . The curve  $C_\alpha$  intersects the Bruhat cell  $X_{s_\alpha}$  in the unique point  $x_{s_\alpha} = s_\alpha B^- / B^-$ . The order of vanishing of  $s_\alpha$  along  $X_{s_\alpha}$  is necessarily the order of vanishing of  $s_\alpha|_{C_\alpha}$  at  $x_{s_\alpha}$ . Now  $\mathcal{O}_X(\lambda)$  restricts to the line bundle  $\mathcal{O}_{\mathbb{P}^1}(\langle \lambda, h_\alpha \rangle)$  on  $C_\alpha$ , and the section  $s_\lambda$  restricts to the highest weight section on  $C_\alpha - x_{s_\alpha}$ . It extends over  $x_\alpha$  iff  $\langle \lambda, h_\alpha \rangle \geq 0$ , by the discussion for the  $SL(2, \mathbb{C})$  case.

Now  $G/B^-$  fibers over  $G/P_\lambda^-$  with projective fibers and so

$$H^0(G/B^-, \mathcal{O}_{G/B^-}(\lambda)) = H^0(G/P_\lambda^-, \mathcal{O}_{G/P_\lambda^-}(\lambda)).$$

Since the result is proved for  $G/B^-$ , this completes the proof.  $\square$

From the point of view of symplectic geometry, the Borel-Weil theorem says that the geometric quantization of a coadjoint orbit equipped with an integral symplectic form (that is, one that is the curvature of some line bundle) is a simple  $K$ -module. Indeed, let  $\Phi$  denote the moment map induced by the action of  $K$  on  $\mathcal{O}_X(\lambda)$ . Since the weight of  $T$  on the fiber of  $\mathcal{O}_X(\lambda)$  over  $B^-/B^-$  is  $-\lambda$ ,  $\Phi$  maps  $X$  onto the coadjoint orbit  $K\lambda$  through  $\lambda$ , see Proposition 3.2.10. Thus in the notation introduced in Section 3.5,  $\mathcal{H}(K\lambda) = V_\lambda$ .

**6.2. The Schur-Horn-Kostant problem.** The Schur-Horn theorem [94], [54] reads:

**Theorem 6.2.1.** *The set of possible diagonal entries of a Hermitian operator with eigenvalues  $\lambda = (\lambda_1, \dots, \lambda_n)$  is the hull of the set of permutations of  $\lambda$ .*

*Example 6.2.2.* If  $K = SO(3)$  then by Proposition 3.2.2 the coadjoint orbit through  $\text{diag}(\lambda, -\lambda)$  may be identified with the sphere of radius  $\lambda$  via the isomorphism  $\mathfrak{k}^\vee = \mathfrak{so}(3)^\vee \rightarrow \mathbb{R}^3$ , and the moment map for the maximal torus action is projection onto the  $z$ -axis, and so has moment image  $[-\lambda, \lambda]$ . The action of the Weyl group  $W = \mathbb{Z}_2$  on  $\mathfrak{t}$  is identified with the sign representation, and so  $[-\lambda, \lambda] = \text{hull}\{-\lambda, \lambda\} = \text{hull}(W\lambda)$  as claimed.

Kostant [71] generalized this result to arbitrary compact connected groups:

**Theorem 6.2.3.** *Let  $K$  be a compact connected group. The projection of a coadjoint orbit  $K\lambda$  of an element  $\lambda \in \mathfrak{t}^\vee$  is the convex hull of the orbit  $W\lambda$  of  $\lambda$  under the Weyl group  $W$ .*

Using the Kempf-Ness and Borel-Weil theorems 5.2.1, 6.1.1, the Schur-Horn-Kostant theorem is equivalent to the following well-known fact in representation theory:

**Theorem 6.2.4.** *With  $K$  as above, let  $\lambda$  be a dominant weight. The set of  $\mu/d$  such that the weight space  $V_{d\lambda, (\mu)} \subset V_{d\lambda}$  is non-trivial for some  $d \in \mathbb{Z}_+$  is the rational convex hull of  $W\lambda$ .*

*Proof.* We identify  $X = K\lambda = G/P_\lambda^-$  and  $\mathbb{C}_\mu$  the trivial bundle over  $X$  with  $T$ -weight  $\mu$  so that  $V_{d\lambda, (\mu)} = H^0(X, \mathbb{C}_\mu^* \otimes \mathcal{O}_X(d\lambda))^T$  by Borel-Weil 6.1.1, which is the space of sections over the quotient  $(X//T)_\mathbb{C}$  by Mumford's Theorem 4.2.6. We may use the Hilbert-Mumford criterion to determine whether there are any semistable points: Given a one-parameter subgroup generated by some  $\xi \in \mathfrak{t}_+$ , a point  $x \in X$  flows under  $\exp(t\xi)$  to  $y_w$  as  $t \rightarrow -\infty$  where  $x \in Y_w := B^- w B^- / B^-$  is the opposite Bruhat cell, see (4.1). The weight of  $T$  on the fiber over  $y_w$  is  $\mu - w\lambda$ . Thus  $x \in Y_w$  is semistable for  $\xi$  iff  $\langle w\lambda - \mu, \xi \rangle \leq 0$  iff  $\mu \in w\lambda - \mathfrak{t}_+^\vee$ . In particular  $Y_1$  is contained in the semistable locus for the one-parameter subgroup generated by  $-\xi$  with  $\xi$  dominant iff  $\mu \in \lambda - (\mathfrak{t}_+^\vee)$ . The semistable locus for the torus action is non-empty iff a generic point is semistable for all one-parameter subgroups iff

$$(6.1) \quad \mu \in \bigcap_{w \in W} w(\lambda - (\mathfrak{t}_+^\vee)).$$

The dual cone to  $\text{hull}(w\lambda, w \in W)$  at  $w\lambda$  is generated by  $(s_\alpha - 1)w\lambda$  where  $\alpha$  ranges over simple roots, which is equal to  $w(\mathfrak{t}_+^\vee)$ . It follows that (6.1) is equivalent to  $\mu \in \text{hull}(w\lambda, w \in W)$  as claimed.  $\square$



*Proof of Theorem 6.2.3.* Let  $X = K\lambda$  be as above. The moment map corresponding to the projective embedding  $K\lambda \rightarrow \mathbb{P}(V_\lambda^\vee)$  is the projection  $\pi$  of  $X$  onto  $\mathfrak{t}^\vee$  by Proposition 3.2.3 (d). Hence the moment map for the projective embedding  $K\lambda \rightarrow \mathbb{P}(V_\lambda^\vee \otimes \mathbb{C}_\mu)$  is  $\pi - \mu$ . By Kempf-Ness  $X//T_{\mathbb{C}} \cong X//T$ , where  $T_{\mathbb{C}}$  is the complexification of  $T$ . Finally  $X//T$  non-trivial iff 0 is in the image of  $\pi - \mu$  iff  $\mu$  is contained in the image of  $\pi$ .  $\square$

**6.3. The Horn-Klyachko problem.** In the previous section we investigated the existence of semistable points for an action of a torus. Horn [55] deals with the following question, which we will rephrase in terms of existence of semistable points for the action of a non-abelian group:

**Question 6.3.1.** *Given the eigenvalues of Hermitian matrices  $H_1, \dots, H_{n-1}$ , what are the possible eigenvalues of  $H_1 + \dots + H_{n-1}$ ?*

Since the eigenvalues are real, we may order them in non-increasing order

$$\lambda_1(H_j) \geq \lambda_2(H_j) \dots \geq \lambda_r(H_j).$$

The eigenvalues must satisfy the trace equality

$$\sum_{i,j} \lambda_i(H_j) = \sum_i \lambda_i(H_1 + \dots + H_{n-1}).$$

After that there are a finite set of linear inequalities, for example the well-known

$$\lambda_1(H_1 + H_2) \leq \lambda_1(H_1) + \lambda_1(H_2).$$

We will describe the complete list. Before we give the answer, we note that this question has a symplectic reformulation as follows. Taking  $H_n = -H_1 - \dots - H_{n-1}$ , obtain a tuple  $(H_1, \dots, H_n)$  with  $H_1 + \dots + H_n = 0$ . Thus the problem is a special case of the *generalized Horn problem*:

**Question 6.3.2.** *Let  $K$  be a compact Lie group. For which  $\lambda_1, \dots, \lambda_n \in \mathfrak{t}_+^\vee$  is the symplectic quotient  $(K\lambda_1 \times \dots \times K\lambda_n)//K$  non-empty?*

By the Kempf-Ness and Borel-Weil theorems, this problem is equivalent to the following

**Question 6.3.3.** *Let  $K$  be a compact Lie group. For which dominant weights  $\lambda_1, \dots, \lambda_n \in \mathfrak{t}_+^\vee$  is space of invariants  $(V_{d\lambda_1} \otimes \dots \otimes V_{d\lambda_n})^K$  non-trivial for some  $d \geq 0$ ?*

In the case  $K = SU(2)$  this question was answered in Section 5.2. The connection between Questions 6.3.2 and 6.3.3 was investigated in the more general setting of projections of coadjoint orbits by Heckman [48]:

**Theorem 6.3.4.** *Suppose that  $L$  is a compact connected group containing a compact connected subgroup  $K$ . The projection of any  $L$ -coadjoint orbit  $L\mu \subset \mathfrak{l}^\vee$  onto  $\mathfrak{k}^\vee$  intersects  $\mathfrak{t}_+^\vee$  in a convex polytope.*

In the case  $L = K^{n-1}$  containing  $K$  via the diagonal embedding, the projection  $(\mathfrak{k}^\vee)^{n-1} \rightarrow \mathfrak{k}^\vee$  is the sum map and the Theorem 6.3.4 implies that the for any given  $\lambda_1, \dots, \lambda_{n-1}$ , the set of possible  $\lambda_n$  in 6.3.2 is a convex polytope.

Next we give a partial answer for which inequalities occur in 6.3.2 in the case  $K = SU(n)$  using max-min description of eigenvalues; this implies inequalities on the invariant theory problem. Then we give a necessary and sufficient answer using the Hilbert-Mumford criterion, following an argument of Klyachko [63]. Finally we give a brief description of works of Belkale [12], Knutson-Tao [66], and Knutson-Tao-Woodward [68] giving a *minimal set* of inequalities. Generalizations to groups of arbitrary type and other actions are described in Berenstein-Sjamaar [14], Kapovich-Leeb-Millson [59] and Ressayre [91].

We begin with the elementary max-min approach for  $K = U(n)$ . If  $H$  is a Hermitian matrix with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r$  then

$$\lambda_j = \max_{\substack{v \in \mathbb{C}^r \\ \dim(V)=j}} \min_{v \in V - \{0\}} \frac{(v, Hv)}{(v, v)}, \quad j \in \{1, \dots, r\}.$$

This has a generalization to partial sums of eigenvalues as follows: For every subspace  $E \subset \mathbb{C}^r$  and Hermitian operator  $H$  we denote by  $H_E$  the operator on  $E$  given by composing  $H$  with restriction and projection. Then for any  $J = \{j_1 < \dots < j_s\} \subset \{1, \dots, r\}$  we have

$$\sum_{j \in J} \lambda_j = \max_{\substack{F_1 \subset \dots \subset F_s \\ \dim(F_l) = j_l}} \min_{\substack{E \in G(s, n) \\ \dim(E \cap F_l) \geq l}} \text{Tr}(H_E).$$

Suppose that  $J_1, \dots, J_n$  are such that for every set of flags  $F_1, \dots, F_n$ , there exists a space  $E \in G(s, r)$  such that  $\dim(E \cap F_{i,l}) \geq j_{i,l}$  for  $i = 1, \dots, n$  and  $l = 1, \dots, s$ . Then

$$\begin{aligned} \sum_{i=1}^n \sum_{j \in J_i} \lambda_{i,j} &= \sum_{i=1}^n \max_{\substack{F_{i,1} \subset \dots \subset F_{i,s} \\ \dim(F_{i,l}) = j_{i,l}}} \min_{\substack{E_i \in G(s, r) \\ \dim(E_i \cap F_{i,l}) \geq l}} \text{Tr}(H_{i,E_i}) \\ &\leq \sum_{i=1}^n \text{Tr}(H_{i,E}) = \text{Tr}\left(\sum_{i=1}^n H_i|E\right) = 0. \end{aligned}$$

*Example 6.3.5.* Suppose that  $J_1 = \{1\}$ ,  $J_2 = \{r\}$ ,  $J_3 = \{r\}$ . Since every subspace of dimension 1 intersects  $\mathbb{C}^r$  in a subspace of dimension 1, namely itself, we obtain the inequality  $\lambda_{1,1} + \lambda_{2,r} + \lambda_{3,r} \leq 0$ . In terms of sums of matrices, this translates to the fact that  $\lambda_r(H_1) + \lambda_r(H_2) \leq \lambda_r(H_1 + H_2)$  for any Hermitian matrices  $H_1, H_2$ .

The existence of such an  $E$  for generic flags is implied by the non-vanishing of the *Schubert coefficient*  $\#[\bar{Y}_{J_1}] \cap \dots \cap [\bar{Y}_{J_n}]$  in the homology  $H(\text{Gr}(s, r))$  of the Grassmannian  $\text{Gr}(s, r)$ , where  $\bar{Y}_{J_i}$  are the *Schubert varieties* of (4.3). (The singular homology has no torsion and with real coefficients is isomorphic to the de Rham cohomology, so there is no conflict with notation.) Thus

**Theorem 6.3.6.** *If the Horn problem for  $\lambda_1, \dots, \lambda_n$  has a solution, then  $\sum_{l=1}^n \sum_{j \in J_i} \lambda_{i,j} \leq 0$  for all  $s < r$  and  $J_1, \dots, J_n$  of size  $s$  such that  $\#[\bar{Y}_{J_1}] \cap \dots \cap [\bar{Y}_{J_n}] > 0$  in  $H(\text{Gr}(s, r))$ .*

Unfortunately, from this point of view it is very difficult to see whether the list of all such inequalities is sufficient. Klyachko [63] noticed that this follows from the Hilbert-Mumford criterion. (See Fulton [35] for a more detailed discussion.) Let  $O_{\lambda_j} = K\lambda_j \cong G/P_{\lambda_j}^-$  for some *dominant*  $\lambda_1, \dots, \lambda_n$ ; for simplicity we assume that  $\lambda_j$  are generic. The quotient  $(O_{\lambda_1} \times \dots \times O_{\lambda_n})//K$  is non-empty iff the semistable locus in  $O_{\lambda_1} \times \dots \times O_{\lambda_n}$  is non-empty, iff a generic point  $F = (F_1, \dots, F_n)$  in  $O_{\lambda_1} \times \dots \times O_{\lambda_n}$  is semistable for all one-parameter subgroups. Let  $\xi \in \mathfrak{k}$  generate a one-parameter subgroup. Under the action of  $\exp(z\xi)$ ,  $z \rightarrow 0$ , the point  $F_j \in O_{\lambda_j}$  flows to a  $T$ -fixed point  $x_{w_j}$  where  $Y_{w_j}$  contains  $F_j$ . Thus  $F$  is  $\xi$ -semistable iff

$$(6.2) \quad \sum_{j=1}^n \langle \lambda_j, w_j^{-1}\xi \rangle \leq 0.$$

So  $F$  is  $\text{Ad}(g)\xi$ -semistable iff the same inequalities hold for  $w_j$  such that  $F_j \in gY_{w_j}$ . Let  $g_j \in G$  be such that  $F_j = g_j B/B$ . Then  $F_j$  lies in  $gY_{w_j}$  iff  $g^{-1}B/B \in g_j^{-1}Y_{w_j^{-1}}$ . Hence the semistable locus for the diagonal action of  $G$  is non-empty iff the inequalities (6.2) hold for dominant  $\xi$  whenever  $(w_1, \dots, w_n)$  are such that the intersection of the varieties  $g_j^{-1}Y_{w_j^{-1}}$  is non-empty for generic  $(g_1, \dots, g_n)$ . This gives a necessary and sufficient set of inequalities. From now on we drop the inverses on the Weyl group elements  $w_j$ , since they appear in both the inequalities and the intersection condition.

The next step is to reduce to inequalities for which the intersection number  $\#[\bar{Y}_{w_1}] \cap \dots \cap [\bar{Y}_{w_n}]$  is non-zero. If the intersection is positive dimensional for generic  $(g_1, \dots, g_n)$  then it represents a non-zero homology class of positive degree, and by Poincaré duality there exists an element  $w_{n+1} \in W$  such that  $\#[\bar{Y}_{w_1}] \cap \dots \cap [\bar{Y}_{w_{n+1}}] \neq 0$ . Then expanding the product of the last two  $[\bar{Y}_{w_n}] \cap [\bar{Y}_{w_{n+1}}]$  and choosing  $w'_n$  so that  $[\bar{Y}_{w'_n}]$  has positive coefficient in  $[\bar{Y}_{w_n}] \cap [\bar{Y}_{w_{n+1}}]$  one obtains  $w'_n$  such that  $\#[\bar{Y}_{w_1}] \cap \dots \cap [\bar{Y}_{w'_n}] \neq 0$ . Then  $w_n \lambda - w'_n \lambda \in \mathfrak{t}_+$  and so the inequality for  $(w_1, \dots, w'_n)$  implies that for  $(w_1, \dots, w_n)$ . The conclusion is that a generic point is semistable iff

$$\#[\bar{Y}_{w_1}] \cap \dots \cap [\bar{Y}_{w_n}] > 0 \implies \sum_{l=1}^n \langle \lambda_l, w_l \xi \rangle \leq 0 \quad \forall \xi \in \mathfrak{t}_+.$$

It suffices to check the inequalities for  $\xi$  in a set of generators for  $\mathfrak{t}_+$ . In particular, for  $K$  semisimple it suffices to check them for  $\xi$  equal to a fundamental coweight  $\omega_j^\vee$ , that is, for a generator of  $\mathfrak{t}_+$ . An argument similar to the one above shows that these inequalities correspond to non-zero intersection numbers in the corresponding generalized partial flag varieties:

**Theorem 6.3.7.** *Let  $K$  be a compact connected semisimple group with complexification  $G$ . A necessary and sufficient set of inequalities for the Horn-Klyachko problem is given by*

$$\#[\bar{Y}_{w_1}] \cap \dots \cap [\bar{Y}_{w_n}] > 0 \implies \sum_{l=1}^n \langle \lambda_l, w_l \omega_l^\vee \rangle \leq 0 \quad \forall \xi \in \mathfrak{t}_+.$$

as  $\omega_l^\vee$  ranges over fundamental coweights,  $[w_1], \dots, [w_n]$  range over elements of  $W/W_{\omega_j}$ ,  $Y_{w_1}, \dots, Y_{w_n} \subset G/P_{\omega_j}$  are the corresponding opposite Bruhat cells in the partial flag variety  $G/P_{\omega_j}$ , with the condition that  $\#[\bar{Y}_{w_1}] \cap \dots \cap [\bar{Y}_{w_n}] \neq 0$  in  $H(G/P_{\omega_j})$ .

For example, suppose that  $K = U(r)$  (and Klyachko’s argument was restricted to this case) so that  $\mathfrak{t}$  is naturally identified with  $\mathbb{R}^n$  and the  $j$ -th fundamental weight is identified with  $\omega_j = e_1 + \dots + e_j$ , where  $e_j$  is the  $j$ -th standard basis vector. In this case one obtains that  $(O_{\lambda_1} \times \dots \times O_{\lambda_n}) // K$  is non-empty iff for each  $j \in \{1, \dots, r\}$  and subsets  $J_1, \dots, J_n \subset \{1, \dots, r\}$  of size  $k$ ,

$$\#[\bar{Y}_{J_1}] \cap \dots \cap [\bar{Y}_{J_n}] > 0 \implies \sum_{l=1}^n \sum_{j \in J_l} \lambda_{l,j} \leq 0$$

c.f. Theorem 6.3.6. So the Hilbert-Mumford approach implies the sufficiency as well as the necessity of these inequalities.

The cohomology of the Grassmannian  $G(s, r)$  has a number of combinatorial models, for example, the famous Littlewood-Richardson rule. A recent “puzzles” model introduced by Knutson and Tao, see [68], is simple enough that we give a brief description. The *puzzle board* is the diagram shown in Figure 6.1. There are  $r$  little triangles along each big edge in the board. The *puzzle*




Figure 6.1: Puzzle board

pieces are shown in Figure 6.2. together with their rotations. A *puzzle* is a way of filling in the




Figure 6.2: Puzzle pieces

puzzle board with puzzle pieces so that all of the edges match.

*Example 6.3.8.* An example of a puzzle is shown in Figure 6.3.

For each puzzle, let  $I$  denote the positions of the 1’s on the northwest boundary,  $J$  the positions of the 1’s on the northeast boundary, and  $K$  the positions of the edge along the southern boundary, reading left to right.

*Example 6.3.9.* For Figure 6.3,  $I = \{2, 4\}$ ,  $J = \{2, 4\}$ ,  $K = \{2, 3\}$ .




Figure 6.3: An example of a puzzle

**Theorem 6.3.10.** [68] *The coefficient of  $[\overline{Y}_K]$  in  $[\overline{Y}_I] \cap [\overline{Y}_J] \in H(G(s, r))$  is the number of puzzles  $n_{I,J}^K$  with boundary data  $I, J, K$ .*

There are several possible proofs: one given by Knutson and Tao checks the equivalence with the Littlewood-Richardson rule. A second proof [69], joint with the author, proves associativity of the product defined by the puzzle numbers by a simple combinatorial trick, and then checks equality with the Schubert coefficients on generators. The formula generalizes to intersection numbers of arbitrary numbers of Schubert varieties, by considering puzzle boards with arbitrary numbers of “large boundaries”. For example, for  $n = 4$  one can take a diamond-shaped puzzle board.

Combining this combinatorial description with Klyachko’s argument gives the following:

**Corollary 6.3.11.** *If there is a puzzle whose 1’s on the boundary are in positions  $I, J, K$  then the inequality*

$$\sum_{i \in I} \lambda_i(H_1) + \sum_{j \in J} \lambda_j(H_2) \leq \sum_{k \in K} \lambda_k(H_1 + H_2)$$

holds for any Hermitian matrices  $A, B$ , and these inequalities together with the trace equality

$$\sum_{i=1}^n \lambda_i(H_1) + \sum_{j=1}^n \lambda_j(H_2) = \sum_{k=1}^n \lambda_k(H_1 + H_2)$$

give sufficient conditions for a triple  $(\lambda(H_1), \lambda(H_2), \lambda(H_1 + H_2))$  to occur.

*Example 6.3.12.* The puzzle in Example 6.3.8 gives the inequality  $\lambda_2(H_1) + \lambda_4(H_1) + \lambda_2(H_2) + \lambda_4(H_2) \leq \lambda_2(H_1 + H_2) + \lambda_3(H_1 + H_2)$ .

The following theorem of Knutson, Tao, and the author [68] (see also the review [67]), extending previous work of Belkale [12], describes a *minimal* set of inequalities:

**Theorem 6.3.13.** *The inequalities corresponding to  $I, J, K$  with  $n_{I,J}^K = 1$  together with the trace equality form a complete and irredundant set of necessary and sufficient conditions for the Horn problem for the sum of two Hermitian matrices.*

Many other problems of this type can be solved in the same way; for example see Agnihotri-Woodward [2] for a discussion of the possible eigenvalues of a product of unitary matrices, and relations with the invariant theory of *quantum* groups. In this case the existence of a good combinatorial model computing the eigenvalue inequalities is still open.

## 7. THE STRATIFICATIONS OF HESSELINK, KIRWAN, AND NESS

According to work of Kirwan [61] and Ness [84], the semistable locus of a  $G$ -variety  $X \subset \mathbb{P}(V)$  can be considered the open stratum in a Morse-type stratification of  $X$ . A theorem of Ness describes the equivalence of this stratification with one introduced by Hesselink [52], which measures the *slope of instability* of a point by its maximal Hilbert-Mumford weight.

**7.1. The Kirwan-Ness stratification.** Let  $X$  be a Hamiltonian  $K$ -manifold with proper moment map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ . Let  $(\cdot, \cdot) : \mathfrak{k} \rightarrow \mathfrak{k} \rightarrow \mathbb{R}$  be an invariant metric on  $\mathfrak{k}$  inducing an identification  $\mathfrak{k} \rightarrow \mathfrak{k}^\vee$ . Let

$$\phi = \frac{1}{2}(\Phi, \Phi) : X \rightarrow \mathbb{R}$$

denote the norm-square of the moment map. The notation  $\Phi(x)_X \in \text{Vect}(X)$  denotes the vector field determined by  $\Phi(x)$ , and  $\Phi(x)_X(x) \in T_x X$  its evaluation at  $x$ .

**Lemma 7.1.1.**  $\text{crit}(\phi) = \{x \in X, \Phi(x)_X(x) = 0\}$ .

*Proof.* We have  $d\phi(x) = (\Phi(x), d\Phi(x)) = -\iota_{\Phi(x)_X(x)}\omega_x$ . Since  $\omega$  is non-degenerate,  $d\phi(x)$  vanishes iff  $\Phi(x)_X(x) \in T_x X$  does.  $\square$

*Example 7.1.2.* Let  $X = \mathbb{P}^2$  and  $K = U(1)^2$  acting by  $(g_1, g_2)[z_0, z_1, z_2] = [z_0, g_1^{-1}z_1, g_2^{-1}z_2]$ . Consider the moment map

$$\Phi([z_0, z_1, z_2]) \mapsto (|z_1|^2/2, |z_2|^2/2) - (1/4, 1/4),$$

which has image the convex hull

$$\Delta(X) = \text{hull}\{(-1/4, -1/4), (-1/4, 3/4), (3/4, -1/4)\}.$$

The critical sets are the level sets of  $\Phi$  at  $(0, 0), (-1/4, 0), (0, -1/4), (1/4, 1/4), (-1/4, -1/4), (-1/4, 3/4), (3/4, -1/4)$ , see Figure 7.1.




Figure 7.1: Critical values for  $X = \mathbb{P}^2$

**Lemma 7.1.3.**  $\Phi(\text{crit}(\phi))$  is a discrete union of  $K$ -orbits in  $\mathfrak{k}^\vee$ , called the set of types for  $X$ .

*Proof.* Suppose first that  $K$  is abelian. Consider the orbit-type decomposition

$$X = \bigcup_{H \subset K} X_H, \quad X_H = \{x \in X | K_x = H\}.$$

where the union is over subgroups  $H \subset K$ . It follows from standard slice theorems that each  $X_H$  is a smooth manifold. Let  $\mathfrak{h}$  denote the Lie algebra of  $H$ . By Lemma 3.3.2,  $\Phi(X_H)$  is an open subset of an affine subspace parallel to  $\text{ann}(\mathfrak{h})$ . Thus  $\Phi(X_H \cap \text{crit}(\phi)) = \{\lambda \in \Phi(X_H) | \lambda \in \mathfrak{h}\}$  which is the set containing the unique point in  $\Phi(X_H)$  closest to 0, if it exists, and empty, otherwise. Since  $\Phi$  is proper, the pre-image of any compact set under  $\Phi$  contains only finitely many orbit-types, which proves the theorem in the abelian case.

Suppose that  $K$  is possibly non-abelian with maximal torus  $T$ . The action of  $T$  on  $X$  is also Hamiltonian with moment map  $\Phi_T$  obtained by composing  $\Phi$  with the projection of  $\mathfrak{k}^\vee$  onto  $\mathfrak{t}^\vee$ . Let  $\phi_T = (\Phi_T, \Phi_T)/2$ . Since  $\phi$  is  $K$ -invariant, any critical point is conjugate to a point  $x \in \text{crit}(\phi)$  with  $\Phi(x) \in \mathfrak{t}^\vee$ . Then  $x \in \text{crit}(\phi)$  iff  $x \in \text{crit}(\phi_T)$  iff  $\Phi(x)$  is a type for the action of  $T$ . Hence the types for  $K$  are locally finite.  $\square$

Choose a compatible  $K$ -invariant metric on  $X$ , and let  $\text{grad}(\phi) \in \text{Vect}(X)$  denote the gradient of  $\phi$ .

**Lemma 7.1.4.** The gradient of  $\phi$  is  $\text{grad}(\phi)(x) = J(x)\Phi(x)_X(x)$ .

*Proof.* Using the proof of Lemma 7.1.1, for  $v \in T_x X$

$$g_x(\text{grad}(\phi)(x), v) = D_x \phi(v) = -\omega_x(\Phi(x)_X(x), v) = g_x(J(x)\Phi(x)_X(x), v).$$

The claim follows.  $\square$

Let  $\varphi_t : X \rightarrow X$  be the flow of  $-\text{grad}(\phi)$ ; since  $\Phi$  is proper, so is  $\phi$  and so  $\varphi_t$  exists for all times  $t \in [0, \infty)$ .

**Proposition 7.1.5** (Duistermaat, see [74], [113]). *Any trajectory of  $\varphi_t$  has a limit.*

For the construction of the Kirwan stratification the actual convergence of  $\varphi_t$  is not needed. For each type  $\lambda$ , let  $C_\lambda = \Phi^{-1}(K\lambda) \cap \text{crit}(\phi)$  denote the corresponding component of the critical set of  $\phi$ . Since the set of types is discrete, any two limit points are contained in some  $C_\lambda \subset \text{crit}(\phi)$ , and in fact in the same connected component of  $\text{crit}(\phi)$ . Let  $X_\lambda$  denote the set of points  $x \in X$  flowing to  $C_\lambda$ ,

$$X_\lambda := \overline{\{\varphi_t(x), t \in [0, \infty)\}} \cap C_\lambda \neq \emptyset.$$

Note that since  $\phi$  is not Morse-Bott in general, there is no guarantee that  $X_\lambda$  is smooth. The *Kirwan-Ness stratification* is the decomposition [61], [84]:

$$X = \bigcup_{\lambda} X_\lambda.$$

**Theorem 7.1.6** (Kirwan). *There exists an invariant metric on  $X$  so that each stratum  $X_\lambda$  is smooth. The spectral sequence for the equivariant stratification  $X = \cup_{\lambda} X_\lambda$  collapses at the second page, so that*

$$H_K(X) \cong \bigoplus_{\lambda} H_K(X_\lambda).$$

In particular the canonical map  $H_K(X) \rightarrow H_K(\Phi^{-1}(0))$  (which is isomorphic to  $H(X//K)$  if  $K$  acts freely on  $\Phi^{-1}(0)$ ) is a surjection and the equivariant Poincaré polynomial of  $X$

$$p_X^K(t) = \sum t^j \text{rank } H_K^j(X)$$

is given by

$$p_X^k(t) = \sum_{\lambda} (-1)^{\text{codim}(X_\lambda)} p_{X_\lambda}^K(t).$$

If  $X$  acts freely on  $\Phi^{-1}(0)$  this means that the difference  $p_X^K(t) - p_{X//K}(t)$  is a finite sum of contributions from fixed point sets of one-parameter subgroups. We will see a version of this formula for sheaf cohomology in the last chapter.

In the case that  $X$  is a Kähler Hamiltonian  $K$ -manifold with proper moment map, the Kirwan-Ness stratification has a more explicit description. For each type  $\lambda$  let  $\varphi_{\lambda,t}$  denote the time  $t$  flow of  $-\text{grad}\langle \Phi, \lambda \rangle$ ,  $Z_\lambda$  the component of the fixed point set  $X^\lambda$  of the action of  $\lambda$  containing  $C_\lambda$ ,  $Y_\lambda$  the subset of  $X$  flowing to  $Z_\lambda$  under  $\varphi_{\lambda,t}$ ,  $K_\lambda$  the centralizer of  $\lambda$ , and  $U(1)_\lambda$  the one-parameter subgroup generated by  $\lambda$ . Then  $K_\lambda/U(1)_\lambda$  acts naturally on  $Z_\lambda$  in Hamiltonian fashion with moment map denoted  $\Phi_\lambda$ , obtained by restricting  $\Phi$  to  $Z_\lambda$  and projecting out the direction generated by  $\lambda$ . We denote by  $Z_\lambda^{\text{ss}}$  the set of points flowing to  $\Phi_\lambda^{-1}(0)$  under the flow of minus the gradient of the norm-square of  $\Phi_\lambda$ . Let  $Y_\lambda^{\text{ss}}$  denote the inverse image of  $Z_\lambda^{\text{ss}}$  in  $Y_\lambda$ .

**Theorem 7.1.7** (Kirwan [61]). *Let  $X$  be a compact Kähler Hamiltonian  $K$ -manifold with proper moment map  $\Phi : X \rightarrow \mathfrak{k}^\vee$ . For the Kähler metric each  $X_\lambda$  is a  $G$ -invariant complex submanifold, each  $Y_\lambda$  is a  $P_\lambda$ -invariant complex submanifold, and  $G \times_{P_\lambda} Y_\lambda^{\text{ss}} \rightarrow X_\lambda$ ,  $[g, y] \mapsto gy$  is an isomorphism of complex  $G$ -manifolds.*

We give a proof, and explain the relation with a theorem of Ness [84], in the following section. In the point of view we will present, a key fact is that the gradient flow of the norm-square of the moment map is essentially equivalent to the gradient flow of the Kempf-Ness function, as was pointed out in Donaldson-Kronheimer [30, Section 6]. Let  $X$  be a Kähler Hamiltonian  $K$ -manifold with proper moment map. For any  $x \in X$ , let  $x_t$  denote the trajectory of the gradient flow of  $-\phi$  starting at  $x$ . On the other hand, let  $\psi : K \backslash G \rightarrow \mathbb{R}$  be the Kempf-Ness function for  $x$ . We may also consider the gradient flow of  $\psi$ , with respect to the given metric on  $\mathfrak{k}$ .

**Proposition 7.1.8.** *Let  $X, x, \psi$  be as above. The map*

$$K \backslash G \rightarrow X/K, [g] \mapsto [gx]$$

*maps the gradient trajectories of  $\psi$  onto the image of the gradient trajectories of  $\phi \bmod K$ .*

*Proof.* Using Corollary 5.2.3,

$$\begin{aligned} J_X((\text{grad}(\psi))([g]))_X(gx) &= J_X(\Phi(gx))_X(gx) \\ &= (\text{grad}(\phi))(gx). \end{aligned}$$

The vector field on  $G$  given by  $g \mapsto i \text{grad}(\psi)([g])$  has trajectories that map to the gradient trajectories of  $\text{grad}(\psi)$  under  $G \rightarrow K \backslash G$ , and to the trajectories of  $\text{grad}(\phi)$  under  $G \rightarrow X$ , which gives the result.  $\square$

In particular, since the trajectories of  $\psi$  exist for all time by the bound on  $\Phi$ , any trajectory of  $-\text{grad}(\phi)$  is contained in a single  $G$ -orbit:  $x_t \in Gx, \forall x \in X, t \in \mathbb{R}$ .

**Corollary 7.1.9.**  *$\psi$  is bounded from below iff the gradient flow for  $-\phi$  converges to  $\Phi^{-1}(0)$ .*

*Proof.* In the algebraic case, this is nothing but a reformulation of 4.3.4. For the Kähler case, note that if  $\psi$  is bounded from below then  $\text{grad}(\psi)$  converges to zero along any gradient trajectory, and by equivalence of gradient flows 7.1.8 it follows that  $\Phi$  must converge to zero. The converse follows as in the proof of Theorem 5.5.9, using that  $\text{grad}(\psi)$  converges to zero exponentially fast along any one-parameter subgroup whose limit corresponds to a polystable point.  $\square$

One obtains an analytic proof of the Hilbert-Mumford criterion Theorem 4.3.1 by combining Corollary 7.1.9 and Theorem 5.4.2.

*Remark 7.1.10.* A rather confusing point is that if  $\psi : K \backslash G \rightarrow \mathbb{R}$  is a Kempf-Ness function, then the slope function  $(K \backslash G)_\infty \rightarrow \mathbb{R}$  of Theorem 5.4.2 is not continuous in the topology on  $(K \backslash G)_\infty$  induced by the identification with the unit sphere in the Lie algebra  $\mathfrak{k}$ , but rather only in the topology induced by the Tits metric. This happens already for the action of  $SL(2, \mathbb{C})$  on  $\mathbb{P}^1$ : the asymptotic slope for the Kempf-Ness function for  $[0, 1]$  is 1 for every direction except that generated by  $\text{diag}(i, -i)$ , where it is  $-1$ ; the topology induced by the Tits metric in this case is discrete.

**7.2. The Hesselink stratification.** Let  $X \subset \mathbb{P}(V)$  be a projective  $G$ -variety, or more generally a compact Kähler Hamiltonian  $K$ -manifold. The Hesselink stratification uses the weights appearing in the Hilbert-Mumford criterion to construct a stratification on  $X$ : Define for any non-zero  $\lambda \in \mathfrak{k}$  the *Hilbert-Mumford slope*

$$\mu_\lambda(x) = \langle \Phi(x_\lambda), \lambda \rangle / \|\lambda\|.$$

By Corollary 5.2.3, the Hilbert-Mumford slope is equal to the asymptotic slope of the Kempf-Ness function studied in Theorem 5.4.2.

**Definition 7.2.1.** A point  $x \in X$  is

- (a) *slope semistable* iff  $\mu_\lambda(x) \leq 0$  for all  $\lambda$ ,
- (b) *slope stable* iff  $\mu_\lambda(x) < 0$  for all  $\lambda$ ,
- (c) *slope unstable* iff  $x$  is not semistable, and
- (d) *slope polystable* iff it is slope semistable and  $\mu_\lambda(x) = 0$  implies  $\mu_{-\lambda}(x) = 0$  for all  $\lambda$ .

Slope semistability might also be called Hilbert-Mumford semistability, but this seems a little unwieldy. We have already seen in the proof of the Kempf-Ness theorem that slope semistability is equivalent to semistability. The equivalence of slope polystability with polystability is proved in Mundet [56]. It follows from Section 5.5 that a point  $x \in X$  is polystable but not stable iff its Jordan-Hölder cone contains a line.

The set of destabilizing one-parameter subgroups is studied by Hesselink in the algebraic case [52], [53], see also Ramanan-Ramanathan [90]. For any  $\lambda$  we denote by  $G_\lambda$  the centralizer of  $\lambda$  and by  $\mathbb{C}_\lambda^*$  the one-parameter subgroup generated by  $\lambda$ . Obviously  $\mathbb{C}_\lambda^* \subset G_\lambda$ . Let  $x \in X$  and  $Z_\lambda$  denote the component of  $X^\lambda$  containing  $x_\lambda$ . Then the action of  $G_\lambda$  on  $Z_\lambda$  descends to an action of  $G_\lambda / \mathbb{C}_\lambda^*$ . Furthermore, the inner product on  $\mathfrak{k}$  determines a splitting  $\mathfrak{g}_\lambda = \mathbb{C}\lambda \oplus \mathfrak{g}_\lambda / \mathbb{C}\lambda$  which defines a lift of  $G_\lambda / \mathbb{C}_\lambda^*$  to the polarizing line bundle, at least up to finite cover. So we may consider

$Z_\lambda$  as a polarized  $G_\lambda/\mathbb{C}_\lambda^*$ -variety, with the caveat that the polarization depends on the choice of inner product on  $\mathfrak{k}$ .

**Theorem 7.2.2.** *Any unstable  $x$  has a unique (up to scalar multiple) maximally destabilizing one-parameter subgroup generated by  $\lambda \in \mathfrak{k}$  with the property that  $x_\lambda$  is a semistable point for the action of  $G_\lambda/\mathbb{C}_\lambda^*$  on  $Z_\lambda$  and  $\lambda$  is maximally destabilizing:  $\mu_\nu(x) \leq \mu_\lambda(x)$  for all  $\nu \in \mathfrak{k} - \{0\}$  and equality holds iff  $\mathbb{R}_+\nu = \mathbb{R}_+\lambda$ .*

*Proof.* We already proved in Theorem 5.4.2 the existence of a maximally destabilizing one-parameter subgroup generated by some  $\lambda \in \mathfrak{k}$ . It remains to show that for any  $x \in X$ ,  $x_\lambda$  is semistable for  $G_\lambda/\mathbb{C}_\lambda^*$ , or what is equivalent,  $\Phi((x_\lambda)_t)$  converges to  $K\lambda$ . Since  $(\Phi(x_\lambda), \lambda) = (\lambda, \lambda)$ , the function  $\psi$  goes to  $-\infty$  along  $\exp(-i\lambda t)x$  as fast as  $x_t$ . If  $\lim \|\Phi((x_\lambda)_t)\| > \lambda$ , then  $\psi((\exp(-i\lambda t_1)x)_{t_2})$  as  $t_1 \gg t_2 \rightarrow \infty$  goes to  $-\infty$  faster than  $\psi(x_t)$ , which contradicts convexity of  $\psi$  as in the proof of Theorem 5.4.2. Hence  $\|\Phi((x_\lambda)_t)\| \rightarrow \lambda$ . Now the gradient trajectories for  $\exp(-i\lambda t)x$  converge to a broken gradient trajectory for  $x_\lambda$ ; since each piece in the broken gradient trajectory must decrease  $\phi$ , the limit has only one piece. That is,  $\Phi((x_\lambda)_t) \rightarrow K\lambda$ .  $\square$

Let  $\Lambda$  denote the set of equivalence classes of one-parameter subgroups appearing in Hesselink's theorem (with equivalence given by the adjoint action) we call the decomposition  $X = \cup_\lambda X_\lambda$  the *Hesselink stratification* of  $X$ .

*Remark 7.2.3.* The Hesselink stratification is the finite-dimensional analog of the Shatz stratification [97] of the moduli stack of vector bundles on a curve by the type of the Harder-Narasimhan filtration.

The following is proved in the algebraic case by Ness [84].

**Theorem 7.2.4.** *The Hesselink and Kirwan-Ness stratifications agree.*

*Proof.* By Theorem 5.4.2 part (c).  $\square$

*Proof of Kirwan's theorem 7.1.7.* Let  $\psi = \psi_\nu$  be a Kempf-Ness function, and  $X_\lambda$  a Hesselink stratum (or equivalently, a Kirwan-Ness stratum.) Let  $U_\lambda$  denote the set of points in  $x$  with direction of maximally negative slope  $\lambda$ . Uniqueness of  $\lambda$  implies that if  $x \in U_\lambda$  and  $g \in G$  is such that  $gx \in U_\lambda$ , then  $g \in P_\lambda$ . Indeed, note  $G = KP_\lambda$  and  $U_\lambda$  is  $P_\lambda$ -stable. Hence it suffices to consider the case  $g \in K$ , and then  $g\lambda$  is also a direction of maximal descent. Hence  $g\lambda = \lambda$  which implies that  $g \in K_\lambda$ , hence  $g \in P_\lambda$ . This implies  $X_\lambda = G \times_{P_\lambda} U_\lambda$ . To see that  $U_\lambda = Y_\lambda^{\text{ss}}$  of Section 7.1, note that  $U_\lambda \subset Y_\lambda^{\text{ss}}$  by Theorem 7.2.2. On the other hand,  $Y_\lambda^{\text{ss}}$  is contained in  $U_\lambda$ : any point in  $Y_\lambda^{\text{ss}}$  has a point in  $\Phi^{-1}(\lambda)$  in its orbit-closure, and  $\|\lambda\|^2$  minimizes  $\|\Phi\|^2$  on  $Y_\lambda^{\text{ss}}$  which implies that  $\lambda$  is the direction of maximally negative slope. This completes the proof.  $\square$

*Remark 7.2.5.* Suppose  $\omega \in \Omega^2(X)$  is a closed two form that is not symplectic, but satisfies  $\omega(\xi_X, J\xi_X) > 0$  for any  $\xi \in \mathfrak{k}$ . The proof above works equally well for moment maps associated to such two-forms. That is, only non-degeneracy of the two-form on the directions generated by the action is used in the proof.

## 8. MOMENT POLYTOPES

According to work of Atiyah, Guillemin-Sternberg, and Kirwan, the quotient of the image of the moment map is convex. (This section could have been placed before that on Schur-Horn convexity.)

**8.1. Convexity theorems for Hamiltonian actions.** Let  $X$  be a Hamiltonian  $K$ -manifold with moment map  $\Phi$ . The *moment image* of  $X$  is  $\Phi(X) \subset \mathfrak{k}$ . The quotient

$$\Delta(X) := \Phi(X)/K \subset \mathfrak{k}^\vee/K$$

can be identified with a subset of the convex cone  $\mathfrak{t}_+^\vee \cong \mathfrak{k}^\vee/K$ .

*Example 8.1.1.* If  $X = \mathbb{P}^{n-1}$  and  $G = U(1)^n$  acts by the standard representation, then the moment image is the *standard  $n$ -simplex*

$$\Phi(X) = \{(\nu_1, \dots, \nu_n) \in \mathbb{R}_{\geq 0}^n \mid \nu_1 + \dots + \nu_n = 1\},$$

see (3.3). The coordinate hyperplane  $\{z_j = 0\} \subset X$  maps to the  $j$ -th facet  $\{\nu_j = 0\} \subset \Phi(X)$ .



Another description of the moment polytope  $\Delta(X)$  involves the *shifted* symplectic quotients: for  $\lambda \in \mathfrak{k}^\vee$ , the quotient

$$X//_\lambda K := \Phi^{-1}(K\lambda)/K = (\mathcal{O}_\lambda^- \times X)//K$$

is the *symplectic quotient of  $X$  at  $\lambda$* . The shifted symplectic quotient is the classical analog of the multiplicity space  $\text{Hom}_K(V_\lambda, V)$  of a representation  $V$  in the following sense:

**Proposition 8.1.2.** *Let  $X$  be a polarized projective  $G$ -variety and  $\lambda$  a dominant weight. Then  $R(X//_\lambda G)_d = \text{Hom}_G(V_{d\lambda}, R(X)_d)$  for any  $d \geq 0$ .*

*Proof.* Combining the Borel-Weil and Kempf-Ness theorems gives

$$\begin{aligned} R(X//_\lambda K)_d &= R(K\lambda^- \times X)_d^K \\ &= (V_{d\lambda}^\vee \otimes R(X)_d)^K \\ &= \text{Hom}_K(V_{d\lambda}, R(X)_d) \end{aligned}$$

□

The following is immediate from the definitions:

**Lemma 8.1.3.**  $\Delta(X) = \{\lambda \mid X//_\lambda K \neq \emptyset\}$  is the set of  $\lambda$  for which the shifted symplectic quotient  $X//_\lambda K$  is non-empty.

The set  $\Delta(X)$  is the “classical analog” of the set of simple modules appearing in a  $G$ -module. Let  $\Delta_{\mathbb{Q}}(X) := \Lambda_{\mathbb{Q}}^\vee \cap \Delta(X)$  denote the set of rational points in  $\Delta(X)$ ; furthermore  $\Delta_{\mathbb{Q}}(X)$  is dense in  $\Delta(X)$ , see for example [73].

**Theorem 8.1.4.**  $\Delta_{\mathbb{Q}}(X) = \Delta(X) \cap \Lambda_{\mathbb{Q}}^\vee$  is equal to the set of points  $\lambda/d$  such that  $V_\lambda \subset R(X)_d$ .

*Proof.* By Lemma 8.1.3 and Proposition 8.1.2. □

Recall that a *convex polyhedron* is the intersection of a finite number of half spaces, while a *convex polytope* is the convex hull of a finite number of points. The fundamental theorem of convex geometry says that any *compact* convex polyhedron is a convex polytope and vice-versa.

**Theorem 8.1.5** (Atiyah [7], Guillemin-Sternberg [37] for the abelian case, Kirwan [62] for the non-abelian case). *Let  $K$  be a compact, connected Lie group and  $X$  a compact connected Hamiltonian  $K$ -manifold. Then  $\Delta(X)$  is a convex polytope. If  $K$  is abelian, then  $\Delta(X)$  is the convex hull of the image  $\Phi(X^K)$  of the fixed point set  $X^K$  of  $K$ .*

$\Delta(X)$  is the *moment polytope* of  $X$ . The arguments of Atiyah and Guillemin-Sternberg in [7], [37] are Morse-theoretic. The equivariant version of Darboux’s theorem implies that the functions  $\langle \Phi, \xi \rangle$  have only critical sets of even index, and this implies that the level sets  $\langle \Phi, \xi \rangle^{-1}(c)$  are connected. Using an inductive procedure one shows that for any subtorus  $K_1 \subset K$ , the level sets of the moment map for  $\Phi_1$  are connected as well. Taking  $K_1$  of codimension one, this shows that the intersection of  $\Phi(X)$  with any rational line is connected and it follows that  $\Phi(X)$  is convex. The reader is referred to the original papers for details. Kirwan’s non-abelian version uses the Morse theory of the norm-square of the moment map. See Lerman-Meinrenken-Tolman-Woodward [73] for a derivation of non-abelian convexity from the abelian case.

Brion [18], following earlier work of Mumford [84, Appendix], pointed out the following proof of convexity, which in language of geometric quantization would be called a “quantum” proof: Suppose  $\lambda_j/d_j \in \Delta_{\mathbb{Q}}(X)$ ,  $j = 0, 1$ . Let  $v_j \in R(X)_{d_j}$  be corresponding highest weight vectors. Then for any  $n_0, n_1 \in \mathbb{N}$ ,  $v_0^{n_0} v_1^{n_1} \in R(X)_{n_0 d_0 + n_1 d_1}$  is a highest weight vector, so

$$\frac{n_0 \lambda_0 + n_1 \lambda_1}{n_0 d_0 + n_1 d_1} = \frac{d_0 n_0}{d_0 n_0 + d_1 n_1} (\lambda_0/d_0) + \frac{d_1 n_1}{n_0 d_0 + n_1 d_1} (\lambda_1/d_1) \in \Delta_{\mathbb{Q}}(X).$$

This implies that  $\Delta_{\mathbb{Q}}(X)$  is convex.

The inequalities of the previous section (for example, the Horn-Klyachko problem) can now be seen as the inequalities describing the moment polytopes of products of coadjoint orbits.

**8.2. Convexity theorems for orbit-closures.** In the case that  $X$  is Kähler, Atiyah [7] also described the images of orbit-closures under the moment map, in the case that  $K$  is abelian. Of course if the orbit-closure is smooth, then this falls under the assumptions of the previous convexity theorem, but Atiyah’s theorem also includes the case of singular orbit-closures:

**Theorem 8.2.1.** [9, Theorem 2] *Let  $K$  be a torus,  $G$  its complexification, and  $X$  a Kähler Hamiltonian  $K$ -manifold. Let  $Y \subset X$  be a  $G$ -orbit. Then*

- (a)  $\Delta := \Phi(\overline{Y})$  is a convex polytope with vertices  $\Phi(\overline{Y} \cap X^G)$ ;
- (b) For each open face  $F \subset \Delta$ ,  $\Phi^{-1}(F) \cap \overline{Y}$  is a single  $G$ -orbit.
- (c)  $\Phi$  induces a homeomorphism of  $\overline{Y}/G$  onto  $\Delta$ .

We will describe Atiyah’s arguments since they are brief and are closely related to the one-parameter subgroups of Hesselink as well as the Jordan-Hölder subgroups of Section 5.5.9. The proof depends on the following

**Lemma 8.2.2.** *Let  $Y \subset X$  be a  $G$ -orbit and  $y \in Y$ . Then*

- (a)  $y_\lambda = \lim_{t \rightarrow \infty} (\exp(it\lambda)y)$  exists and lies in the fixed point set  $X^\lambda$ ;
- (b)  $\lim_{t \rightarrow \infty} \langle \Phi(\exp(it\lambda)y), \lambda \rangle$  exists and is a constant  $d_\lambda$  independent of  $y$ .
- (c)  $d_\lambda = \sup_{y \in Y} \langle \Phi(y), \lambda \rangle$  is the asymptotic slope in Theorem 5.4.2.

Suppose that  $\lambda$  is generic so that  $X^G = X^\lambda$ . The Lemma implies

$$\sup_{y \in Y} \langle \Phi(y), \lambda \rangle = \sup_{y \in X^G \cap \overline{Y}} \langle \Phi(y), \lambda \rangle.$$

Hence  $\Phi(Y)$  is contained in the convex hull of  $\Phi(X^G \cap \overline{Y})$ . To see that  $\Phi(\overline{Y}) = \Delta$ , Atiyah notes that for any  $y \in Y$  and direction  $\xi \in \mathfrak{k}$  of unit length, there exists a time  $t(\xi)$  such that

$$\langle \Phi(\exp(it(\xi)\xi)y), \xi \rangle = \frac{1}{2}(\Phi(y) + d(\xi)).$$

The set of points  $\exp(i\xi)y$  with  $\|\xi\| \leq t(\xi)/\|\xi\|$  defines a neighborhood  $U$  of  $y$  in  $Y$  with  $\Phi(\overline{U}) = \Phi(y) + \frac{1}{2}(\Delta - \Phi(y))$ ; this immediately implies that  $\Phi(\overline{Y})$  is both open and closed in  $\Delta$  and hence equal to  $\Delta$ .

To prove the third part of the Theorem, Atiyah considers for any  $\lambda \in \mathfrak{k}$  and fixed point component  $Z \subset X^\lambda$ , the unstable manifold  $Z^u$  consisting of all points that flow to  $Z$  under  $\exp(it\lambda)$ . By the stable manifold theorem  $Z^u$  is a smooth manifold and the limit of the flow defines a smooth  $G$ -equivariant projection  $Z^u \rightarrow Z$ . In particular, if  $Z$  is any component of  $X^\lambda$  containing a limit point of  $Y$  then  $Y \subset Z^u$  and  $\overline{Y} \cap Z$  is a single  $G$ -orbit. From this it follows that  $\Phi(Z \cap \overline{Y})$  is a face of  $\Delta$  with fibers the orbits of the compact torus  $K$ , see [9, p. 10], and this completes the proof.

*Remark 8.2.3.* Atiyah’s theorem makes the theory of polystable points and Jordan-Hölder vector described in Section 5.5 substantially easier in the abelian case. One sees that the “Jordan-Hölder” cone of Theorem 5.5.9 is the dual cone to the face of the polytope containing 0, in the case that  $Y$  is a semistable orbit.

Atiyah’s convexity theorem for orbit-closures has been generalized to Borel subgroups by Guillemin and Sjamaar [44].

## 9. MULTIPLICITY-FREE ACTIONS

In certain cases Hamiltonian or algebraic actions may be classified by combinatorial data related to the moment map. In this section we discuss an example of this, the *multiplicity-free case*, in which the symplectic and git quotients are points.

**9.1. Toric varieties and Delzant’s theorem.** A *toric variety* is an irreducible normal  $G$ -variety  $X$  such that  $G$  is an algebraic torus and  $X$  contains an open  $G$ -orbit. Affine toric varieties are naturally classified by monoids  $M$  in the group  $\Lambda^\vee$  of weights of  $G$ , with the corresponding toric variety given by  $\text{Spec}(\mathbb{C}[M])$ . Each such monoid spans a rational cone in  $\Lambda_\mathbb{Q}^\vee$ , and defines a dual cone in  $\Lambda_\mathbb{Q}$ . Toric varieties with trivial generic stabilizer are classified by *fans* in  $\Lambda_\mathbb{Q}$ , that is, collections of cones such that any intersection of a cone is again a cone in the fan, see Oda [87] or Fulton [34].

*Example 9.1.1.* Suppose that  $X = \mathbb{P}^2$  with action given by  $(w_1, w_2)[z_0, z_1, z_2] = [z_0, w_1^{-1}z_1, w_2^{-1}z_2]$ . There are seven orbits, given by non-vanishing of various coordinates, and in particular, three closed orbits  $[1, 0, 0]$ ,  $[0, 1, 0]$ ,  $[0, 0, 1]$ , whose cones are generated by pairs of vectors  $(1, 1), (-1, 0), (1, 1), (0, -1)$ , and  $(0, -1), (0, -1)$ . The fan contains these three cones, and their intersections; this is the *dual fan* to the moment polytope  $\text{hull}((0, 0), (1, 0), (0, 1))$ .

A Hamiltonian torus action is *multiplicity free* or *completely integrable* if all the symplectic quotients are points, or equivalently, each fiber of the moment map is an orbit of the torus.

*Example 9.1.2.* The  $U(1)^n$  action on  $\mathbb{P}^n$  is multiplicity-free, since the fibers of the moment map are given by  $[z_0, \dots, z_n]$  with  $|z_1|, \dots, |z_n|$  fixed, which are orbits of  $U(1)^n$ .

Multiplicity-free Hamiltonian torus actions are classified by a theorem of Delzant.

**Definition 9.1.3.** A polytope  $\Delta \subset \mathfrak{k}^\vee$  is called *Delzant* if the normal cone at any vertex is generated by a basis of the weight lattice  $\Lambda^\vee \subset \mathfrak{k}^\vee$ .

**Theorem 9.1.4** (Delzant [28]). *There exists a one-to-one correspondence between Delzant polytopes and multiplicity-free torus actions on compact connected manifolds with trivial stabilizer, given by  $X \mapsto \Phi(X)$ . Any compact connected multiplicity-free Hamiltonian torus action has the structure of a smooth projective toric variety.*

Note that any compatible complex structure is unique up to isomorphism, but not up to Kähler isomorphism. That is, any toric variety has many non-equivalent Kähler structures, see Guillemin [43].

There are “local” and “local-to-global” parts of the proof; the local part follows from the equivariant Darboux theorem, while the “local-to-global” part uses the vanishing of a certain sheaf cohomology group over the polytope.

Existence of a smooth projective toric variety with a given polytope follows from, for example, Lerman’s method of *symplectic cutting* [72] which we now describe. We begin with the simplest case, when  $X$  is a Hamiltonian  $S^1$ -manifold with moment map  $\Phi : X \rightarrow \mathbb{R}$ . The diagonal  $S^1$ -action on  $X \times \mathbb{C}$  is Hamiltonian with moment map

$$\Phi_{X \times \mathbb{C}} : (x, z) \mapsto \Phi(x) - |z|^2/2.$$

Its symplectic quotient at any value  $\lambda$

$$X_{\geq \lambda} := (X \times \mathbb{C}) //_{\lambda} S^1$$

is called the *symplectic cut* of  $X$  at  $\lambda$  admits a decomposition

$$(X \times \mathbb{C}) //_{\lambda} S^1 \cong X //_{\lambda} S^1 \cup (X \times \mathbb{C}^*) //_{\lambda} S^1 \cong X //_{\lambda} S^1 \cup \Phi^{-1}((\lambda, \infty)).$$

It follows from the definitions that the inclusion of  $\Phi^{-1}((\lambda, \infty))$  in  $X_{\geq \lambda}$  is symplectic and so  $X_{\geq \lambda}$  is obtained by removing  $\Phi^{-1}((-\infty, \lambda))$  and “closing off” the boundary by quotienting it by  $S^1$ .

More generally, suppose that  $K$  is a torus,  $\xi \in \mathfrak{k}$  any rational vector, and  $\lambda \in \mathbb{R}$ . Let  $U(1)_{\lambda}$  denote the one-parameter subgroup generated by  $\lambda$ , with moment map  $\langle \Phi, \lambda \rangle$ . Then the symplectic cut  $X_{\geq \lambda} = (X \times \mathbb{C}) //_{\lambda} U(1)_{\lambda} \cong X //_{\lambda} U(1)_{\lambda} \cup \{ \langle \Phi, v \rangle > \lambda \}$  admits the structure of a Hamiltonian  $K$ -manifold with moment polytope  $\Phi(X_{\geq \lambda}) = \Phi(X) \cap \{ \langle \nu, v \rangle \geq \lambda \}$ .

*Example 9.1.5.* Let  $X = \mathbb{P}^2$  equipped with  $U(1)^2$ -action given by with weights  $(0, 0), (-2, 0), (0, -2)$ . The moment polytope is then the convex hull of  $(0, 0), (2, 0), (0, 2)$ . Let  $\lambda = (0, -1)$  so that the one-parameter subgroup generated by  $\lambda$  acts with moment map

$$[z_0, z_1, z_2] \mapsto -2|z_1|^2 / (|z_0|^2 + |z_1|^2 + |z_2|^2).$$

The symplectic cut at  $-1$  is then a toric variety with polytope the convex hull of  $(0, 0), (0, 2), (1, 0), (1, 1)$ , see Figure 9.1.

Suppose that  $\Delta$  is a Delzant polytope defined by inequalities

$$\Delta = \{ \nu \in \mathfrak{k}^\vee \mid \langle \nu, v_j \rangle \geq \lambda_j, j = 1, \dots, m \}$$

for some vectors  $v_j \in \mathfrak{k}$  and some constants  $\lambda_j \in \mathbb{R}, j = 1, \dots, m$ . Let  $X = T^\vee K$ , with moment image  $\mathfrak{k}^\vee$  and the standard Kähler structure. Performing a symplectic cut for each inequality gives a Kähler manifold with Hamiltonian  $K$  action and moment polytope  $\Delta$ .




Figure 9.1: Effect of cutting on a moment polytope

Alternatively any smooth projective toric variety is a symplectic or geometric invariant theory quotient of affine space  $X = \mathbb{C}^m$ . There is an explicit description of the semistable locus given by Audin [10] and Cox [27].

**9.2. Multiplicity-free actions and spherical varieties.** Let  $K$  be a compact connected Lie group. Recall that a  $K$ -module  $V$  is *multiplicity-free* iff  $\dim \text{Hom}_K(V_\lambda, V)$  is at most one, for any simple  $K$ -module  $V_\lambda$  iff  $\text{End}_K(V)$  is abelian, using Schur's lemma. The definition in part (a) of the following was introduced in Guillemin-Sternberg [45]:

**Theorem 9.2.1.** (see [111, Appendix]) *The following conditions are equivalent, and if they hold the action is multiplicity-free:*

- (a)  $C^\infty(X)^K$  is an abelian Poisson algebra.
- (b) The symplectic quotient  $X//_\lambda K := \Phi^{-1}(K\lambda)/K$  is a point for all  $\lambda$ .

*Proof.* We denote by  $r_\lambda : C^\infty(X)^K \rightarrow C^\infty(X//_\lambda K)$  the map of Poisson algebras induced by the symplectic quotient construction, if  $\lambda$  is free. In general, we define

$$C^\infty(X//_\lambda K) := C^\infty(X)^K / \{f, f | \Phi^{-1}(\lambda) = 0\}.$$

A lemma of Arms, Cushman, and Gotay [6], see Sjamaar-Lerman [100], says that this quotient is a non-degenerate Poisson algebra, that is, the bracket vanishes only on constant functions. Suppose (a). Since  $r_\lambda$  is surjective,  $C^\infty(X//_\lambda K)$  is abelian as well, and so  $X//_\lambda K$  must be discrete, hence a point by Kirwan's results. Conversely, if all the reduced spaces are points and  $f, g \in C^\infty(X)^K$  then  $r_\lambda(\{f, g\}) = 0$  for all  $\lambda$  implies that  $\{f, g\} = 0$ .  $\square$

The complex analogs of multiplicity-free Hamiltonian actions are called *spherical varieties*. Let  $G$  be a connected complex reductive group. For the following, see Brion-Luna-Vust [20], the review [65], or the second part of Brion's review in this volume.

**Theorem 9.2.2.** *The following conditions for a normal  $G$ -variety  $X$  are equivalent; if they hold  $X$  is called spherical:*

- (a) some (hence any) Borel subgroup  $B$  has an open orbit;
- (b) the space of rational functions  $\mathbb{C}(X)$  is a multiplicity-free  $G$ -module;
- (c) some (hence any) Borel subgroup  $B$  has finitely many orbits.

*Remark 9.2.3.* For an arbitrary group action, existence of a dense orbit does not imply finitely many orbits. For example, consider the action of  $SL(n, \mathbb{C})$  on the space of  $n \times n$  matrices on the left: any two invertible matrices are related by an element of  $SL(n, \mathbb{C})$ , but there are infinitely many orbits of degenerate matrices distinguished by their kernels.

The classification of toric varieties is generalized to spherical varieties as a special case of a theorem of Luna-Vust [77] which gives a classification of spherical varieties by their generic isotropy group and a *colored fan*, see the contribution of Pezzini in this volume or Knop [65]. Each colored fan is a collection of *colored cones*, convex cones in the space  $\Lambda_X$  dual to the space  $\Lambda_X^\vee$  of characters corresponding to  $B$ -semiinvariant functions  $\mathbb{C}(X)^{(B)}$ , together with a finite set of  $B$ -stable divisors, satisfying certain conditions. The classification of generic isotropy groups that appear, which are called *spherical subgroups*, is the subject of an open conjecture of Luna, see the contribution of Bravi in this volume. The relation between multiplicity-free Hamiltonian actions and spherical varieties is given by the following, which is a consequence of the Kempf-Ness theorem:

**Proposition 9.2.4.** *A smooth  $G$ -variety  $X \subset \mathbb{P}(V)$  is spherical if and only if it is a multiplicity-free Hamiltonian  $K$ -manifold.*

*Proof.* By Proposition 8.1.2  $X//_\lambda K = \text{pt}$  iff  $\text{Hom}_G(V_\lambda, H^0(X, \mathcal{O}_X(d)))$  is dimension one or zero for all  $d \geq 0$ . This holds for all  $\lambda$  and  $d \geq 0$  iff  $\mathbb{C}(X)_d$  is a multiplicity-free  $G$ -module for all  $d \geq 0$  iff  $X$  is spherical.  $\square$

In contrast to the toric case, not every multiplicity-free Hamiltonian action admits the structure of a spherical variety [112].

**9.3. Moment polytopes of spherical varieties.** We have already seen several examples of the following:

**Theorem 9.3.1.** *Let  $X$  be a smooth polarized spherical  $K$ -variety with moment polytope  $\Delta$  and trivial generic stabilizer. Then  $H^0(X, \mathcal{O}_X(1))$  is the multiplicity-free  $K$ -module whose weights are the integral points  $\Delta \cap \mathfrak{k}^\vee$  of  $\Delta$ .*

*Proof.* By Proposition 8.1.2 and the fact that the symplectic quotients are points.  $\square$

The moment polytope of a spherical variety  $X$  is described by a finite set of linear inequalities corresponding to the  $B$ -stable divisors of  $X$  which was described by Brion [19] in the non-abelian case. Let  $X$  be a spherical  $G$ -variety and  $L \rightarrow X$  a  $G$ -equivariant line bundle. First some notation: Let  $\mathbb{C}(X)$  denote the space of rational functions on  $X$ , and  $\mathbb{C}(X)^{(B)}$  the space of  $B$ -semiinvariant vectors. Let  $\Lambda_X^\vee \subset \Lambda^\vee$  denote the group of weights appearing in  $\mathbb{C}(X)^{(B)}$ . Let  $\mathcal{D}(X)$  denote the set of prime  $B$ -stable divisors of  $X$ . Each  $D \in \mathcal{D}(X)$  defines a valuation  $\mathbb{C}(X)^{(B)} \rightarrow \mathbb{Z}$  and so a vector  $v_D$  in the dual  $\Lambda_X$  of  $\Lambda_X^\vee$ . Let  $\mathbb{C}(X, L)$  denote the space of rational sections of  $L$ , and  $s \in \mathbb{C}(X, L)^{(B)}$  with weight  $\mu(s)$ . Let  $n_D(s)$  denote the order of vanishing of  $s$  at  $D$ . Consider the identification  $\mathbb{C}(X)^{(B)} \rightarrow \mathbb{C}(X, L)^{(B)}, f \mapsto fs$ . The section  $fs$  is global iff  $fs$  vanishes to at least zeroth order on each  $D \in \mathcal{D}(X)$ , iff  $f$  vanishes at least to order  $-n_D$ . Thus

**Proposition 9.3.2.** *Let  $X$  be a spherical  $G$ -variety, and  $L \rightarrow X$  a  $G$ -line-bundle. The space of weights for elements of  $\mathbb{C}(X, L)^{(B)}$  is*

$$\Delta(X, L) = \{\mu \in \Lambda_X^\vee \mid v_D(\mu) \geq -n_D(s)\} + \mu(s).$$


*Example 9.3.3.* Here is a typical application which appears in Brion [19] and seems to be due to Macdonald [78]:

*Theorem 9.3.4.* *Let  $V_\lambda$  be a simple  $GL(r)$  module with highest weight  $\lambda = (\lambda_1 \geq \dots \geq \lambda_r)$ . Then  $V_\lambda \otimes \text{Sym}(\mathbb{C}^r)$  admits a multiplicity-free decomposition into simple modules  $V_\mu$  with highest weights  $\mu = (\mu_1, \dots, \mu_r)$  satisfying*

$$\mu_1 \geq \lambda_1 \geq \mu_2 \geq \dots \mu_r \geq \lambda_r.$$

*Proof.* We prove only the case  $r = 2$ ; the general case is similar.  $V_\lambda \otimes \text{Sym}(\mathbb{C}^2)$  is isomorphic to the space of holomorphic sections of the line bundle  $\pi_1^* L_\lambda$  over  $X = \mathbb{P}^1 \times \mathbb{C}^2 = \{([w_0, w_1], (z_0, z_1))\}$ , where  $\pi_1 : \mathbb{P}^1 \times \mathbb{C}^2 \rightarrow \mathbb{P}^1$  is projection on the first factor. We take  $B$  to be the subgroup of upper-triangular invertible matrices. The  $B$ -invariant divisors are given by a single  $G$ -invariant divisor  $D_1 = \{(w, z) \mid z \in w\}$  and two  $B$ -stable divisors  $D_2 = \{w = [1, 0]\}$  and  $D_3 = \{z \in \mathbb{C} \oplus 0\}$ . The space of singular vectors  $\mathbb{C}(X)^{(B)}$  is generated by  $z_1 - w_1 z_0 / w_0$  and  $z_0$  with highest weights  $(0, 1)$  resp.  $(1, 0)$ . The  $B$ -stable divisors are defined by  $D_1 = \{z_1 / z_0 = w_1 / w_0\}$ ,  $D_2 = \{w_1 = 0\}$ ,  $D_3 = \{z_1 = 0\}$  respectively. Hence  $z_1 - w_1 z_0 / w_0$  vanishes to order 1 resp.  $-1, 0$  on  $D_1$  resp.  $D_2, D_3$ ;  $z_1$  vanishes to order 0 resp.  $0, 1$  on  $D_1$  resp.  $D_2, D_3$ . So  $v_{D_1} = (0, 1)$ ,  $v_{D_2} = (0, -1)$ ,  $v_{D_3} = (1, 0)$ . Taking  $s$  to be the section of  $\mathbb{P}^1$  with weight  $(\lambda_1, \lambda_2)$ , which vanishes to order 0 on  $D_1$ ,  $\lambda_1 - \lambda_2$  in  $D_2$ , and 0 on  $D_3$  one obtains  $n_{D_1} = 0, n_{D_2} = \lambda_1 - \lambda_2, n_{D_3} = 0$ . This yields the inequalities  $\mu_2 \geq \lambda_2, -\mu_2 \geq -\lambda_2 - (\lambda_1 - \lambda_2) = -\lambda_1, \mu_1 \geq \lambda_1$  as claimed. See Figure 9.2.  $\square$

*Remark 9.3.5.* Not every  $B$ -stable divisor defines a facet of the moment polytope. This is already apparent in the case of the Borel-Weil theorem, where for a group of rank  $r$  there are  $r$   $B$ -stable divisors (the Schubert varieties of codimension one) but the moment polytope is simply a point.


 Figure 9.2: Decomposition of  $V_\lambda \otimes \text{Sym}(\mathbb{C}^2)$  via Brion's method

Based on his work on the toric case, Delzant asked the question of whether compact multiplicity-free actions are classified by their moment polytopes and generic stabilizers, and answered the question affirmatively in the rank two case [29]. A result of Knop [64] reduces this to the question of whether affine spherical varieties are classified by their moment polytopes and generic stabilizers of the compact group actions; this conjecture has recently been proved by Losev [75], see also his review in this volume.

In the torus case we have

**Corollary 9.3.6.** *With  $X, K, \mathcal{O}_X(1)$  as above, if  $K$  is a torus then the dimension of  $H^0(X, \mathcal{O}_X(1))$  is the number of integral points  $\Delta \cap \mathfrak{k}^\vee$  of  $\Delta$ .*

The dimension of  $H^0(X, \mathcal{O}_X(k))$  can be computed by Riemann-Roch for sufficiently large  $k$ , since  $\mathcal{O}_X(1)$  is by assumption positive. This led to an interesting series of papers on formulas for the number of lattice points in a convex polytope which generalize the Euler-Maclaurin formula and were later proved combinatorially, see [21] for references.

## 10. LOCALIZATION VIA SHEAF COHOMOLOGY

In this section we review various “fixed point methods” for computing moment polytopes, in the context of sheaf cohomology. These include not only the “localization” methods which take as input fixed point data for a one-parameter subgroup, but also the “non-abelian localization” principle which uses the Kirwan-Ness stratification.

**10.1. Grothendieck's local cohomology.** A powerful technique for computing cohomology groups, and therefore for computing moment polytopes, is Grothendieck's local cohomology theory, exposed in [36] and Hartshorne [46]. Let  $X$  be a  $G$ -variety and  $Y \subset X$  a  $G$ -subvariety. Let  $E \rightarrow X$  be a  $G$ -equivariant coherent sheaf. Denote by  $\Gamma_Y(X, E)$  the group of sections whose support is contained in  $Y$ . We denote by  $H_Y^i$  the  $i$ -th derived functor of  $\Gamma_Y$ , so that the *local cohomology group*  $H_Y^i(X, E)$  is a  $G$ -module. These modules have the following properties:

**Theorem 10.1.1.** (a) *(Long Exact Sequence) There is an exact triangle*

$$\dots H_Y(X, E) \rightarrow H(X, E) \rightarrow H(X - Y, E|_{X - Y}) \rightarrow \dots$$

(b) *(Gysin isomorphism) Suppose  $Y \subset X$  is smooth. Then*

$$H_Y^j(X, E) \cong H^{j - \text{codim}(Y)}(Y, E|_Y \otimes \text{Eul}(N)^{-1})$$

where  $N$  is the normal bundle of  $Y$  in  $X$  and  $\text{Eul}(N)^{-1} := \det(N) \otimes \text{Sym}(N)$  (this is an inverse of the  $K$ -theory Euler class  $\text{Eul}(N) = \Lambda(N^\vee)$  although we do not discuss  $K$ -theory here)

(c) *(Spectral sequence associated to a stratification) Let  $X_1 \subset X_2 \subset \dots \subset X_m = X$  be a filtration of  $X$ . There is a spectral sequence*

$$\bigoplus_{i=1}^m H_{X_i - X_{i-1}}(X_i, E|_{X_i}) \implies H(X, E).$$

Let  $\chi(X, E) = \bigoplus (-1)^i H^i(X, E)$  be the Euler characteristic, considered as a virtual  $G$ -representation, and  $\chi_Y(X, E)$  the Euler characteristic of the local cohomology along  $Y$ . These will generally not be finite-dimensional, but rather in our cases of interest the multiplicity of each simple module is finite. Thus the formula below holds in the completion of the representation ring, as an immediate consequence of the spectral sequence:

**Corollary 10.1.2.** *Suppose that  $X_1 \subset \dots \subset X_m = X$  is a filtration of  $X$  such that the differences  $X_i - X_{i-1}$  are smooth with normal bundles  $N_i \rightarrow X_i - X_{i-1}$ . Then*

$$(10.1) \quad \chi(X, E) = \sum_i (-1)^{\text{codim}(X_i - X_{i-1})} \chi(X_i - X_{i-1}, E|_{X_i - X_{i-1}} \otimes \text{Eul}(N_i)^{-1})$$

if both sides are well-defined in the sense that the multiplicity of any simple module is finite.

This formula applies to various filtrations associated to group actions to give ‘‘localization’’ formulae.

*Example 10.1.3.* (Weyl character formula and Borel-Weil-Bott, c.f. Atiyah-Bott [8]) Let  $X = G/B^-$  and  $E = \mathcal{O}_X(\lambda)$  so that if  $\lambda$  is dominant then  $H^0(X, E) = V_\lambda$  by Borel-Weil 6.1.1. The Bruhat decomposition  $X = \cup_{w \in W} X_w$  gives a filtration  $X_i = \cup_{w \in W, l(w) \geq i} X_w$ . Each cell  $X_w$  fibers over  $x_w = wB/B$  with fiber  $X_w \cong M_w := \mathfrak{b} \cap \text{Ad}(w)\mathfrak{b}$ . The normal bundle  $X_w$  has restriction to  $x_w$  given by  $N_w = (\mathfrak{b}/\mathfrak{b} \cap \text{Ad}(w)\mathfrak{b})^\vee$ . The formula (10.1) gives

$$\begin{aligned} \chi(X, \mathcal{O}_X(\lambda)) &= \bigoplus_{w \in W} (-1)^{l(w)} \chi(X_w, E|_{X_w} \otimes \text{Sym}(N_w) \otimes \det(N_w)) \\ &= \bigoplus_{w \in W} (-1)^{l(w)} \chi(x_w, E \otimes \text{Sym}(N_w) \otimes \det(N_w) \otimes \text{Sym}(M_w^\vee)|_{x_w}) \\ &= \bigoplus_{w \in W} (-1)^{l(w)} \mathbb{C}_{w\lambda} \otimes \text{Sym}(\mathfrak{b}^-) \otimes \mathbb{C}_{w\rho - \rho} \end{aligned}$$

where  $\rho$  is the half-sum of positive roots. Thus its character is

$$(10.2) \quad \sum_{w \in W} (-1)^{l(w)} \frac{t^{w(\lambda + \rho) - \rho}}{\prod_{\alpha > 0} (1 - t^{-\alpha})}.$$

Thus if  $\lambda$  is dominant then

*Proposition 10.1.4. (Weyl character formula)* *The character of the action of  $T$  on  $V_\lambda$  is given by (10.2).*

In general, suppose that  $w$  is such that  $w(\lambda + \rho) - \rho$  is dominant. From the spectral sequence we see that the only contribution to  $\chi(X, \mathcal{O}_X(\lambda))$  comes from  $H^{l(w)}(X, \mathcal{O}_X(\lambda))$ , since  $l(w) = \text{codim}(X_w)$ . This is a simple  $G$ -module of highest weight  $w(\lambda + \rho) - \rho$ , since it has the same character as that of  $V_{w(\lambda + \rho) - \rho}$  by the Weyl character formula. If no such  $w$  exists, then the Fourier expansion of the character vanishes on dominant weights and is  $W$ -invariant and so  $H(X, \mathcal{O}_X(\lambda))$  is trivial. Thus:

*Proposition 10.1.5. (Borel-Weil-Bott [17])* *Let  $X = G/B^-$ .  $H^j(X, \mathcal{O}_X(\lambda)) \cong V_{w(\lambda + \rho) - \rho}$  if  $w(\lambda + \rho) - \rho$  is dominant for some (unique)  $w \in W$  and  $j = l(w)$ , and is zero otherwise.*

**10.2. One-parameter localization.** The derivation of the Weyl character formula given in the previous section generalizes to varieties with circle actions as follows. Let  $X$  be a compact  $G \times \mathbb{C}^*$ -variety, and  $X^{\mathbb{C}^*}$  its  $\mathbb{C}^*$ -fixed point set. Let  $\mathcal{F}$  be the set of components of  $X^{\mathbb{C}^*} = \{x \in X \mid zx = x \forall z \in \mathbb{C}^*\}$ . For each  $F \in \mathcal{F}$ , define

$$X_F := \{x \in X \mid \lim_{z \rightarrow 0} zx \in F\}.$$

Let  $N_F$  denote the normal bundle of  $F$  in  $X$ . It admits a decomposition  $N_F = N_F^+ \oplus N_F^-$  into positive and negative weight spaces for the  $\mathbb{C}^*$ -action.

**Proposition 10.2.1.** *(Bialynicki-Birula decomposition [15])* *Suppose that  $X$  is smooth. Then each  $X_F$  is a smooth  $G \times \mathbb{C}^*$ -stable subvariety, equipped with a morphism  $\pi_F : X_F \rightarrow F$ ,  $x \mapsto \lim_{z \rightarrow 0} zx$  which induces on  $X_F$  the structure of a vector bundle whose fibers are isomorphic to the fibers of the normal bundle  $N_F^+ \rightarrow F$  of  $F$  in  $X$ .*

By filtering by the dimension of  $N_F^+$ , applying the localization formula (10.1), and pushing forward with  $\pi_F$  one obtains

**Theorem 10.2.2** (Localization for one-parameter subgroups). *Let  $E \rightarrow X$  be any  $G \times \mathbb{C}^*$ -equivariant coherent sheaf. Then*

$$\chi(X, E) = \sum_{F \subset X^{\mathbb{C}^*}} \chi(F, E|_F \otimes \text{Sym}(N_F^{+, \vee}) \otimes \text{Sym}(N_F^-) \otimes \det(N_F^-))$$

in the completion of the representation ring of  $G$ .

One could equally well choose the stratification for the inverted  $\mathbb{C}^*$ -action, which would lead to the same formula with  $N_F^+, N_F^-$  inverted. In the equivariant cohomology literature such a choice of direction is called a choice of *action chamber*, see Duistermaat [31].

The spectral sequence contains more information than the localization formula, namely, information about the individual cohomology groups. For example,

*Example 10.2.3.* Let  $X = \mathbb{P}^2$  equipped with the  $G = (\mathbb{C}^*)^2$  action by  $(g_1, g_2)[z_0, z_1, z_2] = [z_0, g_1^{-1}z_1, g_2^{-1}z_2]$ . Then  $H^0(X, \mathcal{O}_X(d))$  is spanned by homogeneous polynomials of degree  $d$ . Its Euler characteristic has character

$$(\chi(X, \mathcal{O}_X(d)))(g) = \sum_{d_1+d_2 \leq d, d_1, d_2 \geq 0} g_1^{d_1} g_2^{d_2}.$$

One can also see this easily from the localization formula, which gives (for the  $\mathbb{C}^*$ -action induced by the map  $z \mapsto (z, z^2)$ ) three fixed points with normal weights  $(1, 0)$ ,  $(0, 1)$ , resp.  $(-1, 0)$ ,  $(-1, 1)$  resp.  $(1, -1)$ ,  $(0, -1)$  and so

$$(10.3) \quad (\chi(X, \mathcal{O}_X(d)))(g) = (1 - g_1)^{-1}(1 - g_2)^{-1} - g_1^{d+1}(1 - g_1)^{-1}(1 - g_1^{-1}g_2)^{-1} \\ + g_2^{d+1}g_1^{-1}(1 - g_1^{-1}g_2)^{-1}(1 - g_2)^{-1}.$$

Now suppose that  $X'$  is the blow-up of  $X$  at  $[1, 0, 0]$ . Let  $\pi : X' \rightarrow X$  denote the projection,  $\mathcal{O}_{X'}(d, e) = \pi^*\mathcal{O}_X(d) \otimes E^e$ . The action of  $\mathbb{C}^*$  on  $X'$  has four fixed points (the point at  $[1, 0, 0]$  is replaced by two fixed points in the exceptional divisor with fiber weights  $(e, 0)$ ,  $(0, e)$ ). Hence

$$(10.4) \quad (\chi(X', \mathcal{O}_{X'}(d, e)))(g) = g_1^e(1 - g_1)^{-1}(1 - g_1^{-1}g_2)^{-1} \\ - g_2^{e+1}g_1^{-1}(1 - g_1g_2^{-1})^{-1}(1 - g_2)^{-1} - g_1^d(1 - g_1)^{-1}(1 - g_1^{-1}g_2)^{-1} \\ + g_2^d(1 - g_1^{-1}g_2)^{-1}(1 - g_2)^{-1}.$$

Its Fourier transform is shown below in Figure 10.1. The contributions with weights  $g_1^e$  contributes




Figure 10.1: Euler characteristic of a line bundle on blow-up of  $\mathbb{P}^2$

only to  $H^0$ , while the contribution with weight  $g_2^{e+1}g_1^{-1}$  contributes only to  $H^1$ . The former is the only term whose Fourier transform has support in the larger triangle, while the latter is the only term whose Fourier transform has support in the smaller. Hence the dots in the smaller triangle correspond to vectors in  $H^1$  while those in the larger correspond to  $H^0$ . Very similar results are obtained by a deformation method introduced by Witten [110], and studied by a number of other authors since then, see for example [114].



**10.3. Localization via orbit stratification.** Other stratifications lead to interesting but less well-known localization formulae. For example, suppose that  $G$  acts on  $X$  with only finitely many orbits  $Y$ . We then obtain a formula

$$\chi(X, E) = \sum_{Y \subset X} (-1)^{\text{codim}(Y)} \chi(Y, E|_Y \otimes \text{Eul}(Y)^{-1})$$

assuming that each simple module appears with finite multiplicity as before. In particular, suppose that  $X$  is a toric variety and  $E = \mathcal{O}_X(1)$  a polarization. Indexing the orbits  $Y_F$  by faces  $F$  of the moment polytope  $\Delta$  we see that

$$\chi(Y, E|_Y \otimes \text{Eul}(Y)^{-1}) = \sum_{\mu \in \Delta^\vee \cap C_F} t^\mu \det(N_F)$$

where the sum is over  $\mu$  is the outward normal cone  $C_F$  to  $\Delta$  at  $F$ , and  $\det(N_F)$  is the determinant  $N_F$  of the normal bundle to  $Y_F$ . This is closely related to the *Brianchon-Gram formula*: for any convex polytope  $\Delta$ ,

$$\chi_\Delta = \sum (-1)^{\text{codim}(F)} \chi_{C_F}$$

where  $\chi_{C_F}$  is the characteristic function of  $C_F$  [98].

**10.4. Non-abelian localization.** Let  $X$  be a polarized smooth  $G$ -variety and  $E \rightarrow X$  a  $G$ -equivariant coherent sheaf. Combining the Kirwan-Hessellink-Ness stratification with the Euler characteristic formula (10.1) gives

$$\chi(X, E) = \sum_{\lambda} \chi(X_\lambda, E|_{X_\lambda} \otimes \text{Eul}(N_{X_\lambda})^{-1})$$

where the sum is over types  $\lambda$  or equivalently critical sets for the norm-square of the moment map. This is a sheaf cohomology version of a “non-abelian localization principle” suggested by Witten in the setting of equivariant de Rham cohomology [109]. In fact, this terminology in the sheaf cohomology setting is somewhat confusing: the formula is already quite interesting in the abelian case (non-abelian should read “not necessarily abelian”) and the formula is not really a localization formula, since there is a contribution from the (dense) open stratum. Since  $X_\lambda = G \times_{P_\lambda} Y_\lambda^{\text{ss}}$ , we have

$$\chi(X_\lambda, E|_{X_\lambda} \otimes \text{Eul}(N_{X_\lambda})^{-1}) = \text{Ind}_{G_\lambda}^G \chi(Y_\lambda^{\text{ss}}, E|_{Y_\lambda^{\text{ss}}} \otimes \text{Eul}(N_{X_\lambda}|_{Y_\lambda^{\text{ss}}})^{-1}).$$

(Here  $\text{Ind}$  denotes holomorphic induction, that is, if  $V$  is a  $G_\lambda$ -module then  $\text{Ind}_{G_\lambda}^G(V) = \chi(G \times_{P_\lambda}^- V)$ .) Since  $Y_\lambda^{\text{ss}}$  fibers over  $Z_\lambda^{\text{ss}}$  with affine fibers,

$$(10.5) \quad \begin{aligned} \chi(Y_\lambda^{\text{ss}}, E|_{Y_\lambda^{\text{ss}}} \otimes \text{Eul}(N_{X_\lambda}|_{Y_\lambda^{\text{ss}}})^{-1}) \\ = \chi(Z_\lambda^{\text{ss}}, E|_{Z_\lambda^{\text{ss}}} \otimes \text{Sym}(N_{X_\lambda} X|_{Z_\lambda^{\text{ss}}}) \otimes \det(N_{X_\lambda} X|_{Z_\lambda^{\text{ss}}}) \otimes \text{Sym}(N_{Z_\lambda}^{\text{ss}} Y_\lambda^{\text{ss}})^\vee). \end{aligned}$$

This can be put into a more understandable form if we recognize that  $N_{X_\lambda} X|_{Z_\lambda^{\text{ss}}}$  resp.  $N_{Z_\lambda}^{\text{ss}} Y_\lambda^{\text{ss}}$  is the positive resp. negative part of the normal bundle of  $Z_\lambda^{\text{ss}}$  in  $Y_\lambda^{\text{ss}}$ . One obtains a formula due to Teleman [104] in the algebraic case and Paradan [89] in the general symplectic setting; the latter proof uses techniques of transversally elliptic operators:

**Theorem 10.4.1.**

$$\chi(X, E) = \sum_{\lambda} \text{Ind}_{G_\lambda}^G (\chi(Z_\lambda^{\text{ss}}, E|_{Z_\lambda^{\text{ss}}} \otimes \text{Eul}(N_{Z_\lambda}^{\text{ss}} Y_\lambda^{\text{ss}})_+^{-1}))$$

where the  $+$  indicates the particular choice of (formal) inverse to the  $K$ -theory Euler class given in the previous formula.

*Example 10.4.2.* Let  $X = \mathbb{P}^1$  and  $E = \mathcal{O}(d)$  so  $\chi(X, E)$  has character  $z^{-d} + z^{-d+2} + \dots + z^d$ . The stratification  $\mathbb{P}^1 = \{0\} \cup \mathbb{C}^* \cup \{\infty\}$  leads to the formula

$$z^{-d} + \dots + z^d = \left( \sum_{n \in \mathbb{Z}} z^{d+2n} \right) - z^{d+2}/(1 - z^2) - z^{-d-2}/(1 - z^{-2}).$$

*Example 10.4.3.* We describe the non-abelian localization formula for the action of  $SL(3, \mathbb{C})$  on a partial flag variety for the exceptional group of type  $G_2$ , corresponding to the decomposition of a simple  $G_2$ -module into  $SL(3, \mathbb{C})$ -modules. Let  $\omega_1, \omega_2$  denote the fundamental weights for  $SL(3, \mathbb{C})$ . The dual positive Weyl chamber for  $G_2$  is the span of  $\omega_1$  and  $\omega_1 + \omega_2$ . Let  $P_{\omega_1 + \omega_2}$  denote the maximal parabolic of  $G_2$  corresponding to  $\omega_1 + \omega_2$ , and  $X = G_2/P_{\omega_1 + \omega_2}^-$ , that is, the coadjoint orbit through  $\omega_1 + \omega_2$ . The action is spherical and moment polytope the convex hull of  $\omega_1, \omega_2, \omega_1 + \omega_2$ . We leave the computation of the moment polytope to the reader; it can be computed using one-parameter localization. By Borel-Weil and the computation of the moment polytope,

$$\chi(\mathcal{O}_X(k)) = \sum_{\lambda \in k\Delta \cap Q} \chi_\lambda = \text{Res}_{SL(3, \mathbb{C})}^{G_2}(\chi_{k(\omega_1 + \omega_2)})$$

the character of the irreducible  $G_2$ -representation with highest weight  $k(\omega_1 + \omega_2)$ , restricted to  $SL(3, \mathbb{C})$ ; here  $Q$  is the lattice generated by the long roots shifted by  $k(\omega_1 + \omega_2)$ .

We compute the Kirwan-Ness stratification as follows. Let  $F_1$  be the open face connecting  $\omega_2, \omega_1 + \omega_2$ ,  $F_2$  the open face connecting  $\omega_1, \omega_1 + \omega_2$ , and  $F_3$  the open face connecting  $\omega_1, \omega_2$ . Let  $F_{ij} = F_i \cap F_j$ . The inverse image  $\Phi^{-1}(F_{12})$  contains a unique point,  $x_1$ , which is  $T$ -fixed. None of the other  $T$ -fixed points map to  $\mathfrak{t}_+^\vee$ . Therefore, the remaining points in  $\Phi^{-1}(\text{int}(\mathfrak{t}_+^\vee))$  (the interior of the positive Weyl chamber) have one-dimensional stabilizers. Since  $\Phi^{-1}(\text{int}(\mathfrak{t}_+^\vee))$  has dimension  $2 \dim(T)$ , it is a multiplicity free action, so the inverse image of any face  $F \subset \text{int} \mathfrak{t}_+^\vee$  has infinitesimal stabilizer the annihilator of the tangent space of  $F$ . The stabilizers of the faces  $F_1, F_2, F_3$  are

$$\mathfrak{t}_1 = \text{span}(h_1), \mathfrak{t}_2 = \text{span}(h_2), \mathfrak{t}_3 = \text{span}(h_3)$$

where  $h_1, h_2, h_3$  are the coroots of  $SL(3, \mathbb{C})$ . The level set  $\Phi^{-1}((\omega_1 + \omega_2)/2)$  is a critical set of  $\phi$  with type  $\lambda = ((\omega_1 + \omega_2)/2)$ . The fixed point component  $Z_\xi$  has moment image  $\Phi(Z_\xi) =$




Figure 10.2: Critical values of the norm-square of the moment map for  $X = G_2/P_{\omega_1 + \omega_2}$

$\text{hull}(2\omega_2 - \omega_1, 2\omega_1 - \omega_2)$ . The unstable manifold  $Y_\xi$  has image under the moment map for  $T$  (that is, for the maximal torus of the compact group  $SU(3)$ )

$$\pi_T^G \Phi(\overline{Y_\xi}) = \text{hull}(2\omega_2 - \omega_1, 2\omega_1 - \omega_2, \omega_1 + \omega_2).$$

None of the other facets  $F_j$  contain points  $\xi$  with  $\xi \in \mathfrak{t}_j$ . Therefore, there are no other critical points of  $\phi$  in  $\Phi^{-1}(\text{int}(\mathfrak{t}_+^\vee))$ . Finally consider the inverse image of the vertices  $F_{13}, F_{23}$ . Any  $x \in \Phi^{-1}(F_{jk})$  has  $G_x \neq T$ , hence  $G_x$  cannot intersect the semisimple part  $[G_{\Phi(x)}, G_{\Phi(x)}]$ . Therefore,  $G_x$  is one-dimensional. Let  $Z_x$  denote the fixed point component of  $G_x$  containing  $x$ . Since  $G_x$  is one-dimensional, the image  $\Phi(Z)$  is codimension one, and so meets  $\Phi^{-1}(\text{int}(\mathfrak{t}_+^\vee))$ . But this implies that the  $\mathfrak{g}_x$  is conjugate to either  $\mathfrak{t}_j$  or  $\mathfrak{t}_k$ , and so  $\mathfrak{g}_x$  cannot equal the span of  $F_{jk}$ . Therefore, set of types for the action is  $\{\omega_1 + \omega_2, \frac{1}{2}(\omega_1 + \omega_2)\}$ . (In fact the Kirwan-Ness stratification coincides with the orbit stratification for  $G_{\mathbb{C}}$ . That is,  $X$  is a two-orbit variety, with one open orbit and one of complex codimension two [33].)

We now compute the contributions from the Kirwan-Ness strata. For  $\xi = \omega_1 + \omega_2$ ,  $Z_\xi^{\text{ss}}$  is equal to a point, and the bundle  $N_\xi$  is the representation with weights  $\beta_5, \beta_6$ . Hence

$$\chi_{G_\xi}(Z_\xi^{\text{ss}}, E \otimes \text{Eul}(N_\xi)_+^{-1}) = \sum_{(\lambda, \alpha_1) > k, (\lambda, \alpha_2) > k} z^\lambda.$$

Its induction to  $G$  is

$$\text{Ind}_{G_\xi}^G \chi_{G_\xi}(Z_\xi^{\text{ss}}, E \otimes \text{Eul}(N_\xi)_+^{-1}) = \sum_{(\lambda, \alpha_1) > k, (\lambda, \alpha_2) > k} \chi_\lambda.$$

For  $\xi = (\omega_1 + \omega_2)/2$ , we have  $Z_\xi^{\text{ss}} \cong \mathbb{C}^*$  and  $N_\xi$  trivial. Therefore,

$$\chi_{G_\xi}(Z_\xi^{\text{ss}}, E \otimes \text{Eul}(N_\xi)_+^{-1}) = \sum_{(\lambda, \xi) \geq k(\xi, \xi)} z^\lambda$$

where the sum is over vectors  $\lambda$  such that  $\lambda - k(\omega_1 + \omega_2)$  is in some lattice  $\Lambda_1^\vee$ , and satisfying the inequality above. Hence

$$\text{Ind}_{G_\xi}^G (\chi_{G_\xi}(Z_\xi^{\text{ss}}, E \otimes \text{Eul}(N_\xi)_+^{-1})) = \sum_{\lambda \in k\Delta} \chi_\lambda - \sum_{(\lambda, \alpha_1) > k, (\lambda, \alpha_2) > k} \chi_\lambda.$$

Since the contributions from  $\xi = (\omega_1 + \omega_2), \frac{1}{2}(\omega_1 + \omega_2)$  must have finite sum, the lattice  $\Lambda_1^\vee$  must be the long root lattice. The contribution (for  $k = 6$ ) is shown in Figure 10.3.




Figure 10.3:  $\text{Ind}_T^G \chi_{Z_{(\omega_1 + \omega_2)/2}^{\text{ss}}, T}(E)$

The positive contribution of the open stratum is finite (6 representations, for  $k = 6$ ) and the negative contribution infinite, that is  $\dim(H^{\text{odd}}(M_\xi, L^k)) = \infty$ , for any  $k$ . One can show that the higher cohomology lies in  $H^1$ , using the spectral sequence. The sum of the contributions is  $\chi(\mathcal{O}_X(k)) = \sum_{\lambda \in k\Delta} \chi_\lambda$  as claimed. This completes the example.

Taking invariants in Theorem 10.4.1 gives a formula expressing the difference between  $\chi(X, E)^G$  and  $\chi(X//G, E//G)$ :

**Theorem 10.4.4.**

$$\chi(X, E)^G - \chi(X//G, E//G) = \sum_{\lambda \neq 0} \chi(Z_\lambda^{\text{ss}}, E|_{Z_\lambda^{\text{ss}}} \otimes \text{Eul}(N_{Z_\lambda^{\text{ss}}} Y_\lambda^{\text{ss}})^{-1} \otimes \text{Eul}(\mathfrak{g}/\mathfrak{p}_\lambda^-))^{G_\lambda}$$

The spectral sequence also contains information about the individual cohomology groups. For example, let  $\mathbb{C}_\lambda^* \subset G_\lambda$  denote the one-parameter subgroup generated by  $\lambda$ . The weight of  $\mathbb{C}_\lambda^*$  on  $\det(N_{X_\lambda} X|_{Z_\lambda^{\text{ss}}})$  is positive, if  $\lambda$  is non-trivial. Indeed,  $N_{X_\lambda} X|_{Z_\lambda^{\text{ss}}}$  is the negative part of the tangent bundle. Furthermore,  $\mathfrak{g}/\mathfrak{p}_\lambda^-$  has positive weights under  $\mathbb{C}_\lambda^*$ . Thus

**Corollary 10.4.5** (Teleman [104]). *Suppose that the weights of  $\mathbb{C}_\lambda^*$  on  $E|_{Z_\lambda}$  are positive for all types  $\lambda$ . (This is automatically the case if  $E = \mathcal{O}_X(d)$  is the  $d$ -th tensor product of a polarization  $\mathcal{O}_X(1)$  of  $X$ ). Then  $H^j(X, E)^G = H^j(X//G, E//G)$  for all  $j$ .*

In particular, if the higher cohomology of  $E$  vanishes then so does that of  $E//G$ .

The index maps naturally induce a diagram in  $K$ -theory

$$\begin{array}{ccc} K_G(X) & \longrightarrow & K(X//G) \\ & \searrow & \swarrow \\ & \mathbb{Z} & \end{array}$$

which fails to commute by the above explicit sum of fixed point contributions for one-parameter subgroups. There are similar results in the equivariant cohomology of  $X$  due to Paradan [88] and the author [113], based on earlier work of Witten [109]: a natural diagram of equivariant cohomology groups

$$\begin{array}{ccc} H_G(X) & \longrightarrow & H(X//G) \\ & \searrow & \swarrow \\ & \mathbb{R} & \end{array}$$

fails to commute by an explicit sum of fixed point contributions from one-parameter subgroups. The first explicit version of non-abelian localization is due to Jeffrey-Kirwan [57], and expresses the difference as a sum over certain fixed point sets of the maximal torus. The versions of Paradan, myself [113], and Beasley-Witten [11] express the difference as a sum over critical points of the norm-square of the moment map. The left hand arrow in the diagram above takes some work to define: morally speaking it is defined by  $\alpha \mapsto \int_{X \times \mathfrak{g}} \alpha$ , but this is not well-defined for polynomial equivariant classes. Rather, the left-hand side must be defined by a suitable limit procedure, either by taking the leading term in Riemann-Roch, or (in the context of equivariant de Rham cohomology with smooth coefficients) shifting by equivariant Liouville form and taking the zero limit of the shift, see [113]. From this point of view, the  $K$ -theory approach is more natural.

REFERENCES

- [1] R. Abraham and J. Marsden. *Foundations of Mechanics*. Benjamin/Cummings, Reading, 1978.
- [2] S. Agnihotri and C. Woodward. Eigenvalues of products of unitary matrices and quantum Schubert calculus. *Math. Res. Lett.*, 5(6):817–836, 1998.
- [3] D. N. Akhiezer. *Lie group actions in complex analysis*. Aspects of Mathematics, E27. Friedr. Vieweg & Sohn, Braunschweig, 1995.
- [4] Werner Ballmann, Mikhael Gromov, and Viktor Schroeder. *Manifolds of nonpositive curvature*, volume 61 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1985.
- [5] B. H. Bowditch. A class of incomplete non-positively curved manifolds. *Pacific J. Math.*, 172(1):1–39, 1996.
- [6] J. Arms, R. Cushman, and M. Gotay. A universal reduction procedure for Hamiltonian group actions. In T. Ratiu, editor, *The Geometry of Hamiltonian Systems*, volume 22 of *Mathematical Sciences Research Institute Publications*, Berkeley, 1989, 1991. Springer-Verlag, Berlin-Heidelberg-New York.
- [7] M. F. Atiyah. Convexity and commuting Hamiltonians. *Bull. London Math. Soc.*, 14:1–15, 1982.
- [8] M. F. Atiyah and R. Bott. A Lefschetz fixed point formula for elliptic complexes. II. Applications. *Ann. of Math. (2)*, 88:451–491, 1968.
- [9] M. F. Atiyah and R. Bott. The moment map and equivariant cohomology. *Topology*, 23(1):1–28, 1984.
- [10] M. Audin. *The Topology of Torus Actions on Symplectic Manifolds*, volume 93 of *Progress in Mathematics*. Birkhäuser, Boston, 1991.
- [11] C. Beasley and E. Witten. Non-abelian localization for Chern-Simons theory. *J. Differential Geom.*, 70(2):183–323, 2005.
- [12] P. Belkale. Local systems on  $\mathbb{P}^1 - S$  for  $S$  a finite set. *Compositio Math.*, 129(1):67–86, 2001.

- [13] P. Belkale and S. Kumar. Eigenvalue problem and a new product in cohomology of flag varieties. *Invent. Math.*, 166(1):185–228, 2006.
- [14] A. Berenstein and R. Sjamaar. Coadjoint orbits, moment polytopes, and the Hilbert-Mumford criterion. *J. Amer. Math. Soc.*, 13(2):433–466 (electronic), 2000.
- [15] A. Bialynicki-Birula. Some theorems on actions of algebraic groups. *Ann. of Math. (2)*, 98:480–497, 1973.
- [16] A. M. Bloch and T. S. Ratiu. Convexity and integrability. In *Symplectic geometry and mathematical physics (Aix-en-Provence, 1990)*, volume 99 of *Progr. Math.*, pages 48–79. Birkhäuser Boston, Boston, MA, 1991.
- [17] R. Bott. Homogeneous vector bundles. *Ann. of Math. (2)*, 66:203–248, 1957.
- [18] M. Brion. Sur l’image de l’application moment. In M.-P. Malliavin, editor, *Séminaire d’algèbre Paul Dubreuil et Marie-Paule Malliavin*, volume 1296 of *Lecture Notes in Mathematics*, pages 177–192, Paris, 1986, 1987. Springer-Verlag, Berlin-Heidelberg-New York.
- [19] M. Brion. Groupe de Picard et nombres caractéristiques des variétés sphériques. *Duke Math. J.*, 58(2):397–424, 1989.
- [20] M. Brion, D. Luna, and Th. Vust. Espaces homogènes sphériques. *Invent. Math.*, 84:617–632, 1986.
- [21] M. Brion and M. Vergne. Lattice points in simple polytopes. *J. Amer. Math. Soc.*, 10:371–392, 1997.
- [22] L. Bruasse and A. Teleman. Harder-Narasimhan filtrations and optimal destabilizing vectors in complex geometry. *Ann. Inst. Fourier (Grenoble)*, 55(3):1017–1053, 2005.
- [23] A. Cannas da Silva. *Introduction to symplectic and Hamiltonian geometry*. Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2003.
- [24] Pierre-Emmanuel Caprace and Alexander Lytchak. At infinity of finite-dimensional CAT(0) spaces. *Math. Ann.*, 346(1):1–21, 2010.
- [25] J. S. Carter, D. E. Flath, and M. Saito. *The classical and quantum 6j-symbols*. Princeton University Press, Princeton, NJ, 1995.
- [26] X. Chen and S. Sun. Calabi flow, Geodesic rays, and uniqueness of constant scalar curvature Kähler metrics. arXiv:1004.2012.
- [27] D. A. Cox. The homogeneous coordinate ring of a toric variety. *J. Algebraic Geom.*, 4(1):17–50, 1995.
- [28] T. Delzant. Hamiltoniens périodiques et images convexes de l’application moment. *Bull. Soc. Math. France*, 116:315–339, 1988.
- [29] T. Delzant. Classification des actions Hamiltoniennes des groupes de rang 2. *Ann. Global Anal. Geom.*, 8(1):87–112, 1990.
- [30] S. K. Donaldson and P. Kronheimer. *The geometry of four-manifolds*. Oxford Mathematical Monographs. Oxford University Press, New York, 1990.
- [31] J. J. Duistermaat. Equivariant cohomology and stationary phase. In *Symplectic geometry and quantization, (Sanda and Yokohama, 1993)*, volume 179 of *Contemp. Math.*, pages 45–62, Providence, RI, 1994. Amer. Math. Soc.
- [32] P. B. Eberlein. *Geometry of nonpositively curved manifolds*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1996.
- [33] D. Feldmüller. Two-orbit varieties with smaller orbit of codimension two. *Arch. Math. (Basel)*, 54(6):582–593, 1990.
- [34] W. Fulton. *Introduction to Toric Varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, 1993.
- [35] W. Fulton. Eigenvalues, invariant factors, highest weights, and Schubert calculus. *Bull. Amer. Math. Soc. (N.S.)*, 37(3):209–249 (electronic), 2000.
- [36] A. Grothendieck. *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 4. Société Mathématique de France, Paris, 2005. Séminaire de Géométrie Algébrique du Bois Marie, 1962, Augmenté d’un exposé de Michèle Raynaud. [With an exposé by Michèle Raynaud], With a preface and edited by Yves Laszlo, Revised reprint of the 1968 French original.
- [37] V. Guillemin and S. Sternberg. Convexity properties of the moment mapping. *Invent. Math.*, 67:491–513, 1982.
- [38] V. Guillemin and S. Sternberg. Geometric quantization and multiplicities of group representations. *Invent. Math.*, 67:515–538, 1982.
- [39] V. Guillemin and S. Sternberg. Homogeneous quantization and multiplicities of group representations. *J. Funct. Anal.*, 47:344–380, 1982.
- [40] V. Guillemin and S. Sternberg. *Geometric Asymptotics*, volume 14 of *Mathematical Surveys and Monographs*. Amer. Math. Soc., Providence, R. I., revised edition, 1990.
- [41] V. Guillemin and S. Sternberg. *Symplectic Techniques in Physics*. Cambridge Univ. Press, Cambridge, 1990.
- [42] V. W. Guillemin and S. Sternberg. *Supersymmetry and equivariant de Rham theory*. Springer-Verlag, Berlin, 1999. With an appendix containing two reprints by Henri Cartan [MR 13,107e; MR 13,107f].
- [43] V. Guillemin. Kaehler structures on toric varieties. *J. Differential Geom.*, 40(2):285–309, 1994.
- [44] V. Guillemin and R. Sjamaar. Convexity theorems for varieties invariant under a Borel subgroup. *Pure Appl. Math. Q.*, 2(3, part 1):637–653, 2006.
- [45] V. Guillemin and S. Sternberg. Multiplicity-free spaces. *J. Differential Geom.*, 19(1):31–56, 1984.
- [46] R. Hartshorne. *Residues and duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin, 1966.

- [47] J.-C. Hausmann and A. Knutson. The cohomology ring of polygon spaces. *Ann. Inst. Fourier (Grenoble)*, 48(1):281–321, 1998.
- [48] G. J. Heckman. Projections of orbits and asymptotic behavior of multiplicities for compact Lie groups. *Invent. Math.*, 67:333–356, 1982.
- [49] P. Heinzner and F. Loose. Reduction of complex Hamiltonian  $G$ -spaces. *Geom. Funct. Anal.*, 4(3):288–297, 1994.
- [50] P. Heinzner and A. Huckleberry. Kählerian structures on symplectic reductions. In *Complex analysis and algebraic geometry*, pages 225–253. de Gruyter, Berlin, 2000.
- [51] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*. Academic Press, New York, 1978.
- [52] Wim H. Hesselink. Uniform instability in reductive groups. *J. Reine Angew. Math.*, 303/304:74–96, 1978.
- [53] Wim H. Hesselink. Desingularizations of varieties of nullforms. *Invent. Math.*, 55(2):141–163, 1979.
- [54] A. Horn. Doubly stochastic matrices and the diagonal of a rotation matrix. *Amer. J. Math.*, 76:620–630, 1954.
- [55] A. Horn. Eigenvalues of sums of Hermitian matrices. *Pacific J. Math.*, 12:225–241, 1962.
- [56] Ignasi Mundet i Riera. A Hilbert–Mumford criterion for polystability in Kaehler geometry, 2008. arXiv.org:0804.1067.
- [57] L. C. Jeffrey and F. C. Kirwan. Localization for nonabelian group actions. *Topology*, 34:291–327, 1995.
- [58] V. A. Kaimanovich. Lyapunov exponents, symmetric spaces and a multiplicative ergodic theorem for semisimple Lie groups. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 164(Differentsialnaya Geom. Gruppy Li i Mekh. IX):29–46, 196–197, 1987.
- [59] M. Kapovich, B. Leeb, and J. Millson. Convex functions on symmetric spaces, side lengths of polygons and the stability inequalities for weighted configurations at infinity. *J. Differential Geom.*, 81(2):297–354, 2009.
- [60] G. Kempf and L. Ness. The length of vectors in representation spaces. In K. Lonsted, editor, *Algebraic Geometry*, volume 732 of *Lecture Notes in Mathematics*, pages 233–244, Copenhagen, 1978, 1979. Springer-Verlag, Berlin-Heidelberg-New York.
- [61] F. C. Kirwan. *Cohomology of Quotients in Symplectic and Algebraic Geometry*, volume 31 of *Mathematical Notes*. Princeton Univ. Press, Princeton, 1984.
- [62] F. C. Kirwan. Convexity properties of the moment mapping, III. *Invent. Math.*, 77:547–552, 1984.
- [63] A. A. Klyachko. Equivariant vector bundles on toric varieties and some problems of linear algebra. In *Topics in algebra, Part 2 (Warsaw, 1988)*, pages 345–355. PWN, Warsaw, 1990.
- [64] F. Knop. Automorphisms of multiplicity free Hamiltonian manifolds. arXiv:1002.4256.
- [65] F. Knop. The Luna-Vust theory of spherical embeddings. In *Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989)*, pages 225–249, Madras, 1991. Manoj Prakashan.
- [66] A. Knutson and T. Tao. The honeycomb model of  $\mathfrak{gl}_n(\mathbb{C})$  tensor products. I. Proof of the saturation conjecture. *J. Amer. Math. Soc.*, 12(4):1055–1090, 1999.
- [67] A. Knutson and T. Tao. Honeycombs and sums of Hermitian matrices. *Notices Amer. Math. Soc.*, 48(2):175–186, 2001.
- [68] A. Knutson, T. Tao, and C. Woodward. The honeycomb model of  $GL_n(\mathbb{C})$  tensor products. II. Puzzles determine facets of the Littlewood–Richardson cone. *J. Amer. Math. Soc.*, 17(1):19–48 (electronic), 2004.
- [69] A. Knutson, T. Tao, and C. Woodward. A positive proof of the Littlewood–Richardson rule using the octahedron recurrence. *Electron. J. Combin.*, 11(1):Research Paper 61, 18 pp. (electronic), 2004.
- [70] B. Kostant. Quantization and unitary representations. In C. T. Taam, editor, *Lectures in Modern Analysis and Applications III*, volume 170 of *Lecture Notes in Mathematics*, pages 87–208, Washington, D.C., 1970. Springer-Verlag, Berlin-Heidelberg-New York.
- [71] Bertram Kostant. On convexity, the Weyl group and the Iwasawa decomposition. *Ann. Sci. École Norm. Sup. (4)*, 6:413–455 (1974), 1973.
- [72] E. Lerman. Symplectic cuts. *Math. Res. Letters*, 2:247–258, 1995.
- [73] E. Lerman, E. Meinrenken, S. Tolman, and C. Woodward. Non-abelian convexity by symplectic cuts. *Topology*, 37:245–259, 1998.
- [74] E. Lerman. Gradient flow of the norm squared of a moment map. *Enseign. Math. (2)*, 51(1-2):117–127, 2005.
- [75] I. V. Losev. Proof of the Knop conjecture. *Ann. Inst. Fourier (Grenoble)*, 59(3):1105–1134, 2009.
- [76] D. Luna. Slices étales. *Sur les groupes algébriques, Mém. Soc. Math. France*, 33:81–105, 1973.
- [77] D. Luna and Th. Vust. Plongements d’espaces homogènes. *Comment. Math. Helv.*, 58(2):186–245, 1983.
- [78] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford University Press, New York, 1995. With contributions by A. Zelevinsky.
- [79] J. Marsden and A. Weinstein. Reduction of symplectic manifolds with symmetry. *Rep. Math. Phys.*, 5:121–130, 1974.
- [80] E. Meinrenken. Symplectic surgery and the Spin<sup>c</sup>-Dirac operator. *Adv. in Math.*, 134:240–277, 1998.
- [81] K. Meyer. Symmetries and integrals in mathematics. In M. M. Peixoto, editor, *Dynamical Systems*, Univ. of Bahia, 1971, 1973. Academic Press, New York.
- [82] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric Invariant Theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete, 2. Folge*. Springer-Verlag, Berlin-Heidelberg-New York, third edition, 1994.
- [83] M. S. Narasimhan and C. S. Seshadri. Stable and unitary vector bundles on a compact Riemann surface. *Ann. of Math. (2)*, 82:540–567, 1965.
- [84] L. Ness. A stratification of the null cone via the moment map. *Amer. J. Math.*, 106(6):1281–1329, 1984. with an appendix by D. Mumford.

- [85] P. E. Newstead. *Introduction to moduli problems and orbit spaces*, volume 51 of *Tata Institute of Fundamental Research Lectures on Mathematics and Physics*. Tata Institute of Fundamental Research, Bombay, 1978.
- [86] P. E. Newstead. Geometric invariant theory. In *Moduli spaces and vector bundles*, volume 359 of *London Math. Soc. Lecture Note Ser.*, pages 99–127. Cambridge Univ. Press, Cambridge, 2009.
- [87] T. Oda. *Convex bodies and algebraic geometry*, volume 15 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1988. An introduction to the theory of toric varieties, Translated from the Japanese.
- [88] P.-E. Paradan. The moment map and equivariant cohomology with generalized coefficients. *Topology*, 39(2):401–444, 2000.
- [89] P.-E. Paradan. Localization of the Riemann-Roch character. *J. Funct. Anal.*, 187(2):442–509, 2001.
- [90] S. Ramanan and A. Ramanathan. Some remarks on the instability flag. *Tohoku Math. J. (2)*, 36(2):269–291, 1984.
- [91] N. Ressayre. Geometric invariant theory and generalized eigenvalue problem. arXiv:0704.2127.
- [92] J. Roberts. Asymptotics and 6j-symbols. *Geom. Topol. Monogr.*, 4:245–261, 2002. math.QA/0201177.
- [93] A. H. W. Schmitt. *Geometric invariant theory and decorated principal bundles*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [94] I. Schur. über eine klasse von mittelbindungen mit anwendungen auf der determinanten theorie. *S. B. Berlin Math. Ges.*, 22:9–20, 1923.
- [95] J. -P. Serre. Représentations linéaires et espaces homogènes kählériens des groupes de Lie compacts (d’après Armand Borel et André Weil). In *Séminaire Bourbaki, Vol. 2*, pages Exp. No. 100, 447–454. Soc. Math. France, Paris, 1995.
- [96] C. S. Seshadri. *Fibrés vectoriels sur les courbes algébriques*, volume 96 of *Astérisque*. Société Mathématique de France, Paris, 1982. Notes written by J.-M. Drezet from a course at the École Normale Supérieure, June 1980.
- [97] S. S. Shatz. The decomposition and specialization of algebraic families of vector bundles. *Compositio Math.*, 35(2):163–187, 1977.
- [98] G. C. Shephard. An elementary proof of Gram’s theorem for convex polytopes. *Canad. J. Math.*, 19:1214–1217, 1967.
- [99] R. Sjamaar. Holomorphic slices, symplectic reduction and multiplicities of representations. *Ann. of Math. (2)*, 141:87–129, 1995.
- [100] R. Sjamaar and E. Lerman. Stratified symplectic spaces and reduction. *Ann. of Math. (2)*, 134:375–422, 1991.
- [101] P. Slodowy. Die Theorie der optimalen Einparameteruntergruppen für instabile Vektoren. In *Algebraische Transformationsgruppen und Invariantentheorie*, volume 13 of *DMV Sem.*, pages 115–131. Birkhäuser, Basel, 1989.
- [102] G. Székelyhidi. Extremal metrics and K-stability (PhD thesis). arxiv:math/0611002.
- [103] A. Teleman. Symplectic stability, analytic stability in non-algebraic complex geometry. *Internat. J. Math.*, 15(2):183–209, 2004.
- [104] C. Teleman. The quantization conjecture revisited. *Ann. of Math. (2)*, 152(1):1–43, 2000.
- [105] R. P. Thomas. Notes on GIT and symplectic reduction for bundles and varieties. In *Surveys in differential geometry. Vol. X*, volume 10 of *Surv. Differ. Geom.*, pages 221–273. Int. Press, Somerville, MA, 2006.
- [106] G. Tian. On a set of polarized Kähler metrics on algebraic manifolds. *J. Differential Geom.*, 32(1):99–130, 1990.
- [107] Katrin Wehrheim and Chris T. Woodward. Functoriality for Lagrangian correspondences in Floer theory. *Quantum Topol.*, 1:129–170, 2010.
- [108] A. Weinstein. The symplectic “category”. In *Differential geometric methods in mathematical physics (Clausthal, 1980)*, volume 905 of *Lecture Notes in Math.*, pages 45–51. Springer, Berlin, 1982.
- [109] E. Witten. Two-dimensional gauge theories revisited. *J. Geom. Phys.*, 9:303–368, 1992.
- [110] E. Witten. Holomorphic Morse inequalities. In *Algebraic and differential topology—global differential geometry*, volume 70 of *Teubner-Texte Math.*, pages 318–333. Teubner, Leipzig, 1984.
- [111] C. Woodward. The classification of transversal multiplicity-free group actions. *Ann. Global Anal. Geom.*, 14:3–42, 1996.
- [112] C. Woodward. Multiplicity-free Hamiltonian actions need not be Kähler. *Invent. Math.*, 131(2):311–319, 1998.
- [113] C. T. Woodward. Localization via the norm-square of the moment map and the two-dimensional Yang-Mills integral. *J. Symp. Geom.*, 3(1):17–55, 2006.
- [114] S. Wu. Equivariant holomorphic Morse inequalities. II. Torus and non-abelian group actions. *J. Differential Geom.*, 51(3):401–429, 1999.

Mathematics-Hill Center, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, U.S.A. •  
ctw@math.rutgers.edu