



*J*ournal de l'École polytechnique *Mathématiques*

BORIS BYCHKOV, PETR DUNIN-BARKOWSKI, MAXIM KAZARIAN,
& SERGEY SHADRIN

Explicit closed algebraic formulas for Orlov–Scherbin n -point functions

Tome 9 (2022), p. 1121–1158.

<https://doi.org/10.5802/jep.202>

© Les auteurs, 2022.



Cet article est mis à disposition selon les termes de la licence
LICENCE INTERNATIONALE D'ATTRIBUTION CREATIVE COMMONS BY 4.0.
<https://creativecommons.org/licenses/by/4.0/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques »
(<http://jep.centre-mersenne.org/>) implique l'accord avec les conditions générales
d'utilisation (<http://jep.centre-mersenne.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique



Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org

EXPLICIT CLOSED ALGEBRAIC FORMULAS FOR ORLOV–SCHERBIN n -POINT FUNCTIONS

BY BORIS BYCHKOV, PETR DUNIN-BARKOWSKI, MAXIM KAZARIAN
& SERGEY SHADRIN

ABSTRACT. — We derive a new explicit formula in terms of sums over graphs for the n -point correlation functions of general formal weighted double Hurwitz numbers coming from the Kadomtsev–Petviashvili tau functions of hypergeometric type (also known as Orlov–Scherbin partition functions). Notably, we use the change of variables suggested by the associated spectral curve, and our formula turns out to be a polynomial expression in a certain small set of formal functions defined on the spectral curve.

RÉSUMÉ (Formules algébriques closes explicites pour les fonctions à n points d'Orlov-Scherbin)

Nous présentons une nouvelle formule explicite en termes de sommes sur les graphes pour les fonctions de corrélation à n points des nombres de Hurwitz doubles pondérés formels généraux provenant des fonctions tau de Kadomtsev-Petviashvili de type hypergéométrique (également connues sous le nom de fonctions de partition d'Orlov-Scherbin). Nous utilisons notamment le changement de variables suggéré par la courbe spectrale associée, et notre formule s'avère être une expression polynomiale dans un certain petit ensemble de fonctions formelles définies sur la courbe spectrale.

CONTENTS

1. Introduction.....	1122
2. Operators on the Fock space.....	1129
3. Preliminary computation of $H_{g,n}$	1131
4. Computation of $D_1 \cdots D_n H_n$	1136
5. General formula.....	1143
6. Exceptional cases.....	1149
7. Applying general formula.....	1153
References.....	1156

MATHEMATICAL SUBJECT CLASSIFICATION (2020). — 05A15, 37K10, 14H30, 14N10, 37K30.

KEYWORDS. — Hurwitz numbers, KP tau functions, Fock space.

S. S. was supported by the Netherlands Organization for Scientific Research. M.K. and B.B. wrote Sections 2, 3, 4 as part of the project supported by the International Laboratory of Cluster Geometry NRU HSE, RF Government grant, ag. № 075-15-2021-608 dated 08.06.2021. The work of B.B. on other sections was also partially funded by the grant Basis. P.D.-B. was partially supported by RFBR grant 20-01-00644.

1. INTRODUCTION

1.1. HURWITZ NUMBERS AND KP TAU FUNCTIONS OF HYPERGEOMETRIC TYPE. — Hurwitz numbers enumerate topologically distinct ramified coverings of the sphere S^2 by Riemann surfaces with prescribed ramification data. Different types of Hurwitz numbers are distinguished by the way the ramification data is specified. These data can be encoded by the values of parameters $c_k, s_k, k = 1, 2, \dots$, collected into two formal power series

$$(1) \quad \psi(y) := \sum_{k=1}^{\infty} c_k y^k, \quad y(z) := \sum_{k=1}^{\infty} s_k z^k.$$

We do not reproduce here the precise combinatorial definition of the Hurwitz numbers we are interested in, instead, we identify them as the Taylor coefficients of the corresponding generating function $F(p_1, p_2, \dots)$ introduced below. Namely, its exponential $Z = \exp F$ is a *Kadomtsev–Petviashvili tau function of hypergeometric type* (also known as an *Orlov–Scherbín partition function*) [KMMM95, OS01a, OS01b] given explicitly by its expansion in the basis of Schur functions

$$(2) \quad Z = e^F = \sum_{\lambda} e^{\sum_{(i,j) \in \lambda} \psi(\hbar(j-i))} s_{\lambda}(p) s_{\lambda}(s/\hbar).$$

We regard Z and F as formal power series in the variables p_1, p_2, \dots depending on additional parameters c_k, s_k , and \hbar . The summation runs over the set of all partitions (Young diagrams) λ including the empty one, s_{λ} denotes the corresponding Schur symmetric function represented as a polynomial in the power sums p_k . The parameters c_k are involved as the coefficients of the series ψ while s_k are substituted as the arguments of s_{λ} via $s/\hbar = (s_1/\hbar, s_2/\hbar, \dots)$. We regard a Young diagram λ as a table of rows of lengths $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$, and for a cell of this table with coordinates (i, j) its *content* is defined as the difference $j - i$ of coordinates. For that reason, the exponent $e^{\sum_{(i,j) \in \lambda} \psi(\hbar(j-i))}$ is sometimes referred to as the *content product*.

The (formal) Hurwitz numbers $h_{g,(m_1, \dots, m_n)}$ associated with the series F are defined by the expansion

$$\frac{\partial^n F}{\partial p_{m_1} \dots \partial p_{m_n}} \Big|_{p=0} = \sum_{g=0}^{\infty} \hbar^{2g-2+n} h_{g,(m_1, \dots, m_n)}.$$

The generating functions for many particular families of Hurwitz numbers (e.g. simple, monotone, Bousquet–Mélou–Schaeffer numbers, Grothendieck’s dessins d’enfants, and many others numbers of similar nature both of single, orbifold or double types) are included in F for particular values of the parameters, see Table 1 (cf. [ACEH18b, ALS16, Har16, KL15]).

In the most general case, when ψ and y are arbitrary power series, the Taylor coefficients $h_{g,(m_1, \dots, m_n)}$ have the combinatorial meaning of *weighted double Hurwitz numbers* (see e.g. [Har16]). Roughly speaking, when regarding F as a generating series for Hurwitz numbers, these Hurwitz numbers correspond to a weighted count of coverings with the ramification over the point $\infty \in S^2 = \mathbb{C}P^1$ being encoded by a monomial in the p -variables, the ramification over 0 corresponding to a monomial in

TABLE 1. Types of Hurwitz numbers

Hurwitz numbers	$e^{\psi(y)}$
usual	e^y
atlantes	e^{y^r}
monotone	$1/(1-y)$
strictly monotone	$1+y$
hypermaps	$(1+uy)(1+vy)$
BMS numbers	$(1+y)^m$
polynomial weighted	$1 + \sum_{k=1}^d c_k y^k$
general weighted	$\exp(\sum_{k=1}^{\infty} c_k y^k)$

Variations	$y(z)$
simple	z
orbifold	z^q
double	$\sum_{k=1}^{\infty} s_k z^k$

the s -variables, and the ramification of various types over the points different from 0 and ∞ giving contribution to the weight according to the explicit form of the series ψ . The exponent of the variable \hbar is the negative Euler characteristic $2g - 2 + n$ of the covering surface punctured at the preimages of ∞ , where g is the genus of the covering surface and n is the number of preimages of ∞ .

1.2. n -POINT FUNCTIONS. — Formula (2) is quite explicit and efficient for the numerical computation of particular Hurwitz numbers. Therefore, the main interest is related not to the computation of a single Hurwitz number but to the study of analytical and integrable properties of their generating functions. These properties are often formulated in terms of the *connected* and *disconnected*, respectively, n -point correlation functions defined by

$$(3) \quad H_n = \sum_{k_1, \dots, k_n=1}^{\infty} \frac{\partial^n F}{\partial p_{k_1} \dots \partial p_{k_n}} \Big|_{p=0} X_1^{k_1} \dots X_n^{k_n},$$

$$(4) \quad H_n^\bullet = \sum_{k_1, \dots, k_n=1}^{\infty} \frac{\partial^n Z}{\partial p_{k_1} \dots \partial p_{k_n}} \Big|_{p=0} X_1^{k_1} \dots X_n^{k_n}.$$

These are infinite power series in X_1, \dots, X_n serving as an alternative way of collecting Hurwitz numbers enumerating connected and disconnected, respectively, coverings of the sphere. Connected and disconnected n -point functions are related to one another by the inclusion-exclusion relations

$$(5) \quad \begin{aligned} H_n^\bullet(X_{\{1, \dots, n\}}) &= \sum_{I \vdash \{1, \dots, n\}} \prod_{i=1}^{|I|} H_{|I_i|}(X_{I_i}), \\ H_n(X_{\{1, \dots, n\}}) &= \sum_{I \vdash \{1, \dots, n\}} (-1)^{|I|-1} (|I|-1)! \prod_{i=1}^{|I|} H_{|I_i|}^\bullet(X_{I_i}), \end{aligned}$$

where the sums run over all unordered partitions of the set $\{1, \dots, n\}$, and for $I = \{i_1, i_2, \dots\}$ we denote $X_I := (X_{i_1}, X_{i_2}, \dots)$. The connected n -point function admits a *genus decomposition*

$$H_n = \sum_{g=0}^{\infty} \hbar^{2g-2+n} H_{g,n},$$

where $H_{g,n}$ is independent of \hbar :

$$H_{g,n} = \sum_{m_1, \dots, m_n=1}^{\infty} h_{g, (m_1, \dots, m_n)} X_1^{m_1} \dots X_n^{m_n}.$$

This follows from the combinatorial interpretation of Hurwitz numbers, but it is also a formal corollary of the computation of H_n of the present paper.

One of the main discoveries of last years in the theory of Hurwitz numbers is the fact that in many cases the n -point functions $H_{g,n}$ are governed by the *topological recursion*, a formalism allowing to compute $H_{g,n}$ inductively in g and n . One of the most general cases for which the topological recursion relations had been proved by the time we wrote the first version of the present paper is the one when both $e^{\psi(y)}$ and $y(z)$ are polynomials [ACEH18b, ACEH20]. Consider the following power series

$$(6) \quad X(z) = z e^{-\psi(y(z))},$$

where ψ and y are given by (1), and apply the local change of coordinates $X_i = X(z_i)$ to each of the arguments of $H_{g,n}$. One of the corollaries of the topological recursion is that the *function $H_{g,n}$ written in z -coordinates is rational*. Do note, however, that for the approach of [ACEH18b, ACEH20] the polynomiality of $e^{\psi(y)}$ and $y(z)$ was crucial. See also Remark 1.7 below.

In this paper we show that the function $H_{g,n}$ simplifies considerably after the change (6) even without any assumption of polynomiality or rationality (or even convergence) for the series $\psi(y)$ and $y(z)$. The main result of the paper is an explicit closed formula for $H_{g,n}$ for each pair (g, n) . Through the change (6) we have

$$(7) \quad D := X \frac{\partial}{\partial X} = \frac{1}{Q} z \frac{\partial}{\partial z},$$

where

$$(8) \quad Q = \frac{z}{X} \frac{dX}{dz} = 1 - D\psi(y(z)) = 1 - z y'(z) \psi'(y(z)).$$

For brevity, we will often use the notation of the form $\partial_z := \partial/\partial z$ below.

THEOREM 1.1. — *In the unstable cases $2g-2+n \leq 0$ the n -point functions are given by*

$$(9) \quad \begin{aligned} D_1 H_{0,1} &= y(z_1), \\ H_{0,2} &= \log \left(\frac{z_1^{-1} - z_2^{-1}}{X_1^{-1} - X_2^{-1}} \right), \end{aligned}$$

and for all (g, n) with $2g - 2 + n > 0$ the function $H_{g,n}$ written in z -coordinates admits a closed expression of the form

$$(10) \quad H_{g,n} = \sum_{j_1, \dots, j_n=0}^{\infty} D_1^{j_1} \dots D_n^{j_n} \frac{P_{g;j_1, \dots, j_n}}{Q_1 \dots Q_n} + c_{g,n}$$

with finitely many nonzero summands, where $Q_i = Q(z_i)$, $D_i = D(z_i) = (1/Q_i)z_i \partial_{z_i}$, and $P_{g;j_1, \dots, j_n}$ is a polynomial combination of functions $z_j/(z_i - z_j)$ and derivatives $\psi^{(k)}(y(z_i))$ and $(z_i \partial_{z_i})^k y(z_i)$, $k \geq 1$, $i = 1, \dots, n$. Finally, $c_{g,n}$ is a constant explicitly given by

$$c_{g,n} = (-1)^n \psi^{(2g-2+n)}(0) [u^{2g}] \left(\frac{u}{e^{u/2} - e^{-u/2}} \right)^2,$$

where $[u^{2g}]$ denotes the coefficient in front of u^{2g} in the series expansion.

In particular, an immediate corollary of this theorem is the following statement:

COROLLARY 1.2. — *If both $y'(z)$ and $\psi'(y)$ are rational functions then, for $2g - 2 + n > 0$, $H_{g,n}$ is a rational function in z_1, \dots, z_n .*

An explicit description of the terms entering the formula for $H_{g,n}$ (i.e., a formula where all polynomials $P_{g;j_1, \dots, j_n}$ are given explicitly) is presented in Theorem 5.3 in the case $n > 2$ and in Section 6 in the exceptional cases $n = 1$ and $n = 2$. It might look somewhat complicated but it is actually quite explicit and can be used for practical computations. The formula holds true even in those cases when $\psi(y)$ and $y(z)$ are just formal series with no assumption of rationality or convergence and the topological recursion is not applicable in principle. Moreover, even in those cases when $e^{\psi(y)}$ and $y(z)$ are such that the topological recursion can be applied (e.g. when they are polynomial, as in [ACEH18b, ACEH20], or in a more general case as referred to in Remark 1.7) our formula is more efficient (as a way to compute the n -point functions) since the number of its terms does not depend on the degrees of those polynomials and it does not require finding roots of algebraic equations determining critical points of the function $X(z)$.

REMARK 1.3. — The left-hand side of (10) is a formal power series in z_1, \dots, z_n while the individual summands of the right-hand side have poles on the diagonals $z_i = z_j$ and their interpretation requires additional comments. First note that if both $\psi'(y)$ and $y'(z)$ are rational functions then all terms of (10) are also rational, and the equality implies, in particular, that all poles on the diagonals on the right-hand side cancel out (see Corollary 5.7).

In the general case, one of the possibilities to interpret Equation (10) is to consider the asymptotic Laurent expansion of all of its terms in the sector $|z_1| \ll |z_2| \ll \dots \ll |z_n| \ll 1$. This power expansion involves monomials in z_1, \dots, z_n containing both positive and negative powers of the variables z_i .

It is much more advisable, however, to treat the terms of (10) in a different way. Namely we consider them as elements of the ring

$$R = \mathbb{C}[[z_1, \dots, z_n]][\{(z_i - z_j)^{-1}; i, j \in \{1, \dots, n\}, i < j\}]$$

of ‘formal power series with finite order poles on the diagonals’. It follows that for each $d \geq 0$ the term of homogeneous degree d of each summand in (10) is expressed as a degree d homogeneous rational function in z_1, \dots, z_n with possible poles on the diagonals. After summation, all these poles cancel out and the result is a homogeneous polynomial representing degree d homogeneous term of the Taylor expansion of $H_{g,n}$.

In this paper, we first deal with formal series in z_1, \dots, z_n (from definitions (3)–(4), where we substitute X_i with $X(z_i)$ from (6), itself understood as a formal series in z_i). Then, starting with Proposition 3.4, we introduce the functions $z_i z_j / (z_i - z_j)^2$ understood as their Laurent expansions in the sector $|z_1| \ll |z_2| \ll \dots \ll |z_n| \ll 1$. Finally, in Proposition 4.8 and in what follows after it, we understand all terms as elements of the ring R (which does not make sense prior to that proposition).

1.3. FURTHER REMARKS

REMARK 1.4. — Our results can be naturally extended to the case where $\psi(y)$ and $y(z)$ depend on \hbar^2 , i.e., where c_k and s_k are formal series in \hbar^2 rather than just constants. This is done in [BDBKS20]. See also Remarks 4.9 and 5.6. This means that our statement, in addition to the cases listed in Table 1, also covers e.g. the cases of r -spin Hurwitz numbers [KLPS19] and the coefficients of the extended Ooguri–Vafa partition functions of colored HOMFLY polynomials of torus knots [DBPSS19, DBKP⁺20]; see Table 2, which is an extension of Table 1 to these cases.

TABLE 2. Types of Hurwitz-like numbers requiring \hbar -extension

Hurwitz numbers	$e^{\psi(y)}$	$y(z)$
r -spin q -orbifold	$\exp\left(\frac{(y + \hbar/2)^{r+1} - (y - \hbar/2)^{r+1}}{(r + 1)\hbar}\right)$	z^q
ext. Ooguri-Vafa	$e^{(P/Q)y}$	$\sum_{k=1}^{\infty} \frac{\hbar(A^k - A^{-k})z^k}{e^{k\hbar/2} - e^{-k\hbar/2}}$

REMARK 1.5. — Note that for usual simple Hurwitz numbers [DBKO⁺15, KLS19], for orbifold Hurwitz numbers [DBLPS15, KLS19], for monotone and strictly monotone orbifold Hurwitz numbers [KLS19], for r -spin (and r -spin orbifold) Hurwitz numbers [KLPS19], for the numbers of maps and hypermaps (dessins d’enfants) [KZ15], for the Bousquet–Mélou–Schaeffer numbers [BDBS20], for the coefficients of the extended Ooguri–Vafa partition function of the colored HOMFLY polynomials of torus knots [DBPSS19], and for double Hurwitz numbers [BDK⁺20], there exist

combinatorial-algebraic proofs of the so-called quasi-polynomiality property. This property, in particular, implies the linear loop equation and the projection property of [BS17] for the respective n -point functions. We remark that the results of the present paper, in particular, serve as an independent proof of linear loop equations for all these cases (and, indeed, in the whole generality of the formal weighted double Hurwitz numbers context). We discuss this in more detail in our subsequent publication [BDBKS20].

REMARK 1.6. — One way to interpret the statements of Theorem 1.1 and Theorem 5.3 is to say that they give a conceptual explanation why the change of variables (6) is so ubiquitous in the weighted Hurwitz theory. This change of variables was suggested by Alexandrov–Chapuy–Eynard–Harnad in [ACEH18b] based on the explicit computation of $H_{0,1}$ and the idea that the $(0,1)$ -function should determine the spectral curve for the topological recursion, in the cases when the spectral curve topological recursion is applicable. Specific cases of this sort of change of variables were also used in [ACEH18a], in the combinatorial-algebraic papers mentioned in Remark 1.5 and in other combinatorial papers, see e.g. [Cha09].

Our present paper clarifies the general and unconditional meaning of this change of variables for the n -point functions in the \hbar -expansion in the semi-infinite wedge formalism.

REMARK 1.7. — The results of the present paper have very strong corollaries for the theory of topological recursion for various types of Hurwitz numbers, including all the ones mentioned in Remark 1.5. Specifically, in our subsequent paper [BDBKS20], based on the results of the present paper, we prove the *blobbed* topological recursion (defined in [BS17]) for generalized weighted double Hurwitz numbers basically in full generality, and we prove the regular topological recursion for two very general families of generalized weighted double Hurwitz numbers. These families include as special cases all the cases of Hurwitz-type numbers for which topological recursion was known from the literature (in particular, all the ones mentioned in Remark 1.5), and are actually quite a bit more general than that. Importantly, while previously in the literature the topological recursion for various types of Hurwitz-like numbers has been proved on a case-by-case basis with complicated techniques which differed between the cases, our technique of [BDBKS20] (based on the results of the present paper) gives a clear and uniform way to do this and highlights the underlying common structure.

Moreover, the results of the present paper are also applicable beyond Hurwitz numbers. In particular, we applied them for *maps* and *stuffed maps* and their generalizations: in our another subsequent paper [BDBKS21], based on the results of the present paper, we prove a general duality for the generalized stuffed maps which we call the *ordinary vs fully simple* duality, which also allowed us in that same paper to prove the Borot–Garcia-Failde conjecture on the topological recursion for fully simple maps ([BGF20, Conj. 5.3]).

1.4. **PRIOR WORK OF THE THIRD NAMED AUTHOR.** — The main result of this paper resolves a slightly weaker conjecture of the third named author that he posed in various talks in 2019, see e.g. [Kaz19]. Namely, he conjectured the existence of universal formulas for the Orlov–Scherbin n -point functions $H_{g,n}$ which should represent them as expressions polynomial in

$$\begin{aligned} \psi^{(j)}(y(z_k)), & \quad j \geq 1, \quad k = 1, \dots, n, \\ (z_k \partial_{z_k})^j y(z_k), & \quad j \geq 1, \quad k = 1, \dots, n, \\ z_\ell / (z_k - z_\ell), & \quad 1 \leq k < \ell \leq n, \\ Q(z_k)^{-1}, & \quad k = 1, \dots, n \end{aligned}$$

(cf. the statement of Theorem 1.1). Moreover, using a variety of deformation techniques he later proved his conjecture in [Kaz21], and his proof gave an algorithm to produce the universal formulas inductively (see also [Kaz20]).

It is important to stress that although this paper resolves the conjecture of the third named author in a different way than in [Kaz21], and the formulas for $H_{g,n}$ given in Theorem 5.3 have closed form (as opposed to their inductive algorithmic derivation in [Kaz21]), the present paper is both ideologically and technically very much dependent on [Kaz21]. In particular, many lemmas and computational ideas that we use below are shared directly from [Kaz21].

1.5. **ORGANIZATION OF THE PAPER.** — In Section 2 we recall the basic formalism of the operators on the bosonic Fock space that we use throughout the paper. In Section 3 we compute $H_{g,n}$ as a series in X_1, \dots, X_n , which, in particular, leads to formula giving each particular formal weighted double Hurwitz number $h_{g,(m_1, \dots, m_n)}$ in a closed form. Strictly speaking, this section is not necessary for the rest of the paper, but it sets up the notation and illuminates the logic of computations in the subsequent parts of the paper.

In Section 4 we derive an explicit closed formula for $D_1 \cdots D_n H_{g,n}$. In Section 5 we prove the main theorem of the present paper, which explicitly represents $H_{g,n}$ for given g and n in a closed form. Section 6 deals with the slightly exceptional cases of $n = 1$ for any g and $(g, n) = (0, 2)$. Finally, in Section 7 we give examples of the application of our main general formula, deriving explicit closed formulas for $H_{g,n}$ for particular small g and n .

Acknowledgments. — This project has started when S. S. was visiting the Faculty of Mathematics at the National Research University Higher School of Economics, and S. S. would like to thank the Faculty for warm hospitality and stimulating research atmosphere.

We would like to thank A. Alexandrov, J. van de Leur, and the anonymous referees for helpful remarks.

2. OPERATORS ON THE FOCK SPACE

By the (bosonic) *Fock space* we mean the space of infinite power series $\mathcal{F} = \mathbb{C}[[p_1, p_2, \dots]]$. It has a distinguished element 1 called *vacuum vector* and denoted sometimes by $|0\rangle$, and a distinguished linear function $\mathcal{F} \rightarrow \mathbb{C}$ called *covacuum vector* that takes a series to its free term (the value at $p = 0$) and is denoted by $\langle 0|$.

We will consider some operators acting on the Fock space. In particular, we set $J_m = m \partial_{p_m}$ if $m > 0$, $J_0 = 0$, and $J_m = p_{-m}$ (the operator of multiplication by p_{-m}), if $m < 0$. Note that

$$(11) \quad [J_k, J_\ell] = k\delta_{k+\ell,0}.$$

Introduce also the operator $\mathcal{D}(\hbar)$ acting diagonally in the basis of Schur functions by

$$\mathcal{D}(\hbar) s_\lambda = e^{\sum_{(i,j) \in \lambda} \psi(\hbar(j-i))} s_\lambda.$$

With these notations, and using the Cauchy identity $\sum_\lambda s_\lambda(p)s_\lambda(s) = e^{\sum_{i=1}^\infty s_i p_i/i}$ for Schur polynomials (see, e.g., [Sta99, p. 386]), the definitions of the Orlov-Scherbin partition function and the disconnected n -point functions can be rewritten as follows

$$(12) \quad \begin{aligned} Z &= \mathcal{D}(\hbar) e^{\sum_{i=1}^\infty s_i J_{-i}/i \hbar} |0\rangle, \\ H_n^\bullet &= \sum_{m_1, \dots, m_n=1}^\infty \frac{X_1^{m_1} \cdots X_n^{m_n}}{m_1 \cdots m_n} \langle 0| J_{m_1} \cdots J_{m_n} \mathcal{D}(\hbar) e^{\sum_{i=1}^\infty s_i J_{-i}/i \hbar} |0\rangle. \end{aligned}$$

The introduced standard terminology and notations come from physics. It might look as an unnecessary complication at first glance; its benefit will be seen later.

A bigger set of operators of our interest is constructed as follows.

DEFINITION 2.1. — The Lie algebra A_∞ is the \mathbb{C} -vector space of infinite matrices $(A_{i,j})_{i,j \in \mathbb{Z}+1/2}$ with only finitely many non-zero diagonals (that is, $A_{i,j}$ is not equal to zero only for finitely many possible values of $i - j$), together with the commutator bracket. The standard basis is formed by the matrix units $\{E_{i,j} \mid i, j \in \mathbb{Z} + 1/2\}$ such that $(E_{i,j})_{k,\ell} = \delta_{i,k} \delta_{j,\ell}$.

There is a remarkable projective representation of this algebra in the Fock space by means of differential operators. It is denoted by the hat symbol and defined by the following generating function for the action of the matrix units (see e.g. [MJD00, §6.2]):

$$(13) \quad \sum_{k,\ell \in \mathbb{Z}+1/2} x^\ell y^{-k} \widehat{E}_{k,\ell} = x^{1/2} y^{1/2} \frac{e^{\sum_{i=1}^\infty (y^{-i} - x^{-i}) p_i/i} e^{\sum_{i=1}^\infty (x^i - y^i) \partial_{p_i}} - 1}{x - y}.$$

The expansion of the exponents on the right-hand side enlists all possible monomial differential operators in p -variables. The coefficient of any such monomial differential operator, after cancellation, is a polynomial in the half-integer powers of x and y . The contribution of this operator to $\widehat{E}_{k,\ell}$ is equal to the coefficient of $x^\ell y^{-k}$ in that polynomial.

The term ‘projective representation’ means that the commutator of matrices from A_∞ corresponds to the commutator of their action on the Fock space up to a scalar operator. More explicitly, we have:

$$(14) \quad [\widehat{E}_{a,b}, \widehat{E}_{c,d}] = \delta_{b,c} \widehat{E}_{a,d} - \delta_{a,d} \widehat{E}_{c,b} + \delta_{b,c} \delta_{a,d} (\delta_{b>0} - \delta_{d>0}) \text{Id}.$$

Equivalently, we have actually a representation of the central extension $A_\infty + \mathbb{C}\text{Id}$.

The actual definition of the action of A_∞ in the Fock space goes through fermionic realization of the Fock space and the boson-fermion correspondence, see [MJD00, §§5 & 6] for the details. But as long as the formula (13) is established it can be taken as a definition and most parts of the underlying formalism can be omitted. The profit of using this representation is that while manipulating with operators it is much easier to make computations directly in the algebra A_∞ rather than in its more complicated representation in the Fock space.

However, we will need one more relation that does not follow immediately from (13). Namely, any diagonal matrix $\sum_{k \in \mathbb{Z}+1/2} w_k E_{k,k} \in A_\infty$ acts diagonally in the Schur basis and the corresponding eigenvalue is determined by

$$(15) \quad \sum_{k \in \mathbb{Z}+1/2} w_k \widehat{E}_{k,k} s_\lambda = \sum_{i=1}^{\ell(\lambda)} (w_{\lambda_i - i + 1/2} - w_{-i + 1/2}) s_\lambda = \sum_{(i,j) \in \lambda} v_{j-i} s_\lambda,$$

where

$$v_k = w_{k+1/2} - w_{k-1/2},$$

see [KL15] for details. In particular, for the operator $\mathcal{D}(\hbar)$ introduced above we have

$$\mathcal{D}(\hbar) = \exp\left(\sum_{k \in \mathbb{Z}+1/2} w_k \widehat{E}_{k,k}\right),$$

where w_k is determined from relations $w_{k+1/2} - w_{k-1/2} = \psi(\hbar k)$, $k \in \mathbb{Z}$ (this determines the factors w_k up to a common additive constant, but this constant is unimportant, as (14) implies that $\sum_{k \in \mathbb{Z}+1/2} \widehat{E}_{k,k}$ vanishes on any vector of our Fock space; this follows from taking the limit $y \rightarrow x$ in the RHS of (14) and then taking the free term in the resulting x -series).

Define

$$(16) \quad \mathcal{E}(u, z) := \sum_{m \in \mathbb{Z}} z^m \sum_{k \in \mathbb{Z}+1/2} e^{u(k-m/2)} \widehat{E}_{k-m,k}.$$

Set

$$(17) \quad \mathcal{S}(z) := \frac{e^{z/2} - e^{-z/2}}{z}.$$

Then, setting $x = z e^{u/2}$, $y = z e^{-u/2}$ in (13), we obtain:

PROPOSITION 2.2. — *We have*

$$(18) \quad \mathcal{E}(u, z) = \frac{e^{\sum_{i=1}^{\infty} u \mathcal{S}(u i) J_{-i} z^{-i}} e^{\sum_{i=1}^{\infty} u \mathcal{S}(u i) J_i z^i} - 1}{u \mathcal{S}(u)}.$$

An independent proof of the equality of the coefficients of z^0 of both sides can be found in [SSZ12]. For example, comparing the coefficients of $z^m u^0$ on both sides we find

$$J_m = \sum_{k \in \mathbb{Z} + 1/2} \widehat{E}_{k-m, k}.$$

The commutation relation (11) for these operators also implies the following formula:

PROPOSITION 2.3

$$(19) \quad e^{\sum_{i=1}^{\infty} a_i J_i} e^{\sum_{i=1}^{\infty} b_i J_{-i}} = e^{\sum_{i=1}^{\infty} i a_i b_i} e^{\sum_{i=1}^{\infty} b_i J_{-i}} e^{\sum_{i=1}^{\infty} a_i J_i}$$

for any collection of constants a_i, b_i such that the corresponding infinite sums make sense.

Proof. — This is just a very well-known common special case of the Baker–Campbell–Hausdorff formula, but it is illuminating to see how in this particular case it is just a manifestation of the Taylor formula. Namely, by the Taylor formula, the action of the operator $e^{\sum_{i=1}^{\infty} a_i J_i} = e^{\sum_{i=1}^{\infty} i a_i \partial_{p_i}}$ on a series $f(p_1, p_2, \dots)$ results in a shift of the arguments,

$$e^{\sum_{i=1}^{\infty} a_i J_i} f(p_1, p_2, \dots) = f(p_1 + 1 a_1, p_2 + 2 a_2, \dots).$$

Therefore, we have

$$\begin{aligned} e^{\sum_{i=1}^{\infty} a_i J_i} e^{\sum_{i=1}^{\infty} b_i J_{-i}} f(p_1, p_2, \dots) &= e^{\sum_{i=1}^{\infty} b_i (p_i + i a_i)} f(p_1 + 1 a_1, p_2 + 2 a_2, \dots) \\ &= e^{\sum_{i=1}^{\infty} i a_i b_i} e^{\sum_{i=1}^{\infty} b_i p_i} e^{\sum_{i=1}^{\infty} i a_i \partial_{p_i}} f(p_1, p_2, \dots), \end{aligned}$$

which proves the commutation relation formulated above. □

3. PRELIMINARY COMPUTATION OF $H_{g,n}$

In this section we compute $H_{g,n}$ as a series in X_1, \dots, X_n . In particular, this leads to a computation of each particular weighted double Hurwitz number $h_{g, (m_1, \dots, m_n)}$ in a closed form.

3.1. VACUUM EXPECTATION EXPRESSION FOR H_n^\bullet . — Let us define

$$\mathbb{J}_m := \mathcal{D}(\hbar)^{-1} J_m \mathcal{D}(\hbar).$$

This allows us to rewrite (12) as

$$(20) \quad H_n^\bullet = \sum_{m_1, \dots, m_n=1}^{\infty} \frac{X_1^{m_1} \dots X_n^{m_n}}{m_1 \dots m_n} \langle 0 | \mathbb{J}_{m_1} \dots \mathbb{J}_{m_n} e^{\sum_{i=1}^{\infty} s_i J_{-i} / i \hbar} | 0 \rangle.$$

PROPOSITION 3.1. — The operators $\mathbb{J}_m(\hbar)$ belong to A_∞ for all $m \in \mathbb{Z}$, namely,

$$(21) \quad \mathbb{J}_m(\hbar) = \sum_{k \in \mathbb{Z} + 1/2} \phi_m(\hbar(k - m/2)) \widehat{E}_{k-m, k},$$

where

$$(22) \quad \phi_m(y) := \exp\left(\sum_{i=1}^m \psi\left(y + \frac{2i - m - 1}{2} \hbar\right)\right), \quad m > 0,$$

$$(23) \quad \begin{aligned} \phi_0(y) &:= 1, \\ \phi_m(y) &:= (\phi_{-m}(y))^{-1}, \end{aligned} \quad m < 0.$$

More explicitly, we have

$$(24) \quad \mathbb{J}_m = \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [u^r z^m] \frac{e^{\sum_{i=1}^{\infty} u \hbar \mathcal{S}(u \hbar i) J_{-i} z^{-i}} e^{\sum_{i=1}^{\infty} u \hbar \mathcal{S}(u \hbar i) J_i z^i}}{u \hbar \mathcal{S}(u \hbar)}.$$

Note that $\phi_m(y)$ also depends on \hbar but here and below we omit this argument for brevity.

NOTATION 3.2. — Here and below $[x^k]f(x)$ stands for the coefficient in front of x^k in the series expansion of $f(x)$.

Proof of Proposition 3.1. — For $m = 0$ the statement is evident: from (15), the operator $\sum_{k \in \mathbb{Z}+1/2} \widehat{E}_{k,k}$ annihilates the whole Fock space. Let $m \neq 0$. Recall that $J_m = \sum_{k \in \mathbb{Z}+1/2} \widehat{E}_{k-m,k}$ and $\mathcal{D}(\hbar) = \exp(W)$, where $W = \sum_{k \in \mathbb{Z}+1/2} w_k \widehat{E}_{k,k}$ is represented by a diagonal matrix whose diagonal entries w_k are determined from the relations $w_k - w_{k-1} = \psi(\hbar(k - 1/2))$, $k \in \mathbb{Z}$. Therefore using (14) and Hadamard's formula $e^X Y e^{-X} = e^{\text{ad}_X}(Y)$, where $\text{ad}_X(\cdot) = [X; \cdot]$, we get

$$\begin{aligned} \mathbb{J}_m &= e^{-W} \left(\sum_{k \in \mathbb{Z}+1/2} \widehat{E}_{k-m,k} \right) e^W = \sum_{k \in \mathbb{Z}+1/2} e^{w_k - w_{k-m}} \widehat{E}_{k-m,k} \\ &= \sum_{k \in \mathbb{Z}+1/2} \phi_m(\hbar(k - m/2)) \widehat{E}_{k-m,k}. \end{aligned}$$

For the proof of (24) we compute:

$$\begin{aligned} \mathbb{J}_m &\stackrel{(21)}{=} \sum_{k \in \mathbb{Z}+1/2} \phi_m(\hbar(k - m/2)) \widehat{E}_{k-m,k} \\ &= \sum_{k \in \mathbb{Z}+1/2} \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} \frac{(\hbar(k - m/2))^r}{r!} \widehat{E}_{k-m,k} \\ &\stackrel{(16)}{=} \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [u^r z^m] \mathcal{E}(u \hbar, z) \\ &\stackrel{(18)}{=} \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [u^r z^m] \frac{e^{\sum_{i=1}^{\infty} u \hbar \mathcal{S}(u \hbar i) J_{-i} z^{-i}} e^{\sum_{i=1}^{\infty} u \hbar \mathcal{S}(u \hbar i) J_i z^i}}{u \hbar \mathcal{S}(u \hbar)}. \end{aligned}$$

In the second line we have simply expanded $\phi_m(\hbar(k - m/2))$ in its Taylor series at zero; and in the last line we omit the -1 summand in the numerator coming from (18) since we have set $m \neq 0$ and it vanishes upon applying $[z^m]$. \square

3.2. COMPUTATION OF H_n^\bullet . — Now we can obtain the following expression for the disconnected n -point functions. Let

$$\mathcal{S}(u) = \frac{e^{u/2} - e^{-u/2}}{u} = \sum_{k=0}^{\infty} \frac{u^{2k}}{2^{2k}(2k+1)!}.$$

DEFINITION 3.3. — Denote by U^+ the transformation that takes a Laurent series $f(u, z)$ in u and z to the series in X given by

$$(25) \quad (U^+ f)(X) = \sum_{m=1}^{\infty} \frac{X^m}{m} \sum_{r=0}^{\infty} \partial_y^r \phi_m(y) \Big|_{y=0} [z^m u^r] \frac{e^{u \mathcal{S}(u \hbar z \partial_z) y(z)}}{u \hbar \mathcal{S}(u \hbar)} f(u, z).$$

This formula describes explicitly the coefficients of $U^+ f$ as a power series in X . It makes sense if f is polynomial in u or if f is a series in \hbar whose coefficients are polynomial in u . Remark that $U^+ f$ is a regular series in X (i.e., containing only positive powers of X) even though the series f might have a pole in z at the origin: the non-positive powers of z in the expansion of f are just ignored.

Denote also by U_k^+ a similar transformation applied to u_k and z_k instead of u and z (the output of U_k^+ is a power series in X_k).

In all relations of this section the functions on the right-hand sides are understood as power asymptotic expansion in the sector $|z_1| \ll \dots \ll |z_n| \ll 1$.

PROPOSITION 3.4. — We have

$$(26) \quad H_n^\bullet = U_n^+ \dots U_1^+ \prod_{1 \leq k < \ell \leq n} e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2},$$

where the expression in the product on the right-hand side is understood as its power asymptotic expansion in the sector $|z_1| \ll \dots \ll |z_n| \ll 1$.

Proof. — Let us substitute expressions (24) for \mathbb{J} -operators into (20). We get

$$H_n^\bullet = \sum_{m_1, \dots, m_n=1}^{\infty} \sum_{r_1, \dots, r_n=0}^{\infty} \left(\prod_{k=1}^n \partial_y^{r_k} \phi_{m_k}(y) \Big|_{y=0} \frac{X_k^{m_k}}{m_k} \right) \times \left[\prod_{i=1}^n z_i^{m_i} u_i^{r_i} \right] \left\langle 0 \Big| \prod_{k=1}^n \frac{e^{\sum_{i=1}^{\infty} u_k \hbar \mathcal{S}(u_k \hbar i) J_{-i} z_k^{-i}} e^{\sum_{i=1}^{\infty} u_k \hbar \mathcal{S}(u_k \hbar i) J_i z_k^i}}{u_k \hbar \mathcal{S}(u_k \hbar)} e^{\sum_{i=1}^{\infty} s_i J_{-i} / i \hbar} \Big| 0 \right\rangle.$$

Then we apply commutation relations (19) for the exponentials of J -operators moving the $J_{>0}$ -factors to the right and the $J_{<0}$ -factors to the left. Since $J_{>0}$ is killed by the vacuum vector and $J_{<0}$ is killed by the covacuum, we get

$$\begin{aligned} & \left\langle 0 \Big| \prod_{k=1}^n \frac{e^{\sum_{i=1}^{\infty} u_k \hbar \mathcal{S}(u_k \hbar i) J_{-i} z_k^{-i}} e^{\sum_{i=1}^{\infty} u_k \hbar \mathcal{S}(u_k \hbar i) J_i z_k^i}}{u_k \hbar \mathcal{S}(u_k \hbar)} e^{\sum_{i=1}^{\infty} s_i J_{-i} / i \hbar} \Big| 0 \right\rangle \\ &= \prod_{k=1}^n \frac{\exp\left(\sum_{i=1}^{\infty} u_k \mathcal{S}(u_k \hbar i) s_i z_k^i\right)}{u_k \hbar \mathcal{S}(u_k \hbar)} \prod_{1 \leq k < \ell \leq n} \exp\left(\sum_{i=1}^{\infty} u_k \hbar \mathcal{S}(u_k \hbar i) u_\ell \hbar \mathcal{S}(u_\ell \hbar i) i \left(\frac{z_k}{z_\ell}\right)^i\right). \end{aligned}$$

Recall that

$$\sum_{i=1}^{\infty} s_i z_k^i = y(z_k) =: y_k.$$

Also note that

$$\sum_{i=1}^{\infty} i \left(\frac{z_k}{z_\ell}\right)^i = z_k \partial_{z_k} \frac{z_\ell}{z_k - z_\ell} = \frac{z_k z_\ell}{(z_k - z_\ell)^2},$$

if we assume that $|z_k| \ll |z_\ell|$. Noting all that we finally obtain

$$\begin{aligned} H_n^\bullet &= \sum_{m_1, \dots, m_n=1}^{\infty} \sum_{r_1, \dots, r_n=0}^{\infty} \left(\prod_{k=1}^n \partial_y^{r_k} \phi_{m_k}(y) \Big|_{y=0} \frac{X_k^{m_k}}{m_k} \right) \\ &\quad \times [z_1^{m_1} \cdots z_n^{m_n} u_1^{r_1} \cdots u_n^{r_n}] \prod_{k=1}^n \frac{e^{u_k \mathcal{S}(u_k \hbar z_k \partial_{z_k}) y_k}}{u_k \hbar \mathcal{S}(u_k \hbar)} \\ &\quad \prod_{1 \leq k < \ell \leq n} e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2}, \end{aligned}$$

where the expression in the second line is understood as its power asymptotic expansion in the sector $|z_1| \ll \cdots \ll |z_n| \ll 1$. This formula is equivalent to that of the proposition. \square

REMARK 3.5. — Note that the argument of the U^+ -operators in (26) involves both positive and negative powers of the variables z_k but the left-hand side is determined by those monomials of the right-hand side that contain positive powers of all variables only.

3.3. FROM DISCONNECTED TO CONNECTED n -POINT FUNCTIONS. — With notations of the previous section, we have:

PROPOSITION 3.6

$$(27) \quad H_n = U_n^+ \cdots U_1^+ \sum_{\gamma \in \Gamma_n} \prod_{\{v_k, v_\ell\} \in E_\gamma} \left(e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2} - 1 \right),$$

where Γ_n is the set of all connected simple (i.e., without multiple edges and loops) graphs over n vertices v_1, \dots, v_n , and E_γ is the set of edges of $\gamma \in \Gamma_n$.

Proof. — Let us denote

$$w_{k,\ell} = e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2} - 1$$

and consider the product

$$\prod_{1 \leq k < \ell \leq n} e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2} = \prod_{1 \leq k < \ell \leq n} (1 + w_{k,\ell}).$$

Expanding the brackets we obtain $2^{\binom{n}{2}}$ summands. These summands are labeled by simple graphs on n numbered vertices: the vertices k and ℓ are connected or not connected by an edge if the factor corresponding to the pair of indices k and ℓ is equal to $w_{k,\ell}$ or 1, respectively.

Then, Equation (26) for the disconnected n -point functions attains the following form

$$H_n^\bullet = U_n^+ \cdots U_1^+ \sum_{\gamma} \prod_{\{v_k, v_\ell\} \in E_\gamma} w_{k, \ell},$$

where the summation carries over the set of *all* simple graphs γ on n labeled vertices. The inclusion-exclusion procedure applied to this sum over all simple graphs singles out exactly the terms corresponding to the connected ones. \square

It is sometimes convenient to rearrange the insertion of \hbar in Equation (27) in the following way:

$$\hbar^{2-n} H_n = (\hbar U_n^+) \cdots (\hbar U_1^+) \sum_{\gamma \in \Gamma_n} \hbar^{2(|E_\gamma| - n + 1)} \prod_{\{v_k, v_\ell\} \in E_\gamma} \frac{w_{k, \ell}}{\hbar^2}.$$

Since any connected graph on n vertices has at least $n - 1$ edges, the right-hand side involves only non-negative even powers of the variable \hbar . Indeed, it is easy to see from definition that the series $w_{k, \ell}/\hbar^2$ and the coefficients of the transformation $\hbar U^+$ involve nonnegative even powers of \hbar only. This justifies in a formal way the mentioned *genus decomposition*

$$\hbar^{2-n} H_n = \sum_{g=0}^{\infty} \hbar^{2g} H_{g, n} \quad \text{or} \quad H_n = \sum_{g=0}^{\infty} \hbar^{2g-2+n} H_{g, n},$$

where $H_{g, n}$ is independent of \hbar .

Finally, note that the operators U_i^+ describe explicitly the Taylor coefficients of the resulting series. Therefore, we can regard (27) as an explicit expression for the corresponding Hurwitz numbers:

$$(28) \quad m_1 \cdots m_n h_{g, (m_1, \dots, m_n)} = [\hbar^{2g-2+n}] \sum_{r_1, \dots, r_n=0}^{\infty} \left(\prod_{k=1}^n \partial_y^{r_k} \phi_{m_k}(y) \Big|_{y=0} \right) [z_1^{m_1} \cdots z_n^{m_n} u_1^{r_1} \cdots u_n^{r_n}] \prod_{k=1}^n \frac{e^{u_k \mathcal{S}(u_k \hbar z_k \partial_{z_k}) y_k}}{u_k \hbar \mathcal{S}(u_k \hbar)} \sum_{\gamma \in \Gamma_n} \prod_{\{v_k, v_\ell\} \in E_\gamma} \left(e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2} - 1 \right).$$

REMARK 3.7. — Formulas (27) and (28) provide closed expressions for the connected n -point functions and connected formal weighted double Hurwitz numbers as sums over graphs, respectively. However, note that our main aim, as explained in the introduction, is to express the connected n -point functions as finite polynomials in certain formal functions on the spectral curve, and formula (27) does not achieve that. Indeed, note that in the definition (25) of the operator U^+ we have an infinite sum over m . It turns out that, roughly speaking, it is possible to take these m -sums to arrive at finite expressions, and this is what is done in the two following sections. However, the precise path to arriving at these finite expressions, while being inspired by the contents of the present section, does not explicitly rely on Proposition 3.6 and is, strictly speaking, independent of this section. We do use the notation introduced in the present section in what follows; notably, the ϕ_m 's will play an important role.

4. COMPUTATION OF $D_1 \cdots D_n H_n$

Set

$$D_i = X_i \partial_{X_i}.$$

Denote, for shortness,

$$DH_n^\bullet = \left(\prod_{i=1}^n D_i \right) H_n^\bullet \quad \text{and} \quad DH_n = \left(\prod_{i=1}^n D_i \right) H_n = \sum_{g=0}^{\infty} \hbar^{2g-2+n} DH_{g,n}.$$

In this section we compute these functions in a closed form. Remark that the operator $D_1 \cdots D_n$ multiplies a monomial $X_1^{m_1} \cdots X_n^{m_n}$ by the factor $m_1 \cdots m_n$. Since both H_n and H_n^\bullet only involve monomials with $m_i > 0$, the series DH_n and DH_n^\bullet determine uniquely the original series H_n and H_n^\bullet , respectively.

4.1. COMPLETED n -POINT FUNCTION. — We have from (12)

$$(29) \quad DH_n^\bullet = \sum_{m_1, \dots, m_n=1}^{\infty} X_1^{m_1} \cdots X_n^{m_n} \langle 0 | J_{m_1} \cdots J_{m_n} \mathcal{D}(\hbar) e^{\sum_{i=1}^{\infty} s_i J_{-i}/i \hbar} | 0 \rangle.$$

Define the *completed* version of this function by

$$(30) \quad \widehat{DH}_n^\bullet = \sum_{m_1, \dots, m_n=-\infty}^{\infty} X_1^{m_1} \cdots X_n^{m_n} \langle 0 | J_{m_1} \cdots J_{m_n} \mathcal{D}(\hbar) e^{\sum_{i=1}^{\infty} s_i J_{-i}/i \hbar} | 0 \rangle$$

and the corresponding completed connected functions $\widehat{DH}_n = \sum_{g=0}^{\infty} \hbar^{2g-2+n} \widehat{DH}_{g,n}$ through exactly the same inclusion-exclusion relations as the ones in (5).

These are infinite power series that involve both positive and negative powers of the variables X_i . The advantage of using completed versions of n -point functions is that they are better adapted to convolving in a closed form, as we shall see below.

PROPOSITION 4.1. — *We have*

$$(31) \quad \widehat{DH}_n = DH_n + \delta_{2,n} \frac{X_1 X_2}{(X_1 - X_2)^2},$$

where the last summand is considered as its power expansion over X_1/X_2 :

$$\frac{X_1 X_2}{(X_1 - X_2)^2} = \sum_{m=1}^{\infty} m (X_1/X_2)^m.$$

In other words, for $(g, n) \neq (0, 2)$ we have

$$\widehat{DH}_{g,n} = DH_{g,n}$$

and

$$(32) \quad \widehat{DH}_{0,2} = DH_{0,2} + \frac{X_1 X_2}{(X_1 - X_2)^2}.$$

As a corollary, for $(g, n) \neq (0, 2)$ the series $\widehat{DH}_{g,n}$ only involves positive powers of the variables X_i .

Proof. — Denote

$$\nabla_i^+ = \sum_{m=1}^{\infty} X_i^m J_m = \sum_{m=1}^{\infty} m X_i^m \partial_{p_m}, \quad \nabla_i^- = \sum_{m=1}^{\infty} X_i^{-m} J_{-m} = \sum_{m=1}^{\infty} X_i^{-m} p_m.$$

Using these operators, we can rewrite (29) and (30) as

$$\begin{aligned} DH_n^\bullet &= \nabla_1^+ \cdots \nabla_n^+ Z \Big|_{p=0}, \\ \widehat{DH}_n^\bullet &= (\nabla_1^+ + \nabla_1^-) \cdots (\nabla_n^+ + \nabla_n^-) Z \Big|_{p=0}. \end{aligned}$$

Let us expand brackets in the last equation. By the Leibniz rule, the partial derivatives entering ∇_i^+ are applied to either the linear functions entering ∇_j^- for some $j > i$ or to Z . Therefore, we obtain

$$(33) \quad \widehat{DH}_n^\bullet = \sum_{\{1, \dots, n\} = \sqcup_k \{i_k, j_k\} \sqcup K} \left(\prod_k \frac{X_{i_k} X_{j_k}}{(X_{i_k} - X_{j_k})^2} \right) DH_{|K|}^\bullet(X_K),$$

where the factor $X_i X_j / (X_i - X_j)^2$ for $i < j$ is considered as a power expansion

$$\nabla_i^+ \sum_{m=1}^{\infty} X_j^{-m} p_m = \sum_{m=1}^{\infty} m (X_i / X_j)^m = \frac{X_i X_j}{(X_i - X_j)^2}.$$

By the inclusion-exclusion relations, Equation (33) is equivalent to the relations of the proposition. In order to see this, we observe that if we *define* the connected functions \widehat{DH}_n by (31) then the corresponding disconnected functions are given exactly by (33). \square

4.2. COMPUTATION OF COMPLETED n -POINT FUNCTIONS. — The computation of H_n^\bullet and H_n of the previous section can be extended to the computation of the completed n -point functions \widehat{DH}_n^\bullet and \widehat{DH}_n . The proofs of the corresponding statements for \widehat{DH}_n^\bullet and \widehat{DH}_n are similar to the ones from the previous section, one just needs to extend all summations over $m_i \geq 1$ to summations over $m_i \in \mathbb{Z}$.

Define the transformation U taking a Laurent series $f(u, z)$ in u and z to the Laurent series

$$(Uf)(X) = \sum_{m=-\infty}^{\infty} X^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y) \Big|_{y=0} [z^m u^r] \frac{e^{u\mathcal{S}(u\hbar z \partial_z)y(z)}}{u\hbar\mathcal{S}(u\hbar)} f(u, z).$$

It differs from the transformation U^+ of Definition 3.3 by an extra factor m of the summands and by the summation range of the integer index m . Thus Uf is a Laurent series and might involve negative powers of X . We denote also by U_k a similar transformation applied to u_k and z_k instead of u and z (the output of U_k is a Laurent series in X_k).

Then, similarly to the computation of H_n^\bullet we obtain

$$\begin{aligned} \widehat{DH}_n^\bullet &= \sum_{m_1, \dots, m_n = -\infty}^{\infty} X_1^{m_1} \cdots X_n^{m_n} \langle 0 | J_{m_1} \cdots J_{m_n} \mathcal{D}(\hbar) e^{\sum_{i=1}^{\infty} s_i J_{-i} / i \hbar} | 0 \rangle \\ &= \sum_{m_1, \dots, m_n = -\infty}^{\infty} X_1^{m_1} \cdots X_n^{m_n} \langle 0 | \mathbb{J}_{m_1} \cdots \mathbb{J}_{m_n} e^{\sum_{i=1}^{\infty} s_i J_{-i} / i \hbar} | 0 \rangle \\ &= U_n \cdots U_1 \prod_{1 \leq k < \ell \leq n} e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2}, \end{aligned}$$

where the expression in the product on the right-hand side is understood as its power asymptotic expansion in the sector $|z_1| \ll \cdots \ll |z_n| \ll 1$.

Next, the analogue of the computation of H_n of the previous section is the following equation:

$$(34) \quad \widehat{DH}_n = U_n \cdots U_1 \sum_{\gamma \in \Gamma_n} \prod_{\{v_k, v_\ell\} \in E_\gamma} \left(e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2} - 1 \right),$$

where Γ_n is the set of all connected simple graphs over n vertices v_1, \dots, v_n , and E_γ is the set of edges of $\gamma \in \Gamma_n$.

Since, by Proposition 4.1, \widehat{DH}_n differs from DH_n by a small correction for $n = 2$, we conclude:

COROLLARY 4.2. — For $(g, n) \neq (0, 2)$ we have

$$(35) \quad DH_{g,n} = [\hbar^{2g-2+n} \left(U_n \cdots U_1 \sum_{\gamma \in \Gamma_n} \prod_{\{v_k, v_\ell\} \in E_\gamma} \left(e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) z_k z_\ell / (z_k - z_\ell)^2} - 1 \right) \right)].$$

In particular, all the terms on the right-hand side containing non-positive powers of the variables X_i cancel out.

4.3. PRINCIPAL IDENTITY. — Recall that the transformation U entering the formulas of the previous section acts on a Laurent series $f(u, z)$ in z and u by

$$(Uf)(X) = \sum_{m=-\infty}^{\infty} X^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y) \Big|_{y=0} [z^m u^r] \frac{e^{u \mathcal{S}(u \hbar z \partial_z) y(z)}}{u \hbar \mathcal{S}(u \hbar)} f(u, z).$$

The result of this transformation is a function in X . Up to this point we regarded X and z as independent variables. From now on we assume that they are related by the change $X = X(z)$, where

$$(36) \quad X(z) = z e^{-\psi(y(z))}.$$

Through this change we have

$$D := X \frac{\partial}{\partial X} = \frac{1}{Q} z \frac{\partial}{\partial z},$$

where

$$Q := \frac{z}{X dz/dX} = \frac{z}{X} \frac{dX}{dz} = 1 - D\psi(y) = 1 - z\psi'(y)y'(z).$$

Thus we have

$$z \frac{\partial}{\partial z} = QD.$$

Having this change in mind we treat the result of the transformation U as a function (a Laurent series) in z . We claim that U acts on the coefficients of positive powers of u as a differential operator. More explicitly, define

$$(37) \quad \begin{aligned} L_0(v, y, \hbar) &:= e^{v \left(\frac{\mathcal{S}(v\hbar\partial_y)}{\mathcal{S}(\hbar\partial_y)} - 1 \right) \psi(y)}, \\ L_r(v, y, \hbar) &:= e^{-v\psi(y)} \partial_y^r e^{v\psi(y)} L_0(v, y, \hbar) = (\partial_y + v\psi'(y))^r L_0(v, y, \hbar). \end{aligned}$$

The function $L_r(v, y, \hbar)$ is a series in \hbar^2 whose coefficients are polynomials in v and the higher order derivatives of $\psi(y)$.

The following *principal identity* plays a central role in the proof of the main theorems 4.8 and 5.3 below.

PROPOSITION 4.3. — *Let $H(u, z)$ be arbitrary Laurent series in z whose coefficients are either polynomials in u or infinite series in \hbar such that the coefficient of any power of \hbar is a polynomial in u . Then the following identity holds true:*

$$(38) \quad \begin{aligned} \sum_{m=-\infty}^{\infty} \sum_{r=0}^{\infty} \partial_y^r \phi_m(y) \Big|_{y=0} X^m [z^m u^r] e^{u y(z)} H(u, z) \\ = \sum_{j,r=0}^{\infty} D^j \left(\frac{[v^j] L_r(v, y(z), \hbar)}{Q} [u^r] H(u, z) \right), \end{aligned}$$

where $X = X(z)$ on the left-hand side is given by (36).

Applying this identity to a function of the form

$$H(u, z) = \frac{e^{u(\mathcal{S}(u\hbar Q D) - 1)y(z)}}{u\hbar \mathcal{S}(u\hbar)} f(u, z),$$

we conclude:

COROLLARY 4.4. — *Assume that $f(u, z)$ is a Laurent series in z whose coefficients are polynomials in u of bounded degree and with zero free term. Then the action of the transformation U on f is given by*

$$(39) \quad (Uf)(z) = \sum_{j,r=0}^{\infty} D^j \left(\frac{[v^j] L_r(v, y(z), \hbar)}{Q} [u^r] \frac{e^{u(\mathcal{S}(u\hbar Q D) - 1)y(z)}}{u\hbar \mathcal{S}(u\hbar)} f(u, z) \right).$$

4.4. PROOF OF THE PRINCIPAL IDENTITY. — The proof of the principal identity is split into several lemmas.

LEMMA 4.5. — *Let $\Phi(y)$ and $H(u)$ be two arbitrary regular series. Then*

$$(40) \quad \sum_{r=0}^{\infty} \partial_y^r \Phi(y) \Big|_{y=0} [u^r] e^{uy} H(u) = \sum_{r=0}^{\infty} \partial_y^r \Phi(y) [u^r] H(u).$$

Proof. — We have:

$$\begin{aligned} \sum_{r=0}^{\infty} \partial_y^r \Phi(y) \Big|_{y=0} [u^r] e^{uy} H(u) &= \sum_{r,k=0}^{\infty} \partial_y^{r+k} \Phi(y) \Big|_{y=0} ([u^k] e^{uy}) ([u^r] H(u)) \\ &= \sum_{r,k=0}^{\infty} \partial_y^{r+k} \Phi(y) \Big|_{y=0} \frac{y^k}{k!} [u^r] H(u) \\ &= \sum_{r=0}^{\infty} \partial_y^r \Phi(y) [u^r] H(u). \quad \square \end{aligned}$$

LEMMA 4.6 ([Kaz21]). — *We have:*

$$(41) \quad \phi_m(y) = e^{m\psi(y)} L_0(m, y, \hbar),$$

$$(42) \quad \partial_y^r \phi_m(y) = e^{m\psi(y)} L_r(m, y, \hbar).$$

Proof. — Note that $e^{a\partial_y} f(y) = f(y + a)$. We have (for $m \in \mathbb{Z}_{>0}$):

$$\begin{aligned} e^{m\psi(y)} L_0(m, y, \hbar) &\stackrel{(37)}{=} e^{m\psi(y)} e^{m\left(\frac{\mathcal{S}(m\hbar\partial_y)}{\mathcal{S}(\hbar\partial_y)} - 1\right)\psi(y)} = \exp\left(\frac{e^{(m/2)\hbar\partial_y} - e^{-(m/2)\hbar\partial_y}}{e^{(1/2)\hbar\partial_y} - e^{-(1/2)\hbar\partial_y}} \psi(y)\right) \\ &= \exp\left(e^{-((m-1)/2)\hbar\partial_y} \frac{e^{m\hbar\partial_y} - 1}{e^{\hbar\partial_y} - 1} \psi(y)\right) = \exp\left(e^{-((m-1)/2)\hbar\partial_y} \left(\sum_{i=1}^m e^{(i-1)\hbar\partial_y}\right) \psi(y)\right) \\ &= \exp\left(\sum_{i=1}^m \psi\left(y + \left(i - 1 - ((m - 1)/2)\right)\hbar\right)\right) \stackrel{(22)}{=} \phi_m(y). \end{aligned}$$

The $m = 0$ case is trivial and the $m < 0$ case is analogous. Thus we have proved (41), and (42) is evident from the definition (37) of L_r . \square

We also need a certain form of what is known as the Lagrange-Bürmann formula:

LEMMA 4.7. — *For any Laurent series H in z and for any $m \in \mathbb{Z}$ we have*

$$[z^m] e^{m\psi(y)} H = [X^m] \frac{1}{Q} H,$$

and, therefore,

$$(43) \quad \sum_{m=-\infty}^{\infty} X^m [z^m] e^{m\psi(y)} H = \frac{1}{Q} H,$$

where $y = y(z)$ and the function on the right-hand side is regarded as a Laurent series in X through the change inverse to (36).

Proof. — We have:

$$[z^m] e^{m\psi(y)} H = \operatorname{res}_{z=0} \frac{e^{m\psi(y)} H}{z^{m+1}} dz = \operatorname{res}_{z=0} \frac{H}{z X^m} dz = \operatorname{res}_{z=0} \frac{H}{Q X^{m+1}} dX = [X^m] \frac{1}{Q} H. \quad \square$$

Now we are ready to prove the principal identity.

Proof of Proposition 4.3. — We have:

$$\begin{aligned} & \sum_{m=-\infty}^{\infty} X^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y) \Big|_{y=0} [z^m u^r] e^{u y(z)} H(u, z) \\ & \stackrel{(40)}{=} \sum_{m=-\infty}^{\infty} X^m [z^m] \sum_{r=0}^{\infty} [u^r] \partial_y^r \phi_m(y) \Big|_{y=y(z)} H(u, z) \\ & \stackrel{(42)}{=} \sum_{m=-\infty}^{\infty} X^m [z^m] \sum_{r=0}^{\infty} [u^r] e^{m\psi(y(z))} L_r(m, y(z), \hbar) H(u, z) \\ & = \sum_{j=0}^{\infty} D^j \sum_{m=-\infty}^{\infty} X^m [z^m] \sum_{r=0}^{\infty} [u^r] e^{m\psi(y(z))} [v^j] L_r(v, y(z), \hbar) H(u, z) \\ & \stackrel{(43)}{=} \sum_{j=0}^{\infty} D^j \sum_{r=0}^{\infty} \frac{[v^j] L_r(v, y(z), \hbar)}{Q} [u^r] H(u, z) \end{aligned}$$

(here we consider X and z as independent variables). □

4.5. A CLOSED FORMULA FOR $D_1 \cdots D_n H_n$. — The principal identity together with Corollary 4.2 lead to our first theorem, which is just one step away from the main result formulated in the next section:

THEOREM 4.8. — For $n \geq 2$, $(g, n) \neq (0, 2)$ we have

$$(44) \quad D_1 \cdots D_n H_{g,n} = [\hbar^{2g-2+n}] U_n \cdots U_1 \sum_{\gamma \in \Gamma_n} \prod_{\{v_k, v_\ell\} \in E_\gamma} w_{k,\ell}$$

where

$$(45) \quad w_{k,\ell} = e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar Q_k D_k) \mathcal{S}(u_\ell \hbar Q_\ell D_\ell) z_k z_\ell / (z_k - z_\ell)^2} - 1$$

and U_i is the operator of Proposition 4.3 acting on a function f in u_i and z_i by

$$U_i f = \sum_{j,r=0}^{\infty} D_i^j \left(\frac{[v^j] L_r(v, y(z_i), \hbar)}{Q_i} [u_i^r] \frac{e^{u_i (\mathcal{S}(u_i \hbar Q_i D_i) - 1) y(z_i)}}{u_i \hbar \mathcal{S}(u_i \hbar)} f(u_i, z_i) \right).$$

As before, the sum is over all connected simple graphs on n labeled vertices.

For fixed g and n , after taking the coefficient $[\hbar^{2g-2+n}]$, all sums in this formula for $(\prod_{i=1}^n D_i) H_{g,n}$ become finite, and it becomes a rational expression in z_1, \dots, z_n and the derivatives of the functions $y_i = y(z_i)$ and $\psi(y_i)$.

The coefficient of any power of \hbar in $w_{i,j}$ is a polynomial in u_i and u_j vanishing at $u_i = 0$ and at $u_j = 0$ so that Corollary 4.4 can be applied. The restriction $n > 1$ is imposed because in the case $n = 1$ the operator U_1 is applied to the constant function 1, which is not divisible by u_1 , so the conclusion of Corollary 4.4 does not hold. The requirement $(g, n) \neq (0, 2)$ is a consequence of Corollary 4.1. The cases $n = 1$ and $(g, n) = (0, 2)$ are treated in Section 6 separately.

The (very important) finiteness statement is evident from the way \hbar and u_i enter the expression.

A nice property of the equality of Theorem 4.8 (that does not hold for earlier equalities of Proposition 3.6 and Corollary 4.2) is that it can be applied without expanding the involved functions in Laurent series and is valid in the ring

$$R := \mathbb{C}[[z_1, \dots, z_n]][\{(z_i - z_j)^{-1}; i, j \in \{1, \dots, n\}, i < j\}]$$

of functions with finite order poles on the diagonals $z_i = z_j$. Note that $\mathbb{C}[[z_1, \dots, z_n]]$ can be considered as a subring of this ring, corresponding to the expressions where all factors $(z_i - z_j)^{-1}$ have degree zero.

REMARK 4.9. — Note that the statement of Theorem 4.8 still holds if one allows $\psi(z)$ and $y(z)$ to also be formal series in \hbar^2 . More precisely, an analogous statement can be proved in a very similar way if one puts

$$\psi(\hbar^2, y) := \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} c_{k,m} y^k \hbar^{2m}, \quad y(\hbar^2, z) := \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} s_{k,m} z^k \hbar^{2m},$$

while still keeping the formula for $X(z)$ free of \hbar , i.e., using

$$X(z) = z e^{-\psi(y(z))} \Big|_{\hbar=0}$$

in place of (6), see [BDBKS20, §2].

Theorem 4.8 has an important corollary:

COROLLARY 4.10. — All diagonal poles (i.e., poles at $z_i = z_j$ for $i \neq j$) on the right-hand side of (44) cancel out; in other words, the expression in the right-hand side of (44) actually belongs to the subring $\mathbb{C}[[z_1, \dots, z_n]]$ of the ring R .

Proof. — Indeed, the left-hand side of (44), by definition, is a formal series in X_1, \dots, X_n (containing only positive powers of X_i), into which we plug $X_i = X(z_i)$, where $X(z)$ is a formal series in z (again, containing only positive powers of z). Thus, the expression in the left-hand side of (44) manifestly belongs to the subring $\mathbb{C}[[z_1, \dots, z_n]]$ of the ring R , and thus, by Theorem 4.8, so does the right-hand side. \square

At the end of this section we state several reformulations of Theorem 4.8. First, substituting the definitions of $w_{k,\ell}$ and U_k to (44), we get, explicitly,

$$(46) \quad D_1 \cdots D_n H_{g,n} = [\hbar^{2g-2+n}] \sum_{j_1, \dots, j_n, r_1, \dots, r_n=0}^{\infty} \left(\prod_{i=1}^n D_i^{j_i} \right) \times \left(\prod_{i=1}^n \frac{[v^{j_i}]_{L_{r_i}(v, y(z_i), \hbar)}}{Q_i} [\prod_{i=1}^n u_i^{r_i}] \prod_{i=1}^n \frac{e^{u_i(\mathcal{S}(u_i \hbar Q_i D_i) - 1)y(z_i)}}{u_i \hbar \mathcal{S}(u_i \hbar)} \sum_{\gamma \in \Gamma_n} \prod_{\{v_k, v_\ell\} \in E_\gamma} \left(e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar Q_k D_k) \mathcal{S}(u_\ell \hbar Q_\ell D_\ell) z_k z_\ell / (z_k - z_\ell)^2} - 1 \right) \right).$$

Next, expanding the exponential in a series we can represent the last formula in the following even more explicit form:

$$(47) \quad D_1 \cdots D_n H_{g,n} = [\hbar^{2g-2+n}] \sum_{j_1, \dots, j_n, r_1, \dots, r_n=0}^{\infty} \left(\prod_{i=1}^n D_i^{j_i} \right) \times \left(\prod_{i=1}^n \frac{[v^{j_i}] L_{r_i}(v, y(z_i), \hbar)}{Q_i} \prod_{i=1}^n [u_i^{r_i}] \prod_{i=1}^n \frac{e^{u_i(\mathcal{S}(u_i \hbar Q_i D_i - 1)y(z_i))}}{u_i \hbar \mathcal{S}(u_i \hbar)} \right) \sum_{\gamma \in \tilde{\Gamma}_n} \frac{\hbar^{2|E_\gamma|}}{|\text{Aut}_\gamma|} \prod_{\{v_k, v_\ell\} \in E_\gamma} u_k u_\ell \mathcal{S}(u_k \hbar z_k \partial_{z_k}) \mathcal{S}(u_\ell \hbar z_\ell \partial_{z_\ell}) \frac{z_k z_\ell}{(z_k - z_\ell)^2}.$$

Here $\tilde{\Gamma}_n$ is the set of all connected graphs on n labeled vertices v_1, \dots, v_n , with multiple edges allowed but no loops (i.e., no edges connecting a vertex to itself). Both Equations (46) and (47) hold for $n > 1$ and $(g, n) \neq (0, 2)$.

5. GENERAL FORMULA

In this section we prove the main theorem of the present paper, which explicitly represents $H_{g,n}$ for given g and n in a closed form. What remains is to get rid of $D_1 \cdots D_n$ which are applied in the LHS in Theorem 4.8. Let us introduce in a formal way the operator $D_i^{-1} U_i$ acting on a function $f(u_i, z_i)$ by

$$(48) \quad D_i^{-1} U_i f = \sum_{j,r=0}^{\infty} D_i^{j-1} \left(\frac{[v^j] L_r(v, y(z_i), \hbar)}{Q_i} [u_i^r] \frac{e^{u_i(\mathcal{S}(u_i \hbar Q_i D_i - 1)y(z_i))}}{u_i \hbar \mathcal{S}(u_i \hbar)} f(u_i, z_i) \right),$$

where we define the action of D_i^{-1} on a function $w(z_i)$ by

$$(49) \quad (D_i^{-1} w)(z_i) = \int_0^{z_i} \frac{Q(z)}{z} w(z) dz.$$

Note that this formal definition of the operator $D_i^{-1} U_i$ implies that

$$(50) \quad D_i (D_i^{-1} U_i f) = U_i f.$$

Then we set

$$(51) \quad \tilde{H}_{g,n} = [\hbar^{2g-2+n}] \left(\prod_{i=1}^n D_i^{-1} U_i \right) \sum_{\gamma \in \tilde{\Gamma}_n} \prod_{\{v_i, v_j\} \in E_\gamma} w_{i,j},$$

where $w_{i,j}$ is given by (45).

Now we formulate the propositions needed to prove the theorem; their proofs are given below at the end of this section.

PROPOSITION 5.1. — *Assume that $n \geq 2$ and $(g, n) \neq (0, 2)$. Then each time when the operator D_i^{-1} defined by (49) is applied in the expression (51) for $\tilde{H}_{g,n}$ the corresponding integrated differential form $(Q(z)/z)w(z) dz$ is rational in z with possible poles at $z = z_j$ for $j \neq i$ with zero residues. It follows that its primitive is well-defined as a rational function in z_i and the whole function $\tilde{H}_{g,n}$ is well-defined and has the form as in Theorem 1.1 up to an additive constant.*

By construction, we have $D_1 \cdots D_n H_{g,n} = D_1 \cdots D_n \tilde{H}_{g,n}$. This does not imply that the functions $H_{g,n}$ and $\tilde{H}_{g,n}$ are equal.

PROPOSITION 5.2. — For $n \geq 2$ and $(g, n) \neq (0, 2)$, the difference between $H_{g,n}$ and $\tilde{H}_{g,n}$ is the following constant:

$$(52) \quad H_{g,n} = \tilde{H}_{g,n} - (-1)^{n-1} \psi^{(2g+n-2)}(0) [u^{2g}] \frac{1}{\mathcal{S}^2(u)}.$$

Proposition 5.2 directly implies the main theorem:

THEOREM 5.3. — For $n \geq 2$ and $(g, n) \neq (0, 2)$ we have:

$$(53) \quad H_{g,n} = [\hbar^{2g-2+n}] \left(\prod_{i=1}^n D_i^{-1} U_i \right) \sum_{\gamma \in \Gamma_n} \prod_{\{v_i, v_j\} \in E_\gamma} w_{i,j} + (-1)^n \psi^{(2g+n-2)}(0) [u^{2g}] \frac{1}{\mathcal{S}^2(u)},$$

where Γ_n is the set of simple graphs on n vertices v_1, \dots, v_n with edges E_γ , $w_{i,j}$ is given by (45), and $D_i^{-1} U_i$ is given by (48)–(49).

For fixed g and n , after taking the coefficient $[\hbar^{2g-2+n}]$, this formula turns into a rational expression in z_1, \dots, z_n and the derivatives of the functions $y_i = y(z_i)$ and $\psi(y_i)$.

REMARK 5.4. — Note that the structure of the obtained answer agrees with that suggested by Theorem 1.1. Thus, we have proved Theorem 1.1 in the case $n \geq 2$, $(g, n) \neq (0, 2)$. The special cases $n = 1$ and $(g, n) = (0, 2)$ are treated in the next section.

REMARK 5.5. — Let us also provide another form of the statement of the main theorem (narrowing it slightly to $n \geq 3$), where all integrals (49) are taken explicitly. Namely, for $n \geq 3$ we have:

$$(54) \quad H_{g,n} = [\hbar^{2g-2+n}] \sum_{\gamma \in \Gamma_n} \prod_{v_i \in \mathcal{S}_\gamma} \bar{U}_i \prod_{\{v_i, v_k\} \in E_\gamma \setminus \mathcal{K}_\gamma} w_{i,k} \\ \times \prod_{\{v_i, v_k\} \in \mathcal{K}_\gamma} \left(\bar{U}_i w_{i,k} + \hbar u_k \mathcal{S}(u_k \hbar Q_k D_k) \frac{z_i}{z_k - z_i} \right) + (-1)^n \psi^{(2g+n-2)}(0) [u^{2g}] \frac{1}{\mathcal{S}^2(u)},$$

where Γ_n is the set of simple graphs on n vertices v_1, \dots, v_n , E_γ is the set of edges of a graph γ , \mathcal{S}_γ is the subset of vertices which are not leaves and \mathcal{K}_γ is the subset of edges with one end v_i of valency 1 and another end v_k , and where

$$\bar{U}_i f = \sum_{r=0}^{\infty} \sum_{j=1}^{\infty} D_i^{j-1} \left(\frac{[v^j] L_r(v, y(z_i), \hbar)}{Q_i} [u_i^r] \frac{e^{u_i (\mathcal{S}(u_i \hbar Q_i D_i) - 1) y(z_i)}}{u_i \hbar \mathcal{S}(u_i \hbar)} f \right), \\ w_{k,\ell} \stackrel{(45)}{=} e^{\hbar^2 u_k u_\ell \mathcal{S}(u_k \hbar Q_k D_k) \mathcal{S}(u_\ell \hbar Q_\ell D_\ell) z_k z_\ell / (z_k - z_\ell)^2} - 1, \\ D_i \stackrel{(7)}{=} \frac{1}{Q_i} z \frac{\partial}{\partial z}, \quad L_r \stackrel{(37)}{=} (\partial_y + v \psi'(y))^r e^{v \left(\frac{\mathcal{S}(v \hbar \partial_y)}{\mathcal{S}(\hbar \partial_y)} - 1 \right) \psi(y)}, \\ Q_i \stackrel{(8)}{=} 1 - z y'(z) \psi'(y(z)), \quad \mathcal{S}(u) \stackrel{(17)}{=} \frac{e^{u/2} - e^{-u/2}}{u}.$$

(to help the reader, we included in this list the notation and definitions of some functions introduced earlier in the paper). For $n = 2, g > 0$, the form of statement (53) analogous to (54) is obtained in Section 6.4. For brevity we do not provide the proof of the general case (54) in this text, but it is rather similar to the $(g, 2)$ case of Section 6.4.

REMARK 5.6. — Note that a statement similar to the statement of Theorem 5.3, as in the case of Theorem 4.8, still holds if one allows $\psi(z)$ and $y(z)$ to also be formal series in \hbar^2 . More precisely, it still holds in a very similar form, if one puts

$$\psi(\hbar^2, y) := \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} c_{k,m} y^k \hbar^{2m}, \quad y(\hbar^2, z) := \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} s_{k,m} z^k \hbar^{2m},$$

while still keeping the formula for $X(z)$ free of \hbar , i.e., using

$$X(z) = z e^{-\psi(y(z))} \Big|_{\hbar=0}$$

in place of (6), see [BDBKS20, §3].

Analogously to the case of Theorem 4.8 and Corollary 4.10, Theorem 5.3 has a similar corollary (which is proved via exactly the same reasoning):

COROLLARY 5.7. — *All diagonal poles (i.e., poles at $z_i = z_j$ for $i \neq j$) on the right-hand side of (53) (and of (54)) cancel out; in other words, the respective expressions actually belong to the subring $\mathbb{C}[[z_1, \dots, z_n]]$ of the ring R .*

Now we provide the proofs of the propositions of the present section.

Proof of Proposition 5.1. — The operator D_i^{-1} appears in the summand with $j = 0$ in the definition of $D_i^{-1}U_i$. In the case $j = 0$ we have $[v^0]L_0(v, y, \hbar) = 1$ and, for $r > 0$, $[v^0]L_r(v, y, \hbar) = 0$. Therefore, the summand with $j = 0$ in (48) can be written as

$$(55) \quad D_i^{-1} \frac{1}{Q_i} [u_i^0] \frac{e^{u_i(\mathcal{S}(u_i \hbar Q_i D_i) - 1)y(z_i)}}{u_i \hbar \mathcal{S}(u_i \hbar)} f = D_i^{-1} \frac{1}{\hbar Q_i} [u_i^1] f.$$

Recall that we are interested in (51). Let us check for which graphs γ the product $\prod_{\{v_i, v_j\} \in E_\gamma} w_{i,j}$ has a non-vanishing linear term in u_i . By definition, $w_{i,j}$ is divisible by $u_i u_j$. It follows that if the vertex v_i has valency greater than 1 then the contribution of such graph to the sum has a vanishing linear term in u_i .

If the vertex i has valency 1 and is connected to the vertex k then up to a factor that does not depend on z_i the linear term in u_i is the following:

$$(56) \quad [u_i^1] w_{i,k} = \hbar^2 u_k \mathcal{S}(u_k \hbar Q_k D_k) \frac{z_i z_k}{(z_i - z_k)^2}.$$

The contribution of this term to the $j = 0$ part in the expression one obtains after substituting (48) in (51) is given (again, up to a factor not dependent on z_i)

by applying (55) to (56):

$$\begin{aligned}
 D_i^{-1} \frac{1}{\hbar Q_i} [u_i^1] w_{i,k} &= D_i^{-1} \frac{1}{Q_i} \hbar u_k \mathcal{S}(u_k \hbar Q_k D_k) \frac{z_i z_k}{(z_i - z_k)^2} \\
 (57) \qquad \qquad \qquad &= \hbar u_k \mathcal{S}(u_k \hbar Q_k D_k) z_k \int_0^{z_i} \frac{dz}{(z - z_k)^2} \\
 &= \hbar u_k \mathcal{S}(u_k \hbar Q_k D_k) \frac{z_i}{z_k - z_i}.
 \end{aligned}$$

This function is rational in z_i , as required.

This proves the main part of the statement of Proposition 5.1 in the case $n > 2$. Indeed, we assumed implicitly in the above arguments that the leaf v_i is connected to a vertex v_k which is not a leaf so that D_i^{-1} and D_k^{-1} are not applied simultaneously. This is always the case for a connected graph with the number of vertices $n > 2$. If $n = 2$ then there could be summands linear both in u_1 and u_2 but these summands contribute to the case $g = 0$ only. Therefore, the conclusion of the proposition holds in the case $n = 2$ as well if $g > 0$.

The fact that $\tilde{H}_{g,n}$ has the form as in Theorem 1.1 then follows from the way \hbar and u_i enter the expression (51) (similarly to what happened in Theorem 4.8). The coefficient $[\hbar^{2g-2+n}]$ becomes a finite rational expression of the form described in Theorem 1.1, up to an additive constant. \square

Proof of Proposition 5.2. — We regard all considered functions as elements of the ring R . Let us denote by I the ideal $(z_1 \cdots z_n)$ generated by the product of coordinate functions. $H_{g,n}$ itself lies in I , and we have, by construction (from Equations (44), (51), and (50)),

$$D_1 \cdots D_n H_{g,n} = D_1 \cdots D_n \tilde{H}_{g,n}.$$

Therefore, it suffices to show that the right-hand side of (52) belongs to I . Let us compute $\tilde{H}_{g,n}$ modulo I .

Let $n \geq 3$. From the proof of Proposition 5.1 it follows that each internal edge $\{v_i, v_k\}$ of a graph γ in the sum (51) brings a factor of $z_i z_k$. Indeed, $w_{i,k}$ itself belongs to the ideal $(z_i z_k)$, and the $j = 0$ terms in the sums (48) for $D^{-1}U_i$ and $D^{-1}U_k$ vanish, while the $j > 0$ terms cannot affect the property of divisibility by $z_i z_k$.

On the other hand, if v_i is a leaf (connected to some v_k of valence greater than 1), then $\{v_i, v_k\} \in E_\gamma$ brings a factor of z_i , since, as above, $w_{i,k}$ is divisible by $z_i z_k$ and the $j > 0$ terms of (48) cannot affect this property, while the $j = 0$ term takes the form $\hbar u_k \mathcal{S}(u_k \hbar Q_k D_k) z_i / (z_k - z_i)$ (from Equation (57)), which is divisible by z_i .

Note that since $n \geq 3$ and the graphs are connected all edges belong to one of the above two cases.

Thus the contribution of the whole graph γ is not divisible by z_k for some k only if the vertex k is internal and all adjacent vertices are leaves. In this case the graph is the star with one vertex (labeled by k) of valency $n - 1 \geq 2$ and $n - 1$ vertices of valency 1. We conclude that the contribution of all but the star graphs belong to I .

The star graphs produce the following contributions:

$$\begin{aligned} \tilde{H}_{g,n} + I &= [\hbar^{2g-2+n}] \sum_{k=1}^n D_k^{-1} U_k \prod_{i \neq k} D_i^{-1} U_i \prod_{i \neq k} w_{i,k} + I \\ &= [\hbar^{2g-2+n}] \sum_{k=1}^n \sum_{j,r=0}^{\infty} D_k^{j-1} \frac{[v^j] L_r(v, y(z_k), \hbar)}{Q_k} \\ &\quad \times [u_k^r] \frac{e^{u_k(\mathcal{S}(u_k \hbar Q_k D_k) - 1)y(z_k)}}{u_k \hbar \mathcal{S}(u_k \hbar)} \prod_{i \neq k} D_i^{-1} U_i w_{i,k} + I. \end{aligned}$$

Note that all $j = 0$ terms vanish since v_k is an internal vertex (with valence ≥ 2), as discussed in the proof of Proposition 5.1. Thus we have

$$\begin{aligned} \tilde{H}_{g,n} + I &= [\hbar^{2g-2+n}] \sum_{k=1}^n \sum_{\substack{j \geq 1 \\ r \geq 0}} D_k^{j-1} \frac{[v^j] L_r(v, y(z_k), \hbar)}{Q_k} [u_k^r] \frac{e^{u_k(\mathcal{S}(u_k \hbar Q_k D_k) - 1)y(z_k)}}{u_k \hbar \mathcal{S}(u_k \hbar)} \\ &\quad \prod_{i \neq k} \sum_{j_i, r_i=0}^{\infty} D_i^{j_i-1} \left(\frac{[v^{j_i}] L_{r_i}(v, y(z_i), \hbar)}{Q_i} [u_i^{r_i}] \frac{e^{u_i(\mathcal{S}(u_i \hbar Q_i D_i) - 1)y(z_i)}}{u_i \hbar \mathcal{S}(u_i \hbar)} w_{i,k} \right) + I. \end{aligned}$$

Now note that if any of $j_i > 0$ then the corresponding term is divisible by z_k since $w_{i,k}$ is divisible by z_k and it gets acted upon only by operators of the sort D_k^m and D_i^m for $m \geq 0$ which do not spoil this property. Thus, we can factor out all these terms and we get, applying also formula (57),

$$\begin{aligned} (58) \quad \tilde{H}_{g,n} + I &= [\hbar^{2g-2+n}] \sum_{k=1}^n \sum_{\substack{j \geq 1 \\ r \geq 0}} D_k^{j-1} \frac{[v^j] L_r(v, y(z_k), \hbar)}{Q_k} \\ &\quad \times [u_k^r] \frac{e^{u_k(\mathcal{S}(u_k \hbar Q_k D_k) - 1)y(z_k)}}{u_k \hbar \mathcal{S}(u_k \hbar)} \prod_{i \neq k} \hbar u_k \mathcal{S}(u_k \hbar Q_k D_k) \frac{z_i}{z_k - z_i} + I. \end{aligned}$$

Now we note that all summands with $j \geq 2$ are also divisible by z_k since $D_k = (1/Q_k) z_k \partial/\partial z_k$ and thus only the $j = 1$ term remains. Also note that $Q_k \equiv \mathcal{S}(u_k \hbar Q_k D_k) \equiv 1 \pmod{z_k}$. Taking this into account, we obtain

$$\tilde{H}_{g,n} + I = [\hbar^{2g+n-2}] \sum_{k=1}^n \sum_{r=0}^{\infty} [v^1] L_r(v, y_k, \hbar) [u_k^r] \frac{1}{u_k \hbar \mathcal{S}(u_k \hbar)} \prod_{i \neq k} u_k \hbar \frac{z_i}{z_k - z_i} + I.$$

We have

$$\begin{aligned} (59) \quad [v^1] L_r(v, y_k, \hbar) &\equiv [v^1] (\partial_{y_k} + v \psi'(y_k))^r \left(1 + v \left(\frac{1}{\mathcal{S}(\hbar \partial_{y_k})} - 1 \right) \psi(y_k) \right) \Big|_{y_k=0} \\ &\equiv \frac{\partial_{y_k}^r}{\mathcal{S}(\hbar \partial_{y_k})} \psi(y_k) \Big|_{y_k=0} \pmod{z_k}. \end{aligned}$$

Using the fact

$$\sum_{k=1}^n \prod_{i \neq k} \frac{z_i}{z_k - z_i} = (-1)^{n-1}$$

we finally obtain:

$$\tilde{H}_{g,n} + I = (-1)^{n-1} [\hbar^{2g+n-2}] \sum_{r=0}^{\infty} \frac{\partial_y^r}{\mathcal{S}(\hbar \partial_y)} \psi(y)|_{y=0} [u^r] \frac{(u\hbar)^{n-2}}{\mathcal{S}(u\hbar)} + I.$$

Note that we can reexpand the last sum in \hbar :

$$\begin{aligned} (-1)^{n-1} \sum_{r=0}^{\infty} \frac{\partial_y^r}{\mathcal{S}(\hbar \partial_y)} \psi(y)|_{y=0} [u^r] \frac{(u\hbar)^{n-2}}{\mathcal{S}(u\hbar)} &= (-1)^{n-1} \frac{1}{\mathcal{S}(\hbar \partial_y)} \frac{(\hbar \partial_y)^{n-2}}{\mathcal{S}(\hbar \partial_y)} \psi(y)|_{y=0} \\ &= (-1)^{n-1} \hbar^{n-2} \frac{1}{\mathcal{S}^2(\hbar \partial_y)} \psi^{(n-2)}(y)|_{y=0} \\ &= (-1)^{n-1} \sum_{g=0}^{\infty} \hbar^{2g+n-2} \psi^{(2g+n-2)}(0) [u^{2g}] \frac{1}{\mathcal{S}^2(u)}, \end{aligned}$$

thus

$$(60) \quad \tilde{H}_{g,n} + I = (-1)^{n-1} \psi^{(2g+n-2)}(0) [u^{2g}] \frac{1}{\mathcal{S}^2(u)} + I.$$

This concludes the proof for the $n \geq 3$ case.

For $n = 2, g > 0$, we have only one graph:

$$\begin{aligned} (61) \quad \tilde{H}_{g,2} + I &= [\hbar^{2g}] D_1^{-1} U_1 D_2^{-1} U_2 w_{1,2} + I \\ &= [\hbar^{2g}] \sum_{j_1, r_1=0}^{\infty} \frac{D_1^{j_1-1} [v^{j_1}] L_{r_1}(v, y(z_1), \hbar)}{Q_1} \\ &\quad \times [u_1^{r_1}] \frac{e^{u_1(\mathcal{S}(u_1 \hbar Q_1 D_1) - 1)y(z_1)}}{u_1 \hbar \mathcal{S}(u_1 \hbar)} \sum_{j_2, r_2=0}^{\infty} \frac{D_2^{j_2-1} [v^{j_2}] L_{r_2}(v, y(z_2), \hbar)}{Q_2} \\ &\quad \times [u_2^{r_2}] \frac{e^{u_2(\mathcal{S}(u_2 \hbar Q_2 D_2) - 1)y(z_2)}}{u_2 \hbar \mathcal{S}(u_2 \hbar)} w_{1,2} + I. \end{aligned}$$

Note that if both j_1 and j_2 are positive, then the corresponding terms are divisible by $z_1 z_2$, analogously to what happened above. For $j_1 = j_2 = 0$ we apply (55) and get

$$\begin{aligned} &[\hbar^{2g}] D_1^{-1} \frac{1}{\hbar Q_1} [u_1^1] D_2^{-1} \frac{1}{\hbar Q_2} [u_2^1] w_{1,2} \\ &= [\hbar^{2g}] D_1^{-1} \frac{1}{\hbar Q_1} [u_1^1] D_2^{-1} \frac{1}{\hbar Q_2} [u_2^1] \left(e^{\hbar^2 u_1 u_2 \mathcal{S}(u_1 \hbar Q_1 D_1) \mathcal{S}(u_2 \hbar Q_2 D_2) z_1 z_2 / (z_1 - z_2)^2} - 1 \right) \\ &= [\hbar^{2g}] D_1^{-1} \frac{1}{\hbar Q_1} D_2^{-1} \frac{1}{\hbar Q_2} \hbar^2 \frac{z_1 z_2}{(z_1 - z_2)^2}, \end{aligned}$$

which clearly vanishes for $g > 0$.

Thus the sum in (61) can be represented as combination of two sums, one for $j_1 = 0, j_2 > 0$, and the other for $j_1 > 0, j_2 = 0$. This is actually precisely formula (58) where one substitutes $n = 2$. Thus we have reduced this case to the case of arbitrary $n \geq 2$, so formula (60) holds here as well.

This completes the proof of Proposition 5.2. □

Proof of Theorem 5.3. — The proof of the main statement immediately follows from Proposition 5.2, while the rationality statement is implied by the respective rationality statement of Theorem 4.8. □

6. EXCEPTIONAL CASES

Let us remind the definition of the functions L_r :

$$L_0(v, y, \hbar) := e^{v\left(\frac{\mathcal{S}(v\hbar\partial_y)}{\mathcal{S}(\hbar\partial_y)} - 1\right)\psi(y)},$$

$$L_r(v, y, \hbar) := e^{-v\psi(y)}\partial_y^r e^{v\psi(y)}L_0(v, y, \hbar) = (\partial_y + v\psi'(y))^r L_0(v, y, \hbar).$$

In order to simplify the notation, we denote in Addedthe computations of this section

$$L_{r,i}^j = [v^j]L_r(v, y(z_i), \hbar).$$

Note that in the case $j = 0$ we have

$$L_{0,i}^0 = 1, \quad L_{r,i}^0 = 0 \quad (r > 0).$$

6.1. COMPUTATION OF THE $(0, 1)$ -TERM. — Extracting the terms with $g = 0$ in (27) for $n = 1$ and noting $\phi_m(y)|_{\hbar=0} = e^{m\psi(y)}$ and $\mathcal{S}(\hbar u)|_{\hbar=0} = 1$ we get

$$D_1H_{0,1} = [\hbar^{-1}]D_1U_1^+ 1 = \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r e^{m\psi(y)}|_{y=0} [z^m u^r] \frac{e^{u y(z)}}{u}.$$

In order to apply Lemma 4.5 to the right-hand side one needs to get rid of a pole in u at the origin. One of the possibilities to do that is to differentiate this expression:

$$\begin{aligned} D_1^2H_{0,1} &= \sum_{m=1}^{\infty} mX_1^m \sum_{r=0}^{\infty} \partial_y^r e^{m\psi(y)}|_{y=0} [z^m u^r] \frac{e^{u y(z)}}{u} \\ &= \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r e^{m\psi(y)}|_{y=0} [z^m u^r] z\partial_z \frac{e^{u y(z)}}{u} \\ &= \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r e^{m\psi(y)}|_{y=0} [z^m u^r] e^{u y(z)} QDy(z) \\ &= \sum_{m=1}^{\infty} X_1^m [z^m] \sum_{r=0}^{\infty} \partial_y^r e^{m\psi(y)}|_{y=0} \frac{y(z)^r}{r!} QDy(z) \\ &\stackrel{\prime}{=} \sum_{m=1}^{\infty} X_1^m [z^m] e^{m\psi(y(z))} QDy(z) \\ &\stackrel{\prime\prime}{=} \sum_{m=-\infty}^{\infty} X_1^m [z^m] e^{m\psi(y(z))} QDy(z) \\ &\stackrel{(43)}{=} D_1y(z_1). \end{aligned}$$

The equality $\stackrel{\prime}{=}$ is the Taylor series expansion, and the equality $\stackrel{\prime\prime}{=}$ we obtain from the fact that for all $m \in \mathbb{Z}_{\leq 0}$ holds $[z^m]e^{m\psi(y(z))} QDy(z) = 0$. The constant term equals zero (after one performs integration of the equality $D_1D_1H_{0,1} = D_1y(z_1)$) since both $D_1H_{0,1}$ and $y(z_1)$ are divisible by z_1 . This proves the equality

$$D_1H_{0,1} = y(z_1)$$

of Theorem 1.1.

6.2. COMPUTATION OF THE $(g, 1)$ -TERM, $g > 0$. — By (27), we have

$$\begin{aligned}
 \hbar D_1 H_1 &= \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] \frac{e^{u \mathcal{S}(u \hbar QD)y(z)}}{u \mathcal{S}(u \hbar)} \\
 (62) \quad &= \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] \left(\frac{e^{u \mathcal{S}(u \hbar QD)y(z)}}{u \mathcal{S}(u \hbar)} - \frac{e^{u y(z)}}{u} \right) \\
 &\quad + \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] \frac{e^{u y(z)}}{u}.
 \end{aligned}$$

The expression of the first summand is regular in u and we can apply the principal identity (38) to get

$$\begin{aligned}
 &\sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] \left(\frac{e^{u \mathcal{S}(u \hbar QD)y(z)}}{u \mathcal{S}(u \hbar)} - \frac{e^{u y(z)}}{u} \right) \\
 &= \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] e^{u y(z)} \left(\frac{e^{u (\mathcal{S}(u \hbar QD)-1)y(z)}}{u \mathcal{S}(u \hbar)} - \frac{1}{u} \right) \\
 (63) \quad &= \sum_{m=-\infty}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] e^{u y(z)} \left(\frac{e^{u (\mathcal{S}(u \hbar QD)-1)y(z)}}{u \mathcal{S}(u \hbar)} - \frac{1}{u} \right) \\
 &\stackrel{(38)}{=} \sum_{j,r=0}^{\infty} D_1^j \left(\frac{L_{r,1}^j}{Q_1} [u^r] \left(\frac{e^{u (\mathcal{S}(u \hbar QD)-1)y(z_1)}}{u \mathcal{S}(u \hbar)} - \frac{1}{u} \right) \right) \\
 &= \sum_{j,r=0}^{\infty} D_1^j \left(\frac{L_{r,1}^j}{Q_1} [u^r] \frac{e^{u (\mathcal{S}(u \hbar QD)-1)y(z_1)}}{u \mathcal{S}(u \hbar)} \right).
 \end{aligned}$$

In the second equality we used that $\phi_0 = 1$ from the definition (23) and the fact that the expression after $[z^m u^r]$ does not contain negative powers of z (we will use this switch from summation over m starting from 0 to summation over m starting at $-\infty$ again in what follows, where it is applicable, without further commenting on it). In the last equality the term $1/u$ disappears since the sum goes only over nonnegative r . Note that (63) can be obtained if we take formally the right-hand side of (46) for the case $n = 1$.

The second summand in the right-hand side of (62) can be computed by the differentiation trick similar to the case $g = 0$ above. We have:

$$\begin{aligned}
 &D_1 \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] \frac{e^{u y(z)}}{u} \\
 &= \sum_{m=1}^{\infty} X_1^m \sum_{r=0}^{\infty} \partial_y^r \phi_m(y)|_{y=0} [z^m u^r] e^{u y(z)} QDy(z) \\
 (38) \quad &\stackrel{(38)}{=} \sum_{j,r=0}^{\infty} D_1^j \left(\frac{[v^j] L_r(v, y(z), \hbar)}{Q} [u^r] QDy(z) \right) \\
 &= \sum_{j=0}^{\infty} D_1^j (L_{0,1}^j D_1 y(z_1)) = D_1 y(z_1) + D_1 \sum_{j=1}^{\infty} D_1^{j-1} (L_{0,1}^j D_1 y(z_1)).
 \end{aligned}$$

Putting together and using again that the constant of integration equals zero we conclude

$$D_1(\hbar H_1 - H_{0,1}) = \sum_{j=0}^{\infty} D_1^j \left(\sum_{r=0}^{\infty} \frac{L_{r,1}^j}{Q_1} [u^r] \frac{e^{u(\mathcal{S}(u\hbar QD)-1)y(z_1)}}{u\mathcal{S}(u\hbar)} + L_{0,1}^{j+1} D_1 y(z_1) \right),$$

i.e., for $g > 0$ we have

$$D_1 H_{g,1} = [\hbar^{2g}] \sum_{j=0}^{\infty} D_1^j \left(\sum_{r=0}^{\infty} \frac{L_{r,1}^j}{Q_1} [u^r] \frac{e^{u(\mathcal{S}(u\hbar QD)-1)y(z_1)}}{u\mathcal{S}(u\hbar)} + L_{0,1}^{j+1} D_1 y(z_1) \right).$$

Our next step is to invert the operator D_1 on the right-hand side of Equation 6.2. Possible problems can only appear in the case $j = 0$. Observe that the summand with $r = 0$ vanishes, which implies that the first summand in the term $j = 0$ also vanishes. The second summand in the term with $j = 0$ is equal to

$$L_{0,1}^1 D_1 y(z_1) = \left(\frac{1}{\mathcal{S}(\hbar\partial_y)} - 1 \right) \psi(y) \Big|_{y=y(z_1)} D_1 y(z_1) = D_1 \sum_{k=1}^{\infty} [u^{2k}] \frac{1}{\mathcal{S}(u\hbar)} \psi^{(2k-1)}(y(z_1)).$$

If we define for $g > 0$

$$(64) \quad \tilde{H}_{g,1} := [\hbar^{2g}] \sum_{j=1}^{\infty} D_1^{j-1} \left(\sum_{r=1}^{\infty} \frac{L_{r,1}^j}{Q_1} [u^r] \frac{e^{u(\mathcal{S}(u\hbar QD)-1)y(z_1)}}{u\mathcal{S}(u\hbar)} + L_{0,1}^{j+1} D_1 y(z_1) \right) + \left([u^{2g}] \frac{1}{\mathcal{S}(u)} \right) \psi^{(2g-1)}(y(z_1)),$$

then we have $D_1 H_{g,1} - D_1 \tilde{H}_{g,1} = 0$. This means that $H_{g,1}$ and $\tilde{H}_{g,1}$ may differ only by a constant. To determine this constant let us put $z_1 = 0$ in (64). The second term in the brackets in the first line vanishes, as well as all terms in the j -sum for $j > 1$, and the exponential and Q both turn into 1. Let

$$\frac{1}{\mathcal{S}(x)} = 1 + \sum_{k=1}^{\infty} \sigma_k x^{2k}.$$

The first line of (64) for $z_1 = 0$ turns into the following:

$$\begin{aligned} & [\hbar^{2g}] \sum_{r=1}^{\infty} L_{r,1}^1 \Big|_{y=0} [u^r] \frac{1}{u\mathcal{S}(u\hbar)} \stackrel{(59)}{=} [\hbar^{2g}] \sum_{r=1}^{\infty} \frac{\partial_y^r}{\mathcal{S}(\hbar\partial_y)} \psi(y) \Big|_{y=0} [u^r] \frac{1}{u\mathcal{S}(u\hbar)} \\ &= [\hbar^{2g}] \sum_{r=1}^{\infty} \frac{\partial_y^r}{\mathcal{S}(\hbar\partial_y)} \psi(y) \Big|_{y=0} [u^r] \left(\frac{1}{u\mathcal{S}(u\hbar)} - \frac{1}{u} \right) \\ &= [\hbar^{2g}] \sum_{r=2}^{\infty} \frac{\partial_y^{r-1}}{\mathcal{S}(\hbar\partial_y)} \psi(y) \Big|_{y=0} [u^r] \left(\frac{1}{\mathcal{S}(u\hbar)} - 1 \right) \\ &= [\hbar^{2g}] \sum_{r=2}^{\infty} \partial_y^{r-1} \left(1 + \sum_{k=1}^{\infty} \sigma_k \hbar^{2k} \partial_y^{2k} \right) \psi(y) \Big|_{y=0} [u^r] \left(\sum_{m=1}^{\infty} \sigma_m \hbar^{2m} u^{2m} \right) \\ &= [\hbar^{2g}] \left(\sum_{m=1}^{\infty} \sigma_m \hbar^{2m} \partial_y^{2m-1} \right) \left(1 + \sum_{k=1}^{\infty} \sigma_k \hbar^{2k} \partial_y^{2k} \right) \psi(y) \Big|_{y=0} \\ &= \psi^{(2g-1)}(y)(0) [u^{2g}] \left(\frac{1}{\mathcal{S}(u)^2} - \frac{1}{\mathcal{S}(u)} \right). \end{aligned}$$

Setting $z_1 = 0$ in the second line of (64) is trivial and we arrive at

PROPOSITION 6.1. — For $n = 1$ and $g > 0$ we have:

$$(65) \quad H_{g,1} = [\hbar^{2g}] \sum_{j=1}^{\infty} D_1^{j-1} \left(\sum_{r=1}^{\infty} \frac{L_{r,1}^j}{Q_1} [u^r] \frac{e^{u(\mathcal{S}(u\hbar QD)-1)y(z_1)}}{u \mathcal{S}(u\hbar)} + L_{0,1}^{j+1} D_1 y(z_1) \right) + \left([u^{2g}] \frac{1}{\mathcal{S}(u)} \right) \psi^{(2g-1)}(y(z_1)) - \left([u^{2g}] \frac{1}{\mathcal{S}(u)^2} \right) \psi^{(2g-1)}(0).$$

Note that the structure of this formula agrees with the statement of Theorem 1.1.

6.3. COMPUTATION OF THE $(0, 2)$ -TERM. — We have

$$\begin{aligned} D_1 D_2 H_{0,2} + \frac{X_1 X_2}{(X_1 - X_2)^2} &\stackrel{(32)}{=} \widehat{D} \widehat{H}_{0,2} \stackrel{(34)}{=} [\hbar^0] U_2 U_1 w_{1,2} \\ &\stackrel{(39)}{=} [\hbar^0] \sum_{j_1, j_2=0}^{\infty} D_1^{j_1} D_2^{j_2} \sum_{r_1, r_2=0}^{\infty} \frac{L_{r_1,1}^{j_1} L_{r_2,2}^{j_2}}{Q_1 Q_2} [u_1^{r_1} u_2^{r_2}] \frac{z_1 z_2}{(z_1 - z_2)^2} \\ &= [\hbar^0] \sum_{j_1, j_2=0}^{\infty} D_1^{j_1} D_2^{j_2} \frac{L_{0,1}^{j_1} L_{0,2}^{j_2}}{Q_1 Q_2} \frac{z_1 z_2}{(z_1 - z_2)^2} = \frac{1}{Q_1 Q_2} \frac{z_1 z_2}{(z_1 - z_2)^2}. \end{aligned}$$

Thus, we get

$$\begin{aligned} D_1 D_2 H_{0,2} &= \frac{1}{Q_1 Q_2} \frac{z_1 z_2}{(z_1 - z_2)^2} - \frac{X_1 X_2}{(X_1 - X_2)^2} \\ &= D_1 \left(\frac{1}{Q_2} \frac{z_1}{z_2 - z_1} - \frac{X_1}{X_2 - X_1} \right) = D_1 D_2 \log \left(\frac{z_1^{-1} - z_2^{-1}}{X_1^{-1} - X_2^{-1}} \right). \end{aligned}$$

The function $\tilde{H}_{0,2} = \log \left(\frac{z_1^{-1} - z_2^{-1}}{X_1^{-1} - X_2^{-1}} \right)$ represents a regular series vanishing at $z_1 = 0$ and at $z_2 = 0$ and satisfies $D_1 D_2 H_{0,2} = D_1 D_2 \tilde{H}_{0,2}$. Therefore, it coincides with $H_{0,2}$. This proves (9).

This completes the proof of remaining exceptional cases of Theorem 1.1.

6.4. COMPUTATION OF THE $(g, 2)$ -TERM, $g > 0$. — This case is actually already covered by Theorem 5.3, but we can present a more explicit form of the answer (in line with Remark 5.5). We have

$$\begin{aligned} D_1 D_2 H_{g,2} &\stackrel{(35)}{=} [\hbar^{2g}] U_2 U_1 w_{1,2} \\ &\stackrel{(56)}{=} [\hbar^{2g}] U_2 D_1 \left(\bar{U}_1 w_{1,2} + \hbar u_2 \mathcal{S}(u_2 \hbar Q_2 D_2) \frac{z_1}{z_2 - z_1} \right) \\ &= [\hbar^{2g}] D_1 \left(\bar{U}_1 U_2 w_{1,2} + U_2 \hbar u_2 \mathcal{S}(u_2 \hbar Q_2 D_2) \frac{z_1}{z_2 - z_1} \right) = D_1 D_2 \tilde{H}_{g,2}, \end{aligned}$$

where

$$\begin{aligned} \tilde{H}_{g,2} &= [\hbar^{2g}] \left(\bar{U}_1 \left(\bar{U}_2 w_{1,2} + \hbar u_1 \mathcal{S}(u_1 \hbar Q_1 D_1) \frac{z_2}{z_1 - z_2} \right) \right. \\ &\quad \left. + \bar{U}_2 \left(\hbar u_2 \mathcal{S}(u_2 \hbar Q_2 D_2) \frac{z_1}{z_2 - z_1} \right) \right). \end{aligned}$$

One extra term that we omitted here contributes only in the case $g = 0$, which we considered above.

Arguing as in the proof of Proposition 5.2, we conclude that $H_{g,2}$ and $\tilde{H}_{g,2}$ differ by a constant that is given by the same formula as in the general case, and we obtain:

PROPOSITION 6.2. — For $n = 2$ and $g > 0$ we have:

$$H_{g,2} = [\hbar^{2g}] \left(\bar{U}_1 \bar{U}_2 w_{1,2} + \bar{U}_1 \left(\hbar u_1 \mathcal{S}(u_1 \hbar Q_1 D_1) \frac{z_2}{z_1 - z_2} \right) + \bar{U}_2 \left(\hbar u_2 \mathcal{S}(u_2 \hbar Q_2 D_2) \frac{z_1}{z_2 - z_1} \right) \right) + \psi^{(2g)}(0) [u^{2g}] \frac{1}{\mathcal{S}^2(u)}.$$

Remark that the structure of the obtained answer agrees with that suggested by Theorem 1.1 and correlates with Equation (54).

7. APPLYING GENERAL FORMULA

In this section we derive explicit expressions for $H_{g,n}$ for small g and n in terms of small number of basic functions. These functions include:

$$\begin{aligned} \psi_i^{(k)} &= \psi^{(k)}(y(z_i)), \quad k \geq 1, \\ y_i^{[k]} &= (z_i \partial_{z_i})^k y(z_i), \quad k \geq 1, \\ Q_i &= Q(z_i) = 1 - \psi'_i y_i^{[1]}. \end{aligned}$$

If $n = 1$ we set $z_1 = z$ and drop the lower index $i = 1$. In the case $n \geq 2$ we will use also the functions

$$\begin{aligned} \gamma_{i,j} &= \gamma_{j,i} = \frac{z_i z_j}{(z_i - z_j)^2}, \\ \gamma_{i,j}^{[k]} &= (-1)^k \gamma_{j,i}^{[k]} = (z_i \partial_{z_i})^k \gamma_{i,j}, \quad k \geq 0. \end{aligned}$$

Then, according to (the proof of) Proposition 5.1, the application of D_i^{-1} is reduced to

$$D_i^{-1}(Q_i \gamma_{i,j}^{[k]}) = (-1)^k D_i^{-1}(Q_i \gamma_{j,i}^{[k]}) = (z_i \partial_{z_i})^{-1} \gamma_{i,j}^{[k]} := \gamma_{i,j}^{[k-1]}.$$

This formula can be applied also for $k = 0$ if we set, in addition,

$$\gamma_{i,j}^{[-1]} = -1 - \gamma_{j,i}^{[-1]} = \frac{z_i}{z_j - z_i}.$$

7.1. COMPUTATIONS FOR $n = 1$. — Substituting the genus expansions

$$(66) \quad \frac{e^{u(\mathcal{S}(u \hbar z \partial_z) - 1)y(z)}}{u \mathcal{S}(u \hbar)} = \frac{1}{u} + \frac{1}{24} (u^2 y^{[2]} - u) \hbar^2 + O(\hbar^4)$$

to Equation (65) in the case $n = 1$, $g \geq 1$, we obtain

$$\begin{aligned} H_{g,1} &= [\hbar^{2g}] \sum_{j=0}^{\infty} D^j \frac{1}{Q} [v^j] \left(\frac{1}{24} \left(\frac{L_2(v, y, \hbar)}{v} y^{[2]} - \frac{L_1(v, y, \hbar)}{v} \right) \hbar^2 + \frac{L_0(v, y, \hbar)}{v^2} + O(\hbar^4) \right) \\ &\quad + [u^{2g}] \frac{1}{\mathcal{S}(u)} \psi^{(2g-1)}(y(z)) - [u^{2g}] \frac{1}{\mathcal{S}(u)^2} \psi^{(2g-1)}(0). \end{aligned}$$

Then, using explicit expressions for the series L_r ,

$$(67) \quad \begin{aligned} L_0(v, y, \hbar) &= 1 + (v^3 - v) \frac{\psi''(y)}{24} \hbar^2 + O(\hbar^4), \\ L_1(v, y, \hbar) &= v \psi'(y) + O(\hbar^2), \\ L_2(v, y, \hbar) &= v \psi''(y) + v^2 \psi'(y)^2 + O(\hbar^2), \\ L_3(v, y, \hbar) &= v^3 \psi'(y)^3 + 3v^2 \psi'(y) \psi''(y) + v \psi^{(3)}(y) + O(\hbar^2), \end{aligned}$$

we obtain, in the case $g = 1$,

$$H_{1,1} = D \frac{(\psi')^2 y^{[2]} + \psi'' y^{[1]}}{24Q} + \frac{\psi'' y^{[2]} - \psi'}{24Q} - \frac{\psi'}{24} + \frac{\psi'(0)}{12}.$$

Similar computations in the case $(g, n) = (2, 1)$ give

$$\begin{aligned} H_{2,1}(z) &= D^4 \frac{10\psi'' (\psi')^2 y^{[2]} + 5(\psi'')^2 y^{[1]} + 5(\psi')^5 (y^{[2]})^2}{5760Q} + \dots \\ &+ \frac{5\psi^{(5)} (y^{[2]})^2 - 20\psi^{(4)} y^{[2]} + 3\psi^{(4)} y^{[4]} + 17\psi^{(3)} + 5(\psi'')^2 y^{[1]}}{5760Q} + \frac{7\psi^{(3)}}{5760} - \frac{\psi^{(3)}(0)}{240}, \end{aligned}$$

where the dots denote the terms containing D^j with $j = 1, 2, 3$.

7.2. COMPUTATIONS FOR $n = 2$. — If $n \geq 1$, then equation of Theorem 5.3 can be applied. It is convenient to represent the transformation $D^{-1}U$ of Theorem 5.3 acting on a function $f(u, z)$ in u and z as follows

$$D^{-1}Uf = \frac{1}{\hbar} \sum_{r=0}^{\infty} M_r([u^r]f),$$

where M_r is the differential operator acting on a function $f(z)$ in z by

$$M_r f = \sum_{k,j=0}^{\infty} D^{j-1} \left(\frac{[v^j]L_k(v, y(z), \hbar)}{Q} [u^k] u^r \frac{e^{u(\mathcal{S}(u\hbar z\partial_z) - 1)y(z)}}{u\mathcal{S}(u\hbar)} f \right).$$

From (66) and (67) we find, explicitly,

$$\begin{aligned} M_1 f &= D^{-1} \frac{f}{Q} + \left(\frac{(\psi^{(3)} y^{[2]} - 2\psi'') f}{24Q} + D \frac{\psi' (3\psi'' y^{[2]} - \psi') f}{24Q} \right. \\ &\quad \left. + D^2 \frac{((\psi')^3 y^{[2]} + \psi'') f}{24Q} \right) \hbar^2 + O(\hbar^4), \\ M_2 f &= \frac{\psi' f}{Q} + O(\hbar^2), \\ M_3 f &= \frac{\psi'' f}{Q} + D \frac{(\psi')^2 f}{Q} + O(\hbar^2). \end{aligned}$$

We denote by $M_{k,i}$ the transformation M_k applied to the functions in u_i and z_i instead of u and z , respectively. With this notation, the statement of Theorem 5.3

can be written as follows

$$H_{g,n} = [\hbar^{2g}] \sum_{r_1, \dots, r_n=1}^{\infty} M_{r_1,1} \cdots M_{r_n,n} [u_1^{r_1} \cdots u_n^{r_n}] \sum_{\gamma \in \Gamma_n} \hbar^{2(|E_\gamma|-n+1)} \prod_{\{v_i, v_j\} \in E_\gamma} \bar{w}_{i,j} + (-1)^n [u^{2g}] \frac{1}{\mathcal{S}^2(u)} \psi^{(2g+n-2)}(0),$$

where

$$\begin{aligned} \bar{w}_{i,j} &= \frac{w_{i,j}}{\hbar^2} = \frac{e^{\hbar^2 u_i u_j \mathcal{S}(u_i \hbar z_i \partial_{z_i}) \mathcal{S}(u_j \hbar z_j \partial_{z_j}) \gamma_{i,j}} - 1}{\hbar^2} \\ &= u_i u_j \gamma_{i,j} + \left(\frac{u_i^3 u_j + u_i u_j^3}{24} \gamma_{i,j}^{[2]} + \frac{1}{2} u_i^2 u_j^2 (\gamma_{i,j})^2 \right) \hbar^2 + O(\hbar^2). \end{aligned}$$

If $n = 2$, then the sum over graphs is reduced to just $\bar{w}_{1,2}$ and we get

$$\begin{aligned} H_{g,2} &= [\hbar^{2g}] \sum_{r_1, r_2=1}^{\infty} M_{r_1,1} M_{r_2,2} [u_1^{r_1} u_2^{r_2}] \bar{w}_{1,2} + [u^{2g}] \frac{1}{\mathcal{S}^2(u)} \psi^{(2g)}(0) \\ &= [\hbar^{2g}] \left(M_{1,1} M_{1,2} \gamma_{1,2} + \left(M_{3,1} M_{1,2} \frac{\gamma_{1,2}^{[2]}}{24} + M_{1,1} M_{3,2} \frac{\gamma_{1,2}^{[2]}}{24} + M_{2,1} M_{2,2} \frac{(\gamma_{1,2})^2}{2} \right) \hbar^2 \right. \\ &\quad \left. + O(\hbar^4) \right) + [u^{2g}] \frac{1}{\mathcal{S}^2(u)} \psi^{(2g)}(0). \end{aligned}$$

In particular, for $(g, n) = (1, 2)$ we have

$$\begin{aligned} H_{1,2} &= D_1^2 \frac{\gamma_{2,1}^{[-1]} (\psi_1'' + (\psi_1')^3 y_1^{[2]})}{24Q_1} + D_1 \frac{\psi_1' (\psi_1' (\gamma_{2,1}^{[1]} - \gamma_{2,1}^{[-1]}) + 3\gamma_{2,1}^{[-1]} \psi_1'' y_1^{[2]})}{24Q_1} \\ &\quad + \frac{\psi_1'' (\gamma_{2,1}^{[1]} - 2\gamma_{2,1}^{[-1]}) + \psi_1^{(3)} \gamma_{2,1}^{[-1]} y_1^{[2]}}{24Q_1} + D_2^2 \frac{\gamma_{1,2}^{[-1]} (\psi_2'' + (\psi_2')^3 y_2^{[2]})}{24Q_2} \\ &\quad + D_2 \frac{\psi_2' (\psi_2' (\gamma_{1,2}^{[1]} - \gamma_{1,2}^{[-1]}) + 3\gamma_{1,2}^{[-1]} \psi_2'' y_2^{[2]})}{24Q_2} + \frac{\psi_2'' (\gamma_{1,2}^{[1]} - 2\gamma_{1,2}^{[-1]}) + \psi_2^{(3)} \gamma_{1,2}^{[-1]} y_2^{[2]}}{24Q_2} \\ &\quad + \frac{(\gamma_{1,2})^2 \psi_1' \psi_2'}{2Q_1 Q_2} - \frac{1}{12} \psi''(0). \end{aligned}$$

7.3. COMPUTATIONS FOR $n = 3$. — There are four possible connected simple graphs on three labeled vertices, and summing up the contributions of these four graphs we get

$$\begin{aligned} H_{g,3} &= [\hbar^{2g}] \sum_{r_1, r_2, r_3=1}^{\infty} M_{r_1,1} M_{r_2,2} M_{r_3,3} [u_1^{r_1} u_2^{r_2} u_3^{r_3}] (\bar{w}_{1,2} \bar{w}_{1,3} + \bar{w}_{1,2} \bar{w}_{3,3} + \bar{w}_{1,3} \bar{w}_{2,3} \\ &\quad + \hbar^2 \bar{w}_{1,2} \bar{w}_{1,3} \bar{w}_{2,3}) - [u^{2g}] \frac{1}{\mathcal{S}^2(u)} \psi^{(2g+1)}(0). \end{aligned}$$



For instance, for $g = 0$ using $\bar{w}_{i,j} = u_i u_j \gamma_{i,j} + O(\hbar^2)$ we get

$$\begin{aligned} H_{0,3} &= (M_{2,1} M_{1,2} M_{1,3} \gamma_{1,2} \gamma_{1,3} + M_{1,1} M_{2,2} M_{1,3} \gamma_{1,2} \gamma_{2,3} + M_{1,1} M_{1,2} M_{2,3} \gamma_{1,3} \gamma_{2,2}) \Big|_{\hbar=0} - \psi'(0). \end{aligned}$$

This gives the final answer

$$(68) \quad \begin{aligned} H_{0,3} &= \frac{\psi'_1}{Q_1} \gamma_{2,1}^{[-1]} \gamma_{3,1}^{[-1]} + \frac{\psi'_2}{Q_2} \gamma_{1,2}^{[-1]} \gamma_{3,2}^{[-1]} + \frac{\psi'_3}{Q_3} \gamma_{1,3}^{[-1]} \gamma_{2,3}^{[-1]} - \psi'(0) \\ &= \sum_{i=1}^3 \frac{\psi'(y_i)}{Q(z_i)} \prod_{j \neq i} \frac{z_j}{z_i - z_j} - \psi'(0). \end{aligned}$$

REMARK 7.1. — Note that Equation (68) differs from [ACEH20, Prop.10.2 & Eq.(10.4)] produced by means of the spectral curve topological recursion. The formula given in Equation (10.4) in *op. cit.* does not appear to be vanishing on the coordinate axes and seems to have an incorrect overall sign, which are typical bugs that often occur in applications of topological recursion.

7.4. COMPUTATION FOR $(g, n) = (0, 4)$. — In the case $g = 0$ the graphs that contribute to $H_{0,n}$ are trees. For $n = 4$ there are 4 trees on 4 labeled vertices isomorphic to  and 12 more trees isomorphic to . They contribute to the corresponding summands in $H_{0,4}$:

$$\begin{aligned} H_{0,4} &= [\hbar^0] \sum_{r_1, \dots, r_4 \geq 1} \left(\prod_{k=1}^4 M_{r_k, k} \right) \\ &\quad \left[\prod_{k=1}^4 u_k^{r_k} \right] \left((u_1 u_2 \gamma_{1,2} u_1 u_3 \gamma_{1,3} u_1 u_4 \gamma_{1,4} + \dots (4 \text{ terms in total})) \right. \\ &\quad \left. + (u_1 u_2 \gamma_{1,2} u_2 u_3 \gamma_{2,3} u_3 u_4 \gamma_{3,4} + \dots (12 \text{ terms in total})) \right) + \psi''(0) \\ &= \left((M_{3,1} M_{1,2} M_{1,3} M_{1,4} (\gamma_{1,2} \gamma_{1,3} \gamma_{1,4}) + \dots (4 \text{ terms in total})) \right. \\ &\quad \left. + (M_{1,1} M_{2,2} M_{2,3} M_{1,4} (\gamma_{1,2} \gamma_{2,3} \gamma_{3,4}) + \dots (12 \text{ terms in total})) \right) \Big|_{\hbar=0} + \psi''(0), \end{aligned}$$

and we get the final answer

$$\begin{aligned} H_{0,4} &= \left(D_1 \frac{(\psi'_1)^2 \gamma_{2,1}^{[-1]} \gamma_{3,1}^{[-1]} \gamma_{4,1}^{[-1]}}{Q_1} + \frac{\psi''_1 \gamma_{2,1}^{[-1]} \gamma_{3,1}^{[-1]} \gamma_{4,1}^{[-1]}}{Q_1} + \dots (2 \times 4 \text{ terms in total}) \right) \\ &\quad + \left(\frac{\psi'_2 \psi'_3 \gamma_{1,2}^{[-1]} \gamma_{2,3}^{[-1]} \gamma_{4,3}^{[-1]}}{Q_2 Q_3} + \dots (12 \text{ terms in total}) \right) + \psi''(0). \end{aligned}$$

REFERENCES

- [ACEH18a] A. ALEXANDROV, G. CHAPUY, B. EYNARD & J. HARNAD — “Fermionic approach to weighted Hurwitz numbers and topological recursion”, *Comm. Math. Phys.* **360** (2018), no. 2, p. 777–826.
- [ACEH18b] ———, “Weighted Hurwitz numbers and topological recursion: an overview”, *J. Math. Phys.* **59** (2018), no. 8, article no. 081102 (21 pages).
- [ACEH20] ———, “Weighted Hurwitz numbers and topological recursion”, *Comm. Math. Phys.* **375** (2020), no. 1, p. 237–305.
- [ALS16] A. ALEXANDROV, D. LEWANSKI & S. SHADRIN — “Ramifications of Hurwitz theory, KP integrability and quantum curves”, *J. High Energy Phys.* (2016), no. 5, article no. 124 (30 pages).

- [BDK⁺20] G. BOROT, N. DO, M. KAREV, D. LEWANSKI & E. MOSKOVSKY – “Double Hurwitz numbers: polynomiality, topological recursion and intersection theory”, 2020, [arXiv:2002.00900](https://arxiv.org/abs/2002.00900).
- [BGF20] G. BOROT & E. GARCIA-FAILDE – “Simple maps, Hurwitz numbers, and topological recursion”, *Comm. Math. Phys.* **380** (2020), no. 2, p. 581–654.
- [BS17] G. BOROT & S. SHADRIN – “Blobbed topological recursion: properties and applications”, *Math. Proc. Cambridge Philos. Soc.* **162** (2017), no. 1, p. 39–87.
- [BDBKS20] B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN & S. SHADRIN – “Topological recursion for Kadomtsev–Petviashvili tau functions of hypergeometric type”, 2020, [arXiv:2012.14723](https://arxiv.org/abs/2012.14723).
- [BDBKS21] ———, “Generalised ordinary vs fully simple duality for n -point functions and a proof of the Borot–Garcia-Failde conjecture”, 2021, [arXiv:2106.08368](https://arxiv.org/abs/2106.08368).
- [BDBS20] B. BYCHKOV, P. DUNIN-BARKOWSKI & S. SHADRIN – “Combinatorics of Bousquet–Mélou–Schaeffer numbers in the light of topological recursion”, *European J. Combin.* **90** (2020), article no. 103184 (35 pages).
- [Cha09] G. CHAPUY – “Asymptotic enumeration of constellations and related families of maps on orientable surfaces”, *Combin. Probab. Comput.* **18** (2009), no. 4, p. 477–516.
- [DBKO⁺15] P. DUNIN-BARKOWSKI, M. KAZARIAN, N. ORANTIN, S. SHADRIN & L. SPITZ – “Polynomiality of Hurwitz numbers, Bouchard–Mariño conjecture, and a new proof of the ELSV formula”, *Adv. Math.* **279** (2015), p. 67–103.
- [DBLPS15] P. DUNIN-BARKOWSKI, D. LEWANSKI, A. POPOLITOV & S. SHADRIN – “Polynomiality of orbifold Hurwitz numbers, spectral curve, and a new proof of the Johnson–Pandharipande–Tseng formula”, *J. London Math. Soc. (2)* **92** (2015), no. 3, p. 547–565.
- [DBKP⁺20] P. DUNIN-BARKOWSKI, M. KAZARIAN, A. POPOLITOV, S. SHADRIN & A. SLEPTSOV – “Topological recursion for the extended Ooguri–Vafa partition function of colored HOMFLY–PT polynomials of torus knots”, 2020, [arXiv:2010.11021](https://arxiv.org/abs/2010.11021).
- [DBPSS19] P. DUNIN-BARKOWSKI, A. POPOLITOV, S. SHADRIN & A. SLEPTSOV – “Combinatorial structure of colored HOMFLY–PT polynomials for torus knots”, *Commun. Number Theory Phys.* **13** (2019), no. 4, p. 763–826.
- [Har16] J. HARNAD – “Weighted Hurwitz numbers and hypergeometric τ -functions: an overview”, in *String-Math 2014*, Proc. Sympos. Pure Math., vol. 93, American Mathematical Society, Providence, RI, 2016, p. 289–333.
- [Kaz19] M. KAZARIAN – “Topological recursion for generalized Hurwitz numbers”, Talk at Skoltech Center for Advanced Studies, <https://youtu.be/Hq7RVZBA8f8>, 2019.
- [Kaz20] ———, “Quasirationality of weighted Hurwitz numbers”, Talk at the National Research University Higher School of Economics, <https://youtu.be/Yu1dGgBidr0>, 2020.
- [Kaz21] ———, Paper in preparation, Preprint, 2021.
- [KZ15] M. KAZARIAN & P. ZOGRAF – “Virasoro constraints and topological recursion for Grothendieck’s dessin counting”, *Lett. Math. Phys.* **105** (2015), no. 8, p. 1057–1084.
- [KL15] M. E. KAZARYAN & S. K. LANDO – “Combinatorial solutions to integrable hierarchies”, *Uspekhi Mat. Nauk* **70** (2015), no. 3(423), p. 77–106.
- [KMMM95] S. KHARCHEV, A. MARSHAKOV, A. MIRONOV & A. MOROZOV – “Generalized Kazakov–Migdal–Kontsevich model: group theory aspects”, *Internat. J. Modern Phys. A* **10** (1995), no. 14, p. 2015–2051.
- [KLPS19] R. KRAMER, D. LEWANSKI, A. POPOLITOV & S. SHADRIN – “Towards an orbifold generalization of Zvonkine’s r -ELSV formula”, *Trans. Amer. Math. Soc.* **372** (2019), no. 6, p. 4447–4469.
- [KLS19] R. KRAMER, D. LEWANSKI & S. SHADRIN – “Quasi-polynomiality of monotone orbifold Hurwitz numbers and Grothendieck’s dessins d’enfants”, *Doc. Math.* **24** (2019), p. 857–898.
- [MJD00] T. MIWA, M. JIMBO & E. DATE – *Solitons*, Cambridge Tracts in Math., vol. 135, Cambridge University Press, Cambridge, 2000.
- [OS01a] A. Y. ORLOV & D. M. SCHERBIN – “Hypergeometric solutions of soliton equations”, *Teoret. Mat. Fiz.* **128** (2001), no. 1, p. 84–108.
- [OS01b] ———, “Multivariate hypergeometric functions as τ -functions of Toda lattice and Kadomtsev–Petviashvili equation”, *Phys. D* **152–153** (2001), p. 51–65.

- [SSZ12] S. SHADRIN, L. SPITZ & D. ZVONKINE – “On double Hurwitz numbers with completed cycles”, *J. London Math. Soc. (2)* **86** (2012), no. 2, p. 407–432.
- [Sta99] R. P. STANLEY – *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Math., vol. 62, Cambridge University Press, Cambridge, 1999.

Manuscript received 20th September 2021
accepted 4th July 2022

BORIS BYCHKOV, Faculty of Mathematics, National Research University Higher School of Economics
Usacheva 6, 119048 Moscow, Russia

and

Center of Integrable Systems, P.G. Demidov Yaroslavl State University
Sovetskaya 14, 150003, Yaroslavl, Russia

and current affiliation:

Department of Mathematics, University of Haifa
Mount Carmel, 3488838, Haifa, Israel

E-mail : bbychkov@hse.ru

PETR DUNIN-BARKOWSKI, Faculty of Mathematics, National Research University Higher School of Economics
Economics

Usacheva 6, 119048 Moscow, Russia

and

HSE–Skoltech International Laboratory of Representation Theory and Mathematical Physics,
Skoltech

Nobelya 1, 143026, Moscow, Russia

and

NRC “Kurchatov Institute” – ITEP
117218 Moscow, Russia

E-mail : ptdunin@hse.ru

MAXIM KAZARIAN, Faculty of Mathematics, National Research University Higher School of Economics
Usacheva 6, 119048 Moscow, Russia

and

Center for Advanced Studies, Skoltech

Nobelya 1, 143026, Moscow, Russia

E-mail : kazarian@mccme.ru

SERGEY SHADRIN, Korteweg-de Vries Institute for Mathematics, University of Amsterdam
Postbus 94248, 1090 GE Amsterdam, The Netherlands

E-mail : S.Shadrin@uva.nl