

JOURNAL

de Théorie des Nombres
de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Peter BRUIN

The Tate pairing for Abelian varieties over finite fields

Tome 23, n° 2 (2011), p. 323-328.

http://jtnb.cedram.org/item?id=JTNB_2011__23_2_323_0

© Société Arithmétique de Bordeaux, 2011, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

The Tate pairing for Abelian varieties over finite fields

par PETER BRUIN

RÉSUMÉ. Nous décrivons un accouplement arithmétique associé à une isogenie entre variétés abéliennes sur un corps fini. Nous montrons qu'il généralise l'accouplement de Frey et Rück, donnant ainsi une démonstration brève de la perfection de ce dernier.

ABSTRACT. In this expository note, we describe an arithmetic pairing associated to an isogeny between Abelian varieties over a finite field. We show that it generalises the Frey–Rück pairing, thereby giving a short proof of the perfectness of the latter.

1. Introduction

Throughout this note, k denotes a finite field of q elements, and \bar{k} denotes an algebraic closure of k . If n is a positive integer, then μ_n denotes the group of n -th roots of unity in \bar{k}^\times .

Let C be a complete, smooth, geometrically connected curve over k , let J be the Jacobian variety of C , and let n be a divisor of $q - 1$. In [1], Frey and Rück defined a perfect pairing

$$\{ \cdot, \cdot \}_n : J[n](k) \times J(k)/nJ(k) \longrightarrow \mu_n(k)$$

as follows: if D and E are divisors on C with disjoint supports and f is a non-zero rational function with divisor nD , then

$$\{[D], [E] \bmod nJ(k)\}_n = f(E)^{(q-1)/n},$$

where

$$f(E) = \prod_{x \in C(\bar{k})} f(x)^{n_x} \quad \text{if } E = \sum_{x \in C(\bar{k})} n_x x.$$

Remark. We have composed the map as defined in [1] with the isomorphism $k^\times / (k^\times)^n \rightarrow \mu_n(k)$ that raises elements to the power $(q - 1)/n$.

Manuscrit reçu le 25 mai 2010.

I am grateful to Hendrik Lenstra and Ed Schaefer for discussions on this subject, and to Andreas Enge for the suggestion to publish this note. I thank the two referees for useful remarks.

To prove the perfectness of $\{ \ , \}_n$, Frey and Rück used a pairing introduced by Tate [10], which relates certain cohomology groups of an Abelian variety over a p -adic field, and an alternative description of this pairing given by Lichtenbaum [6] in the case of the Jacobian of a curve over a p -adic field. The names ‘Tate pairing’ and ‘Tate–Lichtenbaum pairing’ are therefore often used for what we call the Frey–Rück pairing.

There are now several proofs of the non-degeneracy of the Frey–Rück pairing that no longer use the cited results of Tate and Lichtenbaum; see Heß [2] and Schaefer [8]. In this note, we consider a separable isogeny $\phi: A \rightarrow B$ between Abelian varieties over k such that $\ker \phi$ is annihilated by $q - 1$. We define a pairing

$$[\ , \]_\phi: \ker \hat{\phi}(k) \times \operatorname{coker}(\phi(k)) \longrightarrow k^\times$$

between the groups

$$\ker \hat{\phi}(k) = \{b \in B(k) \mid \hat{\phi}(b) = 0\}$$

and

$$\operatorname{coker}(\phi(k)) = B(k)/\phi(A(k)).$$

We show that this pairing is perfect in the sense that it induces isomorphisms

$$\ker \hat{\phi}(k) \xrightarrow{\sim} \operatorname{Hom}(\operatorname{coker}(\phi(k)), k^\times)$$

and

$$\operatorname{coker}(\phi(k)) \xrightarrow{\sim} \operatorname{Hom}(\ker \hat{\phi}(k), k^\times).$$

Furthermore, we show that the Frey–Rück pairing is the special case of multiplication by n on the Jacobian of a curve over k . The more general pairing appears to be known, but has not to my knowledge appeared in the literature. In any case, I do not make any claim to originality. It seems appropriate to call $[\ , \]_\phi$ the *Tate pairing associated to ϕ* .

Remark. The condition that $\ker \phi$ be annihilated by $q - 1$ can be relaxed somewhat. Let m be the exponent of $\ker \phi$ and let d be the order of q in $(\mathbf{Z}/m\mathbf{Z})^\times$. If m and d are coprime, then one can pass to an extension of degree d of k and consider appropriate eigenspaces for the Frobenius action. This reproduces a result of Frey and Rück [1, Proposition 2.5].

2. Preliminaries

By a *pairing* we mean a bilinear map

$$A \times B \rightarrow C$$

between Abelian groups. It is called *perfect* if the induced group homomorphisms

$$A \rightarrow \operatorname{Hom}(B, C) \quad \text{and} \quad B \rightarrow \operatorname{Hom}(A, C)$$

are isomorphisms.

The Galois group $G_k = \text{Gal}(\bar{k}/k)$ is canonically isomorphic to the profinite group $\widehat{\mathbf{Z}}$, with the element 1 of $\widehat{\mathbf{Z}}$ corresponding to the q -power Frobenius automorphism σ . The only actions of G_k that we consider are continuous actions on Abelian groups with the discrete topology. An Abelian group equipped with such an action of G_k is called a G_k -module.

The fact that G_k is generated as a topological group by σ implies that for any G_k -module M , the Galois cohomology group $H^1(G_k, M)$ has an easy description via the isomorphism

$$(2.1) \quad \begin{aligned} H^1(G_k, M) &\xrightarrow{\sim} M/(\sigma - 1)M \\ [c] &\longmapsto c(\sigma) \bmod (\sigma - 1)M \end{aligned}$$

of Abelian groups, where $[c]$ denotes the class of a cocycle $c: G_k \rightarrow M$; see Serre [9, XIII, §1, proposition 1].

If F is a finite G_k -module, the Cartier dual of F is the Abelian group

$$F^\vee = \text{Hom}(F, \bar{k}^\times)$$

with the G_k -action given by

$$(\sigma a)(x) = \sigma(a(\sigma^{-1}x)) \quad \text{for all } a \in F^\vee, \sigma \in G_k, x \in F.$$

The subgroup $F^\vee(k)$ of G_k -invariants consists of the elements of $\text{Hom}(F, \bar{k}^\times)$ that are homomorphisms of G_k -modules.

Lemma 2.1. *Let F be a finite G_k -module annihilated by $q - 1$, and let F^\vee be its Cartier dual. There is a perfect pairing*

$$F^\vee(k) \times H^1(G_k, F) \longrightarrow k^\times$$

sending $(a, [c])$ to $a(c(\sigma))$ for every $a \in F^\vee(k)$ and every cocycle $c: G_k \rightarrow F$.

Proof. By assumption, we can write

$$\begin{aligned} F^\vee(k) &= \text{Hom}(F, \bar{k}^\times)^{G_k} \\ &= \text{Hom}(F, k^\times)^{G_k} \\ &= \text{Hom}(F/(\sigma - 1)F, k^\times). \end{aligned}$$

Applying the isomorphism (2.1) with $M = F$, we get an isomorphism

$$F^\vee(k) \xrightarrow{\sim} \text{Hom}(H^1(G_k, F), k^\times)$$

such that the induced bilinear map $F^\vee(k) \times H^1(G_k, F) \longrightarrow k^\times$ is given by the formula in the statement of the lemma. Since $H^1(G_k, F)$ is annihilated by the order of k^\times , it is (non-canonically) isomorphic to $F^\vee(k)$. This implies that the map

$$H^1(G_k, F) \longrightarrow \text{Hom}(F^\vee(k), k^\times)$$

is also an isomorphism, so the pairing is perfect. □

3. Definition of the Tate pairing

Let $\phi: A \rightarrow B$ be a separable isogeny between Abelian varieties over k such that $\ker \phi$ is annihilated by $q-1$. We write $\phi(k)$ for the homomorphism $A(k) \rightarrow B(k)$ induced by ϕ . We note that

$$(\ker \phi)(k) = \ker(\phi(k)),$$

and we denote this group from now on by $\ker \phi(k)$. We also note that there is no similar equality for cokernels.

We start with the short exact sequence

$$0 \rightarrow \ker \phi \rightarrow A \rightarrow B \rightarrow 0$$

of commutative group varieties over k . Since A, B and ϕ are defined over k , taking \bar{k} -points in this sequence gives a short exact sequence of G_k -modules. Taking Galois cohomology, we obtain a long exact sequence

$$0 \rightarrow \ker \phi(k) \rightarrow A(k) \xrightarrow{\phi(k)} B(k) \xrightarrow{\delta} H^1(G_k, \ker \phi(\bar{k})) \rightarrow H^1(G_k, A(\bar{k})).$$

The connecting homomorphism δ is given by

$$\delta(b) = [\tau \mapsto \tau(a) - a],$$

where a is any element of $A(\bar{k})$ such that $\phi(a) = b$. A theorem of Lang on Abelian varieties over finite fields [4, Theorem 3] implies that $H^1(G_k, A(\bar{k}))$ vanishes. This means that δ is surjective and hence gives an isomorphism

$$B(k)/\phi(A(k)) \xrightarrow{\sim} H^1(G_k, \ker \phi(\bar{k})).$$

The pairing from Lemma 2.1 becomes a perfect pairing

$$(3.1) \quad (\ker \phi)^\vee(k) \times B(k)/\phi(A(k)) \rightarrow k^\times$$

sending $(f, b \bmod \phi(A(k)))$ to $f(\sigma(a) - a)$, with $a \in A(\bar{k})$ such that $\phi(a) = b$.

Now let $\hat{\phi}: \hat{B} \rightarrow \hat{A}$ denote the isogeny dual to ϕ . The Cartier duality theorem for isogenies of Abelian varieties (see for example Mumford [7, § 15, Theorem 1]) gives a canonical isomorphism

$$\epsilon_\phi: \ker \hat{\phi} \xrightarrow{\sim} (\ker \phi)^\vee$$

of G_k -modules. Combining this with the pairing (3.1) gives a perfect pairing

$$(3.2) \quad [\ , \]_\phi: \ker \hat{\phi}(k) \times \text{coker}(\phi(k)) \rightarrow k^\times$$

$$(x, y) \mapsto (\epsilon_\phi x)(\sigma a - a),$$

where a is any element of $A(\bar{k})$ with $(\phi(a) \bmod \phi(A(k))) = y$. We call this pairing the *Tate pairing* associated to ϕ .

4. Comparison to the Frey–Rück pairing

We now take $A = B = J$, where J is the Jacobian of a complete, smooth, geometrically connected curve C over k , and we take ϕ to be the multiplication by a positive integer n dividing $q - 1$. We identify J with its dual Abelian variety using the canonical principal polarisation. Then the Tate pairing (3.2) becomes a perfect pairing

$$[\ , \]_n : J[n](k) \times J(k)/nJ(k) \longrightarrow k^\times.$$

Moreover, from the Cartier duality isomorphism $\epsilon_n : \hat{J}[n] \xrightarrow{\sim} J[n]^\vee$ and the identification of $J[n]$ with $\hat{J}[n]$ we obtain a perfect pairing

$$\begin{aligned} e_n : J[n] \times J[n] &\longrightarrow \mu_n \\ (x, y) &\longmapsto (\epsilon_n y)(x) \quad \text{for all } x, y \in J[n](\bar{k}), \end{aligned}$$

called the *Weil pairing*. It can be computed as follows. Let D and E be two divisors of degree 0 on $C_{\bar{k}}$ with disjoint supports and such that their classes $[D]$ and $[E]$ in $J(\bar{k})$ are n -torsion points. Then there exist non-zero rational functions f and g on $C_{\bar{k}}$ such that

$$nD = \operatorname{div} f \quad \text{and} \quad nE = \operatorname{div} g,$$

and we have

$$(4.1) \quad e_n([D], [E]) = \frac{g(D)}{f(E)};$$

see Howe [3]. The fact that this depends only on the linear equivalence classes of D and E comes down to *Weil reciprocity*: if f and g are two rational functions on $C_{\bar{k}}$ whose divisors have disjoint supports, then

$$f(\operatorname{div} g) = g(\operatorname{div} f).$$

For a proof, we refer to Lang [5, Chapter VI, corollary to Theorem 10].

Theorem 4.1. *The pairing $[\ , \]_n$ equals the Frey–Rück pairing $\{ \ , \ \}_n$.*

Proof. Consider elements of $J[n](k)$ and $J(k)/nJ(k)$, represented by (k -rational) divisors D and E , respectively, such that the supports of D and E are disjoint. We choose a divisor E_0 on $C_{\bar{k}}$ such that nE_0 is linearly equivalent to E and rational functions f on C and g on $C_{\bar{k}}$ such that

$$nD = \operatorname{div} f \quad \text{and} \quad nE_0 - E = \operatorname{div} g.$$

Then we have

$$\operatorname{div}(\sigma g/g) = n(\sigma E_0 - E_0).$$

Using the definitions of the pairing $[\ , \]_n$ and of the Weil pairing e_n , the formula (4.1), the fact that σ is the q -power map on \bar{k}^\times , and Weil reciprocity, we can compute $[\ , \]_n$ as follows:

$$\begin{aligned}
 [[D], [E] \bmod nJ(k)]_n &= (\epsilon_n[D])([\sigma E_0 - E_0]) \\
 &= e_n([\sigma E_0 - E_0], [D]) \\
 &= \frac{f(\sigma E_0 - E_0)}{(\sigma g/g)(D)} \\
 &= \frac{\sigma(f(E_0))/f(E_0)}{\sigma g(D)/g(D)} \\
 &= (f(E_0)/g(D))^{q-1} \\
 &= (f(nE_0)/g(nD))^{(q-1)/n} \\
 &= (f(nE_0)/f(nE_0 - E))^{(q-1)/n} \\
 &= f(E)^{(q-1)/n}.
 \end{aligned}$$

This equality is what we had to prove. □

References

- [1] G. FREY and H.-G. RÜCK, *A remark concerning m -divisibility and the discrete logarithm in class groups of curves*. Mathematics of Computation **62** (1994), 865–874.
- [2] F. HESS, *A note on the Tate pairing of curves over finite fields*. Archiv der Mathematik **82** (2004), no. 1, 28–32.
- [3] E. W. HOWE, *The Weil pairing and the Hilbert symbol*. Mathematische Annalen **305** (1996), 387–392.
- [4] S. LANG, *Abelian varieties over finite fields*. Proceedings of the National Academy of Sciences of the U.S.A. **41** (1955), no. 3, 174–176.
- [5] S. LANG, *Abelian Varieties*. Interscience, New York, 1959.
- [6] S. LICHTENBAUM, *Duality theorems for curves over p -adic fields*. Inventiones Mathematicae **7** (1969), 120–136.
- [7] D. MUMFORD, *Abelian Varieties*. Tata Institute of Fundamental Research, Bombay, 1970.
- [8] E. F. SCHAEFER, *A new proof for the non-degeneracy of the Frey–Rück pairing and a connection to isogenies over the base field*. In: T. SHASKA (editor), *Computational Aspects of Algebraic Curves* (Conference held at the University of Idaho, 2005), 1–12. Lecture Notes Series in Computing **13**. World Scientific Publishing, Hackensack, NJ, 2005.
- [9] J-P. SERRE, *Corps locaux*. Hermann, Paris, 1962.
- [10] J. TATE, *WC-groups over p -adic fields*. Séminaire Bourbaki, exposé 156. Secrétariat mathématique, Paris, 1957.

Peter BRUIN
 Université Paris-Sud 11
 Département de Mathématiques
 Bâtiment 425
 91405 Orsay cedex
 France
E-mail: Peter.Bruin@math.u-psud.fr