

JOZEF SICIAK

Families of polynomials and determining measures

Annales de la faculté des sciences de Toulouse 5^e série, tome 9, n° 2
(1988), p. 193-211

[<http://www.numdam.org/item?id=AFST_1988_5_9_2_193_0>](http://www.numdam.org/item?id=AFST_1988_5_9_2_193_0)

© Université Paul Sabatier, 1988, tous droits réservés.

L'accès aux archives de la revue « Annales de la faculté des sciences de Toulouse » (<http://picard.ups-tlse.fr/~annales/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Families of polynomials and determining measures

JOZEF SICIĄK⁽¹⁾

RÉSUMÉ. — Soit μ une mesure de probabilité sur une partie borélienne bornée non pluripolaire E de \mathbb{C}^N , on étudie l'allure de croissance des familles de polynômes ponctuellement bornées μ -presque partout sur E . On définit une fonction $\mathcal{M}(t; E, \mu)$ ($0 \leq t \leq 1$) associée au couple (E, μ) . Sous des hypothèses naturelles sur E et μ , on montre que $\mathcal{M}(1; E, \mu) = 1$ si et seulement si le couple (E, μ) satisfait à la condition polynomiale (\mathcal{L}^*) , généralisant la condition polynomiale de Leja dans le cas plan, si et seulement si μ est une mesure déterminante pour E par rapport à la fonction L -extrémale L_E^* .

ABSTRACT. — Given a probability measure μ on a bounded nonpluripolar Borel subset E of \mathbb{C}^N , we study the growth behaviour of polynomial families which are pointwise bounded μ -a.e. on E . We define a function $\mathcal{M}(t, E, \mu)$ ($0 \leq t \leq 1$) associated to the pair (E, μ) . Under natural assumptions on E and μ we prove that $\mathcal{M}(1; E, \mu) = 1$ if and only if the pair (E, μ) satisfies the polynomial condition (\mathcal{L}^*) (a generalization of the Leja's condition in the plane), if and only if μ is determining for E with respect to the L -extremal function L_E^* .

0 - Introduction

Given a domain Ω in \mathbb{C}^N , we denote by $P(\Omega)$ the class of plurisubharmonic (plsh) functions on Ω . Let

$$\mathcal{L} := \{u \in P(\mathbb{C}^N); u(z) \leq \beta + \log(1 + |z|) \text{ in } \mathbb{C}^N\},$$

where β is a real constant depending on u . For a bounded set E in \mathbb{C}^N define

$$L_E(z) := \sup \{u(z); u \in \mathcal{L}, u \leq 0 \text{ on } E\}. \quad (0.1)$$

⁽¹⁾ Jagellonian University, Institute of Mathematics, ul. Reymonta 4, 30-059 Krakow - Poland

The uppersemicontinuous regularization $L_E^*(z) := \limsup_{w \rightarrow z} L_E(w)$ is called the L -extremal function of E . It is known that if E is a compact subset of \mathbb{C} with positive logarithmic capacity then L_E is identical with the Green function for $\mathbb{C} \setminus \hat{E}$ with pole at infinity.

For a bounded set E in \mathbb{C}^N either $L_E^* \equiv \infty$, in which case E is pluripolar (plp), or $L_E^* \in \mathcal{L}$.

DEFINITION 0.1. — We say that a point a in \mathbb{C}^N is an \mathcal{L} -regular point of $E \subset \mathbb{C}^N$, if $L_E^*(a) = 0$. A point $a \in \mathbb{C}^N$ such that $L_E(a) = 0$ and $L_E^*(a) > 0$ is called irregular point of E . It is clear that L_E^* is continuous at each regular point. By Bedford-Taylor theorem on negligible sets the set of irregular points of any subset E of \mathbb{C}^N is plp. If E is a compact set and $L_E = L_E^*$ on E (i.e. if L_E^* is continuous at each point of E) then L_E is continuous in \mathbb{C}^N and $L_E = L_E^*$. A compact set E with $L_E^* = L_E$ is called \mathcal{L} -regular. The set of \mathcal{L} -regular points of a compact \mathcal{L} -regular set E is identical with the polynomially convex hull \hat{E} of E .

DEFINITION 0.2. — A finite positive Borel measure μ on a bounded Borel set E in \mathbb{C}^N is called determining for E , if for every Borel subset F of E with $\mu(F) = \mu(E)$ one has $L_F^* = L_E^*$.

Observe that if $L_E^* = L_E$ and μ is determining for E , then for every $F \subset E$ with $\mu(F) = \mu(E)$ one has $L_F = L_E$ (because $L_F^* = L_E^* = L_E \leq L_F$).

It is known that $L_{E \cup A}^* = L_E^*$, if A is plp. Therefore $L_E^* = L_F^*$ for a subset F of E if and only if $L_F^* = 0$ quasi-almost everywhere (q.a.e.) on E . We say that a property \mathcal{P} holds q.a.e. on E , if it holds for each point of E except at most of a plp subset of E .

We say that a property \mathcal{P} holds quasi-star-almost-everywhere ($q^*.a.e.$) on E , if it holds for each point of a subset F of E with $L_F^* = L_E^*$.

It is clear that if μ is determining for E and \mathcal{P} holds $\mu.a.e.$ on E then it holds $q^*.a.e.$ on E .

DEFINITION 0.3. — Let μ be a finite positive Borel measure on a bounded Borel set in \mathbb{C}^N . We say that the pair (E, μ) satisfies (\mathcal{L}^*) -condition at a point a of \mathbb{C}^N , if for every family \mathcal{F} of polynomials of N -complex variables and for every number $b > 1$ the polynomial family

$$\mathcal{F}_b := \{b^{-\deg f} f; f \in \mathcal{F}\} \quad (0.2)$$

is uniformly bounded on a neighborhood \mathcal{U} of a .

We say that the pair (E, μ) satisfies (\mathcal{L}^*) -condition, if for every $b > 1$ and for every polynomial family \mathcal{F} bounded μ -a.e. on E the family \mathcal{F}_b is uniformly bounded on a neighborhood of E .

It is clear that if E is compact then (E, μ) satisfies (\mathcal{L}^*) -condition, if and only if it satisfies (\mathcal{L}^*) at each point of E .

All these notions are important for applications of the extremal function L_E^* . There are strict relations between them. Also are known important examples of pairs (E, μ) satisfying (\mathcal{L}^*) and of determining measures (e.g. [2], [5], [6], [7], [8], [14]).

In this paper we introduce a new function $\mathcal{M}(t) \equiv \mathcal{M}(t; E, \mu)$ associated to every pair (E, μ) by the formula

$$\log \mathcal{M}(t) := \sup \left\{ \sup_E L_A; A \subset E, \mu(A) \geq t\mu(E) \right\}, \quad 0 \leq t \leq 1.$$

It is clear that \mathcal{M} is a decreasing function and $1 \leq \mathcal{M}(t) \leq +\infty$. The function $\mathcal{M}^*(t) := \lim_{\tau \uparrow t} \mathcal{M}(\tau)$ ($0 < t \leq 1$), $\mathcal{M}^*(0) := \mathcal{M}(0)$, is decreasing and uppersemicontinuous on $[0, 1]$.

In the sequel we shall often assume (without loss of generality) that μ is a probability measure (i.e. $\mu(E) = 1$).

The function \mathcal{M} appears to be a useful notion strictly related to the determining measures and the (\mathcal{L}^*) -condition. For example we have obtained the following results involving the function \mathcal{M} .

THEOREM A. — *If $E \subset \mathbb{C}^N$ is compact and μ vanishes on plp subsets of \mathbb{C}^N then the following conditions are equivalent*

- (i) *The pair (E, μ) satisfies (\mathcal{L}^*) -condition;*
- (ii) *If $u \in \mathcal{L}$ and $u \leq 0$ μ -a.e. on E , then $u \leq 0$ on E ;*
- (iii) $\mathcal{M}^*(1) = 1$;
- (iv) $\mathcal{M}(1) = 1$;
- (v) μ *is determining for E and E is \mathcal{L} -regular.*

THEOREM B. — *Let $A \subset \mathbb{C}^P$, $B \subset \mathbb{C}^Q$ be two bounded Borel sets and μ, ν two probability measures on A and B , respectively. Put $\mathcal{M}_A(t) := \mathcal{M}(t; A, \mu)$, $\mathcal{M}_B(t) := \mathcal{M}(t; B, \nu)$ and $\mathcal{M}_{A \times B}(t) := \mathcal{M}(t, A \times B, \mu \otimes \nu)$. Then*

- (i) $\mathcal{M}_{A \times B}(1) \leq \mathcal{M}_A(1)\mathcal{M}_B(1)$

$$(ii) \mathcal{M}_{A \times B}^*(1) \leq \mathcal{M}_A^*(1) \mathcal{M}_B^*(1)$$

COROLLARY . — *If $A \subset \mathbb{C}^P$, $B \subset \mathbb{C}^P$ are compact sets and μ, ν vanish on plp sets, then if the pairs (A, μ) , (B, ν) satisfy one of the equivalent conditions of Theorem A then the pair $(A \times B, \mu \otimes \nu)$ satisfies each of the conditions.*

The equivalence of the conditions (i) and (v) was earlier obtained by LEVENBERG [6]. NGUYEN THANH VAN formulated the (\mathcal{L}^*) -condition in his paper [7]; his definition was inspired by LEJA's paper [5] containing as a special case so called "Polynomial Lemma", which in the present language reads as follows :

Let Γ be a rectifiable curve in the complex plane and let λ be the length measure on Γ . Then the pair (Γ, μ) satisfies (\mathcal{L}^*) .

It is worthwhile to mention that the LEJA's paper [5] permits immediately to obtain the following estimate for the function $\mathcal{M}(t) \equiv \mathcal{M}(t; [A, b], \lambda)$:

$$\mathcal{M}(t) \leq \mathcal{J} \left(\sqrt{\frac{1-t}{t-9/10}} \right), \quad 9/10 < t < 1,$$

where

$$\mathcal{J}(\alpha) := \exp \int_0^1 \log \frac{x^2 + \alpha^2}{x^2} dx \leq \exp \alpha(\pi + \alpha).$$

The exact formula for the function $\mathcal{M}(t; [a, b], \lambda)$, where $[a, b]$ is a bounded interval of the real line \mathbf{R} and λ is the Lebesgue measure on \mathbf{R} , reads as follows

$$\mathcal{M}(t; [a, t], \lambda) = 2t^{-1} - 1 + 2t^{-1}\sqrt{1-t}, \quad 0 \leq t \leq 1,$$

and may be easily derived from the following inequality due to DUDLEY and RANDOL [4]

$$\|f\|_{[a,b]} / \|f\|_A \leq (2^{-1} - 1 + 2t^{-1}\sqrt{1-t})^{\deg f}$$

true for every polynomial f of a complex variable and for every compact set $A \subset [a, b]$ with $\lambda(A) \geq t(b-a)$, $0 \leq t \leq 1$.

This paper was written during the author's stay at the Université Paul Sabatier, Toulouse (September-October 1987), as an invited profesor. I want to thank the University for the invitation. My warmest thanks go to Profesor Nguyen Thanh Van for his extraordinary hospitality and for the inspiring discussions on the subject of the paper.

1 - Determining measures for arbitrary bounded Borel subsets of \mathbb{C}^N

Let us start with the following.

LEMMA 1.1.— *Let F be a subset of a bounded set E in \mathbb{C}^N . If the set*

$$G := \{z \in E \setminus F; L_F^*(z) > 0\}$$

is not plp, then there exist a nonplp subset G_o of G , a number $b > 1$ and a polynomial family \mathcal{F} such that

- 1) \mathcal{F} is bounded at each point of F .
- 2) \mathcal{F}_b given by (0.2) is unbounded at each point $z \in G_o$.

Proof.— It is known that $F_1 := \{z \in F; L_F^*(z) > 0\}$ is plp, so there exists a function w in the class \mathcal{L} with $w = -\infty$ on F_1 and $w \leq -\log 2$ on E . It is also known [10] that w can be represented in the form

$$w = \left(\limsup_{m \rightarrow \infty} \frac{1}{m} \log |P_m| \right)^*, \quad (1.1)$$

where P_m is a polynomial on \mathbb{C}^N of degree $\leq m$. We shall consider two cases : either $F_1 = F$, or $F_1 \neq F$.

Case $F_1 = F$. By Bedford-Taylor theorem on negligible sets [1] the set

$$\left\{ \limsup_{m \rightarrow \infty} \sqrt[m]{|P_m|} < \left(\limsup_{m \rightarrow \infty} \sqrt[m]{|P_m|} \right)^* \right\}$$

is plp. Hence there exists a non pluripolar subset G' of G such that

$$-\infty < w(z) = \limsup_{m \rightarrow \infty} \frac{1}{m} \log |P_m(z)| \text{ for } z \in G'.$$

There is a real number ϵ with $0 < \epsilon < 1$ such that the set $G_o := \{x \in G'; w(z) \geq \log \epsilon\}$ is not plp. Take any b with $1 < b < 2$. Then the family

$$\mathcal{F} := \left\{ \left(\frac{2}{\epsilon} \right)^m P_m; m \geq 1 \right\}$$

has the required properties. Indeed, 1) is satisfied because

$$\limsup_{m \rightarrow \infty} \sqrt[m]{\left(\frac{2}{\epsilon b} \right)^m |P_m(z)|} = 0 \text{ on } F.$$

If $z \in G_o$, we have

$$\limsup_{m \rightarrow \infty} \sqrt[m]{\left(\frac{2}{\epsilon b}\right)^m |P_m(z)|} = \frac{2}{\epsilon b} \exp w(z) > \frac{2}{b} > 1, \text{ which implies 2).}$$

Case $F_1 \neq F$. Since $F_1 \neq F$, we have $L_F^* \in \mathcal{L}$ and $u_k := \frac{1}{k} w + \frac{k-1}{k} L_F^* \in \mathcal{L}$ for every $k \geq 1$. If $z \in G$ and $w(z) > 0$ the sequence $u_k(z)$ is increasing to the limit $L_F^*(z) > 0$. Therefore there exists k such that the set $G_k := \{z \in G; u_k(z) > 0\}$ is not plp. For such k there is $\epsilon > 0$ such that

$$G' := \{z \in G; u_k(z) \geq \log(1 + \epsilon)\}$$

is not plp. Write u_k in the form

$$u_k = \left(\limsup_{j \rightarrow \infty} \frac{1}{j} \log |P_j| \right)^* \quad (\deg P_j \leq j)$$

By the theorem on negligible sets there is a non pluripolar subset G_o of G' with

$$u_k(z) \equiv \frac{1}{k} w(z) + \frac{k-1}{k} L_F^*(z) = \limsup_{j \rightarrow \infty} \frac{1}{j} \log |P_j(z)|, z \in G_o.$$

The set G_o , any number b with $1 < b < 1 + \epsilon$ and the polynomial family $\mathcal{F} := \{P_j; j \geq 1\}$ have the required property. Indeed $\limsup \sqrt[j]{|P_j(z)|} \leq \exp u_k(z) \leq 2^{-k} < 1$ on F , which gives 1). On the other hand, if $z \in G_o$ then

$$\limsup_{j \rightarrow \infty} \sqrt[j]{b^{-j} |P_j(z)|} = b^{-1} \exp u_k(z) \geq \frac{1 + \epsilon}{b} > 1,$$

which implies 2).

LEMMA 1.2.— *If a polynomial family \mathcal{F} is bounded q^* .a.e. on a subset E of \mathbb{C}^N , then for every $b > 1$ the family \mathcal{F}_b is bounded q.a.e. on E and uniformly on a neighborhood of every \mathcal{L} -regular point a of E . If E is compact and \mathcal{L} -regular, and \mathcal{F} is bounded q^* .a.e. on E then for each $b > 1$ the family \mathcal{F}_b is uniformly bounded on a neighborhood of E .*

Proof.— Without loss of generality we can assume E is not plp. Let \mathcal{F} be a polynomial family bounded at each point of a subset F of E with $L_F^* = L_E^*$. Put

$$E_j := \{z \in E; |f(x)| \leq j, \forall f \in \mathcal{F}\}, j \geq 1 \quad (1.2)$$

Then $E_j \subset E_{j+1}$ and $F \subset E_o := \bigcup_1^\infty E_j$. Hence $L_{E_j}^* \downarrow L_{E_o}^* = L_F^* = L_E^*$. By the definition of the \mathcal{L} -extremal function we have

$$|f(z)| \leq j (\exp L_{E_j}^*(z))^{\deg f}, \quad z \in \mathbb{C}^N, \quad j \geq 1, \quad f \in \mathcal{F} \quad (1.3)$$

which implies that for each $b > 1$ the family \mathcal{F}_b is bounded at every \mathcal{L} -regular point of E . So \mathcal{F}_b is bounded q.a.e. on E . Moreover, if $L_E^*(a) = 0$, then $L_E^*(z) < b$ on a ball $|z-a| \leq r$. By Dini's argument there is j sufficiently large with $L_{E_j}^*(z) < b$ on the ball $|z-a| \leq r$, which implies by (1.3) that the family \mathcal{F}_b is uniformly bounded on a ball $|z-a| < r$, if a is any \mathcal{L} -regular point of E . The proof of the remaining part of Lemma 1.2. is trivial.

THEOREM 1.3. — *Given a probability measure on a bounded Borel set E in \mathbb{C}^N the following conditions are equivalent*

- I. The measure μ is determining for E ;*
- II. If $u \in \mathcal{L}$ and $u \leq 0$ μ -a.e. on E , then $u \leq 0$ q.a.e. on E ;*
- III. If \mathcal{F} is a polynomial family bounded μ -a.e. on E , then for every $b > 1$ the family \mathcal{F}_b is bounded q.a.e. on E .*

Proof. $I \Rightarrow II$. Let u be a fixed function in the class \mathcal{L} with $u \leq 0$ μ -a.e. on E . Put $F := \{z \in E; u(z) \leq 0\}$. Then $u(z) \leq L_F^*(z) = L_E^*(z)$. Hence $u \leq 0$ q.a.e. on E .

$I \Rightarrow III$. Let \mathcal{F} be a polynomial family bounded μ -a.e. on E . Let E_j be given by (1.2). Then $E_j \subset E_{j+1}$ and $\mu(F) = \mu(E)$ for $F := \bigcup_1^\infty E_j$. By I $L_F^* = L_E^*$. It is known [12] that $L_{E_j}^* \downarrow L_F^*$ as $j \rightarrow \infty$. Hence by (1.3) the family \mathcal{F} is bounded q*.a.e. on E , and by Lemma 1.2. the family \mathcal{F}_b is bounded q.a.a. on E for every $b > 1$.

The implication $III \Rightarrow I$ follows directly from Lemma 1.1.

It remains to show that $II \Rightarrow I$. Fix $F \subset E$ with $\mu(F) = \mu(E)$ and let u be a function of the class \mathcal{L} such that $u \leq 0$ on F . Then $u \leq 0$ q.a.e. on E . Hence $u \leq L_E^*$ in \mathbb{C}^N . By the arbitrariness of u we get $L_F^* \leq L_E^*$, which gives $L_F^* = L_E^*$, because $L_E \leq L_F$.

2 - The function $\mathcal{M}(t; E, \mu)$

Given a probability measure μ on a bounded Borel set E in \mathbb{C}^N the function \mathcal{M} is defined by the formula

$$\log \mathcal{M}(t) := \sup \left\{ \sup_E L_A; A \subset E, \mu(A) \geq t \right\}, \quad 0 \leq t \leq 1 \quad (2.1)$$

It is clear that $1 \leq \mathcal{M}(t_2) \leq \mathcal{M}(t_1) \leq +\infty$ if $0 \leq t_2 < t_1 \leq 1$. The function $\mathcal{M}^*(t) := \limsup_{\tau \rightarrow t} \mathcal{M}(\tau)$ is also decreasing. It follows from (0.1) that

$$\log \mathcal{M}(t) = \sup \left\{ \sup_E u; u \in \mathcal{L}, u \leq 0 \text{ on } A, A \subset E, \mu(A) \geq t \right\}, \quad (2.2)$$

which implies

$$\sup_E u - \sup_A u \leq \log \mathcal{M}(\mu(A)), \text{ if } u \in \mathcal{L}, A \subset E, \quad (2.3)$$

$$L_A(z) \leq \log \mathcal{M}(\mu(A)) + L_E(z), z \in \mathbb{C}^N, A \subset E, \quad (2.4)$$

where A is any Borel subset of E .

Remark 2.1.— If μ vanishes on plp sets then

$$\mathcal{M}(t) \equiv \mathcal{M}_1(t) := \sup \left\{ \sup_E (\exp L_A^*); A \subset E, \mu(A) \geq t \right\}.$$

Indeed, it is clear that $\mathcal{M} \leq \mathcal{M}_1$. In order to prove the opposite inequality observe that given t with $0 \leq t \leq 1$ and $m \in \mathbf{R}$ with $m < \mathcal{M}_1(t)$ there exists $A \subset E$ such that $\mu(A) \geq t$ and $\sup_E L_A^* > \log m$. Put $A_o := \{z \in A; L_A(z) = L_A^*(z)\}$. Then $\mu(A_o) = \mu(A) \geq t$ and $L_A^* \geq L_{A_o}$. Hence $\log m < \sup_E L_A^* \leq \sup_E L_{A_o} \leq \mathcal{M}(t)$. By the arbitrariness of m we get $\mathcal{M}_1(t) \leq \mathcal{M}(t)$.

Remark 2.2.— If $\mathcal{M}^*(1) = 1$, then \mathcal{M} is continuous at $t = 1$ and $\mathcal{M}(1) = 1$. Hence, if $\mathcal{M}^*(1) = 1$ then $L_{A_n} \rightarrow L_E$ for every sequence A_n of Borel subsets of E such that $\mu(A_n) \rightarrow \mu(E)$.

Remark 2.3.— If μ vanishes on plp sets and $\mathcal{M}(1) = 1$ then $L_E^* = 0$ on E . In particular, if E is compact and $\mathcal{M}(1) = 1$ then E is \mathcal{L} -regular. Indeed, put $E_o := \{z \in E; L_E^*(z) = 0\}$. Then $\mu(E_o) = 1$ and $L_E^* \leq L_{E_o} \leq \log \mathcal{M}(1) = 0$ on E , i.e. $L_E^* = 0$ on E . It is clear that the pairs (E, μ) for which $\mathcal{M}(1) = 1$ or $\mathcal{M}^*(1) = 1$ are of great importance for applications of the \mathcal{L} -extremal function.

PROPOSITION 2.4.— *Define*

$$\mathcal{M}_{\mathcal{P}}(t) := \sup \left\{ (\|f\|_E / \|f\|_A)^{1/\deg f}; \deg f \geq 1, A \subset \subset E, \mu(A) \geq t \right\}$$

where the sup is taken over all polynomials f of degree ≥ 1 and over all compact sets $A \subset E$. If E is compact, then

$$\mathcal{M}_{\mathcal{P}}(t) \leq \mathcal{M}(t) \leq \mathcal{M}_{\mathcal{P}}^*(t), 0 \leq t \leq 1$$

Proof.— Fix t with $0 \leq t \leq 1$. Given any number m with $m < \mathcal{M}(t)$, take $A \subset E$ with $\mu(A) \geq t$ and $\sup_E L_A > \log m$. Next choose u in the class \mathcal{L} with $u \leq 0$ on A and $\sup_E u > \log m$. By the Approximation Lemma [13] there is a sequence $u_\nu := \max_{1 \leq j \leq s_\nu} \frac{1}{n_j} \log |f_j|$, where f_j is a polynomial of degree $\leq n_j$, such that $u_\nu \downarrow u$ as $\nu \rightarrow \infty$. Given $\epsilon > 0$ there exists a compact set $K \subset A$ with $\mu(K) \geq t - \epsilon$. Take ν so large that $\sup_K u_\nu < \epsilon$ and choose j with $\sup_E \frac{1}{n_j} \log |f_j| > \log m$. Then $m e^{-\epsilon} \leq (\|f_j\|_E / \|f_j\|_K)^{1/n_j} \leq \mathcal{M}_{\mathcal{P}}(\mu(K)) \leq \mathcal{M}_{\mathcal{P}}(t - \epsilon)$. Hence $m \leq \mathcal{M}_{\mathcal{P}}^*(t)$. By the arbitrariness of m we get $\mathcal{M}(t) \leq \mathcal{M}_{\mathcal{P}}^*(t)$. The inequality $\mathcal{M}_{\mathcal{P}}(t) \leq \mathcal{M}(t)$ is obvious.

PROPOSITION 2.5.— *If E is nonplp compact set in \mathbb{C}^N then $\mathcal{M}_{\mathcal{P}}(t) = \lim_{n \rightarrow \infty} B_n^{1/n}(t) = \sup_{n \geq 1} B_n^{1/n}(t)$, where*

$$B_n(t) := \sup \{ \|f\|_E; \deg f \leq n, \|f\|_A = 1, A \subset \subset E, \mu(A) \geq t \}.$$

Proof.— Given $m \geq 1$ and c with $0 < c^m < B_m(t)$, let A be a compact subset of E with $\mu(A) \geq t$ and let f_m be a polynomial of degree $\leq m$ such that $\|f_m\|_A = 1$ and $\|f_m\|_E > c^m$. Every natural number $n \geq m$ can be written in the form $n = km + r$ with $0 \leq r < m$. Observe that

$$c^{km} < \|f_m\|_E^k \leq B_n(t).$$

Hence $\liminf_{n \rightarrow \infty} B_n^{1/n}(t) \geq c$, which implies that $B_m^{1/m}(t) \leq \liminf_{n \rightarrow \infty} B_n^{1/n}(t)$, and consequently we get the required result.

THEOREM 2.6.— *Let $A \subset \mathbb{C}^p$, $B \subset \mathbb{C}^q$ be bounded Borrel sets and let μ, ν be probability measures on A and B , respectively. Put $\mathcal{M}_A(t) := \mathcal{M}(t; A, \mu)$, $\mathcal{M}_B(t) := \mathcal{M}(t; B, \nu)$ and $\mathcal{M}_{A \times B}(t) := \mathcal{M}(t; A \times B, \lambda)$ with $\lambda := \mu \otimes \nu$.*

Then

$$(i) \mathcal{M}_{A \times B}(1) \leq \mathcal{M}_A(1) \mathcal{M}_B(1)$$

$$(ii) \mathcal{M}_{A \times B}^*(1) \leq \mathcal{M}_A^*(1) \mathcal{M}_B^*(1)$$

Proof.— (i) Let $E \subset A \times B$ with $\lambda(E) = 1$ and put $B^* := \{w \in B; (z, w) \in E\}$. Then $\nu(B^*) = 1$ μ -a.e. on A . Let $u \in \mathcal{L}(\mathbb{C}^p \times \mathbb{C}^q)$, $u \leq 0$ on E . Then $u(z, w) \leq \log \mathcal{M}_B(1) + L_B(w)$ for all $z \in A_o$ and $w \in \mathbb{C}^q$, where $A_o \subset A$ and $\mu(A_o) = 1$. Hence

$$u(z, w) \leq \log \mathcal{M}_B(1) + L_B(w) + \log \mathcal{M}_A(1) + L_A(z), (z, w) \in \mathbb{C}^p \times \mathbb{C}^q.$$

Hence by (2.2) one gets (i).

(ii) Let m be a fixed number with $m < \log \mathcal{M}_{A \times B}^*(1)$. There is a sequence E_n of Borel subsets of $A \times B$ such that $\lambda(E_n) \geq 1 - 2^{-n}$ and $\sup_{A \times B} L_{E_n} > m$. Define

$$\begin{aligned} B_n^z &:= \{w \in B; (z, w) \in E_n\}, \quad z \in A, \quad n \geq 1 \\ A_{n\epsilon} &:= \{z \in A; \nu(B_n^z) \geq 1 - \epsilon\}, \quad n \geq 1, \quad 0 < \epsilon < 1 \end{aligned}$$

We claim that $\mu(A_{n\epsilon}) \rightarrow 1$ as $n \rightarrow \infty$. Indeed,

$$\begin{aligned} \lambda(E_n) &= \int_A \nu(B_n^z) d\mu(z) = \int_{A_{n\epsilon}} + \int_{A \setminus A_{n\epsilon}} \leq \mu(A_{n\epsilon}) + (1 - \mu(A_{n\epsilon})) (1 - \epsilon) \\ &= 1 - \epsilon + \epsilon \mu(A_{n\epsilon}). \end{aligned}$$

Hence $\liminf_{n \rightarrow \infty} \mu(A_{n\epsilon}) \geq 1$, which implies the claim. Fix $n \geq 1$ and let u be a function of the class $\mathcal{L}(\mathbb{C}^p \times \mathbb{C}^q)$ with $u \leq 0$ on E_n . Then for every fixed z in A we have $u(z, w) \leq 0$ on B_n^z . Therefore

$$u(z, w) \leq \log \mathcal{M}_B(\nu(B_n^z)) + L_B(w)$$

which implies

$$u(z, w) \leq \log \mathcal{M}_B(1 - \epsilon) + L_B(w) \text{ for } z \in A_{n\epsilon}, \quad w \in \mathbb{C}^q.$$

Hence

$$u(z, w) \leq \log \mathcal{M}_B(1 - \epsilon) + L_B(w) + \log \mathcal{M}_A(\mu(A_{n\epsilon})) + L_A(z)$$

for all $(z, w) \in \mathbb{C}^p \times \mathbb{C}^q$. By the arbitrariness of u we can replace u by $L_{E_n}(z, w)$. Then we get

$$m < \sup_{A \times B} L_{E_n} \leq \log \mathcal{M}_B(1 - \epsilon) + \log \mathcal{M}_A(\mu(A_{n\epsilon})), \quad n \geq 1, \quad 0 < \epsilon < 1.$$

After passing to the limits, first with n to ∞ and next with ϵ to 0, we get $m \leq \log \mathcal{M}_B^*(1) + \log \mathcal{M}_A^*(1)$. By the arbitrariness of m we get (ii).

The following corollary is important for applications of the function \mathcal{M} .

COROLLARY 2.7.— *If $\mathcal{M}_A(1) = 1$, $\mathcal{M}_B(1) = 1$, (resp. $\mathcal{M}_A^*(1) = 1$, $\mathcal{M}_B^*(1) = 1$) then $\mathcal{M}_{A \times B}(1) = 1$ (resp. $\mathcal{M}_{A \times B}^*(1) = 1$).*

Example 2.8. — Let $I = \{a, b\}$ be an interval of the real line \mathbf{R} with end points a, b such that $-\infty < a < b < +\infty$. Then

$$\mathcal{M}(t) \equiv \mathcal{M}(t; I, \lambda_1) = 2t^{-1} - 1 + 2t^{-1}\sqrt{1-t}, \quad 0 \leq t \leq 1,$$

λ_1 denoting the Lebesgue measure on \mathbf{R} .

Proof. — Without loss of generality we may assume that $I = [a, b]$ is closed. By [4] for every polynomial f of degree $\leq n$, $\|f\|_A / \|f\|_I \leq B_n(t)$, if $A \subset \subset I$ and $\lambda_1(A) \geq t(b-a)$, where

$$B_n(t) := \frac{1}{2} \left[(2t^{-1} - 1 + 2t^{-1}\sqrt{1-t})^n + (2t^{-1} - 1 - 2t^{-1}\sqrt{1-t})^n \right].$$

Moreover, if A is a subinterval of I with a common end point, this bound is best possible. Therefore by Proposition 2.5 we have

$$\mathcal{M}_p(t) = \sup_{n \geq 1} B_n^{1/n}(t) = 2t^{-1} - 1 + 2t^{-1}\sqrt{1-t}, \quad 0 \leq t \leq 1.$$

By Proposition 2.4. $\mathcal{M}_p = \mathcal{M}$.

Remark 2.9. — If Ω is a bounded open set in \mathbf{R}^N (resp. in \mathbf{C}^N) then for every determining measure μ for Ω one has $\mathcal{M}(1; \Omega, \mu) = 1$. Indeed, it is known [12] that $L_\Omega^* = L_\Omega$. So if $F \subset \Omega$ and $\mu(F) = 1$, then $L_F = L_\Omega$ which implies that $\mathcal{M}(1; \Omega, \mu) = 1$. As an example of such μ one can take the Lebesgue measure λ_N in \mathbf{R}^N (resp. λ_{2N} in \mathbf{C}^N).

If μ is a probability measure on Ω such that $\mathcal{M}^*(1; \Omega, \mu) = 1$, then the closure E of Ω , $E = \overline{\Omega}$, is an \mathcal{L} -regular compact. Indeed, let K_n be an increasing sequence of \mathcal{L} -regular compact subsets of Ω such that $\mu(\Omega) = \lim \mu(K_n)$ and $\Omega = \bigcup_1^\infty K_n$. Then

$$\log \mathcal{M}(\mu(K_n)) \geq \sup_{\Omega} L_{K_n} = \sup_E L_{K_n} \geq \sup_E L_E^*, \quad n \geq 1,$$

which implies that $L_E^* = 0$ on E , i.e. E is \mathcal{L} -regular.

Example 2.10. — We shall now construct a bounded open subset Ω of \mathbf{C} with the following properties.

- 1) $E := \overline{\Omega}$ is \mathcal{L} -regular.
- 2) For every probability measure μ on Ω , $\mathcal{M}^*(1; \Omega, \mu) > 1$.
- 3) There exists no finite positive Borel measure μ on Ω such that the pair (Ω, μ) satisfies the (\mathcal{L}^*) -condition.

Indeed, let $\{a_n\}$ be a discrete sequence in the upper half plan $\{Im z > 0\}$ such that each point of $I = [0, 1]$ is a limit of a subsequence of $\{a_n\}$ and the sequence $\{a_n\}$ has no other limit points. There exists a sequence of positive real numbers $\{r_n\}$ such that

$$L_{\Omega_n}(z) \geq 5 - (2^{-1} + \dots + 2^{-n}), \quad z \in I, \quad n \geq 1, \quad (*)$$

with $\Omega_n := \bigcup_{j=1}^n \{|z - a_j| < r_j\}$. Namely, it is clear that $L_{\Omega_1}(z) > 5 - 2^{-1}$ on I , if $r_1 > 0$ is sufficiently small. Suppose r_1, \dots, r_n are already chosen so that $(*)$ is satisfied. Put $\Omega(r) := \Omega_n \cup \{|z - a_{n+1}| < r\}$. Then $L_{\Omega(r)} \uparrow L_{\Omega_n}$ in $\mathbb{C} \setminus \{a_{n+1}\}$ as $r \uparrow 0$. By Dini's argument the convergence is uniform on I . Hence $(*)$ is satisfied for $n+1$ with $r = r_{n+1}$ sufficiently small. The open set $\Omega := \bigcup_{n=1}^{\infty} \Omega_n$ has the required properties. It is clear that $E := \bar{\Omega}$ is \mathcal{L} -regular. If μ is a finite positive Borel measure on Ω , then $\log \mathcal{M}(\mu(\Omega_n)) \geq \sup_{\Omega} L_{\Omega_n} = \sup_E L_{\Omega_n} \geq 4$ ($n \geq 1$). Hence $\mathcal{M}^*(1; \Omega, \mu) \geq 4$. The set $G := \{z \in E : L_{\Omega}^*(z) > 0\}$ contains the interval I , so G is not plp. By Lemma 1.1 the pair (Ω, μ) does not satisfy (\mathcal{L}^*) (see also theorem 3.1).

PROPOSITION 2.11. — *If μ is determining for a nonpluripolar bounded Borel set in \mathbb{C}^N , μ vanishes on plp sets and $\mathcal{M}^*(1; E, \mu) = 1$ then (E, μ) satisfies (\mathcal{L}^*) .*

Proof. — Given a polynomial family \mathcal{F} bounded μ -a.e. on E , let E_j be the sequence of subsets of E defined by (1.2). Then $E_j \uparrow F$ with $\mu(F) = 1$. Therefore $L_{E_j}^* \downarrow L_F^* = L_E^* = L_E$ (see Remark 2.3). Given $b > 1$ the set $\Omega_b := \{L_F^* < \sqrt{b}\}$ is an open neighborhood of E . By (1.2) and (2.4)

$$|f(z)| \leq j \left(\mathcal{M}(\mu(E_j)) \exp L_E^*(z) \right)^{\deg f}, \quad f \in \mathcal{F}, \quad j \geq 1.$$

If j is sufficiently large the family \mathcal{F}_b is bounded by j uniformly on Ω_b .

Problem 2.12. — Let $\Delta = \{|z| < 1\}$ be the unit disk on the complex plane \mathbb{C} . Let θ denote the length measure on the boundary $\partial\Delta$ of Δ and let λ_2 be the Lebesgue measure on $\mathbb{C} \equiv \mathbb{R}^2$. Compute the functions $\mathcal{M}(t; \partial\Delta, \theta)$ and $\mathcal{M}(t; \Delta, \lambda_2)$, $0 \leq t \leq 1$.

3 - Determining measures for bounded Borel sets with \mathcal{L} -regular closure

The main result of this section is given by the following.

THEOREM 3.1. — *Let μ be a probability measure on a bounded Borel set E in \mathbb{C}^N such that \overline{E} is \mathcal{L} -regular. Then the following conditions are equivalent.*

- (1) *The pair (E, μ) satisfies (\mathcal{L}^*) -condition;*
- (2) *If $u \in \mathcal{L}$ and $u \leq 0$ μ -a.e. on E then $u \leq 0$ on \overline{E} ;*
- (3) *$\mathcal{M}^*(1) \equiv \mathcal{M}^*(1; E, \mu) = 1$ and $L_E = L_{\overline{E}}$;*
- (4) *$\mathcal{M}(1) \equiv \mathcal{M}(1; E, \mu) = 1$ and $L_E = L_{\overline{E}}$;*
- (5) *If $A \subset E$ and $\mu(A) = 1$, then $L_A = L_{\overline{E}}$;*
- (6) *For every $b > 1$ there exists a neighborhood Ω of \overline{E} such that for every polynomial family \mathcal{F} bounded μ -a.e. on E the family \mathcal{F}_b (given by (0.2)) is uniformly bounded on Ω ;*
- (7) *If \mathcal{F} is a polynomial family bounded μ -a.e. on E then for every number $b > 1$ the family \mathcal{F}_b is bounded q.a.e. on \overline{E} .*

Proof. — (1) \Rightarrow (2). Let u be a function of the class \mathcal{F} with $u \leq 0$ μ -a.e. on E . The function u can be written in the form

$$u = \left(\limsup_{j \rightarrow \infty} \frac{1}{j} \log |f_j| \right)^*$$

where f_j is a polynomial of degree $\leq j$. Given any fixed number $b > 1$ the polynomial family $\mathcal{F} := \{b^{-j} f_j; j \geq 1\}$ is bounded μ -a.e. on E . By (1) there are a constant $M > 0$ and a neighborhood Ω of E such that

$$\|f_j\|_{\Omega} \leq M b^{2j}, \quad j \geq 1,$$

which implies $\|f_j\|_{\overline{E}} \leq M b^{2j}$ ($j \geq 1$). Hence by the definition of $L_{\overline{E}}$ we obtain $\frac{1}{j} \log |f_j(z)| \leq \frac{1}{j} \log M + 2 \log b + L_E(z)$ in \mathbb{C}^N ($j \geq 1$). Therefore $u(z) \leq 2 \log b$ on \overline{E} . By the arbitrariness of $b > 1$ we get $u \leq 0$ on \overline{E} .

(2) \Rightarrow (3). If (2) is satisfied, then $L_E \leq L_{\overline{E}} \leq L_E$, so that $L_E = L_{\overline{E}}$. It remains to show that $\lim_{t \uparrow 1} \mathcal{M}(t) = 1$. Suppose there exists $b > 1$ with $\mathcal{M}(t) > b$ for all t with $0 < t < 1$. Let A_n be Borel subsets of E such that

$$\mu(A_n) \geq 1 - 2^{-n} \text{ and } \sup_E (\exp L_{A_n}) > b \quad (n \geq 1) \quad (*)$$

Put $E_n := A_n \cap A_{n+1} \cap \dots$ and observe that $E_{n+1} \supset E_n$, $E_n \subset A_n$ and

$$\begin{aligned} \mu(E_n) &= \mu(A_n) - \mu(A_n \setminus E_n) \geq \mu(A_n) - \mu(E \setminus E_n) \\ &\geq \mu(A_n) - \sum_{j=0}^{\infty} \mu(E \setminus A_{n+j}) \geq 1 - 2^{-n} - \sum_{j=0}^{\infty} 2^{-n-j} = 1 - 3 \cdot 2^{-n}, \end{aligned}$$

which implies that $\mu(E_n) \rightarrow 1$. Put $F := \cup E_n$. Then $L_{E_n}^* \downarrow L_F^*$ and $\mu(F) = 1$. By (2) $L_F \leq L_{\overline{E}}$ and since $L_{\overline{E}} \leq L_F$, we get $L_F = L_{\overline{E}}$. By Dini's argument $L_{E_n} \leq L_{E_n}^* < \log b$ on \overline{E} , if $n > n_o = n_o(b)$. This however contradicts the second inequality of (*). Therefore $\mathcal{M}^*(1) = \lim_{t \uparrow 1} \mathcal{M}(t) = 1$.

(3) \Rightarrow (4) obvious.

(5) \Rightarrow (6) If $b > 1$, then the set $\Omega_b := \{z \in \mathbb{C}^N; L_E(z) < \sqrt{b}\}$ is by (5) an open neighborhood of \overline{E} . Let \mathcal{F} be a polynomial family bounded μ -a.e. on E . Put $E_k := \{z \in E; |f(z)| \leq k, \forall f \in \mathcal{F}\}$. Then $E_k \subset E_{k+1}$ and $\mu(E_k) \uparrow 1$. Hence by (5) $L_{E_k}^* \downarrow L_A^* = L_{\overline{E}}$ with $A := \cup_1^\infty E_k$, the convergence being uniform on \overline{E} . Hence $L_{E_k} \leq \frac{1}{2} \log b$ on \overline{E} if $k > k_o$. It is clear that

$$\begin{aligned} |f(z)| &\leq k(\exp L_{E_k}(z))^{\deg f} \\ &\leq k \left(\exp \left[\frac{1}{2} \log b + L_E(z) \right] \right)^{\deg f} \\ &\leq k b^{\deg f}, \text{ if } z \in \Omega_b, f \in \mathcal{F}, k > k_o. \end{aligned}$$

(6) \Rightarrow (7) is obvious.

(7) \Rightarrow (1) follows from lemma 1.2.

4 - Determining measures for compact sets in \mathbb{C}^N

THEOREM 4.1. — *If μ is a probability measure on a compact set E in \mathbb{C}^N vanishing on plp subsets of E , then the following conditions are equivalent.*

- (i) *The pair (E, μ) satisfies (\mathcal{L}^*) ;*
- (ii) *If $u \in \mathcal{L}$ and $u \leq 0$ μ -a.e. on E , then $u \leq 0$ on E ;*
- (iii) $\mathcal{M}^*(1, E, \mu) = 1$;
- (iv) $\mathcal{M}(1; E, \mu) = 1$;
- (v) μ *is determining for E and E is \mathcal{L} -regular.*

Proof. — First observe that each of the conditions (i), (ii), (iii), (iv) implies \mathcal{L} -regularity of E , and next apply Theorem 3.1.

Example 4.2 . — (most likely well known to the reader). Let E be a compact subset in the complex plane. Assume E has a positive logarithmic

capacity $c(E)$. By the classical potential theory there exists a unique probability measure λ with support on E such that

$$\log c(E) = \int_E \int_E \log |z - \zeta| d\lambda(z) d\lambda(\zeta) = \sup_{\mu} \int_E \int_E \log |z - \zeta| d\mu(z) d\mu(\zeta)$$

the supremum being taken over all probability measures μ on E . The measure λ is called the *equilibrium measure of E* . We shall show that λ is determining for E . Indeed, if F is a Borel subset of E with $\lambda(F) = 1$ there is (by Choquet capacitability theorem) a sequence F_n of compact subsets of F with $c(F_n) \nearrow c(F)$. Without loss of generality we may assume E is contained in the disk $|z| < 1/2$. Then

$$\begin{aligned} \log c(E) &\geq \log c(F_n) \geq \frac{1}{\lambda^2(F_n)} \int_{F_n} \int_{F_n} \log |z - \zeta| d\lambda(z) d\lambda(\zeta) \\ &\geq \frac{1}{\lambda^2(F_n)} \log c(E). \end{aligned}$$

Therefore $c(F_n) \uparrow c(E) = c(F)$. For all sufficiently large n the function $u_n(z) := L_{F_n}^*(z) - L_E^*(z)$, $u_n(\infty) := \log[c(E)/c(F_n)]$, is harmonic in $\overline{\mathbb{C}} \setminus \widehat{E}$, $u_{n+1} \leq u_n$ and $u_n(\infty) \downarrow 0$. By Harnack's theorem $u_n \downarrow 0$ locally uniformly in $\overline{\mathbb{C}} \setminus \widehat{E}$. The function $u := \lim L_{F_n}^*$ is subharmonic on \mathbb{C} , $u \geq L_E^*$ on \mathbb{C} , and $u = L_E^*$ in $\mathbb{C} \setminus \widehat{E}$ as well as at each regular point of $\partial \widehat{E}$. By the generalized maximum principle for subharmonic function, $u \leq 0$ on \widehat{E} except at most the polar set of irregular points of $\partial \widehat{E}$. On the other hand $u \geq 0$ on \mathbb{C} . Therefore $u = L_E^*$. Observe that $L_E \leq L_F \leq L_{F_n}$ ($n \geq 1$). Hence $L_E^* = L_F^*$. It follows that μ is determining for E . Hence by theorem 3.1, if E is an \mathcal{L} -regular subset of \mathbb{C} and λ is the equilibrium measure of E , then the pair (E, λ) satisfies each of the equivalent conditions of theorem 4.1.

Remark 4.3. — Given a norm \mathcal{N} on \mathbb{C}^N the logarithmic capacity $c(E) \equiv c(E, \mathcal{N})$ of a bounded subset E of \mathbb{C}^N is defined by the formula

$$-\log c(E) := \limsup_{\mathcal{N}(z) \rightarrow \infty} [L_E(z) - \log \mathcal{N}(z)].$$

If E is a probability measure on E with $\mathcal{M}(1; E, \mu) = 1$, then for every $F \subset E$ with $\mu(F) = \mu(E)$ one has $c(F) = c(E)$.

On the plane, if E is bounded and $F \subset E$, then : $c(F) = c(E) \Leftrightarrow L_F^* = L_E^*$, which implies that μ is determining for E iff $F \subset E$, $\mu(F) = 1 \Rightarrow c(F) = c(E)$ (i.e.; iff μ is determining in the sense of ULLMAN [14]).

If $N \geq 2$ and $F \subset E$, it is clear that $L_F^* = L_E^* \Rightarrow \forall \mathcal{N} c(F, \mathcal{N}) = c(E, \mathcal{N})$. But we do not know whether the inverse implication is true.

The aim of the following example is to illustrate an application of theorem 2.6.

Example 4.4. — Let Ω be a bounded open set in \mathbf{R}^N (resp. in \mathbf{C}^N). Then it is known that λ_N (resp. λ_{2N}) is determining for Ω . We can propose the following proof of this result.

It is sufficient to consider the case of \mathbf{R}^N (because by (1.1) for every $u \in \mathcal{L}(\mathbf{C}^N)$ there is $\tilde{u} \in \mathcal{L}(\mathbf{C}^{2N})$ such that $\tilde{u}(x_1, y_1, \dots, x_N, y_N) = u(x_1 + iy_1, \dots, x_N + iy_N)$ for $(x_1 + iy_1, \dots, x_N + iy_N) \in \mathbf{C}^N \equiv \mathbf{R}^{2N}$. Hence, if $u \in \mathcal{L}(\mathbf{C}^N)$ and $u \leq 0$ λ_{2N} -a.e. on $\Omega \subset \mathbf{C}^N$ then $u \leq 0$ on Ω). Let $u \in \mathcal{L}(\mathbf{C}^N)$ and let $u \leq 0$ λ_N -a.e. on Ω . Given a point $a = (a_1, \dots, a_N)$ in Ω , let $Q := \{|x_j - a_j| \leq r \ (j = 1, \dots, N)\}$ be a closed cube with center a contained in Ω . Since by Theorem 4.1 (via example 2.8) λ_1 is determining for $[a_j - r, a_j + r]$, so by theorem 2.6 the measure λ_N is determining for the cube Q . Therefore $u \leq 0$ on Q . By the arbitrariness of Q we get $u \leq 0$ on Ω . Hence $L_\Omega = L_F^*$ for every $F \subset \Omega$ with $\lambda_N(F) = \lambda_N(\Omega)$.

Let $I^N = [0, 1]^N$ be the unit cube in \mathbf{R}^N . If A is a nonsingular affine mapping of \mathbf{R}^N onto itself, then the set $P := A(I^N)$ is called a parallelepiped.

Let Ω be a bounded open subset of \mathbf{R}^N such that for each point $b \in \overline{\Omega}$ there exists a parallelepiped P such that $P \subset \Omega \cup \{b\}$ and $b \in P$. Then $\overline{\Omega}$ is \mathcal{L} -regular and the pair (Ω, λ_N) satisfies each of the equivalent conditions of theorem 3.1.

Indeed, it is easy to see that each parallelepiped P is \mathcal{L} -regular. Therefore $\overline{\Omega}$ is \mathcal{L} -regular, because $L_{\overline{\Omega}} \leq L_P$. We already know that the pair (I^N, λ_N) satisfies (\mathcal{L}^*) . Hence for every parallelepiped P the pair (P, λ_N) satisfies (\mathcal{L}^*) . Therefore the pair (Ω, λ_N) satisfies (\mathcal{L}^*) at each point of $\overline{\Omega}$, which implies that (Ω, λ_N) satisfies (6) of Theorem 3.1.

5 - Polynomial inequality of Bernstein-Markov type and pairs (E, μ) satisfying the (\mathcal{L}^*) -condition

DEFINITION 5.1. — Let p be a positive number, E a bounded Borel set in \mathbf{C}^N and μ a probability measure on E . We say that the triple (p, E, μ) has Bernstein-Markov Property, if for every $b > 1$ there exist a positive constant M and a neighborhood G of E such that for every polynomial f of

N complex variables one has

$$\|f\|_G \leq M b^{\deg f} \|f\|_{\mu p} \quad (BM)$$

with $\|f\|_{\mu p} := (\int_E |f(x)|^p d\mu(z))^{1/p}$.

It was shown in [11] that if (E, μ) satisfies (\mathcal{L}^*) and μ satisfies some density condition, then the triple (p, E, μ) has BMP for every $p > 0$. Due to a remark by A.ZERIAHI the density condition may be dropped and one gets the following.

THEOREM 5.1. — *Let E be a Borel subset of \mathbb{C}^N and let μ be a positive measure on E such that (E, μ) satisfies (\mathcal{L}^*) . Then for every $p > 0$ the triple (p, E, μ) has the Bernstein-Markov Property (BMP).*

Proof. — Let $s(f)$ denote the degree of f . It is sufficient to prove that for every $p > 0$ and for every $b > 1$ there exists a constant $M > 0$ such that for every polynomial f

$$\|f\| := \|f\|_E \leq M b^{s(f)} \|f\|_{\mu p}.$$

Suppose the statement is not true. Then we can find $p > 0$, $b > 1$ and a sequence of polynomials f_k such that

$$\|f_k\| > k^k b^{s(f_k)} \|f_k\|_{\mu p} \text{ for } k \geq 1. \quad (5.1)$$

It follows that $\|f_k\| > 0$ and $0 < \|f_k\|_{\mu p} < +\infty$ ($k \geq 1$). We claim that for every $q > 1$ and every $\eta > 1$ the sequence of polynomials $g_k := \eta^{-k} q^{-s(f_k)} f_k / \|f_k\|_{\mu p}$ is bounded μ -a.e. on E . Indeed, following NGUYEN THANH VAN [8], put $E_{nk} := \{z \in E; |g_k(z)| \geq n\}$, $E_n := \bigcup_{k=1}^{\infty} E_{nk}$ and observe that

$$\mu(E_n) \leq \sum_{k=1}^{\infty} n^{-p} \eta^{-kp} q^{-qs(f_k)} \leq n^{-p} / \eta^{p-1}, \quad n \geq 1,$$

whence it follows that $\{g_k\}$ is bounded μ -a.e. on E . Now by the assumption (E, μ) satisfies (\mathcal{L}^*) , so that we can find $G \supset E$ and $M > 0$ such that $\|g_k\|_G \leq M q^{s(f_k)}$, $k \geq 1$. Hence

$$\|f_k\|_G \leq M \eta^k q^{2s(f_k)} \|f_k\|_{\mu p}, \quad k \geq 1 \quad (5.2)$$

Put $q = b^{1/2}$. Then (5.1) and (5.2) imply

$$k^k < M \eta^k, \quad k \geq 1,$$

which is an absurd.

THEOREM 5.2. — *If $\mathcal{M}^*(1; E, \mu) = 1$ and there is $p > 0$ such that the triple (p, E, μ) has the BMP, then (E, μ) satisfies (\mathcal{L}^*) .*

Proof. — Take $b > 1$ and let \mathcal{F} be a polynomial family bounded μ -a.e. on E . Define E_j by formula (1.2). Then $\mu(E_j) \uparrow 1$ and

$$|f(z)| \leq j \mathcal{M}(\mu(E_j))^{\deg f} \text{ for all } z \in E, f \in \mathcal{F}, j \geq 1.$$

Hence by BMP

$$\|f\|_G \leq j M [b \mathcal{M}(\mu(E_j))]^{\deg f}, \quad f \in \mathcal{F}, j \geq 1,$$

which implies the required result.

COROLLARY . — *If $\mathcal{M}^*(1; E, \mu) = 1$, then the pair (E, μ) satisfies (\mathcal{L}^*) if and only if for every $p > 0$ (for some $p > 0$) the triple (p, E, μ) has the BMP.*

References

- [1] BEDFORD (E.) et TAYLOR (B.A.). — A new capacity for plurisubharmonic functions, *Acta Math.*, t. 49, 1982, p. 1-40.
- [2] BEDFORD (E.) et TAYLOR (B.A.). — Fine topology, Silov boundary and $(dd^c)^n$, *Journal of Functional Analysis*, t. 72, 1987, p. 225-251.
- [3] DUDLEY (R.M.) and RANDOL (B.). — Implications of pointwise bounds on polynomials, *Duke Math. J.*, t. 29(3), 1962, p. 455-458.
- [4] BERNSTEIN (S.). — Sur l'ordre de la meilleure approximation des fonctions continues, *Mémoires de l'Académie Royale de Belgique* (2), t. 4, 1914, p. 1-104.
- [5] LEJA (F.). — Sur les suites des polynômes bornées presque partout sur la frontière d'un domaine, *Math. Ann.*, t. 108, 1933, p. 517-524.
- [6] LEVENBERG (N.). — Monge-Ampère measures associated to extremal plurisubharmonic functions in \mathbb{C}^N , *Trans. Amer. Math. Soc.*, t. 289 (1), 1985, p. 333-343.
- [7] NGUYEN THANH VAN . — Famille de polynômes ponctuellement bornés, *Ann. Pol. Math.*, t. 31, 1975, p. 83-90.
- [8] NGUYEN THANH VAN . — *Inégalités de Bernstein-Markov relative à une fonction poids positive.* — Séminaire d'Analyse Numérique de Toulouse, 1976-1977.

- [9] NGUYEN THANH VAN et ZERIAHI (A.). — Familles des polynômes presque partout bornées, *Bull. Sci. Math. 2ème série*, t. **107**, 1983, p. 81-91.
- [10] SICIĄK (J.). — Extremal plurisubharmonic functions in C^N , *Ann. Pol. Math.*, t. **39**, 1981, p. 175-211.
- [11] SICIĄK (J.). — On some inequalities for polynomials, *Zeszyty Nauk. Univ. Jagiellon*, t. **21**, 1979, p. 7-10.
- [12] SICIĄK (J.). — *Extremal plurisubharmonic functions and capacities in C^N* . — Sophia Kokyuroku in Math.14, Sophia University, Tokio 1982.
- [13] SICIĄK (J.). — *On logarithmic capacities and pluripolar sets C^N* . — preprint, 1986.
- [14] ULLMAN (J.L.). — Orthogonal polynomials for general measures I, *Proc 1983 Tampa Conf. on Rational Approximation and Interpolation, Springer Lectures Notes in Math.*, t. **1105**, 1985, p. 524-528.

(Manuscrit reçu le 14 décembre 1987)