

# ANNALES DE L'I. H. P., SECTION A

LOUIS MICHEL

BUNJI SAKITA

## **Group of invariance of a relativistic supermultiplet theory**

*Annales de l'I. H. P., section A*, tome 2, n° 2 (1965), p. 167-170

[http://www.numdam.org/item?id=AIHPA\\_1965\\_\\_2\\_2\\_167\\_0](http://www.numdam.org/item?id=AIHPA_1965__2_2_167_0)

© Gauthier-Villars, 1965, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## Group of invariance of a relativistic supermultiplet theory (\*)

par

**Louis MICHEL**

Institut des Hautes Études Scientifiques,  
Bures-sur-Yvette (S.-et-O.)

and

**Bunji SAKITA**

Argonne National Laboratory  
Argonne (Illinois).

---

Recently Sakita [1], Gürsey and Radicati [2] [4] and Pais [3] [4] have proposed a generalization of Wigner supermultiplet theory [5] for the nucleus to baryons and mesons [6]. This raises the question: what is a relativistic supermultiplet theory ? In this paper we shall consider only the problem of defining the invariance group  $G$  for such a theory [7].

We denote by  $P$  the connected Poincaré group. It is the semi-direct product  $P = T \times L$  where  $T$  is the translation group and  $L$  is the homogeneous Lorentz group.

CONDITION 1. — The invariance group  $G$  of a relativistic theory contains  $P$ . We shall not discuss here the discrete invariance  $P$ ,  $C$ ,  $T$ , so we shall add.

CONDITION 2. —  $G$  is a connected topological group (with  $P$  as topological subgroup) [8].

Invariance under  $G$  is considered as the largest symmetry for strong coupling physics [1] [2] [3] [4]. The particles of a supermultiplet have

---

(\*) Work performed under the auspices of the U. S. Atomic Energy Commission.

the same mass and for a given energy momentum  $p$ , all possible states (spin, charges, etc.) of these particles form a finite dimensional Hilbert space which is the space of an irreducible unitary representation of a compact group  $S_p$ , the « little group » of  $p$ . From the classical Wigner analysis it is easy to translate this as conditions on  $G$ .

CONDITION 3. — The translation group  $T$  is invariant subgroup of  $G$ . The action of  $G$  on  $T$  (by its inner automorphisms) preserves the Minkowski metric and the little group (mathematicians say stabiliser or isotopy group) in  $G/T$  of a time-like translation  $a \in T$  is a compact group  $S$ .

THEOREM. — If a group  $G$  satisfies condition 1, 2 and 3, then  $G/T$  is a direct product of  $H \times L$ .

*Proof :*

The condition 3 implies that for every  $g \in G$ ,  $a \in T$  and its transformed  $g(a) = gag^{-1}$  have same Minkowski length :  $a \cdot a = g(a) \cdot g(a)$ . In the dual of  $T$  [i. e., the four dimensional vector space of energy momentum] the orbits of  $G$  are the connected sheets of mass hyperboloid. Denote  $f: G \xrightarrow{f} \text{Aut } T$ , the homomorphism of  $G$  which describes its action, by inner automorphisms, on its invariant subgroup  $T$ . It is easy to prove [9] that the connected group of continuous automorphisms of  $G$  which preserves the Minkowski metric is  $L$ . So the image of  $f$  is  $L$ :  $\text{Im } f = L$ . Since  $T$  is abelian  $T \subset K = \text{Ker } f$ , the kernel of  $f$ , and  $f$  is factorized into  $g \circ p$ , where  $p: G \xrightarrow{p} G/T$  and  $g: G/T \xrightarrow{g} L$ . The restriction of  $g$  to the subgroup  $L = P/T \subset G/T$ , is an identity transformation. By definition of the semi-direct product, therefore,  $G/T$  is the semi-direct product  $H \times L$  where  $H = \text{Ker } g = K/T$ . Furthermore, by definition of  $H = \text{Ker } g$ ,  $H$  is an invariant subgroup of every stabilizer (little group)  $S$  for any  $a$ . For a time-like  $a$ ,  $S_a$  is compact, this implies that its invariant subgroup  $H$  is compact, and from a theorem of Iwasawa [10]  $H$  compact and  $G/T$  connected implies that it is a central extension of kernel  $H$ . As we have seen, it is also the semi-direct product  $H \times L$ . Hence it is a direct product:

$$G/T = H \otimes L. \quad (1)$$

The proof of the theorem also gives conditions on the little group  $S$ , for a time-like translation. Indeed it must be isomorphic to the direct product  $H \otimes R$  where  $R$  is the three dimensional rotation group. Of course, this excludes  $SU(6)$  or any simple Lie group for  $S$ .

A possible way to have a relativistic theory with supermultiplets of particles classified by irreducible unitary representation of  $SU(6)$ , is to find a (connected Lie) group  $\bar{G}$  with irreducible unitary representations characterized by  $m > 0$  and those of  $SU(6)$ , and such that  $\bar{G} \supset \bar{P}$ , the covering of the Poincaré group. We proceed now to build such a group  $\bar{G}$ .

Among all subgroups of the linear group with enumerable dimension let us look for the smallest group  $H$  such that:

$$SU(6) \subset H, \quad SL(2, C) \subset H, \quad SU(6) \cap SL(2, C) = SU(2) \quad (2)$$

where  $SU(2)$  is the covering of  $R$  and  $SL(2, C)$  the covering of  $L$ . The smallest group  $H$  is the intersection of all groups which satisfy (2).  $SL(6, C)$  is one of them, so  $SU(6) \subset H \subset SL(6, C)$ . But  $SU(6)$  is maximal subgroup of  $SL(6, C)$ ; this implies:  $H = SL(6, C)$ . The elements  $x \in H$  are  $6 \times 6$  matrices with determinant 1. They can be decomposed in a unique way into the product  $x = hu$  where  $h$  is a  $6 \times 6$  hermitian positive matrix of determinant 1 and  $u$  is a  $6 \times 6$  unitary matrix with determinant 1. The matrices  $u$  generates  $SU(6)$  and the set  $\{h\}$  of matrices  $h$  is the homogenous space  $SL(6, C)/SU(6)$ . The smallest Lie group generated by  $\{h\}$  is the additive group of the  $6 \times 6$  hermitian matrices; it is the 36 real parameter simply connected abelian Lie group. We shall denote it  $T_{36}$ .

The group  $\bar{G}$  is the semi-direct product of  $SL(6, C)$  by  $T_{36}$  with the action:  $h \rightarrow xhx^*$  (indeed this contains the action of  $SL(2, C)$  on  $T_4$ , hence  $\bar{G} \supset \bar{P}$ ). The orbit of  $\bar{G}$  on  $T_{36}$  are characterized by  $\det h$  and the sign of the eigenvalues. If  $h > 0$ , one can take as representative  $h = m1$ . Its little group is that of the matrices with determinant 1 such that  $xhx^* = mx^*x = m1$ ; it is  $SU(6)$ .

Hence the smallest connected Lie group which contains  $P$  and has unitary irreducible linear representations characterized by:  $m > 0$  and the unitary representations of  $SU(6)$ , is the group  $G$  we just defined. It is a 106 parameter Lie group [11]. As we shall explain elsewhere the use of such  $G$  as invariance group for a relativistic supermultiplet theory of elementary particle is possible, but we do not like it.

One of us (L. M.) has benefitted very much from a two-month visit to the Argonne National Laboratory, where this work was done.

We would like to thank Dr. W. D. McGlinn for his reading of the manuscript.

## REFERENCES

- [1] B. SAKITA, *Phys. Rev.*, t. 136, B, 1964, p. 1756.
- [2] F. GÜRSEY and L. RADICATI, *Phys. Rev. Letters*, t. 13, 1964, p. 173.
- [3] A. PAIS, *Phys. Rev. Letters*, t. 13, 1964, p. 175.
- [4] F. GÜRSEY, A. PAIS and L. RADICATI, *Phys. Rev. Letters*, t. 13, 1964, p. 299.
- [5] E. P. WIGNER, *Phys. Rev.*, t. 51, 1937, p. 106.
- [6] More than twenty papers on this subject have been published or mimeographed.
- [7] For physicists who may find our simple rigorous proof too abstract we are writing a more detailed paper on the subject in terms of Lie algebra. We will also show in this paper that the invariance properties of such a theory cannot be reduced to the study of a group.
- [8] This is a purely technical condition; with the work of E. C. ZEEMAN, *J. Math. Phys.*, t. 5, 1964, p. 491, we can obtain that  $G/T$  is the semi-direct product  $H \times L$  without condition 2.
- [9] Indeed ZEEMAN [8], has proven it without the assumptions of continuity and automorphisms.
- [10] K. IWASAWA, *Ann. Math.*, t. 50, 1949, p. 507.
- [11] This group has been mentioned at the end of reference [1].

(Manuscrit reçu le 28 janvier 1965).

---

*Directeur de la publication : P. GAUTHIER-VILLARS.*

---

IMPRIMÉ EN FRANCE

DÉPÔT LÉGAL ÉD. N° 1304a.

IMPRIMERIE BARNÉOUD S. A. LAVAL, N° 5014. 3-1965.  
65-01