

ANNALES DE L'I. H. P., SECTION A

G. MARMO

G. MENDELLA

W. M. TULCZYJEW

Symmetries and constants of the motion for dynamics in implicit form

Annales de l'I. H. P., section A, tome 57, n° 2 (1992), p. 147-166

http://www.numdam.org/item?id=AIHPA_1992__57_2_147_0

© Gauthier-Villars, 1992, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Symmetries and constants of the motion for dynamics in implicit form

by

G. MARMO

Dipartimento di Scienze Fisiche,
Mostra d'Oltremare, pad.19. 80125 Napoli. Italy;
Istituto Nazionale di Fisica Nucleare. Sezione di Napoli. Italy

G. MENDELLA

Dipartimento di Scienze Fisiche,
Mostra d'Oltremare, pad.19. 80125 Napoli. Italy;
Istituto Nazionale di Fisica Nucleare. Sezione di Napoli. Italy

and

W. M. TULCZYJEW

Dipartimento di Matematica e Fisica,
Università di Camerino, 62032 Camerino (MC), Italy,
Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, Italy

ABSTRACT. — Variational principles applied to singular Lagrangians give rise to equations of motion in implicit form.

In the present paper we analyze the concepts of symmetries and constants of the motion for such differential equations.

RÉSUMÉ. — Les principes variationaux appliqués à des lagrangiennes singulières donnent lieu à équations du mouvement en forme implicite.

Dans le présent article nous analysons les concepts des symétries et constantes du mouvement pour ces équations différentielles.

PACS numbers: 0320, 0240.

INTRODUCTION

Many dynamical systems in Physics are described by singular Lagrangians in the sense of Dirac and Bergmann ([12], [13], [2], [7], [22]). This class includes all dynamical systems associated with gauge theories. In the usual Euler-Lagrange approach these theories give rise to dynamical systems (differential equations) which are implicit, *i. e.* they are not in a normal form and, because of the singularity (degeneracy) of the Lagrangian, they cannot be put in a normal form in an intrinsic way.

In the past, one of us ([23], [26], [27], [28], [19]) introduced a geometrical formalism for dealing directly with these dynamical systems in their implicit form. As a matter of fact this formalism allows for the usual Lagrangian, Hamiltonian and Hamilton-Jacobi treatment of these systems directly in their implicit form.

In this paper we wish to extend this approach by including also symmetries and constants of the motion in this setting.

The paper is organized as follows:

In section one we establish notation and recall some definitions and properties of the geometrical structure of tangent bundles. Then we briefly recall the definition of dynamical systems in their implicit form, define symmetries and constants of the motion for these systems, and prove some useful properties.

In section two we add the symplectic structure to the carrier space in order to be able to deal with Hamiltonian dynamical systems. We consider again symmetries and constants of the motion in this context and show that the usual relation between symmetries and constants of the motion is present.

For geometric formulations of dynamics we refer the reader to recent texts ([1], [18], [16]).

SECTION 1

1.1. Notation and derivations of forms

For the geometry of differential manifolds and tensor bundles we refer to standard text-books (for example [9]).

Let P be a differential manifold. We denote by TP the tangent bundle over P , equipped with the projection $\tau_p: TP \rightarrow P$. We recall that a tangent vector in TP is an equivalence class of smooth curves in P . Each curve in this class is called an integral curve of the vector. The equivalence class of a curve $\gamma: \mathbf{R} \rightarrow P$ is denoted by $t\gamma(0)$. We define the tangent prolongation

$t\gamma: \mathbf{R} \rightarrow TP$ of a curve γ by $t\gamma(s) = t\gamma_s(0)$, where $\gamma_s: \mathbf{R} \rightarrow P$ is defined by $\gamma_s(r) = \gamma(s+r)$. One of the properties of the tangent prolongation is expressed by the equality $\langle t\gamma(s), df \rangle = \frac{d}{ds}(f \circ \gamma)$, where f is a function on P .

Let P and R be two differential manifolds. Each differentiable mapping $\varphi: P \rightarrow R$ induces the tangent mapping $T\varphi: TP \rightarrow TR$ defined by $T\varphi(v) = t(\varphi \circ \gamma)(0)$, where γ is an integral curve of v . As a consequence of the definition of the tangent prolongation, we have

$$T\varphi(t\gamma(s)) = t(\varphi \circ \gamma)(s).$$

For more details on the geometry of tangent bundles we refer to [32], [10], [30], [20].

We use symbols $\mathcal{F}(P)$, $\mathcal{X}(P)$ and $\Lambda(P)$ to denote the algebra of smooth functions, the Lie algebra of vector fields and the exterior algebra of differential forms on P respectively. The space of differential k -forms will be denoted by $\Lambda^k(P)$. The space $\Lambda^0(P)$ is the same as $\mathcal{F}(P)$. The pull-back mapping induced by a differentiable mapping $\varphi: P \rightarrow R$ is denoted by $\varphi^*: \Lambda(R) \rightarrow \Lambda(P)$.

The Lie algebra of vector fields can be extended to a graded Lie algebra of derivations on $\Lambda(P)$. Here we follow the approach of Frölicher and Nijenhuis [14]; see also Klein [15].

A linear mapping $a: \Lambda(P) \rightarrow \Lambda(P)$ is called a derivation of $\Lambda(P)$ of degree r if $a\mu \in \Lambda^{k+r}(P)$ and $a(\mu \wedge \nu) = a\mu \wedge \nu + (-1)^{kr} \mu \wedge a\nu$, where k is the degree of μ .

The commutator $[a, b] = ab - (-1)^{rs} ba$ of two derivations of degrees r and s respectively is a derivation of degree $r+s$.

The exterior differential d is a derivation of $\Lambda(P)$ of degree 1.

A derivation a of $\Lambda(P)$ is said to be of type i_* if $af = 0$ for each function $f \in \Lambda^0(P)$. A derivation a of $\Lambda(P)$ is said to be of type d_* if $[a, d] = 0$. The exterior differential d is a derivation of type d_* and the commutator $[a, d]$ is a derivation of type d_* for any derivation a . If a is a derivation of type d_* then there is a unique derivation b of type i_* such that $a = [b, d]$.

A derivation of $\Lambda(P)$ is characterized by its action on $\Lambda^0(P)$ and $d\Lambda^0(P) \subset \Lambda^1(P)$. A derivation of type d_* is characterized by its action on $\Lambda^0(P)$.

For each vector field $X \in \mathcal{X}(P)$ there is a derivation i_X of degree -1 and type i_* characterized by $i_X \mu = \langle X, \mu \rangle$ for each $\mu \in \Lambda^1(P)$. The derivation $d_X = [i_X, d]$ of degree 0 coincides with the well known Lie derivative of differential forms with respect to X , usually denoted by L_X . The equation $[d_X, d_Y] = d_{[X, Y]}$ establishes a relation between the Lie bracket $[X, Y]$ of vector fields X and Y and the commutator of the derivations of type d_* associated with these fields.

The following extension of derivations turns out to be useful to deal with the tangent bundle structure TP on P .

Let $\varphi: P \rightarrow R$ be a differentiable mapping. A linear mapping $a: \Lambda(R) \rightarrow \Lambda(P)$ is called a φ -derivation from $\Lambda(R)$ to $\Lambda(P)$ of degree r if $a\mu \in \Lambda^{k+r}(P)$ and $a(\mu \wedge \nu) = a\mu \wedge \varphi^*\nu + (-1)^{kr} \varphi^*\mu \wedge a\nu$, where k is the degree of μ .

A φ -derivation from $\Lambda(R)$ to $\Lambda(P)$ is characterized by its action on $\Lambda^0(R)$ and $\Lambda^1(R)$.

Let a be a φ -derivation from $\Lambda(R)$ to $\Lambda(P)$ of degree r and let b denote a pair (b_R, b_P) of derivations of degree s of the exterior algebras $\Lambda(R)$ and $\Lambda(P)$ respectively. If the relation $\varphi^* \circ b_R = b_P \circ \varphi^*$ is satisfied, then the commutator $[a, b] = ab_R - (-1)^{rs} b_P a$ is a φ -derivation from $\Lambda(R)$ to $\Lambda(P)$ of degree $r+s$.

A φ -derivation a from $\Lambda(R)$ to $\Lambda(P)$ is said to be of type i_* if $a f = 0$ for each function $f \in \Lambda^0(R)$. The derivation a is said to be of type d_* if $[a, d] = 0$, where d stands for the pair (d, d) of exterior differentials in $\Lambda(R)$ and $\Lambda(P)$.

Properties of φ -derivations are similar to the properties of ordinary derivations of differential forms. For the complete theory of φ -derivations we refer the reader to [21].

A differentiable mapping $U: P \rightarrow TR$ such that $\tau_R \circ U = \varphi$ is called an infinitesimal deformation of φ . For each infinitesimal deformation U of φ we introduce a φ -derivation i_U from $\Lambda(R)$ to $\Lambda(P)$ of degree -1 and type i_* characterized by $i_U \mu = \mu \circ U$, where μ is a 1-form on R . We denote by d_U the φ -derivation $[i_U, d]$.

1.2. Derivations on tangent bundles

Let M be a manifold and TM its tangent bundle. We introduce a derivation which will turn out to be useful.

The identity mapping of TM is obviously an infinitesimal deformation of the projection $\tau_M: TM \rightarrow M$. We denote this deformation by T and introduce the τ_M -derivations i_T and d_T from $\Lambda(M)$ to $\Lambda(TM)$ (see [24], [25]).

We have the following proposition:

PROPOSITION 1.1. — *The kernel of the operator d_T is the space of the locally constant functions on M .*

Proof. — If f is a function on M , then $d_T f(v) = (i_T df)(v) = \langle v, df \rangle$ for each vector $v \in TM$. Hence, if $d_T f = 0$, f is locally constant. If $\mu = \mu_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}$ is a k -form, with $k > 0$, then

$$d_T \mu = (d_T \mu_{i_1 \dots i_k}) dx^{i_1} \wedge \dots \wedge dx^{i_k} + \mu_{i_1 \dots i_k} \sum_j dx^{i_1} \wedge \dots \wedge dx^{i_j} \wedge \dots \wedge dx^{i_k}$$

is a combination of independent k -forms. It follows that $d_T \mu = 0$ implies $\mu = 0$. \square

Remark. – We denote by $\tau_M^* \mathcal{F}(M)$ the subalgebra of the algebra $\mathcal{F}(TM)$ formed by pull-backs of functions on M . We denote by $d_T \mathcal{F}(M)$ the image of the mapping $d_T: \mathcal{F}(M) \rightarrow \mathcal{F}(TM)$. We note that the derivations of $\Lambda(TM)$ of type d_* are completely determined by their action on $\tau_M^* \mathcal{F}(M)$ and $d_T \mathcal{F}(M)$.

Let X be a vector field on M and Y a vector field on TM . If $T \tau_M \circ Y = X \circ \tau_M$ then Y is called a lift of X to TM and Y is said to be τ_M -projectable onto X .

Let $\varphi: \mathbf{R} \times M \rightarrow M$ be a one-parameter group of diffeomorphisms of M . The tangent lift of φ is the one-parameter group $\varphi^T: \mathbf{R} \times TM \rightarrow TM$ defined by $\varphi^T(s, \cdot) = T\varphi_s$, where φ_s denotes the mapping $\varphi(s, \cdot): M \rightarrow M$.

Let $X \in \mathcal{X}(M)$ be a complete vector field and let $\varphi: \mathbf{R} \times M \rightarrow M$ be the flow of X . The tangent lift $X^T \in \mathcal{X}(TM)$ of X is defined as the infinitesimal generator of the tangent lift φ^T of the group φ . This definition is easily extended to arbitrary fields by using local flows. The set $\mathcal{X}^T(TM)$ of all tangent lifts is a subalgebra of $\mathcal{X}(TM)$.

If $X = X^i \frac{\partial}{\partial x^i}$ is the local expression of a field X , then the local expression of the tangent lift is

$$X^T = (\tau_M^* X^i) \frac{\partial}{\partial x^i} + (d_T X^i) \frac{\partial}{\partial \dot{x}^i}.$$

It can be shown that the tangent lift X^T of a vector field $X \in \mathcal{X}(M)$ is the unique lift $Y \in \mathcal{X}(TM)$ of X satisfying:

$$[(d_Y, d_X), d_T] = 0. \tag{1}$$

Proof. – In order to prove this statement it is sufficient to show that $(d_Y d_T - d_T d_X)f = 0$, for all functions $f \in \mathcal{F}(M)$, implies that Y is the tangent lift of X . In local coordinates, with $Y = (\tau_M^* X^i) \frac{\partial}{\partial x^i} + Y^i \frac{\partial}{\partial \dot{x}^i}$, $X^i \in \mathcal{F}(M)$, $Y^i \in \mathcal{F}(TM)$, we have:

$$\begin{aligned} (d_Y d_T - d_T d_X)f &= d_Y \left(\frac{\partial f}{\partial x^i} \dot{x}^i \right) - d_T \left(\frac{\partial f}{\partial x^j} X^j \right) \\ &= \frac{\partial^2 f}{\partial x^i \partial x^j} X^j \dot{x}^i + \frac{\partial f}{\partial x^i} Y^i - \frac{\partial^2 f}{\partial x^i \partial x^j} X^j \dot{x}^i - \frac{\partial f}{\partial x^i} \frac{\partial X^i}{\partial x^j} \dot{x}^j \\ &= \frac{\partial f}{\partial x^i} \left(Y^i - \frac{\partial X^i}{\partial x^j} \dot{x}^j \right). \end{aligned}$$

If $(d_Y d_T - d_T d_X)f = 0$, then $Y^i = \frac{\partial X^i}{\partial x^j} \dot{x}^j$ or $Y = X^T$. \square

The following statements are equivalent to the earlier characterization of the tangent lift:

$$[(i_X^T, i_X), d_T] = 0 \quad (2)$$

$$[(d_X^T, d_X), i_T] = 0 \quad (3)$$

Proof. — It is easy to derive these relations from (1). In $d\Lambda^0(M)$, we find:

$$\begin{aligned} i_X^T d_T(df) &= i_X^T dd_T f = d_X^T d_T f = d_T d_X f = d_T i_X(df) \\ d_X^T i_T(df) &= d_X^T d_T f = d_T d_X f = i_T dd_X f = i_T d_X(df). \quad \square \end{aligned}$$

Having introduced the basic structures of the tangent bundle, we are now prepared to define implicit differential equations on M .

1.3. Implicit differential equations

DEFINITION 1.1. — A *first order differential equation (f.o.d.e.)* on a differential manifold M is a submanifold E of the tangent bundle TM .

DEFINITION 1.2. — A differentiable curve $\gamma: I \subset \mathbf{R} \rightarrow M$ is said to be a *solution* of a differential equation $E \subset TM$ if $\text{Im}(t\gamma) \subset E$.

A solution of E is also called an *integral curve* of E .

DEFINITION 1.3. — A differential equation $E \subset TM$ is said to be *integrable* if for each $v \in E$ there is a solution $\gamma: I \rightarrow M$ of E such that $t\gamma(0) = v$.

DEFINITION 1.4. — A differential equation $E \subset TM$ is said to be *explicit* if there is a vector field $\Gamma: M \rightarrow TM$ such that $E = \text{Im}(\Gamma)$.

Differential equations which are not explicit are said to be *implicit*. Explicit differential equations are integrable.

For a discussion of integrability of implicit differential equations and related problems we refer to [3], [4], [5], [6].

We give simple examples of integrable and non-integrable implicit differential equations.

Example 1:

$$\begin{aligned} M &= \mathbf{R}^2, & TM &= \mathbf{R}^4 \\ E &= \{ (x, y, \dot{x}, \dot{y}) \mid \dot{x}^2 + \dot{y}^2 + x^2 + y^2 = 1 \} \end{aligned}$$

E is a 3-dimensional submanifold; this means that it cannot be put in normal form, not even locally, for this would require E to be 2-dimensional. E is integrable, indeed $\tau_M(E) = 2$ -dimensional disk of radius 1; for $x^2 + y^2 < 1$ there is no problem to exhibit integral curves. For $x^2 + y^2 = 1$, $\dot{x} = \dot{y} = 0$, thus integral curves are simply constant curves.

Example 2:

$$\begin{aligned} M &= \mathbf{R}^2, & TM &= \mathbf{R}^4 \\ E &= \{ (x, 0; 0, \dot{y}) \in \mathbf{R}^4 \} \end{aligned}$$

Here E is 2-dimensional. This differential equation is clearly non integrable, indeed there is no way to get a vector in the direction of y by differentiating a curve along x .

Example 3:

$$\begin{aligned} M &= \mathbf{R}, & TM &= \mathbf{R}^2 \\ E &= \{ (x, \dot{x}) \mid \dot{x}^2 = f^2(x) \} \end{aligned}$$

Here E has the right dimension to be the image of a vector field on \mathbf{R} ; however $\dot{x} = \pm f(x)$ are two possible vector fields on \mathbf{R} with images in E . This system is clearly integrable.

Example 4:

$$\begin{aligned} M &= S^1 = \{ (x, y) \in \mathbf{R}^2 \mid x^2 + y^2 = 1 \}, & TM &= S^1 \times \mathbf{R} \\ \varepsilon : \mathbf{R} &\rightarrow S^1 \times \mathbf{R} : \varphi \mapsto (\cos \varphi, \sin \varphi, \varphi), & E &= \text{Im } \varepsilon \end{aligned}$$

This equation is integrable. Its solutions are motions in a circle S^1 with constant acceleration. For each $\varphi \in S^1$ there is an infinity of solutions γ such that $\gamma(0) = \varphi$. This is due to the fact that E is a helix, which is only locally the image of a vector field.

This brief introduction to implicit differential equations permits us to approach the problem of symmetries and constants of the motion for this class of differential equations.

1.4. Symmetries and constants of the motion for implicit differential equations

DEFINITION 1.5. — A differentiable function $f: M \rightarrow \mathbf{R}$ is said to be a *constant of the motion* for a differential equation $E \subset TM$ if for each solution $\gamma: I \rightarrow M$ of E the composition $f \circ \gamma$ is a constant function. In other words a constant of the motion is a function constant along any solution of E .

PROPOSITION 1.2. — *If $E \subset TM$ is integrable, then a function $f: M \rightarrow \mathbf{R}$ is a constant of the motion for E if and only if $\langle v, df \rangle = 0$ for each $v \in E$.*

Proof. — Let f be a constant of the motion. For each $v \in E$ there is a solution γ of E such that $t\gamma(0) = v$. Hence,

$$\langle v, df \rangle = \langle t\gamma(0), df \rangle = \frac{d}{ds}(f \circ \gamma) \Big|_{s=0} = 0.$$

Conversely, if f is a function on M such that $\langle v, df \rangle = 0$ for each $v \in E$ and if γ is a solution of E then $\frac{d}{ds}(f \circ \gamma) = \langle t\gamma(s), df \rangle = 0$. Hence, f is a constant of the motion. \square

The property $\langle v, df \rangle = 0$ for each $v \in E$, used to characterize a constant of the motion for an integrable system E , is equivalent to $d_T f|_E = 0$. The application of this criterion to a function f does not require the integrability of E . We will refer to a function $f: M \rightarrow \mathbf{R}$ with this property as a constant of the motion for $E \subset TM$ even if the integrability of E has not been established.

DEFINITION 1.6. — A diffeomorphism $\varphi: M \rightarrow M$ is said to be a *symmetry* of a differential equation $E \subset TM$ if the composition $\varphi \circ \gamma$ of φ with a solution $\gamma: I \rightarrow M$ of E is again a solution of E .

PROPOSITION 1.3. — *If $E \subset TM$ is integrable, then a diffeomorphism $\varphi: M \rightarrow M$ is a symmetry of E if and only if $T\varphi(E) = E$.*

Proof. — Let φ be a symmetry. For any $v \in E$ we consider a solution $\gamma: I \rightarrow M$ such that $t\gamma(0) = v$. Since $\varphi \circ \gamma$ is a solution,

$$T\varphi(v) = T\varphi(t\gamma(0)) = t(\varphi \circ \gamma)(0) \in E.$$

It follows that $T\varphi(E) \subset E$. By applying the same reasoning to φ^{-1} we obtain $E \subset T\varphi(E)$. Hence $T\varphi(E) = E$.

Conversely, if $T\varphi(E) = E$ and $\gamma: I \rightarrow M$ is a solution of E , then $t(\varphi \circ \gamma)(s) = T\varphi(t\gamma(s)) \in E$. Hence $\varphi \circ \gamma$ is a solution. \square

In analogy with the terminology adopted for constants of the motion, we refer to a diffeomorphism $\varphi: M \rightarrow M$ such that $T\varphi(E) = E$ as a symmetry of $E \subset TM$ even if the integrability of E has not been established.

DEFINITION 1.7. — A vector field $X: M \rightarrow TM$ is said to be an *infinitesimal symmetry* of $E \subset TM$ if the local diffeomorphisms of M belonging to the local one-parameter groups generated by X are symmetries of E .

PROPOSITION 1.4. — *A vector field $X: M \rightarrow TM$ is an infinitesimal symmetry of an integrable differential equation $E \subset TM$ if and only if $X^T(E) \subset TE$.*

Proof. — The proof is similar to that of proposition 1.3 and is based on the fact that X^T is the infinitesimal generator of φ^T , if X is the infinitesimal generator of φ . \square

The criterion $X^T(E) \subset TE$ can be applied to a vector field $X: M \rightarrow TM$ even if the integrability of E has not been established. We refer to a vector field X which satisfies this criterion as an infinitesimal symmetry of E . We note that $X^T(E) \subset TE$ simply means that the vector field X^T is tangent to E .

This inclusion is obviously equivalent to the inclusion:

$$L_{X^\Gamma} \mathcal{F}_E \subset \mathcal{F}_E$$

where \mathcal{F}_E denotes the ideal $\mathcal{F}_E = \{f \in \mathcal{F}(TM) \mid f|_E = 0\}$.

It is frequently convenient to characterize \mathcal{F}_E as generated locally by a set of independent functions $\phi^i \in \mathcal{F}_E$, in the style of the original works of Dirac ([12], [13]). If the submanifold E is characterized by such a set of functions, infinitesimal symmetries of E satisfy:

$$L_{X^\Gamma} \phi^i = A_j^i \phi^j$$

with $A_j^i \in \mathcal{F}(TM)$.

From the definition of symmetries of E it follows easily that if ϕ_1 and ϕ_2 are symmetries for E , the composition $\phi_1 \circ \phi_2$ is also a symmetry. Thus symmetries for E are a subgroup of $\text{Diff}(M)$. Similarly, infinitesimal symmetries for E are a Lie subalgebra.

1.5. Infinitesimal symmetries and constants of the motion for differential equations in normal form

The case of dynamical systems in normal form is more familiar. For this reason we give explicit proofs of proposition 1.2 and 1.4 in the case of explicit differential equations.

We consider images of vector fields for reasons of simplicity; however what we are going to say is equally valid for sets E which are unions of images of local vector fields. An example of this more general situation was given in the example 4 of section 1.3.

In local coordinates:

$$\Gamma = \Gamma^i(m) \frac{\partial}{\partial x^i}$$

and the submanifold $E \subset TM$ can be defined as:

$$E = \text{Im } \Gamma = \{(m, v) \in TM \mid \phi^i := \dot{x}^i - \Gamma^i(m) = 0\}$$

where $\{\phi^i\}$ is a generating set for E .

Explicitly, an infinitesimal symmetry $X = A^i \frac{\partial}{\partial x^i}$, $A^i \in \mathcal{F}(M)$ for Γ satisfies:

$$Z = [\Gamma, X] = 0.$$

In coordinates:

$$Z^i = \Gamma^j \frac{\partial A^i}{\partial x^j} - A^j \frac{\partial \Gamma^i}{\partial x^j} = 0$$

According to prop. 1.4, an infinitesimal symmetry of E is a vector field satisfying

$$L_{X^T} \phi^i = A_j^i \phi^j$$

where $A_j^i \in \mathcal{F}(TM)$.

The tangent lift of a generic vector field X is given by:

$$X^T = A^i \frac{\partial}{\partial x^i} + \frac{\partial A^i}{\partial x^j} \dot{x}^j \frac{\partial}{\partial \dot{x}^i}$$

Then:

$$\begin{aligned} L_{X^T} \phi^i - A_j^i \phi^j &= i_{X^T} (d\dot{x}^i - d\Gamma^i) - A_j^i \phi^j = \frac{\partial A^i}{\partial x^j} \dot{x}^j - \frac{\partial \Gamma^i}{\partial x^j} A^j - A_j^i \phi^j \\ &= \frac{\partial A^i}{\partial x^j} (\dot{x}^j - \Gamma^j) - A_j^i \phi^j + Z^i = Z^i \end{aligned}$$

It follows that $X \in \mathcal{X}(M)$ is an infinitesimal symmetry for Γ iff its tangent lift $X^T \in \mathcal{X}(TM)$ is tangent to E.

Concerning constants of the motion, if we consider the vector field Γ as a map $\Gamma : M \rightarrow TM$, we have:

$$d_T f|_E = \Gamma^* (d_T f) = \Gamma^i \frac{\partial f}{\partial x^i} = L_{\Gamma} f$$

Thus a function $f \in \mathcal{F}(M)$ is a constant of the motion for Γ iff $d_T f|_E = 0$.

Summarizing the above discussion we have, for $E = \text{Im } \Gamma$:

$$\begin{aligned} X \in \mathcal{X}(M) : [\Gamma, X] = 0 &\Leftrightarrow X^T \text{ tangent to } E \subset TM \\ f \in \mathcal{F}(M) : L_{\Gamma} f = 0 &\Leftrightarrow d_T f|_E = 0 \end{aligned}$$

Until now we have considered generic first order differential equations. To deal with Hamiltonian or Lagrangian equations in implicit form we need to further qualify M to carry a cotangent bundle structure.

SECTION 2

2.1. The tangent bundle over the phase space

We choose the manifold M to be a cotangent bundle, let us say $M = T^*Q$. We describe the geometrical structure of the space TT^*Q and prove some useful properties.

As it is well known, the canonical structure of the cotangent bundle consists of:

- (i) a projection map $\pi_Q : T^*Q \rightarrow Q$
- (ii) a canonical 1-form $\theta_0 = p_i dq^i$

(iii) a symplectic structure $\omega_0 = d\theta_0 = dp_i \wedge dq^i$

To each vector field $Z \in \mathcal{X}(Q)$ there corresponds a vector field Z^* on T^*Q characterized by $L_{Z^*}\theta_0 = 0$ and called the canonical lift of Z .

If the manifold M is a cotangent bundle, the structure of this bundle induces additional structure in the tangent bundle $TM = TT^*Q$. This additional structure includes a canonical isomorphism from TT^*Q to T^*TQ , which, in local coordinates $(q^i, p_j, \dot{q}^k, \dot{p}_l)$ of TT^*Q , is given by:

$$\alpha: (q^i, p_j, \dot{q}^k, \dot{p}_l) \mapsto (q^i, \dot{q}^k, \dot{p}_l, p_j)$$

Other objects belonging to this additional structure are the 1-forms:

$$\begin{aligned}\theta_1 &:= d_T \theta_0 = \dot{p}_i dq^i + p_i d\dot{q}^i, \\ \theta_2 &:= i_T \omega_0 = p_i dq^i - \dot{q}^i dp_i\end{aligned}$$

and the 2-form:

$$\omega := d_T \omega_0 = d\dot{p}_i \wedge dq^i + dp_i \wedge d\dot{q}^i.$$

The following relations

$$\begin{aligned}\theta_2 &= \theta_1 - d(p_i \dot{q}^i) \\ \omega &= d\theta_1 = d\theta_2\end{aligned}$$

are satisfied. The manifold TT^*Q with the 2-form ω is a symplectic manifold.

For an intrinsic construction of the isomorphism α see [23], [27], [28], [30], [11], [8]. Intrinsic definitions of forms θ_1 , θ_2 and ω can be found in [23], [26], [27], [28], [29].

From the definition of ω and the property (1), *i.e.* $L_{X^T} d_T = d_T L_X$, it follows that

$$L_{X^T} \omega = L_{X^T} d_T \omega_0 = d_T L_X \omega_0.$$

Since $d_T: \Lambda^k(T^*Q) \rightarrow \Lambda^k(TT^*Q)$ is a monomorphism for $k > 0$ (see prop. 1.1), we find:

$$L_{X^T} \omega = 0 \Leftrightarrow L_X \omega_0 = 0. \quad (4)$$

As a consequence of the geometric structure of TT^*Q , we have the following

PROPOSITION 2.1. — The tangent lift X^T of a (locally) Hamiltonian vector field $X \in \mathcal{X}(T^*Q)$ preserves the 1-form θ_2 :

$$L_{X^T} \theta_2 = 0$$

and is globally Hamiltonian with respect to the symplectic structure ω . If X is globally Hamiltonian and $i_X \omega_0 = -df$, then $i_{X^T} \omega = -d(d_T f)$.

Proof. — From the definition of θ_2 and the property $L_{X^T} i_T = i_T L_X$ [see (3)], we find that, if $L_X \omega_0 = 0$, then

$$L_{X^T} \theta_2 = L_{X^T} i_T \omega_0 = i_T L_X \omega_0 = 0.$$

Further

$$i_{X^T} \omega + d(i_{X^T} \theta_2) = i_{X^T} d\theta_2 + di_{X^T} \theta_2 = L_{X^T} \theta_2 = 0$$

Hence X^T is globally Hamiltonian. Moreover:

$$\begin{aligned} i_{X^T} \omega &= i_{X^T} d_T \omega_0 = d_T i_X \omega_0 \\ &= i_T di_X \omega_0 + di_T i_X \omega_0 = i_T L_X \omega_0 - d(-i_T i_X \omega_0) = -dF, \end{aligned}$$

where $F = -i_T i_X \omega_0$. We have used the relation (2) and the assumption that X is locally Hamiltonian. When $i_X \omega_0 = -df$, we find $F = d_T f$. \square

We remark that a tangent lift preserving θ_2 is not necessarily the lift of an Hamiltonian vector field. A simple counterexample is provided by a vector field $X = A^i \frac{\partial}{\partial q^i} + B_i \frac{\partial}{\partial p_i}$, with $\frac{\partial A^i}{\partial q^j} = 0$, $\frac{\partial B_i}{\partial p_j} = 0$, which is not in general Hamiltonian, in spite of satisfying $L_{X^T} \theta_2 = 0$.

Symmetries of differential equations in T^*Q are frequently point transformations. Similarly, infinitesimal symmetries are usually canonical lifts of vector fields on Q . Let $Z = Z^i(q) \frac{\partial}{\partial q^i}$ be a vector field on Q . Then:

$$\begin{aligned} Z^* &= Z^i \frac{\partial}{\partial q^i} - \left(p_j \frac{\partial Z^j}{\partial q^i} \right) \frac{\partial}{\partial p_i} \\ Z^{*T} &= Z^i \frac{\partial}{\partial q^i} - \left(p_j \frac{\partial Z^j}{\partial q^i} \right) \frac{\partial}{\partial p_i} + \left(\frac{\partial Z^i}{\partial q^j} \dot{q}^j \right) \frac{\partial}{\partial \dot{q}^i} - \left(\dot{p}_j \frac{\partial Z^j}{\partial q^i} + p_j \frac{\partial^2 Z^j}{\partial q^i \partial q^k} \dot{q}^k \right) \frac{\partial}{\partial \dot{p}_i} \end{aligned}$$

We note that Z^* is an Hamiltonian vector field on T^*Q as a direct consequence of the definition of the canonical lift. It follows that its tangent lift preserves θ_2 .

2.2. Generalized Hamiltonian systems

We recall a few basic facts about symplectic manifolds. We refer the reader for further details to [1], [18], [16], [31].

Let (P, ω) be a symplectic manifold. At each point p of a submanifold $N \subset P$ we consider the tangent space $T_p N \subset T_p P$ and the symplectic polar $(T_p N)^\perp = \{ w \in T_p P \mid \langle w \wedge u, \omega \rangle = 0, \forall u \in T_p N \}$. In terms of these two spaces we have the following definitions:

- (a) N is said to be isotropic at p if $T_p N \subset (T_p N)^\perp$.
- (b) N is said to be coisotropic at p if $T_p N \supset (T_p N)^\perp$.
- (c) N is said to be Lagrangian at p if it is isotropic and coisotropic: $T_p N = (T_p N)^\perp$.

In an equivalent way we can characterize Lagrangian submanifolds as follows.

DEFINITION 2.1. — A submanifold $N \subset P$ is called a Lagrangian submanifold of (P, ω) if $\omega|_N = 0$ and $\dim N = 1/2 \dim P$.

DEFINITION 2.2. — A first order differential equation which is a Lagrangian submanifold of $(TP, d_T \omega)$ is called a *generalized Hamiltonian system*. For more details see [19].

Let $\Gamma_H = \frac{\partial H}{\partial p_i} \frac{\partial}{\partial q^i} - \frac{\partial H}{\partial q^j} \frac{\partial}{\partial p_j}$ be the Hamiltonian vector field on T^*Q associated with the Hamiltonian $H \in \mathcal{F}(T^*Q)$. The image E of Γ_H , described by the equations

$$\begin{aligned} \dot{q}^i &= \partial H / \partial p_i \\ \dot{p}_j &= -\partial H / \partial q^j \end{aligned}$$

is both an integrable differential equation and a Lagrangian submanifold. Hence it is an integrable generalized Hamiltonian system. It is even an ordinary Hamiltonian system. E can be characterized in an equivalent way as the set of all points in TT^*Q on which the forms θ_2 and $(-\tau_{T^*Q}^* dH)$ coincide. In this case we say that E is generated by the function $-H(q, p)$ in the sense of [23], [29].

Also a Lagrangian function $\mathcal{L} \in \mathcal{F}(TQ)$ leads to a generalized Hamiltonian system $E \subset TT^*Q$. It is described by the equations

$$\begin{aligned} \dot{p}_i &= \partial \mathcal{L} / \partial q^i \\ p_i &= \partial \mathcal{L} / \partial \dot{q}^i. \end{aligned}$$

Also in this case E can be characterized as the set of all points in TT^*Q on which the 1-forms θ_1 and $(\pi_{T^*Q}^* d\mathcal{L})$ coincide.

A generalized Hamiltonian system obtained from a Lagrangian is not necessarily an ordinary Hamiltonian system and is not necessarily integrable. An algorithm to extract an integrable part of a generalized Hamiltonian system derived from a Lagrangian function is found in the original papers on constrained Hamiltonian systems [12], [13], [2], [7].

2.3. Infinitesimal symmetries and constants of the motion for Hamiltonian systems

The symplectic structure of a symplectic manifold (P, ω) provides a connection between infinitesimal symmetries and constants of the motion. We recall that if $f \in \mathcal{F}(P)$ is a constant of the motion of an ordinary Hamiltonian system then the associated Hamiltonian vector field $X \in \mathcal{X}(P)$ is an infinitesimal symmetry. The converse is not always true as is seen from the following example.

Let us consider a particle in a constant gravitational field, whose Lagrangian is

$$\mathcal{L} = (1/2)m(\dot{x}_1^2 + \dot{x}_2^2 + \dot{x}_3^2) - kx_3.$$

The Hamiltonian function for this system is

$$H = (1/2m)(p_1^2 + p_2^2 + p_3^2) + kx_3$$

and the dynamical vector field Γ_H is given by:

$$\Gamma_H = \frac{p_1}{m} \frac{\partial}{\partial x_1} + \frac{p_2}{m} \frac{\partial}{\partial x_2} + \frac{p_3}{m} \frac{\partial}{\partial x_3} - k \frac{\partial}{\partial p_3}$$

A well known symmetry for this system is the translation along x_3 -axis, generated by the vector field $X = \frac{\partial}{\partial x_3}$. $X^* = \frac{\partial}{\partial x_3}$ is an infinitesimal symmetry for Γ_H , since $[X^*, \Gamma_H] = 0$, but the Hamiltonian function $f = p_3$ associated with X^* via ω_0 is not a constant of the motion:

$$L_{\Gamma_H} f = L_{\Gamma_H} p_3 = -k \neq 0.$$

As we have seen in Section 1 the natural setting for the analysis of implicit differential equations is the geometry of objects lifted to the tangent bundle. For this reason we will translate the above described relations between infinitesimal symmetries and constants of the motion into relations among the lifted geometrical objects introduced earlier on the tangent bundle TT^*Q , *i. e.* the tangent lift X^T , its Hamiltonian function $F = d_T f$ and the derived symplectic structure $\omega = d_T \omega_0$.

More precisely, let E be the image of the Hamiltonian vector field Γ_H . If the function $F = d_T f \in \mathcal{F}(TT^*Q)$ satisfies $F|_E = 0$, then the Hamiltonian vector field X^T associated to F is tangent to E . However if X^T is tangent to E , then $F|_E$ is constant but not necessarily zero.

In the example described earlier, X^* is an Hamiltonian vector field whose tangent lift is obviously tangent to $E = \text{Im } \Gamma_H$. As expected, the function $d_T f = \dot{p}_3$, associated with $X^{*T} = \frac{\partial}{\partial x_3}$ via ω , satisfies

$$d_T f|_E = \dot{p}_3|_E = -k \neq 0.$$

Having defined a generalized Hamiltonian system, we may further qualify our infinitesimal symmetries to be canonical.

DEFINITION 2.2. — An infinitesimal symmetry $X: T^*Q \rightarrow TT^*Q$ of a generalized Hamiltonian system $E \subset TT^*Q$ is said to be canonical if $L_X \omega_0 = 0$.

We show that the relations between canonical infinitesimal symmetries and constants of the motion described above for ordinary Hamiltonian systems are present also in the case of generalized Hamiltonian systems.

PROPOSITION 2.2. — Let $f \in \mathcal{F}(T^*Q)$ be a constant of the motion for a generalized Hamiltonian system $E \subset TT^*Q$, and let $X \in \mathcal{X}(T^*Q)$ be the Hamiltonian vector field associated with f .

Then X is a canonical infinitesimal symmetry of E .

Proof. — By proposition 2.1, X^T is globally Hamiltonian. The assumption $F|_E=0$ implies $(i_{X^T}\omega)|_E=0$. This relation means that for each $v \in E$, we have $\langle X^T(v) \wedge w, \omega \rangle = 0 \quad \forall w \in T_v E$. Thus $X^T(v) \in (T_v E)^\perp$. Since E is Lagrangian, $(T_v E)^\perp = T_v E$. Hence $X^T(v) \in T_v E$ for each $v \in E$, i.e. X is a canonical infinitesimal symmetry of E . \square

PROPOSITION 2.3. — Let $X \in \mathcal{X}(T^*Q)$ be a canonical infinitesimal symmetry of a generalized Hamiltonian system $E \subset TT^*Q$. Then the Hamiltonian function of X , $f \in \mathcal{F}(T^*Q)$, satisfies $d_T f|_E = \text{Const}$.

Proof. — Since X is Hamiltonian, X^T is globally Hamiltonian and its Hamiltonian function is $d_T f$. The assumption that E is Lagrangian implies $\omega|_E=0$. Since X^T is tangent to E , we have also $(i_{X^T}\omega)|_E=0$. Hence $d(d_T f)|_E=0$, i.e. $d_T f|_E = \text{Const}$. \square

2.4. Implicit systems defined by Lagrangian functions

As seen before, if our system of differential equations is explicit, the submanifold E is the image of a global section Γ_H of the tangent bundle $\tau_{T^*Q}: TT^*Q \rightarrow T^*Q$.

However when, due to the singularity of the Lagrangian, the equations are truly implicit, i.e. they cannot be put in normal form, the submanifold E is not the image of a vector field, not even locally.

One of the possible consequences of the singularity of the Lagrangian is the presence of constraints. The mechanism of the appearance of constraints is clarified by the following construction, which is the geometric version of the construction described in section 2.2.

Given a Lagrangian $\mathcal{L} \in \mathcal{F}(TQ)$, we can associate with it a submanifold of T^*TQ , the image of the differential $d\mathcal{L}$. This submanifold, in turn, can be mapped into TT^*Q via the inverse of the isomorphism (see [23], [26]):

$$\alpha: TT^*Q \rightarrow T^*TQ.$$

As a matter of fact, if the Lagrangian is regular, the submanifold obtained in TT^*Q is the image of a global vector field Γ_H . But in the singular case the situation is quite different.

For regular Lagrangians, the submanifold obtained in TT^*Q projects via τ_{T^*Q} on the whole T^*Q ; on the contrary, in the singular case, the projection of this submanifold is usually only a part of T^*Q , and more precisely it coincides exactly with the image of TQ via the Legendre map $F\mathcal{L}$ it is the submanifold in T^*Q defined by the primary constraints (in the theory of Dirac and Bergmann).

It is instructive to follow the construction described above in local coordinates:

$$\begin{aligned} (x, \dot{x}) \in \text{TQ} &\xrightarrow{d\mathcal{L}} \left(x, \dot{x}, \frac{\partial \mathcal{L}}{\partial x}, \frac{\partial \mathcal{L}}{\partial \dot{x}} \right) \in \text{T}^* \text{TQ} \\ &\xrightarrow{\alpha^{-1}} \left(x, \frac{\partial \mathcal{L}}{\partial \dot{x}}, \dot{x}, \frac{\partial \mathcal{L}}{\partial x} \right) \in \text{TT}^* \text{Q} \\ &\xrightarrow{\tau_{\text{T}^* \text{Q}}} \left(x, \frac{\partial \mathcal{L}}{\partial \dot{x}} \right) \in \text{T}^* \text{Q} \end{aligned}$$

The following diagram illustrates the situation:

$$\begin{array}{ccc} \text{T}^* \text{TQ} & \xrightarrow{\alpha^{-1}} & \text{TT}^* \text{Q} \\ d\mathcal{L} \uparrow & & \downarrow \tau_{\text{T}^* \text{Q}} \\ \text{TQ} & \xrightarrow{F_{\mathcal{L}}} & \text{T}^* \text{Q} \end{array}$$

As a pedagogical aid, it is instructive to follow step by step the above construction for a very simple example; let us consider the relativistic free particle, whose Lagrangian, with the metric tensor $g_{\mu\nu} = \text{diag}(-+++)$, is given by:

$$\mathcal{L} = -m(-\dot{x}^2)^{1/2}$$

We use the homogeneous Lagrangian in order to have a parametrization invariant action functional. Explicitly, we obtain:

$$\text{TQ} \xrightarrow{d\mathcal{L}} \text{T}^* \text{TQ} \xrightarrow{\alpha^{-1}} \text{TT}^* \text{Q} \xrightarrow{\tau_{\text{T}^* \text{Q}}} \text{T}^* \text{Q}$$

$$(x^\mu, \dot{x}^\mu) \mapsto \left(x^\mu, \dot{x}^\mu, 0, \frac{m\dot{x}^\mu}{(-\dot{x}^2)^{1/2}} \right) \mapsto \left(x^\mu, \frac{m\dot{x}^\mu}{(-\dot{x}^2)^{1/2}}, \dot{x}^\mu, 0 \right) \mapsto \left(x^\mu, \frac{m\dot{x}^\mu}{(-\dot{x}^2)^{1/2}} \right)$$

The submanifold $E \subset \text{TT}^* \text{Q}$ can be described by:

$$\begin{aligned} \phi_\mu &:= p_\mu - \frac{m\dot{x}_\mu}{(-\dot{x}^2)^{1/2}} = 0 \\ \phi_{\mu+n} &:= \dot{p}_\mu = 0 \end{aligned}$$

These differential equations cannot be represented by a vector field on $\text{T}^* \text{Q}$; indeed, the projection of E only covers a closed submanifold of $\text{T}^* \text{Q}$, namely the submanifold described by

$$p_\mu p^\mu + m^2 = 0$$

We can easily obtain also the infinitesimal symmetries and constants of the motion for the submanifold E . According to our definitions, a vector

field $X \in \mathcal{X}(\text{TT}^*Q)$ given locally as:

$$X = A^\mu \frac{\partial}{\partial x^\mu} + B_\mu \frac{\partial}{\partial p_\mu} + d_T A^\mu \frac{\partial}{\partial \dot{x}^\mu} + d_T B_\mu \frac{\partial}{\partial \dot{p}_\mu}$$

is the tangent lift of an infinitesimal symmetry for E represented by $\{\phi_\mu, \phi_{\mu+n}\}$ if it satisfies:

$$\begin{aligned} L_X \phi_\mu|_E &= B_\mu - m \frac{\partial}{\partial \dot{x}^\nu} \left(\frac{\dot{x}_\mu}{(-\dot{x}^2)^{1/2}} \right) d_T A^\nu|_E \\ &= B_\mu + \frac{m}{(-\dot{x}^2)^{3/2}} (\dot{x}^2 g_{\mu\nu} - \dot{x}_\mu \dot{x}_\nu) d_T A^\nu|_E = 0 \end{aligned}$$

$$L_X \phi_{\mu+n}|_E = d_T B_\mu|_E = 0.$$

It is possible to solve these equations in full generality. However, we limit ourselves to list the well known symmetries associated with the Poincaré group.

First we can consider the generator of space-time translations:

$$Z = \alpha^\mu \frac{\partial}{\partial x^\mu}$$

which is obviously an infinitesimal symmetry of E ; the associated Hamiltonian function

$$f = \alpha^\mu \frac{m \dot{x}_\mu}{(-\dot{x}^2)^{1/2}} = \alpha^\mu P_\mu$$

satisfies the condition:

$$d_T f|_E = \alpha^\mu \dot{p}_\mu|_E = 0$$

hence it is a constant of the motion for E .

Now we consider spatial rotations, generated by the vector fields:

$$J_i = \varepsilon_{ij}^k x^j \frac{\partial}{\partial x^k}$$

In this case the lift to TT^*Q is given by:

$$W_i := (J_i^*)^T = \varepsilon_{ij}^k x^j \frac{\partial}{\partial x^k} + \varepsilon_{ik}^j p_j \frac{\partial}{\partial p_k} + \varepsilon_{ij}^k \dot{x}^j \frac{\partial}{\partial \dot{x}^k} + \varepsilon_{ik}^j \dot{p}_j \frac{\partial}{\partial \dot{p}_k}$$

and it satisfies:

$$\begin{aligned} L_{W_i} \left(p_k - \frac{m \dot{x}_k}{(-\dot{x}^2)^{1/2}} \right) \Big|_E &= \varepsilon_{ik}^j p_j + \frac{m}{(-\dot{x}^2)^{3/2}} (\dot{x}^2 g_{kl} - \dot{x}_k \dot{x}_l) \varepsilon_{ij}^l \dot{x}^j \Big|_E \\ &= \varepsilon_{ik}^j \left(p_j - \frac{m}{(-\dot{x}^2)^{1/2}} \dot{x}_j \right) \Big|_E = 0 \end{aligned}$$

$$L_{W_i} \dot{p}_k|_E = \varepsilon_{ik}^j \dot{p}_j|_E = 0$$

For the Hamiltonian functions:

$$f_i = \frac{m \varepsilon_{ij}^k x^j \dot{x}_k}{(-\dot{x}^2)^{1/2}} = \varepsilon_{ij}^k x^j p_k$$

we have:

$$d_T f_i|_E = \varepsilon_{ij}^k \frac{m \dot{x}^j \dot{x}_k}{(-\dot{x}^2)^{1/2}} + \varepsilon_{ij}^k x^j \dot{p}_k|_E = \varepsilon_{ij}^k x^j \dot{p}_k|_E = 0$$

In the same way we could analyze the boost generators:

$$K_i = x^0 \frac{\partial}{\partial x^i} + x^i \frac{\partial}{\partial x^0}$$

and thus complete the list of symmetries associated with the Poincaré group.

2.5. Conclusions

The theory of dynamical systems described by explicit differential equations (vector fields) is well established. In particular, relations between symmetries and conservation laws are well known.

Dynamical systems which are not described by explicit equations were first studied by Dirac and Bergmann. Such systems, derived from singular Lagrangians, are common in theoretical physics. They are encountered in gauge theories and relativity. Systems of this kind are usually represented by families of explicit differential equations by working locally in appropriate charts and by “fixing gauges”.

In this paper we have chosen a formulation in terms of implicit differential equations and have extended the concepts of symmetries and conservation laws to this formulation.

An analysis of constrained systems based on the geometry of T^*TQ can be found in [17].

We have not yet dealt with the classification of constraints (primary, secondary, ..., first class and second class) and their relation with symmetries, neither we have described relations between the traditional formulation and the formulation in terms of implicit differential equations.

We have limited the discussion to first order differential equations on phase manifolds, leaving out description of dynamics by second order differential equations on configuration manifolds.

REFERENCES

- [1] R. ABRAHAM and J. E. MARSDEN, *Foundations of Mechanics*, Benjamin, Mass. 1978.
- [2] J. L. ANDERSON and P. G. BERGMANN, Constraints in Covariant Field Theories, *Phys. Rev.*, Vol. **83**, 1951, pp. 1018-1026.
- [3] V. I. ARNOLD, *Geometrical Methods in the Theory of Ordinary Differential Equations*, Springer-Verlag, XI, New York, Heidelberg, Berlin, 1983.
- [4] V. I. ARNOLD, A. VARCHENKO and S. GOUSSEIN-ZADÉ Eds., *Singularités des applications différentiables, I*, Editions Mir, Moscou, 1986.
- [5] V. I. ARNOLD Ed., *Dynamical Systems, III*, Springer-Verlag, New York, Heidelberg, Berlin, 1988.
- [6] D. V. ASONOV and V. I. ARNOLD Eds., *Dynamical Systems, I*, Springer-Verlag, New York, Heidelberg, Berlin, 1987.
- [7] P. G. BERGMANN and J. GOLDBERG, Dirac Bracket Transformations in Phase Space, *Phys. Rev.*, Vol. **98**, 1955, pp. 531-538.
- [8] F. CANTRIJN, M. CRAMPIN, W. SARLET and D. SAUNDERS, The Canonical Isomorphism Between T^*T^*M and T^*T^*M , *C. R. Acad. Sci. Paris*, T. **309**, Serie II, 1989, pp. 1509-1514.
- [9] Y. CHOQUET-BRUHAT, C. DE WITT-MORETTE and M. DILLARD-BLEICK, *Analysis, Manifolds and Physics*, North-Holland, Amsterdam, Oxford, New York, Tokio, 1982.
- [10] M. CRAMPIN, Tangent Bundle Geometry for Lagrangian Dynamics, *J. Phys. A: Math. Gen.*, Vol. **16**, 1983, pp. 3755-3772.
- [11] M. CRAMPIN, F. CANTRIJN and W. SARLET, Lifting Geometric Objects to a Cotangent Bundle, and the Geometry of the Cotangent Bundle of a Tangent Bundle, *J. Geom. Phys.*, Vol. **4**, 1987, pp. 469-492.
- [12] P. A. M. DIRAC, Generalized Hamiltonian Dynamics, *Can. J. Math.*, Vol. **2**, 1950, pp. 129-148.
- [13] P. A. M. DIRAC, *Lectures in Quantum Mechanics*, Belfer Graduate School of Science, Yeshiva University, New York, 1967.
- [14] A. FRÖLICHER and A. NIJENHUIS, Theory of vector valued differential forms, *Nederl. Akad. Wetensch. Proc.*, Vol. **A 59**, 1956, pp. 338-359.
- [15] J. KLEIN, On Lie Algebras of Vector Fields defined by Vector Forms, *Colloquia Mathematica Societatis Janos Bolyai*, 46, Topics in Differential Geometry, Debrecen, Hungary, 1984.
- [16] P. LIBERMANN and C. M. MARLE, *Symplectic Geometry and Analytical Mechanics*, Reidel D, Dordrecht, 1987.
- [17] L. LUSANNA, An Enlarged Phase Space for Finite-dimensional Constrained Systems, Unifying Their Lagrangian, Phase- and Velocity-space Descriptions, *Physics Reports*, Vol. **185**, 1990, pp. 1-54.
- [18] G. MARMO, E. J. SALETAN, A. SIMONI and B. VITALE, *Dynamical Systems. A Differential Geometric Approach to Symmetry and Reduction*, Wiley, New York, 1985.
- [19] M. R. MENZIO and W. M. TULCZYJEW, Infinitesimal Symplectic Relations and Generalized Hamiltonian Dynamics, *Ann. Inst. H. Poincaré*, Vol. **A28**, 1978, pp. 349-367.
- [20] G. MORANDI, C. FERRARIO, G. LO VECCHIO, G. MARMO and C. RUBANO, The Inverse Problem in the Calculus of Variations and the Geometry of the Tangent Bundle, *Phys. Rep.*, Vol. **188**, 1990, pp. 147-284.
- [21] G. PIDELLO and W. M. TULCZYJEW, Derivations of differential forms on Jet Bundles, *Ann. Matem. Pura e Applic.*, Vol. **147**, 1987, pp. 249-265.
- [22] E. C. G. SUDARSHAN and N. MUKUNDA, *Classical Dynamics: A Modern Perspective*, Wiley, New York, 1974.
- [23] W. M. TULCZYJEW, Hamiltonian Systems, Lagrangian Systems and the Legendre Transformation, *Symp. Math.*, Vol. **16**, 1974, pp. 247-258.

- [24] W. M. TULCZYJEW, Sur la différentielle de Lagrange, *C. R. Acad. Sci. Paris*, T. **280**, 1975, pp. 1295-1298.
- [25] W. M. TULCZYJEW, The Lagrange Differential, *Bull. Acad. Pol. des Sciences*, Vol. **24**, 1976, pp. 1089-1096.
- [26] W. M. TULCZYJEW, Le sous-varietes Lagrangiennes et la dynamique Hamiltonienne, *C. R. Acad. Sci. Paris*, T. **283**, 1976, p. 15.
- [27] W. M. TULCZYJEW, Le sous-varietes Lagrangiennes et la dynamique Lagrangienne, *C. R. Acad. Sci. Paris*, T. **283**, 1976, pp. 675-678.
- [28] W. M. TULCZYJEW, The Legendre Transformation, *Ann. Int. H. Poincaré*, Vol. **27**, 1977, pp. 101-114.
- [29] W. M. TULCZYJEW, A Symplectic Formulation of Relativistic Particle Dynamics, *Acta Phys. Pol.*, Vol. **B8**, 1977, pp. 431-447.
- [30] W. M. TULCZYJEW, *Geometrical Formulations of Physical Theories*, Bibliopolis, Napoli, 1989.
- [31] A. WEINSTEIN, *Lectures on Symplectic Manifolds*, American Math. Soc., Providence, R.I., 1977.
- [32] K. YANO and S. ISHIHARA, *Tangent and Cotangent Bundles*, Dekker, New York, 1973.

(Manuscript received February 25, 1991.)