

# ANNALES DE L'I. H. P., SECTION A

V. WÜNSCH

## **Moments and Huygens' principle for conformally invariant field equations in curved space-times**

*Annales de l'I. H. P., section A*, tome 60, n° 4 (1994), p. 433-455

[http://www.numdam.org/item?id=AIHPA\\_1994\\_\\_60\\_4\\_433\\_0](http://www.numdam.org/item?id=AIHPA_1994__60_4_433_0)

© Gauthier-Villars, 1994, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Moments and Huygens' principle for conformally invariant field equations in curved space-times

by

V. WÜNSCH

Fachbereich Mathematik der Pädagogischen Hochschule,  
99089 Erfurt, Germany

---

**ABSTRACT.** – By means of a certain conformal covariant differentiation process we define an infinite sequence of conformally invariant tensors (moments) for Weyl's neutrino equation in a curved space-time. In the cases of the conformally invariant-scalar wave equation and Maxwell's equations such moments were introduced by Günther. We prove some properties of the moments and study the relationship between the moments and the validity of Huygens' principle for these conformally invariant field equations. Using suitable generating systems of conformally invariant tensors we derive the first moment equations and obtain from them results on Huygens' principle.

*Key words* : Conformally invariant field equations, moments, Huygens' principle.

**RÉSUMÉ.** – Grâce à un certain processus de différentiation de covariant conformes, nous définissons une suite infinie de tenseurs invariants conformes (moments) pour l'équation de Weyl dans un espace-temps courbe. Ces moments sont introduits par Günther pour l'équation des ondes scalaires et les équations de Maxwell. Nous prouvons quelques propriétés des moments et étudions le lien entre ces moments et la validité du principe de Huygens pour les équations de champ invariante conformes. En utilisant des tenseurs invariants conformes adaptés, nous déduisons les premières équations de moments et obtenons des résultats à propos du principe de Huygens.

---

## 1. INTRODUCTION

In a four-dimensional pseudo-Riemannian manifold  $(M, g)$  with a smooth metric of Lorentzian signature the following conformally invariant field equations are considered:

$$\begin{aligned} \text{Scalar wave equation } g^{ab} \nabla_a \nabla_b u - \frac{1}{6} R u &= O & E_1 \\ \text{Maxwell's equations } d\omega = O, \delta\omega = O & & E_2 \\ \text{Weyl's neutrino equation } \nabla_X^A \varphi_A &= O & E_3 \end{aligned}$$

P. Günther [G4] defined for the equations  $E_1$  and  $E_2$  an infinite sequence of symmetric, trace-free, conformally invariant tensors

$$I_{i_1 \dots i_\nu}^\sigma(x), \quad \sigma = 1, 2, \quad \nu = 0, 1, 2, \dots$$

which he called moments of the equation  $E_\sigma$  of order  $\nu$ . He derived these moments by means of a certain conformal covariant differentiation process. The moments are of particular importance in the theory of Huygens' principle for  $E_1$  and  $E_2$ . For one of the equations  $E_\sigma$  Huygens' principle (in the sense of Hadamard's "minor premise") is valid if and only if the corresponding tail term vanishes [Ha; G2, 4; W4]. Consequently, if  $(M, g)$  is analytic we have the following relationship between the moments and the validity of Huygens' principle: The equation  $E_1$  or  $E_2$  satisfies Huygens' principle if and only if all corresponding moments vanish on  $M$ . These moment equations  $I_{i_1 \dots i_\nu}^\sigma(x) = O$ ,  $(\sigma = 1, 2)$  are determined explicitly up to now for  $0 \leq \nu \leq 4$  (see [G1, 2; W1, 2, 4; McL 2]). For example, we have  $I_{i_1 i_2}^\sigma = \alpha^{(\sigma)} B_{i_1 i_2}$ , where  $\alpha^{(\sigma)} \in \mathbb{R} \setminus \{0\}$  and  $B_{i_1 i_2}$  denotes the Bach tensor [G1, 2; W4].

In this paper we define such a sequence of conformally invariant tensors also for Weyl's neutrino equation  $E_3$ . Using some results on the theory of conformally invariant tensors [GW 2, 3], in particular, the notion "conformal covariant derivative" and suitable linear independent systems of conformally invariant tensors, we give new information about the general structure of the moment equations  $I_{i_1 \dots i_\nu}^\sigma = O$  for  $\sigma \in \{1, 2, 3\}$  and  $\nu \in \{0, 1, \dots, 6\}$ , which imply some results on Huygens' principle for  $E_\sigma$ .

The paper is organized as follows. After some preliminaries we give in Section 3 necessary and sufficient conditions for the validity of Huygens' principle for the equations  $E_\sigma$  and, further, the transformation law for the corresponding tail terms under conformal transformations of the metric. In Section 4, Günther's and the author's contributions to the theory of polynomial conformally invariant tensors [GW2, 3] are generalized by including the Levi-Civita pseudo-tensor and conformally invariant spinors. Further, some classes of conformally invariant tensors, which are important

for the moments and necessary conditions for  $(M, g)$  to be conformal to an Einstein space-time are given. In Section 5, we introduce moments also for Weyl's neutrino equation and derive some properties of the moments. Using the results of Section 4 on generating systems of conformally invariant tensors we obtain information about the algebraic structure of the first moments. Finally, in Section 6, we show the importance of the moments for the validity of Huygens' principle, especially in the case of a Petrov type N space-time.

## 2. PRELIMINARIES

Let  $(M, g)$  be a space-time, i.e. a 4-manifold together with a smooth metric of Lorentzian signature, and  $g_{ab}, g^{ab}, \nabla_a, R_{abcd}, R_{ab}, R, C_{abcd}$  the local coordinates of the covariant and contravariant metric tensor, the Levi-Civita connection, the curvatur tensor, the Ricci tensor, the scalar curvature and the Weyl curvature tensor, respectively.  $\mathcal{J}$  and  $\Lambda^p$  denote the space of the  $C^\infty$  scalar fields and the  $p$ -forms of class  $C^\infty$ , respectively. On  $\Lambda^p$  the exterior derivative  $d$ , the co-derivative  $\delta$  and  $\Delta := -(d\delta + \delta d)$  are defined. Assuming that  $(M, g)$  can be equipped with a spin structure we denote the complex spinor bundles of covariant and contravariant 1-spinors and their conjugates by  $S, S^*, \bar{S}, \bar{S}^*$ , the set of all cross sections of  $S, S^*, \bar{S}, \bar{S}^*$  by  $\mathcal{S}, \mathcal{S}^*, \bar{\mathcal{S}}, \bar{\mathcal{S}}^*$ , respectively, the coordinates of  $\varphi \in \mathcal{S}, \psi \in \bar{\mathcal{S}}$ , the connection quantities (generalized Pauli-matrices), the Levi-Civita spinor, the connection coefficients and the spinor covariant derivative by [PR]

$$\varphi_A, \psi_{\dot{X}}, \sigma_{A\dot{X}}^a, \varepsilon_{AB}, \Gamma_{aA}^B, \nabla_{A\dot{X}} := \sigma_{A\dot{X}}^a \nabla_a, \quad \begin{matrix} A \in \{1, 2\} \\ \dot{X} \in \{\dot{1}, \dot{2}\}. \end{matrix} \quad (2.1)$$

If we define for  $\varphi \in \mathcal{S}, \psi \in \bar{\mathcal{S}}$

$$(\mathcal{M}\varphi)_{\dot{X}} := \nabla_{\dot{X}}^A \varphi_A, \quad (\mathcal{N}\psi)_A := \nabla_A^{\dot{X}} \psi_{\dot{X}}, \quad (2.2)$$

we have [W4; PR]

$$-2(\mathcal{N}\mathcal{M}\varphi)_A = g^{ab} \nabla_a \nabla_b \varphi_A - \frac{1}{4} R \varphi_A =: (\mathcal{L}^{(1/2)} \varphi)_A. \quad (2.3)$$

In the following we consider the conformally invariant wave equation

$$\mathcal{L}^{(0)} u \equiv g^{ab} \nabla_a \nabla_b u - \frac{1}{6} R u = 0, \quad u \in \mathcal{J} \quad E_1$$

the (source-free) Maxwell equations

$$du = O, \quad \delta u = O, \quad u \in \Lambda^2 \quad E_2$$

and Weyl's neutrino equation

$$\mathcal{M}u = O, \quad u \in \mathcal{S}. \quad E_3$$

Let  $M$  be a causal domain  $[F; G4]$  and  $\Gamma(x, y)$  the square of the geodesic distance of  $x, y \in M$ . For any fixed  $y \in M$  the set  $\{x \in M \mid \Gamma(x, y) > 0\}$  decomposes naturally into two open subsets of  $M$ ; one of them is called the future  $D_+(y)$  and the other one the past  $D_-(y)$  of  $y$ . The characteristic semi null cones  $C_\pm(y)$  are defined as the boundary sets of  $D_\pm(y)$ , respectively.

Let  $G_\pm^{(0)}(y)$ ,  $G_\pm^{(1)}(y)$  and  $G_\pm^{(1/2)}(y)$  be the fundamental solutions of the linear operators  $\mathcal{L}^{(0)}$ ,  $\Delta$ ,  $\mathcal{L}^{(1/2)}$  and  $T^{(\alpha)}(\cdot, y)$ ,  $\alpha = 0, 1, 1/2$  the tail terms of  $G_\pm^{(\alpha)}(y)$  with respect to  $y$ , respectively. The tail term is just the factor of the regular part of the corresponding fundamental solution, which is a distribution supported inside the future of  $y$   $[F; G4]$ . For  $T^{(\alpha)}$  there is an asymptotic expansion in  $\Gamma$

$$T^{(\alpha)} \sim \sum_{k=0}^{\infty} \frac{1}{2^k \cdot k!} U_{(k+1)}^{(\alpha)} \Gamma^k, \quad (2.4)$$

where the Hadamard coefficients  $U_{(k)}^{(\alpha)}$  are determined recursively by the transport equations [F, G4, W4].

### 3. HUYGENS' PRINCIPLE

Let  $F$  be a space-like 3 dimensional submanifold of  $M$ ,  $D_F(y)$  that part of the interior of the past semi null cone  $C_-(y)$ , which is bounded by the hypersurface  $F$  and, finally,  $F(y) := F \cap D_F(y)$ . Cauchy's problem for one of the equations  $E_\sigma$  is the problem of determining a solution which assumes given values  $u$  (and their normal derivative for  $E_1$ ) on the given submanifold  $F$ . These values are called Cauchy data. Local existence and uniqueness of the solution of Cauchy's problem has been proved by Hadamard [Ha] for  $E_1$ , by Günther [G2] for  $E_2$  and by the author [W4] for  $E_3$ .

**DEFINITION 3.1.** – *One of the equations  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$ , is said to satisfy Huygens' principle (in the sense of Hadamard's "minor premise") if and only if for every Cauchy problem and for every  $y \in M$  the solution*

$u$  depends only on the Cauchy data in an arbitrarily small neighbourhood of  $F(y)$ .

Only if Huygens' principle is valid the wave propagation is free of tails [F; G4; McL2; W4], that is the solution depends on the sources distributions on the past null cone of the field point only and not on the sources inside the cone. Huygens' principle plays an important role also in quantum field theory in curved space-times. According to Lichnerowicz [L; B] the support of the commutator – or the anticommutator – distribution, respectively, lies on the null cone if and only if Huygens' principle holds for the corresponding field equation.

In [Ha; F; G2; W4] it was proved:

PROPOSITION 3.1. – *The equation  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$  satisfies Huygens' principle iff*

$$\forall x, y \in M : T^{(0)}(x, y) = O \quad \text{in the case } \sigma = 1 \quad (3.1)$$

$$\forall x, y \in M : K(x, y) : d^{(1)} d^{(2)} T^{(1)}(x, y) = O \text{ in the case } \sigma = 2 \quad (3.2)$$

$$\forall x, y \in M : N(x, y) := \mathcal{M}^{(1)} T^{(1/2)}(x, y) = O \text{ in the case } \sigma = 3. \quad (3.3)$$

Here the superscripts <sup>(1)</sup>, <sup>(2)</sup> indicate whether the derivative is meant with respect to  $x$  or  $y$ .

DEFINITION 3.2. – *The terms*

$$T^{(0)}(x, y), \quad K(x, y), \quad N(x, y) \quad (3.4)$$

are called tail terms of the equation  $E_1, E_2, E_3$ , respectively.

If the metric  $g$  undergoes a conformal transformation

$$\tilde{g}_{ab} = e^{2\Phi} g_{ab}, \quad \phi \in C^\infty(M), \quad (3.5)$$

the tail terms (3.4) transform according to [GW1; McL2; W4, 5]

$$\tilde{T}^{(0)}(x, y) = e^{-[\phi(x)+\phi(y)]} T^{(0)}(x, y) \quad (3.6)$$

$$\tilde{K}(x, y) = K(x, y) \quad (3.7)$$

$$\tilde{N}(x, y) = e^{-3/2[\phi(x)+\phi(y)]} N(x, y). \quad (3.8)$$

Consequently, a conformal transformation (3.5) preserve the Huygens' character of the equation  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$  [G4; W4; Ø]. In particular,

the conditions (3.1)-(3.3) are fulfilled for flat metrics [G4; W4], which implies that if  $g$  is conformally flat for the equations  $E_\sigma$  Huygens' principle is (trivially) valid.

Because the functional relationship between the tail terms (3.4) and the metric is not clear, the problem of determination of all metrics, for which any equation  $E_\sigma$  satisfies Huygens' principle, is not yet completely solved (see [G4; W2, 4; McL3; CM; I]). A step forward is the derivation of suitable infinite sequences of conformally invariant tensors, the so called moments of  $E_\sigma$ .

#### 4. CONFORMALLY INVARIANT TENSORS

We consider polynomial tensors, i.e. tensors, whose coordinates are polynomial in  $g^{ab}$  and the partial derivatives of  $g_{ab}$ . These tensors are just the elements of the tensor algebra  $\mathcal{R}_0$  generated by the tensors.

$$g^{ab}, g_{ab}, \nabla_{(i_1} \cdots \nabla_{i_r} R_{i_{r+1}|ab|i_{r+2}}), \quad r = 0, 1, 2, \dots \quad (4.1)$$

by means of the usual tensor operations. Furthermore, let  $\mathcal{R}$  be the algebra generated by the tensors (4.1) and the Levi-Civita tensor  $e_{abcd}$ .<sup>1)</sup>

DEFINITION 4.1. – A tensor  $T(g) \in \mathcal{R}$  is said to be conformally invariant of weight  $\omega$ , if under the conformal transformation (3.5)  $T(g)$  has the transformation law

$$T(\tilde{g}) = e^{2\omega\phi} T(g). \quad (4.2)$$

It is an important problem to give a survey of all conformally invariant tensors or to give methods for constructing special classes of such tensors [GW2, 3; W3].

LEMMA 4.1. –  $T(g) \in \mathcal{R}$  is conformally invariant iff it is invariant under all infinitesimal conformal transformations, i.e. iff

$$P(g, \Phi) := \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [T(e^{2\varepsilon\Phi} g) - e^{2\varepsilon\omega\Phi} T(g)] = O. \quad (4.3)$$

Let  $\tau$  be the subalgebra of those elements of  $\mathcal{R}$  which contain only *first* derivatives of  $\Phi$  in their transformation law [GW 2]. From Lemma 4.1 it follows that  $T(g) \in \tau$  iff  $P(g, \Phi)$  has the form

$$P(g, \Phi) = X^k(T) \nabla_k \Phi. \quad (4.4)$$

For the linear operator  $X^k$ , defined on  $\tau$  by (4.4), holds Leibniz's rule [GW2].

COROLLARY 4.1. -  $T(g) \in \mathcal{R}$  is conformally invariant iff  $T(g) \in \tau$  and  $X^k(T) = 0$ .

Examples. -  $X^k(g_{ab}) = 0$ ,  $X^k(C_{abcd}) = 0$ ,  $X^k(\nabla_u C^u{}_{.abc}) = C^k{}_{.abc}$ . If  $T(g) \in \tau$  then in general we have  $\nabla_a T \notin \tau$ . Let be

$$L_{ab} := -R_{ab} + \frac{1}{6} Rg_{ab}, \quad *C_{abd} := \frac{1}{2} e_{abkl} C^k{}_{.cd}.$$

DEFINITION 4.2. - For  $T \in \tau$  the tensor

$$\overset{c}{\nabla}_a T := \nabla_a T - \frac{1}{2} X^k(T) L_{ka} \tag{4.5}$$

is called the conformal covariant derivative of  $T$  [GW2].

In [GW 2; GeW 1] it was proved:

PROPOSITION 4.1. - (i) The conformal covariant derivative  $\overset{c}{\nabla}_a$  is linear, obeys Leibniz's rule and commutes with contractions.

(ii)  $\overset{c}{\nabla}_a : \tau \rightarrow \tau$

(iii)  $\tau$  is generated by the tensors

$$g^{ab}, g_{ab}, e_{abcd}, \overset{c}{\nabla}_{(i_1} \dots \overset{c}{\nabla}_{i_r} C_{i_{r+1}|ab|i_{r+2})}, \quad r = 0, 1, 2, \dots \tag{4.6}$$

(iv) If  $T \in \tau$  has the weight  $\omega$ , then

$$X^k(\overset{c}{\nabla}_a T) - \overset{c}{\nabla}_a(X^k(T)) = 2\omega \delta_a^k T + \Delta_a^k T, \tag{4.7}$$

where

$$\begin{aligned} \Delta_a^k(T_{ij\dots lm} \dots) &:= \Delta_{as}^{kl} T_{ij\dots sm\dots} + \Delta_{as}^{km} T_{ij\dots ls\dots} + \dots \\ &- \Delta_{ai}^{ks} T_{sj\dots lm\dots} - \Delta_{aj}^{ks} T_{is\dots lm\dots} - \dots \end{aligned} \tag{4.8}$$

and

$$\Delta_{ai}^{ks} := \delta_a^k \delta_i^s + \delta_i^k \delta_a^s - g_{ai} g^{ks}.$$

Examples. - (i)  $\overset{c}{\nabla}_k C_{abcd} = \nabla_k C_{abcd}$ ,  $\overset{c}{\nabla}_k *C_{abcd} = \nabla_k *C_{abcd}$

(ii)  $B_{i_1, i_2} := \nabla_a \nabla_b C^a{}_{.i_1 i_2}{}^b + \frac{1}{2} C^a{}_{.i_1 i_2}{}^b R_{ab} = \overset{c}{\nabla}_a \overset{c}{\nabla}_b C^a{}_{.i_1 i_2}{}^b$  (Bach tensor)  $X^k(B_{i_1 i_2}) = 0$ . Therefore, the Bach tensor is a conformally invariant tensor of weight  $-1$ .

Whenever in the following latin indices with subindices arise (e. g.  $i_1 \dots i_r$ ) we assume that symmetrization has been carried out over the indices. If  $T$  is any tensor with covariant rank  $r$  ( $r \geq 2$ ), then we denote by  $TS(T)$  the trace-free part of  $T$ . For a symmetric tensor  $T_{i_1 \dots i_r}$  with  $r \geq 2$  we write

$$T_{i_1 \dots i_r} \equiv O \pmod{g} \quad \text{iff} \quad TS(T_{i_1 \dots i_r}) = O.$$

LEMMA 4.2. — *If  $T_{i_1 \dots i_{k-l}}$  is a symmetric, conformally invariant tensor with covariant rank  $(k-l)$  and weight  $\omega$  then*

$$\begin{aligned} & X^\gamma (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} T_{i_{l+1} \dots i_k}) \\ & \equiv l(2\omega + l - 2k + 1) \delta_{i_1}^\gamma \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_l} T_{i_{l+1} \dots i_k} \pmod{g} \end{aligned} \quad (4.9)$$

*Proof.* — By (4.7) one gets (see [G4], p. 510)

$$\begin{aligned} & X^\gamma (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} T_{i_{l+1} \dots i_k}) \\ & \equiv l(2\omega - l + 1) \delta_{i_1}^\gamma \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_l} T_{i_{l+1} \dots i_k} \\ & \quad + l \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_{l-1}} \Delta_{i_l}^\gamma (T_{i_{l+1} \dots i_k}) \pmod{g}. \end{aligned}$$

On account of (4.8) it follows  $\Delta_{i_1 i_2}^{\gamma r} \equiv 2 \delta_{i_1}^\gamma \delta_{i_2}^r \pmod{g}$  and, therefore,

$$\Delta_{i_l}^\gamma (T_{i_{l+1} \dots i_k}) = -2(k-l) \delta_{i_l}^\gamma T_{i_{l+1} \dots i_k}$$

and the assertion (4.9).

A conformally invariant tensor  $T$  is called trivial if  $T$  is generated by

$$\{g^{ab}, g_{ab}, e_{abcd}, C_{abcd}\}.$$

Lemma 4.1, Corollary 4.1, Definition 4.2 and Proposition 4.1 are very useful for the construction of nontrivial conformally invariant tensors.

Let  $\mathcal{S}_{2r}$ ,  $\mathcal{S}_{2r+1}$  ( $r = 0, 1, 2, \dots$ ) be the set of all symmetric, trace-free, conformally invariant tensors contained in  $\mathcal{R}_0$  and  $\mathcal{R}$ , respectively, with weight  $-1$  and covariant rank  $2r$  and  $2r+1$ , respectively. Using the above results, in [GW3; GeW1, 2] linear independent generating systems of  $\mathcal{S}_r$  for  $0 \leq r \leq 6$  were derived.<sup>2)</sup>

PROPOSITION 4.2. - *If  $S_r = \{S_{i_1 \dots i_r}\} \in \mathcal{S}_r$  then one has*

$$S_r = 0 \quad \text{for } r = 0, 1, 3, \quad S_2 = \alpha B$$

$$S_4 = \sum_{m=1}^3 \beta_m W^{(m)}, \quad S_5 = \gamma^{(2)} S_5^{(2)} + \sum_{k=1}^4 \gamma_k^{(3)} S_5^{(3,k)}$$

$$S_6 = \sum_{k=1}^2 \delta_k^{(2)} S_6^{(2,k)} + \sum_{l=1}^6 \delta_l^{(3)} S_6^{(3,l)},$$

where  $B$  is the Bach tensor,

$$W_{i_1 \dots i_4}^{(1)} := TS [\nabla^a C^b_{i_1 i_2}{}^c \nabla_a C_{b i_3 i_4 c} + 16 \nabla_u C^u_{i_1 i_2 a} \nabla_k C^k_{i_3 i_4}{}^a + 4 C^a_{i_1 i_2}{}^b \{2 \nabla_a \nabla_u C^u_{i_3 i_4 b} - C_{a i_3 i_4}{}^c L_{bc}\}],$$

$$W_{i_1 \dots i_4}^{(2)} := TS [2 \nabla_{i_1} C^b_{i_2 i_3}{}^c \nabla_u C^u_{ab i_4} + 2 \nabla_u C^u_{i_1 i_2 a} \nabla_k C^k_{i_3 i_4}{}^a - C^a_{i_1 i_2}{}^b \{2 \nabla_{i_3} \nabla_u C^u_{ab i_4} - C^c_{ab i_3} L_{c i_4}\}],$$

$$W_{i_1 \dots i_4}^{(3)} := TS [C^{abcd} C_{a i_1 i_2 d} C_{b i_3 i_4 c}],$$

$$S_{i_1 \dots i_5}^{(2)} := TS [\nabla^c C^a_{i_1 i_2}{}^b \nabla_c C_{a i_3 i_4 b} + 4 C^a_{i_1 i_2}{}^b \nabla_{i_3} \nabla_a \nabla_u C^u_{i_4 i_5 b} - 6 \nabla_{i_1}^* C^a_{i_2 i_3}{}^b \nabla_a \nabla_u C^u_{i_4 i_5 b} + 26 \nabla_u^* C^u_{i_1 i_2}{}^a \nabla_{i_3} \nabla_k C^k_{i_4 i_5 a} - 5 C^a_{i_1 i_2}{}^b \nabla_{i_3} C^c_{i_4 i_5 b} L_{ac} - 4 C^a_{i_1 i_2}{}^b \nabla_a C^c_{i_3 i_4 b} L_{i_5 c} - 4 C^a_{i_1 i_2}{}^b \nabla_u C^u_{ab i_3} L_{i_4 i_5} + 21 C^a_{i_1 i_2}{}^b \nabla_u C^u_{i_3 i_4 a} L_{i_5 b}],$$

$$S_{i_1 \dots i_5}^{(3,1)} := TS [12 C^a_{i_1 i_2}{}^b C_{a i_3 i_4}{}^c \nabla_u C^u_{bc i_5} + C_{a \dots i_1}{}^{cd} C_{b c d i_2} \nabla_{i_3} C^a_{i_4 i_5}{}^b],$$

$$S_{i_1 \dots i_5}^{(3,2)} := TS [8 C^a_{i_1 i_2}{}^b C_{a i_3 i_4}{}^c \nabla_u C^u_{bc i_5} - C_{a \dots i_1}{}^{cd} C_{b i_2 c d} \nabla_{i_3} C^a_{i_4 i_5}{}^b],$$

$$S_{i_1 \dots i_5}^{(3,3)} := TS [12 C^a_{i_1 i_2}{}^b C_{a i_3 i_4}{}^c \nabla_u C^u_{bc i_5} + C_{a \dots i_1}{}^{cd} C_{b c d i_2} \nabla_{i_3} C^a_{i_4 i_5}{}^b],$$

$$S_{i_1 \dots i_5}^{(3,4)} := TS [8 C^a_{i_1 i_2}{}^b C_{a i_3 i_4}{}^c \nabla_u C^u_{bc i_5} - C_{a \dots i_1}{}^{cd} C_{b i_2 c d} \nabla_{i_3} C^a_{i_4 i_5}{}^b]$$

and  $\alpha, \beta_m, \gamma^{(2)}, \gamma_k^{(3)}, \delta_k^{(2)}, \delta_l^{(3)} \in \mathbb{R}$ .

For the definition of the tensors  $S_6^{(2,k)}$ ,  $S_6^{(3,l)}$ , which contain many monomials see [GeW 2, pp. 127-136].<sup>3)</sup>

*Remark 4.1.* – In the case of  $R_{ab} = 0$  the tensors  $B$ ,  $W^{(2)}$  and  $S_6^{(2,2)}$  vanish. Consequently, if  $(M, g)$  is conformally related to an Einstein space-time with  $R_{ab} = 0$  then [W8]

$$B = 0, W^{(2)} = 0, S_6^{(2,2)} = 0.$$

*Remark 4.2.* – McLenaghan and Leroy investigate in [McL, L] the class of generalized plane wave metrics

$$ds^2 = 2 dx^1 [dx^2 + \{a(z + \bar{z})x^2 + Dz^2 + \bar{D}\bar{z}^2 + ez\bar{z} + Fz + \bar{F}\bar{z}\} dx^1 - 2 [dz + az^2 dx^1] [d\bar{z} + a\bar{z}^2 dx^1], \quad (4.10)$$

where  $z = x^3 + ix^4$  and  $a = \bar{a}$ ,  $D, e, F$  are arbitrary functions of  $x^1$  only. For the special case  $a = 0$  we obtain the important subclass of plane wave metrics [S2, G3, 4; DC; W4]. In [AW; Ge W2] it was shown:

LEMMA 4.3. – *For a metric (4.10) it holds*

$$B = 0; W^{(m)} = 0, S_5^{(2)} = 0, S_5^{(3,k)} = 0; S_6^{(3,l)} = 0 \\ (m = 1, 2, 3; k = 1, \dots, 4; l = 1, \dots, 6)$$

$$S_6^{(2,1)} = 11 S_6^{(2,2)} = 363 Z,$$

where

$$Z_{i_1 \dots i_6} := TS [C_{i_1 i_2}^a \cdot C_{ai_3 i_4 b} \cdot R_{i_5} \cdot R_{i_6 c}].$$

LEMMA 4.4. – *A generalized plane wave metric is a plane wave metric iff  $Z = 0$ .*

Consequently, in virtue of Remark 4.1 a generalized plane wave metric  $g$  is conformally related to an Einstein metric iff  $g$  is a plane wave metric.

It is well known that the equations  $E_1 - E_3$  are conformally invariant [PR; W4]. In particular, for Weyl's neutrino operator under the conformal transformation [PR; W4]

$$\tilde{g}_{ab} = e^{2\Phi} g_{ab}, \tilde{\sigma}_a^{A\dot{X}} = e^\Phi \cdot \sigma_a^{A\dot{X}}, \tilde{\varepsilon}_{AB} = \varepsilon_{AB}$$

we get

$$\tilde{\mathcal{M}} \tilde{\varphi} = e^{-(5/2)\Phi} \mathcal{M} \varphi,$$

where  $\tilde{\varphi} := e^{-(3/2)} \varphi$ . In Section 5 we need a conformal covariant derivative of a conformal invariant spinor  $S_{\dot{X}}$  with the weight  $\omega$ , i.e. a spinor with the transformation law

$$\tilde{S}_{\dot{X}} = e^{2\omega\phi} S_{\dot{X}}, \quad \omega \in \mathbb{R}.$$

Similarly to (4.5) one can define inductively for  $r > 0$  :

$$\begin{aligned} \overset{c}{\nabla}_{i_1} S_{\dot{X}} &= \nabla_{i_1} S_{\dot{X}}, & \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}} \\ &= \nabla_{i_1} (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}}) - \frac{1}{2} X^k (\overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}}) L_{ki_1}. \end{aligned} \tag{4.11}$$

Here  $X^k (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}})$  is again given by (see (4.3), (4.4))

$$\begin{aligned} \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} \tilde{S}_{\dot{X}} &= e^{2\omega\phi} [\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}} \\ &+ X^k (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}}) \nabla_k \phi + \dots], \end{aligned}$$

where we omit terms which are nonlinear in the derivatives of  $\phi$ . The transformation law of  $\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r} S_{\dot{X}}$  contains only first derivatives of  $\phi$ .

LEMMA 4.5. – A conformally invariant spinor  $S_{\dot{X}}$  with weight  $\omega$  satisfies

$$\begin{aligned} X^\gamma (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_k} S_{\dot{X}}) &\equiv k(2\omega - k + 1) \delta_{i_1}^\gamma \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_k} S_{\dot{X}} \\ &- k \sigma_{[i_1}^{Y\dot{Z}} \sigma_{k] \dot{X} Y} g^{\gamma k} \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_k} S_{\dot{Z}} \pmod{g}. \end{aligned}$$

*Proof.* – Induction with respect to  $k$ : by  $X^\gamma (S_{\dot{X}}) = 0$  the formula (4.11) is correct if  $k = 0$ . If (4.11) is fulfilled for any  $k$ , then on account of (see [W 4])

$$\nabla_a S_{\dot{X}} = \partial_a S_{\dot{X}} - \bar{\Gamma}_{a\dot{X}}^{\dot{Y}} S_{\dot{Y}}, \quad \bar{\Gamma}_{aA}^B = \Gamma_{aA}^B + g^{bc} \sigma_{[a}^{B\dot{Y}} \sigma_{b] A\dot{Y}} \nabla_c \phi$$

we have analogues to (4.7)

$$\begin{aligned} X^\gamma (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_{k+1}} S_{\dot{X}}) &\equiv \overset{c}{\nabla}_{i_1} X^\gamma (\overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_{k+1}} S_{\dot{X}}) \\ &+ 2(\omega - k) \delta_{i_1}^\gamma \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_{k+1}} S_{\dot{X}} \\ &- \sigma_{[i_1}^{A\dot{Y}} \sigma_{m] A\dot{X}} g^{\gamma m} \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_{k+1}} S_{\dot{Z}} \pmod{g}. \end{aligned}$$

In order to calculate  $\overset{c}{\nabla}_{i_1} X^\gamma (\overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_{k+1}} S_{\dot{X}})$  we use the induction hypothesis and obtain under consideration of  $\overset{c}{\nabla}_a g_{bd} = 0$ ,  $\overset{c}{\nabla}_a \sigma_b^{A\dot{X}} = O$  the assertion.

## 5. MOMENTS OF THE FIELD EQUATIONS

### 1. The moments of the scalar wave equation

From (2.4) and the transport equations for the Hadamard coefficients it follows, that the coincidence values  $\nabla_{i_1}^{(1)} \cdots \nabla_{i_r}^{(1)} T^{(0)}(x, x)$  are elements of  $\mathcal{R}_0$  [G4]. Here  $\nabla^{(1)}$  denotes the covariant derivative of  $T^{(0)}$  with respect to the first variables. Because of (3.6) and Proposition 4.1 the conformal covariant derivatives  $\overset{c}{\nabla}_{i_1}^{(1)} \cdots \overset{c}{\nabla}_{i_r}^{(1)} T^{(0)}(x, x)$ ,  $r = 0, 1, 2, \dots$  are contained in  $\tau$ . In [G4] it was defined and proved:

DEFINITION 5.1. – *The covariant  $r$ -tensors given recursively by*

$$I_{i_1 \dots i_r}^1(x) := TS \left[ \overset{c}{\nabla}_{i_1}^{(1)} \cdots \overset{c}{\nabla}_{i_r}^{(1)} T^{(0)}(x, x) - \sum_{l=1}^r d_{l,r}^1 \overset{c}{\nabla}_{i_1} \cdots \overset{c}{\nabla}_{i_l} I_{i_{l+1} \dots i_r}^1(x) \right]$$

with

$$d_{l,r}^1 := \binom{r}{l}^2 / \binom{2r-l+1}{l}$$

are called moments of the scalar wave equation  $E_1$  of order  $r$ .

PROPOSITION 5.1. – *It holds  $I_{i_1 \dots i_r}^1 \in \mathcal{S}_r$ ,  $r \geq 0$  and  $I_{i_1 \dots i_r}^1 \equiv 0$  if  $r$  odd.* <sup>4)</sup>

### 2. The moments of Maxwell's equations $E_2$

The tail term  $K(x, y)$  of Maxwell's equations  $E_2$  (see (3.2)) is a double differential form of degree 2:

$$K(x, y) = K_{ij, \alpha\beta}(x, y) dx^i \wedge dx^j dy^\alpha \wedge dy^\beta.$$

One defines for every  $x \in M$  and  $r \geq 2$  the covariant  $r$  tensors [G4; W2]

$$M_{i_1 \dots i_r}(x) := g^{j\beta}(x) \overset{c}{\nabla}_{i_1}^{(1)} \cdots \overset{c}{\nabla}_{i_{r-2}}^{(1)} K_{i_{r-1}j, i_r\beta}(x, x)$$

and for  $r$  even recursively

$$I'_{i_1 \dots i_r} := TS \left[ M_{i_1 \dots i_r} - \sum_{l=1}^{r-2} d_{l,r}^2 \overset{c}{\nabla}_{i_1} \cdots \overset{c}{\nabla}_{i_l} I'_{i_{l+1} \dots i_r} \right]$$

with

$$d_{l,r}^2 := \binom{r-2}{l} \binom{r}{l} / \binom{2r-l+1}{l}$$

On the other hand let be

$$\begin{aligned}
 {}^*K(x, y) &= {}^*K_{ij, \alpha\beta}(x, y) dx^i \wedge dx^j dy^\alpha \wedge dy^\beta \\
 &:= \frac{1}{2} e_{ijkl} K_{.. \alpha\beta}^{lk}(x, y) dx^i \wedge dx^j \wedge dy^\alpha \wedge dy^\beta.
 \end{aligned}$$

Then  ${}^*K(x, y)$  is the left dual of  $K(x, y)$  which is analogues to  $K(x, y)$  (see (3.7)) conformally invariant with weight zero. Now one defines for  $x \in M$  and  $r \geq 2$  [G4; W2]

$${}^*M_{i_1 \dots i_r} := g^{j\beta}(x) \overset{c}{\nabla}_{i_1}^{(1)} \dots \overset{c}{\nabla}_{i_{r-2}}^{(1)} {}^*K_{i_{r-1} j, i_r \beta}(x, x)$$

and for  $r$  odd recursively ♦

$$I''_{i_1 \dots i_r} := TS \left[ {}^*M_{i_1 \dots i_r} - \sum_{l=1}^{r-2} d_{l,r}^2 \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} I''_{i_{l+1} \dots i_r} \right] \tag{5.1}$$

DEFINITION 5.2. – *The covariant  $r$ -tensors given by*

$$I^2_{i_1 \dots i_r} := \begin{cases} I'_{i_1 \dots i_r} & \text{if } r \geq 2, \text{ even} \\ I''_{i_1 \dots i_r} & \text{if } r \geq 3, \text{ odd} \end{cases} \tag{5.2}$$

are called moments of Maxwell's equations of order  $r$ . <sup>5)</sup>

In [G4] the following was shown, using in particular (3.2):

PROPOSITION 5.2. – *The moments (5.2) are elements of  $\mathcal{S}_r$  ( $r \geq 2$ ).*

3. *The moments of Weyl's neutrino equation*

The tail term  $N(x, y)$  of the equation  $E_3$  (see (3.3)) is a spinor of  $\bar{\mathcal{S}}_{(x)} \otimes \mathcal{S}_{(y)}$  [W4]. We denote by  $N_{\dot{X}\underline{A}}(x, y)$  its coordinates. Here the underlined indices refer to  $y$ . In [W4, p. 69] it was proved:

LEMMA 5.1. – *It is  $N(x, y) = -N(y, x)$  for  $x, y \in M$ .*

Now we define for every  $x \in M$  and  $r \geq 1$  the covariant (complex)  $r$ -tensors [W4]

$$N_{i_1 \dots i_r}(x) := \sigma_{i_1}^{\underline{A}\dot{X}}(x) \overset{c}{\nabla}_{i_2}^{(1)} \dots \overset{c}{\nabla}_{i_r}^{(1)} N_{\dot{X}\underline{A}}(x, x) \tag{5.3}$$

and for  $r \geq 1$  recursively

$$I^*_{i_1 \dots i_r} := N_{i_1 \dots i_r} - \sum_{l=1}^{r-1} d_{l,r}^3 \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} I^*_{i_{l+1} \dots i_r} \tag{5.4}$$

with

$$d_{l,r}^3 := \binom{r-1}{l} \binom{r}{l} / \binom{2r-l+1}{l} \tag{5.5}$$

PROPOSITION 5.3. – *The tensor  $i^r I_{i_1 \dots i_r}^*$  is real for  $r \geq 1$ .*

*Proof.* – Let us write in this proof instead of  $I_{i_1 \dots i_r}^*$ ,  $N_{i_1 \dots i_r}$ ,  $\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_r}$ ,  $\overset{c}{\nabla}_{i_1}^{(1)} \dots \overset{c}{\nabla}_{i_r}^{(1)}$ ,  $\overset{c}{\nabla}_{i_1}^{(2)} \dots \overset{c}{\nabla}_{i_r}^{(2)}$  shortly  $I_r$ ,  $N_r$ ,  $\nabla_r$ ,  $\nabla_r^1$ ,  $\nabla_r^2$ , respectively.

Then it may be shown that for  $r \geq 1$

$$I_r + (-1)^{r+1} \bar{I}_r = 0. \tag{5.6}$$

Using the differentiation rule  $\overset{c}{\nabla} = \overset{c}{\nabla}^{(1)} + \overset{c}{\nabla}^{(2)}$  along the diagonal of  $M \times M$  and Lemma 5.1 we obtain

$$\begin{aligned} \nabla_l^1 N_{\dot{X}A}(x, x) &= -\nabla_l^2 \bar{N}_{\dot{X}A}(x, x) = (\nabla_l - \nabla_l^2) N_{\dot{X}A}(x, x) \\ &= \sum_{n=0}^l (-1)^{l-n} \binom{l}{n} \nabla_n \nabla_{l-n}^2 N_{\dot{X}A}(x, x) \\ &= \sum_{n=0}^l (-1)^{l-n+1} \binom{l}{n} \nabla_n \nabla_{l-n}^1 \bar{N}_{\dot{X}A}(x, x), \end{aligned}$$

hence

$$N_{l+1} = \sum_{n=0}^l (-1)^{l-n+1} \binom{l}{n} \nabla_n \bar{N}_{l-n+1}. \tag{5.7}$$

We prove (5.6) by induction with respect to  $r$ . By (5.7) we have for  $r = 1$ :

$$I_1 = N_1 = -\bar{N}_1 = -\bar{I}_1.$$

Now let (5.6) be fulfilled for  $1 \leq r \leq (k-1)$ . Then from (5.4) and (5.7) it follows for  $r = k$

$$\begin{aligned}
 \bar{I}_k + (-1)^{k+1} I_k &= \bar{N}_k + (-1)^{k+1} N_k \\
 &- \sum_{l=1}^{k-1} d_{l,k}^3 [\nabla_l \bar{I}_{k-l} + (-1)^{k+1} \nabla_l I_{k-l}] \\
 &= \sum_{n=1}^{k-1} (-1)^{k-n} \binom{k-1}{n} \nabla_n \left[ I_{k-n} + \sum_{l=1}^{k-n-1} d_{l,k-n}^3 \nabla_l I_{k-n-l} \right] \\
 &+ \sum_{n=1}^{k-1} d_{n,k}^3 (-1)^k (1 - (-1)^n) \nabla_n I_{k-n} \\
 &= \sum_{n=2}^{k-1} (-1)^k \nabla_n I_{k-n} \left[ \sum_{l=0}^n (-1)^l \binom{k-1}{l} d_{n-l,k-l}^3 - (-1)^n d_{n,k}^3 \right] \\
 &= \sum_{n=2}^{k-1} (-1)^k d_{n,k}^3 \nabla_n I_{k-n} [F(-n, n-2k-1, -k, 1) - (-1)^n] = O,
 \end{aligned}$$

where  $F$  is the hypergeometric polynomial at the point 1.

DEFINITION 5.3. – *The covariant real  $r$ -tensors given by*

$$I_{i_1 \dots i_r}^3 = \begin{cases} TS [I_{i_1 \dots i_r}^*] & \text{if } r \geq 2, \text{ even} \\ TS [i I_{i_1 \dots i_r}^*] & \text{if } r \geq 1, \text{ odd} \end{cases}$$

are called moments of Weyl's neutrino equation of order  $r$ .

LEMMA 5.2. – *If  $I_{i_1 \dots i_l}^3(x) = 0$  for every  $x \in M$  and  $1 \leq l \leq r-1$  then*

$$I_{i_1 \dots i_r}^3 = \begin{cases} I_{i_1 \dots i_r} & \text{if } r \geq 2, \text{ even} \\ i I_{i_1 \dots i_r} & \text{if } r \geq 1, \text{ odd,} \end{cases}$$

where

$$I_{i_1 \dots i_r}(x) := \sigma_{i_1}^{A\dot{X}}(x) \partial_{i_2}^{(1)} \dots \partial_{i_r}^{(1)} N_{\dot{X}A}(x, x).$$

*Proof.* – Induction with respect to  $r$ . Obviously, the assertion is true for  $r = 1$ . Now we assume, that the assertion is valid for  $r - 1$ . Then we have

$$\left. \begin{aligned}
 I_{i_1 \dots i_l}(x) &= \sigma_{i_1}^{A\dot{X}}(x) \partial_{i_2}^{(1)} \dots \partial_{i_l}^{(1)} N_{\dot{X}A}(x, x) = O, \\
 0 &\leq l \leq r - 1,
 \end{aligned} \right\} \tag{5.8}$$

from which it follows

$$\sigma_{i_1}^{A\dot{X}} \nabla_{i_2}^{(1)} \dots \nabla_{i_l}^{(1)} N_{\dot{X}A}(x, x) = O, \quad 0 \leq l \leq r - 1. \tag{5.9}$$

Lemma 4.5 and (4.11) imply that

$$\overset{c}{\nabla}_{i_1}^{(1)} \dots \overset{c}{\nabla}_{i_{r-1}}^{(1)} N_{\dot{X}} - \nabla_{i_1}^{(1)} \dots \nabla_{i_{r-1}}^{(1)} N_{\dot{X}}$$

can be represented as linear form in the symmetric covariant derivatives of  $N_{\dot{X}}$  up to the order  $r - 2$ . Consequently, on account of (5.8) and (5.9) we get

$$\overset{c}{\nabla}_{i_1}^{(1)} \dots \overset{c}{\nabla}_{i_{r-1}}^{(1)} N_{\dot{X}A}(x, x) = \partial_{i_1}^{(1)} \dots \partial_{i_{r-1}}^{(1)} N_{\dot{X}A}(x, x)$$

(see also [G4], Lemma 4.10), hence

$$I_{i_1 \dots i_r} \equiv \sigma_{i_1}^{A\dot{X}} \partial_{i_2}^{(1)} \dots \partial_{i_r}^{(1)} N_{\dot{X}A} \pmod{g}. \tag{5.10}$$

In [W4, p. 154] it was proved, that (5.10) is symmetric and trace-free. Hence, the assertion is valid also for  $r$ .

PROPOSITION 5.4. – *The moments  $I_{i_1 \dots i_r}^3$  are elements of  $\mathcal{S}_r$ ,  $r \geq 1$ .*

*Proof.* – The spinor equivalents of the moments are polynomial with respect to the covariant derivatives of the curvature spinors with real coefficients [W4].

Therefore

$$I_{i_1 \dots i_{2r}}^3 \in \mathcal{R}_0, \quad I_{i_1 \dots i_{2r+1}}^3 \in \mathcal{R}, \quad r = 0, 1, 2, \dots, \tag{5.11}$$

where the moments contain the Levi-Civita tensor linearly if  $r$  odd (see [W4]). Now we have to show, that the moments are conformally invariant of weight  $(-1)$ . Obviously, it is sufficient to prove this for the tensors (5.4).

Induction with respect to  $r$ . From (5.3), (5.4), (3.8) and  $\tilde{\sigma}_a^{A\dot{X}} = e^\Phi \sigma_a^{A\dot{X}}$  it follows for  $r = 1$ :

$$\tilde{I}_{i_1}^* (x) = \tilde{N}_{i_1}(x, x) = e^{-2\Phi(x)} N_{i_1}(x, x) = e^{-2\phi(x)} I_{i_1}^* (x).$$

Lemma 4.2 implies for  $l \geq 1$

$$\begin{aligned} X^\gamma (\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} I_{i_{l+1} \dots i_r}^*) \\ \equiv -l(2r - l + 1) \delta_{i_1}^\gamma \overset{c}{\nabla}_{i_2} \dots \overset{c}{\nabla}_{i_l} I_{i_{l+1} \dots i_r}^* \pmod{g}. \end{aligned} \tag{5.12}$$

Further, we get from (5.3), Lemma 4.5, (3.8) and

$$\sigma_{i_1}^{A\dot{X}} \sigma_{[i_2}^{Y\dot{Z}} \sigma_{k]Y\dot{X}} \equiv \frac{1}{2} \sigma_{i_1}^{A\dot{Z}} g_{ki_2} \pmod{g}$$

(see [Sch]) the formula

$$\begin{aligned}
 X^\gamma(N_{i_1 \dots i_r}) &\equiv \sigma_{i_1}^{A\dot{X}} X^\gamma(\overset{c}{\nabla}_{i_2}^{(1)} \dots \overset{c}{\nabla}_{i_r}^{(1)} N_{\dot{X}A}) \\
 &\equiv \sigma_{i_1}^{A\dot{X}} \left[ (r-1) \left( \frac{1}{2} - r \right) \delta_{i_2}^\gamma \overset{c}{\nabla}_{i_3}^{(1)} \dots \overset{c}{\nabla}_{i_r}^{(1)} N_{\dot{X}A} \right. \\
 &\quad \left. + (1-r) \sigma_{[i_2}^{YZ} \sigma_{k]\dot{X}Y} g^{\gamma k} \overset{c}{\nabla}_{i_3}^{(1)} \dots \overset{c}{\nabla}_{i_r}^{(1)} N_{\dot{Z}A} \right] \\
 &\equiv r(1-r) \delta_{i_1}^\gamma N_{i_2 \dots i_r} \pmod{g}.
 \end{aligned}
 \tag{5.13}$$

Finally, from (5.4), (5.5), (5.12), (5.13) it follows

$$\begin{aligned}
 X^\gamma(I_{i_1 \dots i_r}^*) &= X^\gamma(N_{i_1 \dots i_r}) - \sum_{l=1}^{r-1} d_{l,r}^3 X^\gamma(\overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} I_{i_{l+1} \dots i_r}^*) \\
 &\equiv -r(r-1) \delta_{i_1}^\gamma N_{i_2 \dots i_r} + \sum_{l=1}^{r-1} d_{l,r}^3 l(2r-l+1) \delta_{i_r}^\gamma \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_{l-1}} I_{i_l \dots i_{r-1}}^* \\
 &\equiv r(1-r) \delta_{i_1}^\gamma \left[ I_{i_1 \dots i_{r-1}}^* + \sum_{l=1}^{r-2} d_{l,r-1}^3 \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} I_{i_{l+1} \dots i_{r-1}}^* \right] \\
 &\quad + \sum_{l=1}^{r-1} d_{l,r}^3 l(2r-l+1) d_{i_r}^\gamma \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_{l-1}} I_{i_l \dots i_{r-1}}^* \\
 &\equiv \delta_{i_r}^\gamma \sum_{l=1}^{r-2} [r(1-r) d_{l,r-1}^3 + (l+1)(2r-l) d_{l+1,r}^3] \overset{c}{\nabla}_{i_1} \dots \overset{c}{\nabla}_{i_l} I_{i_{l+1} \dots i_{r-1}}^* \\
 &\equiv O \pmod{g}.
 \end{aligned}$$

Consequently  $I_{i_1 \dots i_r}^3$  is conformally invariant. The proof of the weight is easily done.

We summarize the results of this Section:

**THEOREM 5.1.** – *Every field equation  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$  implies an infinite sequence  $\{I_{i_1 \dots i_r}^\sigma\}$  of symmetric, trace-free, conformally invariant  $r$ -tensors of weight  $(-1)$ .*

4. The first moments

The propositions 5.1, 5.2, 5.4 and 4.2 imply

**PROPOSITION 5.5.** – *There are real coefficients  $\alpha^{(\sigma)}$ ,  $\beta_k^{(\sigma)}$ ,  $\gamma^{(2,\sigma)}$ ,  $\gamma_l^{(3,\sigma)}$ ,  $\delta_m^{(2,\sigma)}$ ,  $\delta_p^{(3,\sigma)}$  such that for the moments  $I_{i_1 \dots i_r}^\sigma$  it holds*

$$\begin{aligned}
 I^1 &= O, \quad I_{i_1}^1 = I_{i_1}^3 = O, \quad I_{i_1 i_2}^\sigma = \alpha^{(\sigma)} B_{i_1 i_2}, \quad I_{i_1 i_2 i_3}^\sigma = O, \\
 I_{i_1 \dots i_4}^\sigma &= \sum_{k=1}^3 \beta_k^{(\sigma)} W_{i_1 \dots i_4}^{(k)}, \quad I_{i_1 \dots i_5}^\sigma = \gamma^{(2, \sigma)} S_{i_1 \dots i_5}^{(2)} + \sum_{l=1}^2 \gamma_l^{(3, \sigma)} S_{i_1 \dots i_5}^{(3, l)}, \\
 I_{i_1 \dots i_6}^\sigma &= \sum_{m=1}^2 \delta_m^{(2, \sigma)} S_{i_1 \dots i_6}^{(2, m)} + \sum_{p=1}^6 \delta_p^{(3, \sigma)} S_{i_1 \dots i_6}^{(3, p)} \quad (\sigma = 1, 2, 3). \quad (5.14)
 \end{aligned}$$

**6. HUYGENS’ PRINCIPLE AND MOMENT EQUATIONS**

The following theorem shows the importance of the moments for the validity of Huygens’ principle:

**THEOREM 6.1.** – *Let  $(M, g)$  be an analytic space-time. The field equation  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$  satisfies Huygens’ principle if and only if all the corresponding moments  $I_{i_1 \dots i_r}^\sigma$  of order  $r$  vanish on  $M$ .*

*Proof.* – For the cases  $\sigma = 1, 2$  see [G4, W2]. If for  $E_3$  Huygens’ principle is valid, then because of (3.3) we have for all  $x \in M$  and  $r \geq 0$

$$\partial_{i_1}^{(1)} \dots \partial_{i_r}^{(1)} N_{\dot{X}A}(x, x) = O, \quad (6.1)$$

hence, by Lemma 5.2

$$I_{i_1 \dots i_r}^3 = 0, \quad r \geq 1 \quad (6.2)$$

Conversely, from (6.2) and Lemma 5.2 it follows (6.1). Since  $N_{\dot{X}A}(x, y)$  is analytic the spinor  $N_{\dot{X}A}(x, y)$  can be determined with the help of the Taylor expansion of  $N_{\dot{X}A}(x, y)$  at the point  $y = x$ , which by (6.1) has only zero coefficients. Hence,  $N(x, y) = 0$  and  $E_3$  satisfies Huygen’s principle.

**DEFINITION 6.1.** – *The equations*

$$I_{i_1 \dots i_r}^\sigma = 0, \quad \sigma \in \{1, 2, 3\} \quad (ME)_r^\sigma$$

*are called moment equations for  $E_\sigma$  of order  $r$ .*

During the last forty years in the investigations of Huygens’ principle for  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$ , the first moments were determined, using the transport equations for Hadamard’s coefficients, suitable test metrics and often normal coordinates [A; G1, 2, 4; GW1; McL1, 2; S3; W1, 2, 4]. By complicate

calculations one obtained in this way more detailed information about the coefficients of Proposition 5.5.

LEMMA 6.1. – *It holds*  $\alpha^{(1)} = -1/60$  [G1],  $\alpha^{(2)} = 1/20$  [G2],  $\alpha^{(3)} = -1/80$  [W4],

$$\beta_1^{(1)} = -1/2^3 \cdot 5 \cdot 7, \quad \beta_2^{(1)} = 1/2 \cdot 3 \cdot 5 \cdot 7, \quad \beta_3^{(1)} = O \quad [\text{W1}],$$

$$\beta_1^{(2)} = 1/2^3 \cdot 3 \cdot 7, \quad \beta_2^{(2)} = -2/3 \cdot 5 \cdot 7, \quad \beta_3^{(2)} = O \quad [\text{W2}],$$

$$\beta_1^{(3)} = -1/2^2 \cdot 3 \cdot 5 \cdot 7, \quad \beta_2^{(3)} = 13/2^5 \cdot 3 \cdot 5 \cdot 7, \quad \beta_3^{(3)} = O \quad [\text{W4}],$$

$$\gamma^{(2,\sigma)} \neq 0 \text{ for } \sigma = 2, 3 [\text{AW}, \text{A}]$$

and [RW; GeW2]

$$I_{i_1 \dots i_6}^1 = 1/2^3 \cdot 3^2 \cdot 5 \cdot 7 \left[ \frac{1}{33} S_{i_1 \dots i_6}^{(2,1)} + \frac{7}{66} S_{i_1 \dots i_6}^{(2,2)} + \frac{5}{2} S_{i_1 \dots i_6}^{(3,1)} + 2 S_{i_1 \dots i_6}^{(3,2)} - \frac{5}{2} S_{i_1 \dots i_6}^{(3,3)} - 5 S_{i_1 \dots i_6}^{(3,4)} \right].$$

COROLLARY 6.1. – *If the equation*  $E_\sigma$ ,  $\sigma \in \{1, 2, 3\}$  *satisfies Huygens' principle, then, in particular, the following moment equations* <sup>7)</sup> *must hold*

$$B_{i_1 i_2} = O \quad (ME)_2^\sigma$$

$$W_{i_1 \dots i_4}^{(1)} - k_\sigma W_{i_1 \dots i_4}^{(2)} = O \quad (ME)_4^\sigma$$

$$I_{i_1 \dots i_5}^\sigma = O \quad (ME)_5^\sigma$$

$$I_{i_1 \dots i_6}^\sigma = O \quad (ME)_6^\sigma$$

where

$$k_1 = 4/3, \quad k_2 = 16/5, \quad k_3 = 13/8.$$

PROPOSITION 6.1. – (i) *If*  $g$  *is a plane wave metric, then for all*  $r$  *and*  $\sigma$  *it holds*  $I_{i_1 \dots i_r}^\sigma = 0$ .

(ii) *If*  $g$  *is an Einstein metric, a central symmetric metric a (2,2)-decomposable metric or a conformally recurrent metric and*  $\sigma \in \{1, 2, 3\}$ , *then from*  $\{(ME)_k^\sigma \mid k = 2, 4\}$  *it follows, that*  $g$  *is a conformally flat or a plane wave metric.*

(iii) *Let*  $g$  *be conformally equivalent to a metric with*  $\nabla_a C_{abcd} = 0$  *and*  $\sigma \in \{1, 2, 3\}$ . *Then the equations*  $\{(ME)_k^\sigma \mid k = 2, 4\}$  *imply, that*  $g$  *is of Petrov type*  $N$ .

(iv) Let  $g$  be of Petrov type  $N$  and  $\sigma \in \{2, 3\}$ . The moment equations  $\{(ME)_k^\sigma \mid k = 2, 4, 5\}$  are satisfied if and only if  $g$  is conformally equivalent to a generalized plane wave metric.

(v) Let  $g$  be of Petrov type  $N$ ,  $\sigma \in \{1, 2, 3\}$  and

$$11\delta_1^{(2,\sigma)} + \delta_2^{(2,\sigma)} \neq O. \quad (6.3)$$

Then the equations  $\{(ME)_k^\sigma \mid k = 2, 4, 5, 6\}$  are fulfilled if and only if  $g$  is conformally equivalent to a plane wave metric. <sup>8)</sup>

(vi) Let  $g$  be of Petrov type  $D$  and  $\sigma \in \{1, 2, 3\}$ . There are no metrics, for which the equations  $\{(ME)_k^\sigma \mid k = 2, 4\}$  are valid.

*Proof.* – If  $g$  is a plane wave metric then every equation  $E_\sigma$  satisfies Huygens' principle [G3; S1; W4]. Hence, Theorem 6.1 implies the assertion (i). The assertion (ii) was proved in [W4]. For the case  $R_{ab} = 0$  see [McL 1]. For the proof of (iii) see [W6]. If  $g$  is of Petrov type  $N$  and  $(ME)_k^\sigma$  hold for  $k = 2$  and  $4$ , then  $g$  is conformally equivalent to special cases of complex recurrent metrics [CM]. If in addition  $(ME)_5^\sigma$ ,  $\sigma \in \{2, 3\}$ , is satisfied, then  $g$  is conformally equivalent to a generalized plane wave metric (see [AW], p. 81). Conversely, if  $g$  is conformally equivalent to a generalized plane wave metric, then by Lemma 4.3 the equations  $(ME)_k^\sigma$  are satisfied for  $k = 2, 4, 5$ . In order to prove (v) we remark, that  $g$  is conformally equivalent to a plane wave metric, iff  $g$  is of Petrov type  $N$  and the equations  $(ME)_k^\sigma$  for  $k = 2, 4, 6$  are satisfied [CM]. For  $\sigma \in \{2, 3\}$  by (iv) from  $\{(ME)_k^\sigma \mid k = 2, 4, 5\}$  it follows, that  $g$  is conformally equivalent to a generalized plane wave metric. Now (5.14) and Lemma 4.3 imply

$$I_{i_1 \dots i_6}^\sigma = 33(11\delta_1^{(2,\sigma)} + \delta_2^{(2,\sigma)})Z_{i_1 \dots i_6} = O,$$

from which by Lemma 4.4 and (6.3) it follows, that  $g$  is conformal equivalent to a plane wave metric. The conversion is clear because of (i). Finally, in [W7, CM] it was proved the assertion (vi). <sup>9)</sup>

**COROLLARY 6.2.** – Let  $g$  be of Petrov type  $N$  and the condition (6.3) <sup>8)</sup> fulfilled. Every field equation  $E_\sigma$  satisfies Huygens' principle if and only if  $g$  is conformally equivalent to a plane wave metric.

#### ACKNOWLEDGEMENTS

This paper was prepared during a visit to the University of Waterloo (Canada). I would like to express my appreciation to the Department of Applied Mathematics and the *Deutsche Forschungsgemeinschaft (DFG)*

for financial support. It is a pleasure for me to thank Professor R. G. McLenaghan and the staff of the Department for their hospitality.

## REFERENCES

- [A] M. ALVAREZ, Zum Huygensschen Prinzip bei einigen Klassen spinorieller Feldgleichungen in gekrümmten Raum-Zeit-Mannigfaltigkeiten, *Dissertation A*, Pädagogische Hochschule Erfurt, 1991.
- [AM] M. ALVAREZ, V. WÜNSCH, Zur Gültigkeit des Huygensschen Prinzips bei der Weyl-Gleichung und den homogenen Maxwellschen Gleichungen für Metriken vom Petrow-Typ  $N$ , *Wiss. Zeitschr. d. Päd. Hochschule Erfurt/Mühlhausen, Math.-naturwissensch. Reihe* **27**, 1991, H. 2, pp. 77-91.
- [B] Y. BRUHAT, Sur la théorie des propagateurs, *Ann. Mat. Pura Appl.*, **64**, 1964, pp. 191-228.
- [CM] J. CARMINATI, R. G. MC LENAGHAN, An Explicit Determination of the Space-times on which the Conformally Invariant Scalar Wave Equation Satisfies Huygens' Principle, *Ann. Inst. Henri Poincaré*, Phys. théor. **44**, 1986, pp. 115-153; Part. II, **47**, 1987, pp. 337-354; Part. III, **48**, 1988, pp. 77-96; **54**, 1991, pp. 9-16.
- [DC] L. DEFRISE-CARTER, Conformal Groups and Conformally Equivalent Isometry Groups, *Comm. Math. Phys.* **40**, 1975, pp. 273-282.
- [DP] J.-C. DU PLESSIS, Polynomial Conformal Tensors, *Proc. Cambridge Phil. Soc.*, **68**, 1970, pp. 329-344.
- [F] F. G. FRIEDLANDER, *The Wave Equation on a Curved Space-time*, Cambridge University Press, Cambridge, London, New-York, Melbourne, 1975.
- [GeW1] R. GERLACH, V. WÜNSCH, Über konforminvariante Tensoren ungerader Stufe in gekrümmten Raum-Zeiten, *Wiss. Zeitschr. d. Päd. Hochschule Erfurt-Mühlhausen, Math.-naturwissensch., Reihe* **26**, 1990, H. 1, pp. 20-32.
- [GeW2] R. GERLACH, V. WÜNSCH, Über konforminvariante Tensoren der Stufe sechs in gekrümmten Raum-Zeiten, *Wiss. Zeitschr. d. Päd. Hochschule Erfurt-Mühlhausen, Math.-naturwissensch., Reihe* **27**, 1991, H. 2, pp. 117-143.
- [G1] P. GÜNTHER, Zur Gültigkeit des Huygensschen Prinzips bei partiellen Differentialgleichungen vom normalem hyperbolischen Typ., *Ber. Verh. Sächs. Akad. d. Wiss. Leipzig, M.-nat. Klasse*, **100**, H. 2, 1952.
- [G2] P. GÜNTHER, Einige Sätze über Huygenssche Differentialgleichungen, *Wiss. Zs. Karl-Marx-Univ. Leipzig*, **14**, 1965, pp. 497-507.
- [G3] P. GÜNTHER, Ein Beispiel einer nichttrivialen Huygensschen Differentialgleichung mit vier unabhängigen Veränderlichen, *Arch. Rat. Mech. and Analysis*, **18**, 1965, pp. 103-106.
- [G4] P. GÜNTHER, *Huygen's Principle and Hyperbolic Equations*, Boston, Academic Press, 1988.
- [GW1] P. GÜNTHER, V. WÜNSCH, Maxwellsche Gleichungen und Huygenssches Prinzip, I, *Math. Nachr.*, **63**, 1974, pp. 97-121.
- [GW2] P. GÜNTHER, V. WÜNSCH, Contributions to a Theory of Polynomial Conformal Tensors, *Math. Nachr.*, **126**, 1986, pp. 83-100.
- [GW3] P. GÜNTHER, V. WÜNSCH, On some Polynomial Conformal Tensors, *Math. Nachr.*, **124**, 1985, pp. 217-238.
- [Ha] J. HADAMARD, *Lectures on Cauchy's Problem in Linear Partial Differential Equations*, Yale University Press, New Haven, 1923.

- [He] S. HELGASON, Huygens' Principle for Wave Equations on Symmetric Spaces, *J. Funkt. Anal.*, **107**, 1992, pp. 279-288.
- [I] R. ILLGE, Zur Gültigkeit des Huygensschen Prinzips bei hyperbolischen Differentialgleichungssystemen in statischen Raum-Zeiten, *Zs. für Anal. und Anwendungen*, **6**, 1987, pp. 385-407.
- [L] A. LICHNEROWICZ, Champs spinoriels et propagateurs en relativité générale, *Bull. Soc. math. Fr.*, **92**, 1964, pp. 11-100.
- [McL1] R. G. MC LENAGHAN, An Explicit Determination of the Empty Space-Times on which the Wave Equation Satisfies Huygens' Principle. *Proc. Cambridge Phil. Soc.*, **65**, 1969, pp. 139-155.
- [McL2] R. G. MC LENAGHAN, On the Validity of Huygen's Principle for Second Order Partial Differential Equations with four Independent Variables. Part. I. Derivation of Necessary Conditions, *Ann. Inst. H. Poincaré*, A **20**, 1974, pp. 153-188.
- [McL3] R. G. MC LENAGHAN, Huygens' Principle. *Ann. Inst. H. Poincaré*, A **37**, 1982, pp. 211-236.
- [McL, L] R. G. MC LENAGHAN, J. LEROY, Complex Recurrent Space-times. *Proc. Roy. Soc. London*, A **327**, 1972, pp. 229-249.
- [McL, W] R. G. MC LENAGHAN, G. C. WILLIAMS, An Explicit Determination of the Petrov Type D Space-Times on which Weyl's Neutrino Equation and Maxwell's Equations Satisfy Huygens' Principle. *Ann. Inst. H. Poincaré*, A, **53**, 1990, pp. 217-223.
- [O] B. ORSTED, The Conformal Invariance of Huygens' Principle, *Diff. Geom.*, **16**, 1981, pp. 1-9.
- [PR] R. PENROSE, W. RINDLER, *Spinors and space-time*, Vol. 1, 1984, Vol. 2, 1986, Cambridge Univ. Press.
- [RW] B. RINKE, V. WÜNSCH, Zum Huygensschen Prinzip bei der skalaren Wellengleichung. *Beiträge zur Analysis*, **18**, 1981, pp. 43-75.
- [Sch] E. SCHMUTZER, *Relativistische Physik*, B. G. Teubner-Verlag, Leipzig, 1968.
- [S1] R. SCHIMMING, Zur Gültigkeit des Huygensschen Prinzips bei einer speziellen Metrik. *ZAMM*, **51**, 1971, pp. 201-208.
- [S2] R. SCHIMMING, Riemannsche Räume mit ebenfrontiger und ebener Symmetrie, *Math. Nachr.*, **59**, 1974, pp. 129-162.
- [S3] R. SCHIMMING, Das Huygenssche Prinzip bei hyperbolischen Differentialgleichungen für allgemeine Felder. *Beiträge zur Analysis* **11**, 1978, pp. 45-90.
- [W1] V. WÜNSCH, Über selbstadjungierte huygenssche Differentialgleichungen. *Math. Nachr.* **47**, 1970, pp. 131-154.
- [W2] V. WÜNSCH, Maxwellsche Gleichungen und Huygenssches Prinzip II, *Math. Nachr.*, **73**, 1976, pp. 19-36.
- [W3] V. WÜNSCH, Über eine Klasse konforminvarianter Tensoren, *Math. Nachr.*, **73**, 1976, pp. 37-58.
- [W4] V. WÜNSCH, Cauchy-Problem und Huygenssches Prinzip bei enigen Klassen spinorieller Feldgleichung I, II, *Beiträge zur Analysis*, **12**, 1978, pp. 47-76. **13**, 1979, pp. 147-177.
- [W5] V. WÜNSCH, Konforminvariante Variationsprobleme und Huygenssches Prinzip, *Math. Nachr.* **120**, 1985, pp. 175-193.
- [W6] V. WÜNSCH, C-Räume und Huygenssches Prinzip, *Wiss. Zeitschr. d. Päd. Hochschule Erfurt-Mühlhausen, Math.-naturwissensch., Reihe*, **23**, 1987, H. 1, pp. 103-111.
- [W7] V. WÜNSCH, Huygens' Principle on Petrov Typ D Space-Times, *Ann. d. Physik*, **7**. Folge, Bd. 46, H. 8, 1989, pp. 593-597.
- [W8] V. WÜNSCH, Conformal C- and Einstein Spaces, *Math. Nachr.*, **146**, 1990, pp. 237-245.

## NOTES

- 1) The Levi-Civita tensor  $e_{abcd}$  is strictly speaking a pseudo-tensor.
- 2) In [GW3] generating systems were derived also for higher dimensions.
- 3) It is easy to prove, that there are no trivial tensors in  $S_5$  and  $S_6$ .
- 4) In [G4] moments for a general hyperbolic equation for higher dimensions were defined, too.
- 5) If  $r$  odd then because of (4.12) the moments contain the Levi-Civita pseudo-tensor  $e_{abcd}$  and the moments are strictly speaking pseudo-tensors.
- 6) For this assertion it is sufficient for  $g$  to require to be of class  $C^\infty$ .
- 7) A conjecture is, that the moment equations  $(ME)_r^\sigma$  ( $r = 2, 4, 5, 6$ ) are also sufficient for the validity of Huygens' principle for  $E_\sigma$ ,  $\sigma = 1, 2, 3$ , and that these equations are fulfilled if and only if  $g$  is conformally flat or a plane wave metric [W4, 6; CM].
- 8) For  $\sigma = 1$  by Lemma 6.1 the condition (6.3) is satisfied. For  $\sigma = 2, 3$  we show this in a subsequent paper.
- 9) In order to prove (ii)-(vi) we mainly use the two-component spinor formalism of Penrose and the spin coefficient formalism of Newman and Penrose [PR]

*(Manuscript received May 17, 1993;  
revised version received July 12, 1993)*