

ANNALES DE L'I. H. P., SECTION A

QIHONG FAN

Resonances of relativistic Schrödinger operators with homogeneous electric fields

Annales de l'I. H. P., section A, tome 61, n° 2 (1994), p. 205-221

http://www.numdam.org/item?id=AIHPA_1994__61_2_205_0

© Gauthier-Villars, 1994, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Resonances of relativistic Schrödinger operators with homogeneous electric fields*

by

Qihong FAN

Department of Mathematics, Peking University,
Peking 100871, China

ABSTRACT. – In this paper, we apply the analytic distortion technique and the calculus of pseudodifferential operators to study the resonances of the relativistic Schrödinger operators with homogeneous electric fields. By constructing a suitable approximate operator, we give precise locations for resonances generated by eigenvalues below the bottom of the essential spectra and as a consequence, we obtain an upper bound on the widths of resonances.

RÉSUMÉ. – Dans cet article, nous utilisons la technique de déformation analytique et le calcul pseudodifférentiel pour étudier les résonances d'un opérateur de Schrödinger relativiste en présence d'un champ électrique homogène. En construisant une approximation convenable de cet opérateur, nous fournissons une localisation précise des résonances engendrées par les valeurs propres situées au-dessous du spectre essentiel, ce qui fournit en conséquence une borne supérieure sur la largeur des résonances.

1. INTRODUCTION

The study of resonances in Schrödinger operator theory has been an object of growing interest in last years. There are several mathematical

*This research has been supported by Natural Science Foundation of China.

concepts, but there is not a general theory which can describe all physical phenomena considered as resonances. A very powerful method is the complex scaling which was introduced by Aguilar, Balslev and Combes in [AC], [BC]. It was extended in several directions, *see* [CFKS] for a survey. In [HS], using the calculus of pseudodifferential operators on I-Lagrange manifolds, Fourier integral operators and the microlocal analysis, Helffer and Sjöstrand founded the microlocal theory of resonances for a wide class of h -pseudodifferential operators.

The purpose of this work is to discuss the resonances of relativistic Schrödinger operators with homogeneous electric fields, $H(\beta) = \sqrt{-\Delta + m^2} - m + v(x) + \beta x_1$, where β is a small parameter. In this paper, we assume $v(x)$ satisfies the following conditions.

(H1). There are two constants $C, R_0 > 0$, such that $v(x)$ is an analytic function in variable x_1 in complex region: $\{x_1, |Im x_1| \leq C |Re x_1|, Re x_1 < -R_0\}$.

(H2). $v(x) \rightarrow 0$, as $|x| \rightarrow \infty$, and $|\partial^\alpha v(x)| \leq C_\alpha \langle x \rangle^{-|\alpha|}$, for any $\alpha \geq 0$, and x in the above analyticity region.

One can find many authors who have studied the spectral properties of $H(0)$ in literature. *See* [CMS] and its references there. In general the spectra of $H(0)$ include eigenvalues and the essential spectrum $(0, +\infty)$ (*cf.* [CMS], [Fa2] and [We]). In [HP], Helffer and Parisse studied the decay properties of eigenfunctions and tunneling effect of $H(0)$ in semiclassical limit, and they showed the decays of the eigenfunctions of $H(0)$ are more rapid than that of the Dirac operators. For the operator $H(\beta)$, $\beta \neq 0$, if $v(x)$ is real valued, one can prove $H(\beta)$ is essentially self-adjoint on $C_0^\infty(R^n)$ (*cf.* [Fa1]). Furthermore, in [Fa3], we proved that the essential spectrum of $H(\beta)$ ($\beta \neq 0$) is $(-\infty, +\infty)$. In the following we will study the properties of the discrete eigenvalues of $H(0)$ under the perturbation βx_1 , for β small enough. We will prove that the discrete eigenvalues of $H(0)$ become the resonances of $H(\beta)$, when β is sufficiently small.

The main ideas of this paper are similar to those for usual Schrödinger operators with Stark effect (*cf.* [Wa1], [Wa2] and [Wa3]), but many technical points are quite different. As in [Wa1], we used analytic distortion to define resonances for $H(0)$ and to construct an appropriate Grushin problem to prove the existence and the locations of resonances. The latter is in the spirit of the work of Helffer-Sjöstrand on resonances in semiclassical limit (*cf.* [HS]). The differences between relativistic and non relativistic Schrödinger operators arise from the fact that the former is a pseudodifferential operator (non local operator) while the latter is a differential operator (local operator).

The arrangement of this paper as following. In section 2, we introduce the complex dilation and give the definition of resonances. In section 3, we will study the behavior of eigenfunctions with discrete eigenvalues. In section 4, we will study the location of resonances.

2. COMPLEX DILATION AND THE RESONANCES

In this section, we assume $v(x)$ satisfies the condition (H1), (H2). We will introduce the complex dilation as in [CDKS], [Hu], [Wa1]. Let $\lambda \in \mathbb{R}$, $\lambda < 0$, and $\delta > 0$ sufficiently small. One can choose $\chi \in C^\infty(\mathbb{R})$, such that $\chi(t) = 1$, $t < (\lambda + \delta)/\beta$, $\chi(t) = 0$, $t > (\lambda + 2\delta)/\beta$, and $|\chi(t)| \leq 1$, $\chi^{(k)}(t) \leq C_k \left(\frac{\beta}{\delta}\right)^k$.

For $\theta \in \mathbb{R}$, put $\phi_\theta(x) = (e^{\theta\chi(x_1)} x_1, x')$, $x = (x_1, x')$. Since $\det \phi'_\theta(x) = e^{\theta\chi(x_1)} (1 + \theta\chi'(x_1) x_1)$, one knows that ϕ_θ is invertible when $|\theta|$ is sufficiently small, say $|\theta| < \theta_0$. In this case, let $\psi_\theta = \phi_\theta^{-1}$. In the following, we always assume $|\theta| < \theta_0$. Let $U(\theta)$ be the operator defined by

$$U(\theta) f(x) = f(\phi_\theta(x)) |\phi'_\theta(x)|^{1/2}.$$

In this paper, we will use the Weyl corresponding of symbol $\sigma(x, \xi)$, that is

$$\sigma(x, D) f(x) = (2\pi)^{-n} \int_{\mathbb{R}^{2n}} e^{i(x-y)\xi} \sigma\left(\frac{x+y}{2}, \xi\right) f(y) dy d\xi.$$

The symbol of $H(\beta)$ is given by

$$p(x, \xi) = \sqrt{m^2 + \xi^2} - m + v(x) + \beta x_1$$

The symbol $p_\theta(x, \xi)$, of $U(\theta) p(x, D) U(\theta)^{-1}$, is given by

$$p_\theta(x, \xi) = (2\pi)^{-n} \int \int p\left(\frac{1}{2} \left(\phi_\theta\left(x + \frac{t}{2}\right) + \phi_\theta\left(x - \frac{t}{2}\right)\right), \right. \\ \left. {}^T \Psi_\theta(x, t)^{-1} (\xi + \eta) \right) \cdot J_\theta(x, t) e^{i(\eta, t)} d\eta dt, \quad (2.1)$$

where $J_\theta(x, t) = \left| \phi'_\theta\left(x + \frac{t}{2}\right) \right|^{1/2} \left| \phi'_\theta\left(x - \frac{t}{2}\right) \right|^{1/2} |\det \Psi_\theta(x, t)|^{-1}$, and $\Psi_\theta(x, t)$ satisfies: $\phi_\theta\left(x + \frac{t}{2}\right) - \phi_\theta\left(x - \frac{t}{2}\right) = {}^T \Psi_\theta(x, t) t$ and $\Psi_\theta(x, 0) = \phi'_\theta(x)$.

Let S^k be the symbol classes of Hörmander type $S_{1,0}^k$, that is

$$S^k = \{a(x, \xi) \in C^\infty(R^{2n}), |\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha\beta} \langle \xi \rangle^{k-|\beta|}, \alpha, \beta \geq 0\}.$$

From (2.1) and Taylor expansion in variable t , one can get

$$p_\theta(x, \xi) = p_0(x, \xi) + R_\theta(x, \xi), \quad (2.2)$$

where $p_0(x, \xi) = p(\phi_\theta(x), \psi'_\theta(x)\xi)$, $R_\theta(x, \xi) \in S^0$, and satisfy

$$|\partial_\xi^k \partial_x^\alpha R_\theta(x, \xi)| \leq O(\beta/\delta) C_{k\alpha} \langle \xi \rangle^{-|k|}, \quad \alpha, \beta \geq 0. \quad (2.3)$$

Therefore

$$\|R_\theta(x, \xi)\| \leq C \frac{\beta}{\delta}.$$

Let $\Omega = \{\theta \in C, |\theta| < \rho, \text{Im } \theta > 0\}$. We should notice that in the following, $\rho > 0$ will always be sufficiently small. For $\theta \in \Omega$, we define

$$E(\theta, \delta) = \{e^\theta r + s, s > -4\delta, r < 4\delta\}.$$

For $\beta > 0$ sufficiently small, $p_\theta(x, D)$ defined for $\theta \in R$ and $|\theta| < \rho$ has a natural extension in θ into Ω . In fact, by the analyticity assumption (H1) of $v(x)$, one knows that, for $\beta > 0$ sufficiently small, $U(\theta)v(x)U(\theta)^{-1}$ is analytic in $\theta \in \Omega$. In order to prove that $p_\theta(x, D)$ has an analytic extension in $\theta \in \Omega$, one note that

$$\begin{aligned} U(\theta) \sqrt{m^2 - \Delta} U(\theta)^{-1} f(x) &= \frac{1}{(2\pi)^n} \int \int e^{i(\phi_\theta(x) - \phi_\theta(y)) \cdot \xi} \\ &\times \sqrt{m^2 + \xi^2} |\phi'_\theta(x)|^{1/2} |\phi'_\theta(y)|^{1/2} f(y) d\xi dy, \end{aligned}$$

for $f \in S(R^n)$. $\phi_\theta(x) - \phi_\theta(y) = ((x_1 - y_1) \tilde{\psi}_\theta(x_1, y_1), x' - y')$, where $\tilde{\psi}_\theta(x_1, y_1) = \int_0^1 \phi'_{\theta,1}(x_1 + \tau(y_1 - x_1)) d\tau$ and $\phi_{\theta,1}$ is the first component of ϕ_θ . Since $\phi'_{\theta,1} = e^{\theta \chi(x_1)} (1 + \theta \chi'(x_1) x_1)$, for $\theta \in \Omega$ small enough, one can check that $\tilde{\psi}_\theta(x_1, y_1) \neq 0$, and $|\text{Im } \tilde{\psi}_\theta(x_1, y_1)| \leq C(\theta) |\text{Re } \tilde{\psi}_\theta(x_1, y_1)|$, for $x_1, y_1 \in R$, where $|C(\theta)| < C|\theta|$ and C is a constant. Since $\sqrt{m^2 + \xi^2}$ is an analytic function in variable ξ_1 in the complex region: $|\text{Im } \xi_1| \leq \frac{1}{2} |\text{Re } \xi_1|$, one can make a changement of the integral countor of variable in ξ_1 to show $U(\theta) \sqrt{m^2 - \Delta} U(\theta)^{-1}$

is a standard ps. d. o. and

$$\begin{aligned} U(\theta) \sqrt{m^2 - \Delta} U(\theta)^{-1} f(x) \\ = \frac{1}{(2\pi)^n} \int \int e^{i(x-y) \cdot \xi} \sqrt{m^2 + (\xi_1 / \tilde{\psi}_\theta(x_1, y_1))^2 + \xi'^2} \\ \times \frac{|\phi'_\theta(x)|^{1/2} |\phi'_\theta(y)|^{1/2}}{\tilde{\psi}_\theta(x_1, y_1)} f(y) d\xi dy. \end{aligned} \quad (2.4)$$

The analyticity of $U(\theta) \sqrt{m^2 - \Delta} U(\theta)^{-1}$ in $\theta \in \Omega$ follows from the analyticity of $\tilde{\psi}_\theta(x_1, y_1)$ and (2.4).

THEOREM 2.1. – For $\theta \in \Omega$, $p_\theta(x, D)$ defined on $H^1 \cap D(x_1)$ is closed in $L^2(R^n)$ and is a holomorphic family of type (A).

Proof. – For $\text{Im } z > 0$ sufficiently large, one has

$$|p_\theta(x, \xi) - z| \geq C(\langle \xi \rangle + |\beta x_1| + \text{Im } z).$$

Let $b_\theta(x, \xi) = (p_\theta(x, \xi) - z)^{-1}$. By the calculus of pseudodifferential operators (cf. [CM]), one has

$$(p_\theta(x, D) - z) b_\theta(x, D) = I + R_1(x, D),$$

and

$$b_\theta(x, D) (p_\theta(x, D) - z) = I + R_2(x, D),$$

where $R_i(x, D)$ satisfy the estimates

$$\|R_i(x, D)\| \leq \frac{C}{\text{Im } z}, \quad i = 1, 2.$$

By these estimates, one can prove easily that $p_\theta(x, D)$ is a closed operator. The analyticity of $(p_\theta(x, D) - z)^{-1}$ in $\theta \in \Omega$ follows from the assumption (H1) on $v(x)$ and (2.4). ■

THEOREM 2.2. – For $\theta \in \Omega$, $\sigma_{\text{ess}}(p_\theta(x, D)) \subset E(\theta, \delta)$.

Proof. – One can choose a cut off function χ_t , such that $\text{supp } \chi_t \subset B_{2t}$ and $|v(\phi_\theta(x))| \leq \frac{\delta}{4}$, $x \notin B_t$. Let $\tilde{p}_\theta(x, D) = p_\theta(x, D) - \chi_t v(\phi_\theta(x))$. Then $\sigma_{\text{ess}}(\tilde{p}_\theta(x, D)) = \sigma_{\text{ess}}(p_\theta(x, D))$. In order to characterize the essential spectrum of $\tilde{p}_\theta(x, D)$, we first calculate the numerical range of $\tilde{p}_\theta(x, D)$ on $L^2(R_+^n) \oplus L^2(R_-^n)$, where $R_+^n = \{x = (x_1, x'), x_1 > (\lambda + 2\delta)/\beta\}$, $R_-^n = \{x = (x_1, x'), x_1 < (\lambda + 2\delta)/\beta\}$.

For u with $\text{supp } u \subset \left\{x, x_1 < \frac{\lambda + 2\delta}{\beta}\right\}$, when ρ, β are sufficiently small, one has

$$(\tilde{p}_\theta(x, D)u, u) \in E(\theta, \delta).$$

For u with $\text{supp } u \subset \left\{x, x_1 > \frac{\lambda + 2\delta}{\beta}\right\}$.

$$\begin{aligned} (\tilde{p}_\theta(x, D)u, u) &= \iint \tilde{p}_\theta(x, \xi) W(u)(x, \xi) dx d\xi \\ &= \iint (p_0(x, \xi) - \chi_t v(\phi_\theta(x)) + R_\theta(x, \xi)) W(u)(x, \xi) dx d\xi \\ &= \iint (\sqrt{m^2 + \xi^2} (1 + C_\theta(x, \xi)) + R_\theta(x, \xi) \\ &\quad + (1 - \chi_t(x)) v(\phi_\theta(x)) + e^{\theta x(x)} \beta x_1) W(u)(x, \xi) dx d\xi, \end{aligned}$$

where $W(u)(x, \xi)$ is the Wigner distribution of u (cf. [Fo]), $C_\theta(x, \xi) \in S^0$, and $|C_\theta(x, \xi)| \leq C|\theta|$. Therefore, when ρ, β are sufficiently small, we have $(\tilde{p}_\theta(x, D)u, u) \in E(\theta, \delta)$. We proved that $\sigma_{\text{ess}}(\tilde{p}_\theta(x, D)) \subset E(\theta, \delta)$ on $L^2(R_+^n) \oplus L^2(R_-^n)$. Since the $\sigma_{\text{ess}}(\tilde{p}_\theta(x, D))$ on $L^2(R^n)$ is the same as on $L^2(R_+^n) \oplus L^2(R_-^n)$ (cf. [K]). This completes the proof. ■

The spectra of $p_\theta(x, D)$ in $C \setminus E(\theta, \delta)$ are discrete eigenvalues. We call these eigenvalues the resonances of $p(x, D)$.

THEOREM 2.3. – *The eigenvalues of $p_\theta(x, D)$ are essentially independent of $\theta \in \Omega$. More precisely, for every compact $K \subset \subset \Omega$, there exists $\beta_0 > 0$, such that for $0 < \beta < \beta_0$, $z \in C \setminus E(\theta, \delta)$, $z \in \sigma(p_\theta(x, D))$, for some $\theta \in K$, then $z \in \sigma(p_\theta(x, D))$ for any $\theta \in K$.*

The proof of the theorem is standard, we omit it here.

Remark. – The resonances of $p(x, D)$ are independent of the choice of χ used in the definition of $U(\theta)$ (cf. [Wa1]).

3. DECAY PROPERTIES OF EIGENFUNCTIONS

In this section, we will estimate the decay properties of eigenfunctions of various operators. By the estimates, we can prove the stability theorem of resonances. Some of the results in this section were obtained by Carmona-Master-Simon in [CMS], by probability method. In the following, we will use the commutator method, as in [Wa2], to study the decay properties of eigenfunctions.

Let $H = \sqrt{m^2 - \Delta} - m + v(x)$, and $H(x, \xi) = \sqrt{m^2 + \xi^2} - m + v(x)$.

PROPOSITION 3.1. – Assume that $d(x) \in C^\infty(\mathbb{R}^n)$, $0 \leq d(x) \leq C|x|$, $|d'(x)| < m$. $|\partial^\alpha d(x)| \leq C|x|^{1-|\alpha|}$, for $|x|$ sufficiently large. Then the symbol, $H_d(x, \xi)$, of operator $e^{d(x)} H(x, D) e^{-d(x)}$, is given by

$$H_d(x, \xi) = H(x, \xi + id'(x)) + H_0(x, \xi), \quad (3.1)$$

with $H_0(x, \xi) \in S^0$, and

$$|\partial_x^\alpha \partial_\xi^\beta H_0(x, \xi)| \leq C_{\alpha\beta} \langle x \rangle^{-1} \langle \xi \rangle^{-|\beta|}. \quad (3.2)$$

Proof.

$$\begin{aligned} e^{d(x)} H(x, D) e^{-d(x)} u(x) \\ &= (2\pi)^{-n} \iint e^{i(x-y)\eta + (d(x)-d(y))} H\left(\frac{x+y}{2}, \eta\right) u(y) dy d\eta \\ &= (2\pi)^{-n} \iint e^{i(x-y)(\eta - id'(\tilde{x}))} H\left(\frac{x+y}{2}, \eta\right) u(y) dy d\eta, \end{aligned}$$

where $d(x) - d(y) = (x - y) d'(\tilde{x})$. Since $H(x, \xi)$ is an analytic function in variable η in the complex region $\{\eta \in \mathbb{C}^n, |\operatorname{Im} \eta| < m\}$, one can change the integral contour in η to show

$$\begin{aligned} e^{d(x)} H(x, D) e^{-d(x)} u(x) \\ &= (2\pi)^{-n} \iint e^{i(x-y)\xi} H\left(\frac{x+y}{2}, \xi + id'(\tilde{x})\right) u(y) dy d\xi. \end{aligned}$$

By Taylor formula, one has

$$\begin{aligned} H\left(\frac{x+y}{2}, \xi + id'(\tilde{x})\right) &= H\left(\frac{x+y}{2}, \xi + id'\left(\frac{x+y}{2}\right)\right) \\ &+ H'_\xi\left(\frac{x+y}{2}, \xi + id'\left(\frac{x+y}{2}\right)\right) \left(d'(\tilde{x}) - d'\left(\frac{x+y}{2}\right)\right), \end{aligned}$$

where $\tilde{d}'(x, y) = \tau d'\left(\frac{x+y}{2}\right) + (1 - \tau) d'(\tilde{x})$, $0 < \tau < 1$. Therefore, we have

$$H_d(x, \xi) = H(x, \xi + id'(x)) + H_0(x, \xi),$$

and

$$|\partial_x^\alpha \partial_\xi^\beta H_0(x, \xi)| \leq C_{\alpha\beta} \langle x \rangle^{-1} \langle \xi \rangle^{-|\beta|}.$$

This completes the proof. ■

PROPOSITION 3.2. — Assume that $d(x) \in C^\infty(R^n)$, $0 \leq d(x) \leq C|x|$, $|d'(x)| < C$, $C < m$ and $|\partial^\alpha d(x)| \leq C_\alpha |x|^{1-|\alpha|}$, as $|x|$ sufficiently large. For $E \in R$, we assume that

$$|H(x, \xi + id'(x)) - E| \geq C \langle \xi \rangle, \quad \text{for } \langle x \rangle + \langle \xi \rangle \geq C', \quad (3.3)$$

where C, C' are constants. Then there is a $Q(x, \xi) \in S^{-1}$, such that

$$Q(x, D)(H_d(x, D) - E) = I + R_d(x, D), \quad (3.4)$$

with $R_d(x, \xi) \in S^{-1}$, and satisfy

$$|\partial_\xi^\beta \partial_x^\alpha R_d(x, \xi)| \leq C_{\alpha\beta} \langle x \rangle^{-1} \langle \xi \rangle^{-1-\beta}, \quad (3.5)$$

where the constants $C_{\alpha\beta}$ are uniformly bounded with respect to the C^k semi-norms of $d(x)$.

Proof. — By the assumption (3.3), one can choose $Q(x, \xi) \in S^{-1}$, such that $Q(x, \xi) = (H(x, \xi + id'(x)) - E)^{-1}$, for $\langle x \rangle + \langle \xi \rangle \geq C'$. By the calculus of pseudodifferential operators, one can get easily

$$Q(x, D)(H_d(x, D) - E) = I + R_d(x, D),$$

with $R_d(x, \xi)$ satisfying the estimate (3.5). ■

For $E \in R$, $E < 0$, let $m(E) = \sqrt{2m|E| - E^2}$, for $|E| \leq m$, $m(E) = m$, for $|E| > m$. In the following, we will choose $d^\varepsilon(x) \in C^\infty$, $d^\varepsilon(x) = (1 - \varepsilon)m(E)|x|$, for $\varepsilon > 0$, $|x|$ sufficiently large. One can get the following estimate

$$|\sqrt{m^2 + (\xi + id^{\varepsilon'}(x))^2} - m - E| \geq C \langle \xi \rangle, \quad (3.6)$$

as $|x|$ sufficiently large.

THEOREM 3.3. — Assume $v(x)$ satisfy the conditions (H1) and (H2). Let f be an eigenfunction of operator $\sqrt{m^2 - \Delta} - m + v(x)$ with eigenvalue E , $E \in R$, $E < 0$, and $d^\varepsilon(x)$ chosen as above. Then for any $\varepsilon > 0$, $e^{d^\varepsilon(x)} f(x) \in L^2(R^n)$.

Proof. — If $e^{d^\varepsilon(x)} f(x) \notin L^2(R^n)$, we can choose $\chi_\lambda(t) = t$, for $t \leq \lambda$, and $\chi_\lambda(t) = \lambda + 1$, for $t \geq \lambda + 2$, with $\chi_\lambda \geq 0$, $0 < \chi'_\lambda(t) \leq 1$ and $|\chi_\lambda^{(k)}(t)| \leq C$, for any λ, t . Let $d_\lambda(x) = \chi_\lambda(d^\varepsilon(x))$, and $f_\lambda(x) = e^{d_\lambda(x)} f(x) / \|e^{d_\lambda} f(x)\|$. Then

$$f_\lambda \rightarrow 0, \quad \text{weakly, as } \lambda \rightarrow \infty.$$

For $|x|$ sufficiently large, one has

$$|\sqrt{m^2 + \xi^2} - m - E + v(x)| \geq C \langle \xi \rangle.$$

By the estimate (3.6), one can verify easily

$$|H(x, \xi + id'_\lambda(x)) - E| \geq C \langle \xi \rangle.$$

The constant C is independent of λ . By Proposition (3.2), there is a $Q_\lambda(x, \xi) \in S^{-1}$, such that

$$Q_\lambda(x, D)(H_{d_\lambda}(x, D) - E) = I + R_{d_\lambda}(x, D)$$

and $R_{d_\lambda}(x, \xi)$ satisfies the estimate (3.5) uniformly with respect to λ . Therefore $R_{d_\lambda}(x, D)$ are compact operators and

$$|\partial_\xi^\beta \partial_x^\alpha R_{d_\lambda}(x, \xi)| \leq C_{\alpha\beta} \langle x \rangle^{-1} \langle \xi \rangle^{-\beta}, \quad (3.7)$$

Since $(H_{d_\lambda}(x, D) - E)f_\lambda = 0$, one has

$$(I + R_{d_\lambda}(x, D))f_\lambda = 0 \quad (3.8)$$

From (3.7), one has

$$\lim_{\lambda \rightarrow \infty} R_{d_\lambda}(x, D)f_\lambda = 0.$$

By (3.8), one has $f_\lambda \rightarrow 0$, strongly. This is a contradiction. ■

Remark. – The analyticity assumption on $v(x)$ is not necessary in theorem 3.3. In fact, if $v(x) \rightarrow 0$, as $|x| \rightarrow \infty$, and $|\partial^\alpha v(x)| \leq C_\alpha \langle x \rangle^{-|\alpha|}$, then all results in this section hold. This decay results has been obtained by probability method in [CMS], and in [HP], Helffer and Parrisse studied the decay properties of eigenfunctions of operator $\sqrt{1 - h^2 \Delta} + v(x)$, in the semiclassical limit $h \rightarrow 0$.

For $\eta > 0$, $\eta < -E - \delta$, one can choose $\chi_\eta \in C^\infty(R)$, such that $\chi_\eta(t) = 1$, for $t > -\eta$; $\chi_\eta(t) = 0$, for $t < -2\eta$. Let

$$\tilde{H}(\beta) = \sqrt{m^2 - \Delta} - m + v(x) + \beta x_1 \chi_\eta(\beta x_1).$$

THEOREM 3.4. – Let $f_\beta(x)$ be a normalized eigenfunction of $\tilde{H}(\beta)$ with eigenvalue in $I(\beta) = (E - C\beta^{1/2}, E + C\beta^{1/2})$, $C > 0$ is a constant. Then, for any $\varepsilon > 0$, there exists a $\beta_\varepsilon > 0$, such that

$$\sup_{0 < \beta < \beta_\varepsilon} \|e^{d^\varepsilon(x)} f_\beta(x)\| < \infty \quad (3.9)$$

Proof. – For any $\varepsilon > 0$, one can choose $\eta > 0$, $\beta_\varepsilon > 0$ sufficiently small such that $(1 - \varepsilon)m(E) \leq (1 - \varepsilon')m(E + \eta + \beta_\varepsilon)$, for some $\varepsilon' > 0$. For $|x|$ sufficiently large, one can get easily the following estimate

$$|\sqrt{m^2 + (\xi + id^{\varepsilon'}(x))^2} - m - E(\beta) + v(x) + \beta x_1 \chi_\eta(\beta x_1)| \geq C \langle \xi \rangle.$$

By this estimate and the proof processes in Theorem 3.1 and Theorem 3.3, one can prove (3.9). ■

THEOREM 3.5. – *Let E_0 be a discrete eigenvalue of H with multiplicity N , and $I = (E_0 - C(\beta), E_0 + C(\beta))$, $C(\beta) = C\beta^{1/2}$, $C > 0$. There exists a $\beta_0 > 0$, such that for $0 < \beta < \beta_0$, there are exactly N eigenvalues of $\tilde{H}(\beta)$, repeated according to their multiplicity in I . Let $\mu_1, \dots, \mu_N$, denote these eigenvalues. Then we have $|\mu_j - E_0| < C\beta$, $0 < \beta \leq \beta_0$, $j = 1, \dots, N$, and moreover*

$$\mu_l(\beta) \sim E_0 + \sum_{j=1}^{\infty} \lambda_j^l \beta^j, \quad l = 1, \dots, N. \quad (3.10)$$

Proof. – Let $f_1, \dots, f_N$ be the orthonormalized eigenfunctions of H , $H f_j = E_0 f_j$, $j = 1, \dots, N$. Put

$$g_j(\beta) = \chi_\eta \left(\frac{x_1}{\beta} \right) f_j,$$

where $\chi_\eta \in C^\infty(R)$, such that $\chi_\eta(t) = 1$, for $t > -\eta$; $\chi_\eta(t) = 0$, for $t < -2\eta$ then, from Theorem 3.3, we have $g_j \in D(\tilde{H}(\beta))$ and

$$\tilde{H}(\beta) g_j = E_0 g_j + O(\beta), \quad \text{in } L^2(R^n). \quad (3.11)$$

Now let u_j be the normalized eigenfunctions of $\tilde{H}(\beta)$ associated with the eigenvalue $\mu_j(\beta)$, $j = 1, \dots, \tilde{N}$. Put $\nu_j = \chi_\eta \left(\frac{x_1}{\beta} \right) u_j$. From Theorem 3.4, we obtain

$$H \nu_j = \mu_j \nu_j + O(\beta), \quad \text{in } L^2(R^n). \quad (3.12)$$

By the estimates in Theorem 3.3 and Theorem 3.4, one knows that the matrices $((g_i, g_j))$ and $((\nu_i, \nu_j))$ are close to the identity matrices in $R^{N \times N}$, $R^{\tilde{N} \times \tilde{N}}$ respectively, as $\beta \rightarrow 0$. So $\{g_1, \dots, g_N\}$ and $\{\nu_1, \dots, \nu_{\tilde{N}}\}$ are two linearly independent sets of $L^2(R^n)$, for $0 < \beta < \beta_0$. From this and estimates (3.11), (3.12), one can prove easily that $N = \tilde{N}$, $|\mu_j - E_0| \leq C\beta$. The detailed process can be found in [Wal]. The proof of the asymptotic formula (3.10) is similar to that in [S]. ■

4. LOCATIONS OF RESONANCES

In this section, we will study the locations of resonances of the operator $H(\beta)$. That is to study the positions of eigenvalues of operator $p_\theta(x, D)$. We also assume $v(x)$ satisfies the condition (H1), (H2). Let E_0 be a discrete eigenvalue of H , with multiplicity N . In order to estimate the width of resonances, in the definition of $U(\theta)$ in section 2, we will choose $\lambda = E_0$, $\delta > 0$ sufficiently small, and in the definition of $\tilde{H}(\beta)$ in section 3,

we will choose $\eta < -E_0 - \delta$. By Theorem 3.5, there exist exactly N eigenvalues of $\tilde{H}(\beta)$ near E_0 . In order to study the eigenfunctions of $p_\theta(x, D)$, we will construct an approximate inverse of $(p_\theta(x, D) - z)^{-1}$ out side $\{x, x_1 < (E_0 + \delta)/\beta\}$.

Let $\tilde{p}_\theta(x, \xi) = p_\theta(x, \xi) - \chi_\eta(\beta|x|)v(\phi_\theta(x))$, and $S(E_0) = \{z \in C, |z - E_0| < C\beta^{1/2}\}$. Then for $z \in S(E_0)$, one has the following estimates,

$$|\tilde{p}_\theta(x, \xi) - E| \geq C\delta|\theta|(|\xi| + \beta|x_1| + 1) \quad (4.1)$$

$$|\partial_x^\alpha \tilde{p}_\theta(x, \xi)| \leq C\beta, \quad |\alpha| \geq 1 \quad (4.2)$$

$$|\partial_\xi^\alpha \tilde{p}_\theta(x, \xi)| \leq C\langle \xi \rangle^{1-|\alpha|}, \quad |\alpha| \geq 1 \quad (4.3)$$

The estimate (4.2) comes from $|\chi'(x)| \leq c\frac{\beta}{\delta}$, (2.3) and the property of $v(x)$. By these estimates and the calculus of pseudodifferential operators (cf. [CM]), we have the following proposition.

PROPOSITION 4.1. – *There exists a $\beta_0 > 0$, such that for $\theta \in \Omega^+$, $E \in S(E_0)$, $\tilde{p}_\theta(x, D) - E$ is invertible and moreover we have*

$$\|(\tilde{p}_\theta(x, D) - E)^{-1}\| \leq C\delta^{-k}|\theta|^{-k},$$

for any $E \in S(E_0)$, where k is a constant.

LEMMA 4.2. – *The integral kernel $K_\theta(x, y)$ of the operator $(\tilde{p}_\theta(x, D) - E)^{-1}$ satisfies the estimate*

$$K_\theta(x, y) = O(e^{-\gamma d(x-y)}), \quad (4.4)$$

for some $\gamma > 0$, and uniformly for $E \in S(E_0)$, i.e. for $r > 0$, $y_0 \in R^n$, let $U(y_0) = \{y \in R^n, |y - y_0| < r\}$, one has

$$\|e^{\gamma d(x-y_0)}(\tilde{p}_\theta(x, D) - E)^{-1}g\| \leq C(r)\|g\|_{L^2(U(y_0))}, E \in S(E_0).$$

where $d(x) \in C^\infty(R^n)$, $0 \leq d(x) \leq C\langle x \rangle$, $|\partial^\alpha d(x)| \leq C_\alpha\langle x \rangle^{1-|\alpha|}$.

Proof. – Let $\tilde{p}_{\theta, \gamma}(x, D) = e^{\gamma d(x-y_0)}\tilde{p}_\theta(x, D)e^{-\gamma d(x-y_0)}$, and

$$\tilde{R}_{\theta, \gamma}(x, D) = \tilde{p}_{\theta, \gamma}(x, D) - \tilde{p}_\theta(x, D).$$

By (2.4), one knows that $\tilde{p}_\theta(x, \xi)$ is an analytic function in variable ξ in the complex region: $|\operatorname{Im} \xi| \leq C|\operatorname{Re} \xi|$, where C is a small constant. As in the proof of proposition 3.1, for γ sufficiently small, one can make a

changement of the integral counter of variable in ξ to show that $\tilde{p}_{\theta, \gamma}(x, D)$ is a standard ps. d. o. and

$$\begin{aligned} & \tilde{p}_{\theta, \gamma}(x, D) f(x) \\ &= (2\pi)^{-n} \int \int e^{i(x-y)\xi} \tilde{p}_{\theta} \left(\frac{x+y}{2}, \xi + i\gamma d'_{y_0}(\tilde{x}) \right) f(y) dy d\xi, \end{aligned}$$

where $d_{y_0}(x) = d(x - y_0)$. and $d_{y_0}(x) - d_{y_0}(y) = (x - y) d'_{y_0}(\tilde{x})$. By Taylor expansion, one can get the following estimate easily.

$$\|\tilde{R}_{\theta, \gamma}(x, D)\| \leq C\gamma.$$

When γ sufficiently small one has

$$\begin{aligned} (\tilde{p}_{\theta, \gamma}(x, D) - E)^{-1} &= (1 + (\tilde{p}_{\theta}(x, D) - E)^{-1} \\ &\quad \times \tilde{R}_{\theta, \gamma}(x, D))^{-1} (\tilde{p}_{\theta}(x, D) - E)^{-1} \end{aligned}$$

Let $g \in L^2(R^n)$, and $\text{supp } g \subset U(y_0)$. Put

$$f(x) = \int_{R^n} K_{\theta}(x, y) g(y) dy.$$

Then $(\tilde{p}_{\theta}(x, D) - E) f = g$, and $(\tilde{p}_{\theta, \gamma}(x, D) - E) e^{\gamma d_{y_0}} f = e^{\gamma d_{y_0}} g$. One has

$$e^{\gamma d_{y_0}} f = (\tilde{p}_{\theta, \gamma} - E)^{-1} (e^{\gamma d_{y_0}} g)$$

Therefore

$$\|e^{\gamma d_{y_0}} f\|_{L^2} \leq C(r) \|g\|.$$

This completes the proof. ■

In the following, we will outline the construction of Grushin problem for $p_{\theta}(x, D) - z$, $z \in S(E_0)$, the detailed process, see [HS] and [Wa1]. Let $\mu_1(\beta), \dots, \mu_N(\beta)$ be the eigenvalues of $\tilde{H}(\beta)$ in the interval $I(\beta)$ and $u_1(\beta), \dots, u_N(\beta)$ be the associated normalized eigenfunctions. Define the maps.

$$R_0^+ : L^2(R^n) \rightarrow C^N, \quad \text{by } (R_0^+ u)_j = (u, u_j)$$

$$R_0^- : C^N \rightarrow L^2(R^n), \quad \text{by } (R_0^- \nu) = \sum_{j=1}^N \nu_j u_j$$

For $z \in S(E_0)$, consider

$$\begin{pmatrix} \tilde{H}(\beta) - z & R_0^- \\ R_0^+ & 0 \end{pmatrix} \begin{pmatrix} u \\ u^- \end{pmatrix} = \begin{pmatrix} v \\ v^+ \end{pmatrix} \quad (4.5)$$

with $(v, v^+) \in L^2(R^n) \times C^N$. Let $E' = \text{Span}\{u_1, \dots, u_N\}$, $E'' = E'^\perp$, $\tilde{H}'(\beta) = \tilde{H}(\beta)|_{E''}$. By theorem 3.5, $\tilde{H}'(\beta)$ has no spectrum in $S(E_0)$, one has the following estimate.

$$\|(\tilde{H}'(\beta) - z)^{-1}\| \leq C, \quad z \in S(E_0), \quad 0 < \beta < \beta_0.$$

For $z \in S(E_0)$, then $H(\beta) - z$ is invertible, one has

$$\begin{pmatrix} u \\ u^- \end{pmatrix} = P_0(z) \begin{pmatrix} v \\ v^+ \end{pmatrix},$$

where

$$P_0(z) = \begin{pmatrix} R_0(z) & R_0^+(z) \\ R_0^-(z) & R_0^{-+}(z) \end{pmatrix}$$

$$R_0(z) = (\tilde{H}'(\beta) - z)^{-1}, \quad R_0^{-+}(z) = \text{diag}(z - \mu_j).$$

Let $\rho \in C^\infty(R)$, $\rho(t) = 0$, $t < (E_0 + \delta/2)/\beta$; $\rho(t) = 1$, $t > (E_0 + \delta)/\beta$, and $\psi \in C^\infty(R)$, $\psi(t) = 0$, $t < t_0$, $\psi(t) = 1$, $t > \tilde{t}_0$. Put

$$R^- = \rho R_0^-, \quad R^+ = R_0^+ \rho.$$

Now we consider

$$P(\beta, z) = \begin{pmatrix} p_\theta(x, D) - z & R^- \\ R^+ & 0 \end{pmatrix}$$

Define

$$F(z) = \begin{pmatrix} \rho R_0(z) \psi + (\tilde{p}_\theta(x, D) - z)^{-1} (1 - \psi) & \rho R_0^+(z) \\ R_0^-(z) \psi & R_0^{-+}(z) \end{pmatrix}$$

Then $P(\beta, z) F(z) = I + K(z)$, where $K(z) = (K_{ij}(z))_{2 \times 2}$

$$K_{11} = (p_\theta(x, D) - z) \rho R_0(z) \psi - \psi$$

$$+ \chi_\eta v(x) (\tilde{p}_\theta(x, D) - z)^{-1} (1 - \psi) + \sum (\cdot, \psi u_j) \rho u_j$$

$$K_{12} = (p_\theta(x, D) - z) \rho R_0^+(z) + \rho R_0^+(z) R_0^{-+}(z)$$

$$K_{21} = R_0^+(\rho^2 - 1) R_0(z) \psi + R_0^+(\tilde{p}_\theta(x, D) - z)^{-1} (1 - \psi)$$

$$K_{22} = R_0^+(\rho^2 - 1) R_0^-.$$

In order to estimate the operator $K(z)$, we will need the following lemma.

LEMMA 4.2. – *With the notations above, for any $\varepsilon > 0$, there are $\gamma_0, \gamma_+, \gamma_- > 0$ such that*

$$K_{R_0(z)}(x, y) = O(e^{-\gamma_0 d^\varepsilon(x-y)}), \quad (4.6)$$

$$K_{R_0^+(z)}(x, j) = O(e^{-\gamma_+ d^\varepsilon(x)}), \quad (4.7)$$

$$K_{R_0^-(z)}(j, y) = O(e^{-\gamma_- d^\varepsilon(y)}), \quad (4.8)$$

where the meaning of (4.6) is the same as in the Lemma 4.2.

Proof. – (4.7) and (4.8) follow from (3.8). The proof of (4.6) is similar to the proof of (4.4). ■

LEMMA 4.3. – *One can choose ρ, χ_η, ψ such that the following estimates hold*

$$(p_\theta(x, D) - \tilde{H}(\beta)) R_0(z) \psi = O(\beta^\infty) \quad (4.9)$$

$$(p_\theta(x, D) - z)(\rho - 1) R_0(z) \psi = O(\beta^\infty) \quad (4.10)$$

$$(p_\theta(x, D) - \tilde{H}(\beta)) R_0^+(z) = O(\beta^\infty) \quad (4.11)$$

$$(p_\theta(x, D) - z)(\rho - 1) R_0^+(z) \psi = O(\beta^\infty) \quad (4.12)$$

Proof.

$$(p_\theta(x, D) - \tilde{H}(\beta)) f(x)$$

$$\begin{aligned} &= (p_\theta(x, D) - \tilde{H}(\beta)) (\chi_\eta f)(x) + (p_\theta(x, D) - \tilde{H}(\beta)) (1 - \chi_\eta) f(x) \\ (p_\theta(x, D) - \tilde{H}(\beta)) \chi_\eta f(x) &= U(\theta)^{-1} H(\beta) U(\theta) \chi_\eta f(x) - \tilde{H}(\beta) \chi_\eta f(x) \\ &= U(\theta)^{-1} H(\beta) \chi_\eta f(x) - \tilde{H}(\beta) \chi_\eta f(x) \end{aligned}$$

Therefore $(p_\theta(x, D) - \tilde{H}(\beta)) \chi_\eta f(x) = 0$, for $x_1 > (E_0 + 2\delta)/\beta$. For $x_1 < (E_0 + 2\delta)/\beta$.

$$\begin{aligned} &(p_\theta(x, D) - \tilde{H}(\beta)) \chi_\eta f(x) \\ &= (2\pi)^{-n} \int_{R^{2n}} e^{i(\phi_\theta(x)-y)\cdot\xi} \sqrt{m^2 + \xi^2} \chi_\eta(y) f(y) dy d\xi \\ &\quad - (2\pi)^{-n} \int_{R^{2n}} e^{i(x-y)\cdot\xi} \sqrt{m^2 + \xi^2} \chi_\eta(y) f(y) dy d\xi \\ &= O(\beta^\infty) \end{aligned} \quad (4.13)$$

By (4.6), one can prove that

$$(p_\theta(x, D) - \tilde{H}(\beta))(1 - \chi_\eta) R_0(z) \psi = O(\beta^\infty).$$

From the estimate (4.13) and the estimate above, we get the estimate (4.9). The proofs of the estimates (4.10), (4.11) and (4.12) are similar to this one. ■

Making use of Lemma 4.2 and Lemma 4.3, one can get the following estimate

$$K(z) = O(\beta^\infty), \quad (4.14)$$

in the sense of operator norm on $L^2(R^n) \times C^N$, uniformly for $z \in S(E_0)$.

As a consequence of (4.14), $P(\beta, z)$ is invertible for $\beta > 0$ sufficiently small. We write this inverse, $G(z)$, as

$$G(z) = \begin{pmatrix} R(z) & R^+(z) \\ R^-(z) & R^{-+}(z) \end{pmatrix}.$$

From (4.14) and the formula

$$G(z) = F(z)(I - K(z) + K(z)^2 + \dots)$$

we can prove that

$$R^{-+}(z) = R_0^{-+}(z) + O(\beta^\infty) \quad (4.15)$$

THEOREM 4.4. – *Under the assumption above, $E_0 < 0$ is an eigenvalue of H with multiplicity N . Let $\mu_1(\beta), \dots, \mu_N(\beta)$ be the eigenvalues of $\tilde{H}(\beta)$ in the interval $I(\beta)$. Let $\Gamma(\beta)$ denote the resonances of $H(\beta)$ in $S(E_0)$. Then there exists a bijection $b : \{\mu_1(\beta), \dots, \mu_N(\beta)\} \rightarrow \Gamma(\beta)$ such that:*

$$|b(\mu) - \mu| = O(\beta^\infty), \quad \beta \rightarrow 0_+.$$

Proof. – The proof of this theorem is standard now. The basic point is that $p_\theta(x, D) - z$ is invertible if and only if $R^{-+}(z)$ is bijective on C^N . Then we have the formula:

$$(p_\theta(x, D) - z)^{-1} = R(z) - R^+(z) R^{-+}(z)^{-1} R^-(z) \quad (4.16)$$

(cf. [HS]). By (4.16), we can show that the spectrum of $p_\theta(x, D)$ in $S(E_0)$ is in one-one correspondence with the zero of $\det R^{-+}(z)$, even if we count the multiplicity of these elements. The desired result follows from (4.15). ■

Since the eigenvalues of $\tilde{H}(\beta)$ are real, from Theorem 4.4, one can get easily the following consequence.

COROLLARY 4.5. – *With the notations of Theorem 4.4, for $\beta > 0$ sufficiently small there exist exactly N resonances of $H(\beta)$ in $S(E_0)$. Let $z_1(\beta), \dots, z_N(\beta)$ denote these resonances, repeated according to their multiplicity. Then we have*

$$\begin{aligned} z_j(\beta) &= E_0 + O(\beta), & j &= 1, \dots, N \\ \operatorname{Im} z_j(\beta) &= O(\beta^\infty), & \beta &\rightarrow 0_+ \end{aligned}$$

Note that in the case of non-relativistic Schrödinger operator, the width of resonances in Stark effect is exponential small (cf. [Sig], [Wa1] and [Wa3]).

ACKNOWLEDGEMENT

The author would like to thank professor X. P. Wang for his helpful discussions and suggestions.

REFERENCES

- [AC] J. AGUILLAR, J. M. COMBES, A class of analytic perturbations for one-body Schrödinger Hamiltonians, *Comm. Math. Phys.*, Vol. **22**, 1971, pp. 269-279.
- [BC] E. BALSLEV, J. M. COMBES, Spectral properties of many-body Schrödinger operators with dilation analytic interactions, *Comm. Math. Phys.*, Vol. **22**, 1971, pp. 280-294.
- [CMS] R. CARMONA, W. C. MASTERS, B. SIMON, Relativistic Schrödinger operators: Asymptotic behavior of the eigenfunctions, *J. Funct. Anal.*, Vol. **91**, 1990, pp. 117-142.
- [CM] R. COIFMAN, Y. MEYER, Au-delà des opérateurs pseudodifférentiels, *Astérisque*, Vol. **57**, 1978.
- [CDKS] J. M. COMBES, P. DUCLOS, M. KLEIN, R. SEILER, The shape resonances, *Comm. Math. Phys.*, Vol. **110**, 1987, pp. 215-236.
- [CFKS] H. L. CYCON, R. G. FROESE, W. KIRSCH, B. SIMON, *Schrödinger operators, with application to quantum mechanics and global geometry*, Springer-Verlag, 1986.
- [Fa1] Q. H. FAN, On the essential self-adjointness of pseudodifferential operators, to appear in *Acta Mathematica Sinica*, Vol. **10** (1), 1994.
- [Fa2] Q. H. FAN, On the essential spectrum of pseudodifferential operators I, Nonlinear Analysis and Microlocal Analysis, *Proc. Inter. Confer. at Nankai*, 1991, edited by K. C. Chang etc., World Scientific, Singapore, 1992, pp. 151-159.
- [Fa3] Q. H. FAN, On the essential spectrum of pseudodifferential operators II, to appear in *Chin. Ann. of Math. (Series B)*, 1994.
- [Fo] G. B. FOLLAND, Harmonic analysis in phase space, *Anal. Math. Studies*, Vol. **122**, 1989.
- [HP] B. HELFFER, B. PARISSE, *Comparaison entre la décroissance de fonctions propres pour les opérateurs de Dirac et de Klein-Gordon. Application à l'étude de l'effet tunnel*, preprint, Inst. Fourier, Grenoble, 1993.

- [HS] B. HELFFER, J. SJÖSTRAND, Résonances sur limite semiclassique, *Bull. Soc. Math. France Supp.*, Vol. **114**, 1986.
- [Hu] W. HUNZIKER, Distortion analyticity and molecular resonance curves, *Ann. Inst. Henri Poincaré*, Vol. **45**, 1986, pp. 339-358.
- [K] T. KATO, Perturbation theory for linear operators, Berlin, Heidelberg, New York, Springer, 1966.
- [Sig] I. M. SIGAL, Sharp exponential bounds on resonance states and width of resonances, *Adv. Appl. Math.*, Vol. **9**, 1988, pp. 127-166.
- [S] B. SIMON, Semiclassical analysis of low lying eigenvalue I, non-degenerate minima: Asymptotic expansions, *Ann. Inst. Henri Poincaré*, Vol. **38**, 1983, pp. 295-307.
- [Wa1] X. P. WANG, Bounds on widths of resonances for Stark Hamiltonian, *Acta Math. Sinica* (N. S.), Vol. **6**, 1990, pp. 100-119.
- [Wa2] X. P. WANG, On the magnetic Stark resonances in two dimensional case, *Lecture Notes in Physics*, Vol. **403**, Schrödinger operators, ed. E. Balslev, Springer Verlag, 1992.
- [Wa3] X. P. WANG, Resonances for N-body Schrödinger operators with Stark Hamiltonian, *Ann. Inst. Henri Poincaré*, Vol. **52**, 1990, pp. 1-30.
- [We] R. A. WEDER, Spectral analysis of pseudodifferential operators, *J. Funct. Anal.*, Vol. **20**, 1975, pp. 319-337.

(Manuscript received July 2, 1993;

revised version received December 13, 1993.)