ANNALES DE L'I. H. P., SECTION B

R. W. R. DARLING

Convergence of martingales on manifolds of negative curvature

Annales de l'I. H. P., section B, tome 21, nº 2 (1985), p. 157-175
<http://www.numdam.org/item?id=AIHPB_1985__21_2_157_0>

© Gauthier-Villars, 1985, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section B » (http://www.elsevier.com/locate/anihpb) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Ann. Inst. Henri Poincaré,
Vol. 21, n° 2, 1985, p. 157-175.

Echanges Annales

Probabilités et Statistiques

Convergence of martingales on manifolds of negative curvature

by

R. W. R. DARLING (*)

ABSTRACT. — Let M be a Riemannian manifold whose sectional curvatures are bounded above by a negative constant, and let the stochastic process (X_t) be a martingale with values in M. We give sufficient conditions for (X_t) to have a limit almost surely in the sphere at infinity. We apply the theorem when (X_t) is Brownian motion, proving as a corollary that if the sectional curvatures of M go to minus infinity sufficiently slowly with radial distance, then M admits non-constant bounded harmonic functions. In passing, we obtain an estimate on the passage times of Brownian motion in negatively curved manifolds.

Key-words: Martingale, Brownian motion, negative curvature, Riemannian manifold, stochastic development, escape rates, passage times, bounded harmonic functions.

RÉSUMÉ. — Soit M une variété riemannienne dont les courbures sectionnelles sont majorées par une constante négative, et soit (X_t) une martingale à valeurs dans M. Nous donnons des conditions suffisantes afin que (X_t) ait une direction asymptotique. Nous appliquons le théorème quand (X_t) est le mouvement brownien, et nous prouvons que, si les courbures sectionnelles de M tendent suffisamment lentement vers moins l'infini, alors M admet des fonctions harmoniques bornées et non-constantes. Nous obtenons également une inégalité concernant les temps de passage du mouvement brownien dans les variétés à courbure négative.

(*) Department of Mathematics, University of Southern California, Los Angeles, CA. 90089-1113.

85/02/157/19 /$ 3,90/© Gauthier-Villars

§ 1. INTRODUCTION

Let M be a d-dimensional Riemannian manifold and let Γ denote its Levi-Civita connection. A stochastic process X with values in M is called a Γ-martingale when, roughly speaking, the image of (X_t) under every local Γ-convex function is a local submartingale whenever X_t is in the domain of the function: see Darling [*2*] for details and examples. A convenient method of constructing Γ-martingales is to start with a continuous local martingale on $\mathbb{R}^d$ and « print » it on to M by means of a certain stochastic moving frame: see Darling's thesis [*1*] or Meyer [*10*]. This generalizes the construction of Brownian motion using the orthonormal frame bundle, presented in Elworthy [*5*].

In a previous article [*3*], Darling gave a condition for the almost sure convergence of (X_t), without reference to the curvature of M. The idea was to construct the scalar quadratic variation of (X_t), denoted by $\langle X, X \rangle_t$, by setting

$$d \langle X, X \rangle_t = g_{ij}(X_t) d \langle X^i, X^j \rangle_t$$

where (g_{ij}) is the metric tensor and $(X_t^1, \ldots, X_1^d)$ is the local co-ordinate representation of (X_t). The result is that $X_\infty = \lim_t X_t$ exists almost surely in M, the one-point compactification of M, on the set where $\langle X, X \rangle_\infty < \infty$. Zheng [*15*] proved that on the set where X_∞ exists and lies in M, we have $\langle X, X \rangle_\infty < \infty$ almost surely. These results are put together in the expository paper of Meyer [*11*]. Emery [*6*] attempts to characterize the manifolds on which

$$\{ \lim_{t \to \infty} X_t \text{ exists in M} \} = \{ \langle X, X \rangle_\infty < \infty \}.$$

The present article addresses a different question. We restrict ourselves to manifolds M whose sectional curvatures are bounded above by a negative constant, and which are diffeomorphic to $\mathbb{R}^d$.

We wish to know: when does a Γ-martingale (X_t) converge to a limit in the sphere at infinity (considered as the set of directions of geodesic rays)? Theorem A says that a sufficient condition is that

(1) the process (X_t) has at least two commensurate 'random directions', and

(2) the process (X_t) does not escape too fast from unit balls. These assumptions are made precise in § 3.

In his article of 1975 [*12*], Prat shows that when the sectional curvatures

of M are pinched between two negative constants, Brownian motion on M converges to a limit in the sphere at infinity. To demonstrate that Theorem A is not trivial, we apply it when X is Brownian motion and improve Prat's result, showing that the sectional curvatures may go to minus infinity like $-(\log r(x))^{2-2h}$, where $0 < h < 1$ and $r(x)$ is the distance of x from some fixed point M (Theorem B). As a corollary, we obtain an improvement on Sullivan's result on bounded harmonic functions [*14*]: if the sectional curvatures $K(x)$ of M satisfy

$$-k^2 \geqslant K(x) \geqslant -c^2 (\log r(x))^{2-2h} \quad \text{as} \quad x \to \infty$$

then M admits non-constant bounded harmonic functions.

On the way to Theorem B, we obtain at the end of § 6 a new result about Brownian motion (X_t) when the sectional curvatures of M are pinched between two negative constants: namely, a lower bound on the probability that Brownian motion remains in the ball of radius a up till time t.

§ 2. MARTINGALES IN MANIFOLDS AND ITO'S FORMULA

The idea of martingales on a manifold with a linear connection seems to have originated with Bismut; for an elementary explanation, see Meyer [*10*]. For a more abstract version, see Darling's thesis [*1*]. We shall use the definitions and notations presented in Darling [*3*]. In particular, we shall *not* repeat the definitions of Stratonovitch integration of differential forms, horizontal lifting of a process to the orthonormal frame bundle, or stochastic development. A full account is given in Ikeda and Watanabe [*9*].

Let M be a smooth d-dimensional manifold.

DEFINITION 1. — A *semimartingale* on M will mean a process X with almost surely continuous paths on M, whose image under every C^2 function from M to R is a real-valued semimartingale.

Suppose M has a linear connection Γ on the tangent bundle (for example, the Levi-Civita connection if M is Riemannian). Let ∇ denote covariant differentiation in the cotangent bundle, and for x in M and C^2 functions f on M, define as usual

$$\nabla df(x)(V, W) = (\nabla_V df(x), W), \; V, W \in T_x M$$

(some authors call this Hess $f(x)(V, W)$. In local co-ordinates,

$$\nabla df(x)_{ij} = \nabla df(x)(D_i, D_j) = D_{ij}f(x) - \Gamma_{ij}^k(x)D_k f(x)$$

where the (Γ_{ij}^k) are the Christoffel symbols.

For any semimartingale X on M,

$$\int_0^t \nabla df(X_s)(dX_s, dX_s) \tag{2.1}$$

can be defined by expressing (X_t) in terms of local co-ordinate processes $(X_t^1, \ldots, X_t^n)$ and setting the integrand equal to:

$$\nabla df(X_s)_{ij} d\langle X^i, X^j \rangle_s \tag{2.2}$$

using the angle brackets process of the continuous semimartingales X^i and X^j, whenever X_s is in the domain of the local co-ordinate system.

DEFINITION 2. — (The phrasing is due to Emery [*6*]).

A semimartingale X with values in M is called a Γ-*martingale* if, for all C^2 functions f on M,

$$f(X_t) - 1/2 \int_0^t \nabla df(X_s)(dX_s, dX_s) \tag{2.3}$$

is a real-valued local martingale.

Now assume that M is Riemannian, and Γ is the Levi-Civita connection. Experience in calculating with semimartingales on manifolds shows that it is inefficient to refer constantly to local co-ordinate processes as in expressions (2.1), (2.2) and (2.3). To avoid this, we introduce a process U with values in the orthonormal frame bundle 0(M), called a *horizontal lift* of X to 0(M) through Γ, such that U_t is an isometry from $\mathbb{R}^d$ to the tangent space to M at X_t, with Riemannian inner product. From U we construct a process Z in $\mathbb{R}^d$, called the *stochastic development* of X into $\mathbb{R}^d$. The precise definitions, which are not needed here, are available in Darling [*1*] [*3*]. The crucial relationship of X, U and Z is the Ito formula. For the sake of clarity, take an orthonormal basis $e_1, \ldots, e_d$ for $\mathbb{R}^d$, and write $Z_t = (Z_t^1, \ldots, Z_t^d)$ with respect to this basis. Then $U_s(e_i)$ is a tangent vector at X_s, for each i. Suppose f is a C^2 function from M to $\mathbb{R}$. Then the differential of f is the 1-form

$$df(x) : T_x M \to \mathbb{R}$$

and so the composite process

$$df(X_s) \circ U_s(e_i)$$

is real-valued. Likewise

$$\nabla df(X_s)(U_s(e_i), U_s(e_j))$$

is real-valued. We may now state the

Ito Formula.

For C^2 functions f from M to $\mathbb{R}$,

$$f(X_t) - f(X_0) = \int_0^t (df(X_s) \circ U_s(e_i))dZ_s^i + 1/2 \int_0^t \nabla df(X_s)(U_s(e_i), U_s(e_j))d\langle Z^i, Z^j \rangle_s \qquad (2.4)$$

In fact the use of a basis is not necessary, and we can abbreviate this to:

$$\int_0^t (df(X_s) \circ U_s)dZ_s + 1/2 \int_0^t \nabla df(X_s)(UdZ \otimes UdZ)_s.$$

The virtues of (2.4) are as follows. First, $(U_s(e_1), \ldots, U_s(e_d))$ forms an orthonormal basis of the tangent space X_s, which simplifies many calculations. More importantly, we have:

PROPOSITION 1. — For X to be a Γ-martingale on M, it is necessary and sufficient that any stochastic development Z of X into $\mathbb{R}^d$ is a continuous local martingale; in this case (2.4) is exactly the Doob-Meyer decomposition of the semimartingale $f(X_t) - f(X_0)$.

Proof. — See Darling [*1*], or Meyer [*10*].

Since we refer to Brownian motion on Riemannian manifolds, let us state the following fact:

PROPOSITION 2. — The following there assertions are equivalent:

i) (X_t) is a Markov process on M whose generator is half the Laplacian Δ.

ii) For all C^2 functions f with compact support on M,

$$C_t^f \equiv f(X_t) - f(X_0) - 1/2 \int_0^t \Delta f(X_s)ds$$

is a real-valued martingale on M.

iii) Each stochastic development Z of X into $\mathbb{R}^d$ is a Brownian motion in $\mathbb{R}^d$.

Proof. — The equivalence of (*i*) and (*ii*) follows from the approach to diffusions presented in Stroock and Varadhan [*13*]. Assume (*iii*); then

$$d\langle Z^i, Z^j \rangle_s = \delta^{ij} ds$$

and the left side of (2.4) becomes

$$\int_0^t (df(X_s) \circ U_s) dZ_s + 1/2 \sum_i \int_0^t \nabla df(X_s)(U_s(e_i), U_s(e_i)) ds .$$

Since U_s is an isometry and df is a bounded 1-form the first integrand is bounded; hence the first integral is a martingale. As for the second, since $(U_s(e_i), \ldots, U_s(e_d))$ is an orthonormal basis of the tangent space at X_s, the integrand is $\Delta f(X_s) ds$. Hence

$$f(X_t) - f(X_0) - 1/2 \int_0^t \Delta f(X_s) ds = \text{martingale} .$$

This proves (*ii*). The implication (*ii*) $\Rightarrow$ (*iii*) is easily proved in local coordinates, by the methods presented in Ikeda and Watanabe [*9*].

DÉFINITION 3. — A semimartingale X on a Riemannian manifold M will be called a *Brownian motion* if the assertions of Proposition 2 hold.

§ 3. CONVERGENCE THEOREM

Let $(\Omega, \mathscr{F}, (\mathscr{F}_t), P)$ be a filtered probability space satisfying the usual conditions. All processes will be $(\mathscr{F}_t)$ adapted. Suppose X is a Γ-martingale on a d-dimensional Riemannian manifold (M, g), where Γ is the Levi-Civita connection. Let U be a horizontal lift of X to 0(M) through Γ, and be the corresponding stochastic development of X into $\mathbb{R}^d$.

Our theorem will depend on two main assumptions, (A1) and (A2), about the process X, plus a condition that the sectional curvatures of M are bounded above by a negative constant. The first assumption ensures that X has at least two « random directions » at all times. The second asks that X does not escape too fast from unit balls.

(A1) **Multidirectional assumption.**

This assumption is stated in terms of the stochastic development Z of X into $\mathbb{R}^d$. Actually Z is not unique, because different choices of initial frame U_0

give different processes Z. However if the following assumptions hold for *any* Z, they will hold for *all* stochastic developments Z.

First we make two mild technical assumptions:

$$\mathbb{E}\,|\,Z_t\,|^2 < \infty \quad \text{for all} \quad t \tag{3.1}$$

$$d\langle Z^i, Z^j\rangle_t = A^{ij}(t)dt \tag{3.2}$$

where $\langle Z^i, Z^j\rangle_t$ is the angle-brackets process of Z^i and Z^j, and $(A^{ij}(t))$ is some symmetric $d \times d$ matrix-valued process.

Let $\lambda_1(t) \geqslant \lambda_2(t) \geqslant \ldots \geqslant \lambda_d(t) \geqslant 0$ be the eigenvalues of $A^{ij}(t)$. The crucial assumption is that for some constant c_1 and c_2

$$0 < c_1 < \lambda_1(t) < c_2\lambda_2(t) \text{ a. s.,} \quad \text{for all } t \tag{3.3}$$

Among other things, this ensures that (X_t) moves in at least two 'random directions'. Observe that when (X_t) is Brownian motion on M, then (Z_t) is a Brownian motion in $\mathbb{R}^d$, and all the λ_i are identically 1.

(A2) **Moderated escape rates.**

We need to have a lower estimate for the rate at which (X_t) can escape from unit balls in M. We suppose henceforward that for some $k > 0$, the sectional curvatures of M are bounded *above* by $-k^2 < 0$. Define $c_3 = k(3 - 2\sqrt{2})/c_2$, and define ε_0 to be the value of ε which maximizes

$$h(\varepsilon) = \frac{1}{2}\varepsilon e^{-\varepsilon}(c_3 - \varepsilon)c_1\,; \quad 0 \leqslant \varepsilon \leqslant c_3 \wedge k\,.$$

Define

$$c_4 = c_2 h(\varepsilon_0) \tag{3.4}$$

Let $\delta = 1/c_4$, and define bounded stopping-times as follows:

$$u(0) = 0$$

$$u(n) = \inf\{\, t > u(n-1) \colon d(X_{u(n-1)}, X_t) \geqslant 1 \,\} \wedge (u(n-1) + \delta)\,. \tag{3.5}$$

Note that necessarily $u(n) \leqslant n\delta$. For $n = 0, 1, 2, \ldots$, define $H(n)$ to be the event that (X_t) leaves the unit ball centered on $X_{u(n-1)}$ sometime before $(u(n-1) + \delta)$; more formally,

$$\begin{aligned} H(n) &= \{\, u(n) < u(n-1) + \delta \,\} \\ &= \{\, d(X_{u(n-1)}, X_t) \geqslant 1, \text{ some } u(n-1) < t < u(n-1) + \delta \,\} \end{aligned} \tag{3.6}$$

The assumption is that there exist real numbers $(a(n))$, $n = 1, 2, \ldots$ such that

$$P(H(n) \mid \mathscr{F}_{u(n-1)}) \leqslant a(n)$$

with

$$\sum_{N=1}^{\infty}\left(\prod_{n=1}^{N} a(n)\right) < \infty \tag{3.7}$$

Intuitively speaking, this says that the process (X_t) is not allowed to escape from unit balls too fast.

The theorem below refers to connected manifolds M whose sectional curvatures are bounded above by a negative constant. The Cartan-Hadamard theorem states that for such M, the exponential map at any point x of M provides a diffeomorphism $\exp_x$ from $\mathbb{R}^d$ to M where $d = \dim(M)$. Let us take polar co-ordinates in $\mathbb{R}^d$, so that we now have a diffeomorphism

$$\exp_x : (0, \infty) \times S^{d-1} \to M/\{x\}.$$

The *geometric compactification* of M is the space $M \cup S^{d-1}$ obtained by identifying M with $[0, \infty) \times S^{d-1}$, and adjoining $\{\infty\} \times S^{d-1}$. We say that a sequence $(r(n), \theta(n))$ in M converges to (∞, θ) in $M \cup S^{d-1}$ if $r(n)$ tends to infinity and $\theta(n)$ tends to θ in S^{d-1}. We speak of $\{\infty\} \times S^{d-1}$ as the *sphere at infinity*.

THEOREM A. — Let (M, g) be a simply-connected Riemannian manifold with sectional curvatures bounded above by $-k^2 < 0$. Let (X_t) be a Γ-martingale on M. Suppose that the multidirectional assumption (A1), and the escape rate assumption (A2) are in force. Then we have

i) $r(X_t) \to \infty$ as $t \to \infty$.

ii) $X_\infty = \lim_{t\to\infty} X_t$ exists almost surely in $M \cup S^{d-1}$, and X_∞ lies in S^{d-1} almost surely.

§ 4. PROOF OF THE CONVERGENCE THEOREM

Step 1.

Let $(u(n))$ be the stopping-times defined in § 3.5. The Ito formula (2.4) may be applied to $e^{-\varepsilon r(x)}$ for $\varepsilon > 0$, since (X_t) has zero probability of hitting x_0: thus

$$\begin{aligned} e^{-\varepsilon r(X_{u(n)})} - e^{-\varepsilon r(X_{u(n-1)})} &= \int_{u(n-1)}^{u(n)} (-\varepsilon e^{-\varepsilon r(X_s)} dr(X_s) \circ U_s) dZ_s \\ &\quad - \frac{1}{2}\int_{u(n-1)}^{u(n)} e^{-\varepsilon r(X_s)} \varepsilon(\nabla dr(X_s) - \varepsilon dr \otimes dr(X_s))(U dZ \otimes U dZ)_s \end{aligned} \tag{4.1}$$

Since (X_t) is a Γ-martingale, it follows from Proposition 1 that (Z_t) is a local martingale; by (A1), Z_t is in L^2 and so the first integral, whose integrand is bounded, is a martingale. The second integral can be written, using (A1), as

$$\frac{1}{2}\int_{u(n-1)}^{u(n)} e^{-\varepsilon r(X_s)}\varepsilon\left\{\sum_{i,j}(\nabla dr(X_s)(U_s(e_i), U_s(e_j)) - \varepsilon dr(U_s(e_i))dr(U_s(e_j)))A^{ij}(s)\right\} ds .$$

The right choice of orthonormal basis (e_i) diagonalizes $(A^{ij}(s))$, so that

$$\{\ \dots\ \} = \sum_{i=1}^{d}(\nabla dr(X_s)(U_s(e_i), U_s(e_i)) - \varepsilon(dr(U_s(e_i)))^2)\lambda_i(s) \tag{4.2}$$

Step 2. Hessian comparison theorem.

We shall first perform a small algebraic calculation. Let $(e_1, \dots, e_d)$ denote an orthonormal basis of $\mathbb{R}^d$, and let

$$v = (a_1 e_1 + \dots + a_n e_n), \sum_{i=1}^{d} a_i^2 = 1$$

be a unit vector; define

$$\hat{e}_i = e_i(1 - v.e_i) = (1 - a_i)e_i$$
$$f(a_1, \dots, a_n) = |\hat{e}_1|^2 + |\hat{e}_2|^2 = (1 - a_1)^2 + (1 - a_2)^2 .$$

LEMMA. — $\min\{f(a_1, \dots, a_n) : a_1^2 + \dots + a_n^2 = 1\} = 3 - 2\sqrt{2}$.

Proof. — The minimum is attained when $a_1 = a_2 = 2^{-1/2}$.

We return to the integral obtained in step 1. A (random) tangent vector such as $U_s(e_i)$ can be decomposed into a radial part and an orthogonal part; denote the latter by $U_s(e_i)^{\perp}$. Greene and Wu (*7*, p. 21] show that:

$$\nabla dr(U_s(e_i), U_s(e_i)) = \nabla dr(U_s(e_i)^{\perp}, U_s(e_i)^{\perp}) \geqslant k \| U_s(e_i)^{\perp} \|^2$$

by the Hessian comparison theorem; we have used the fact that the Hessian of the distance function in M is greater (in a sense described precisely by Greene and Wu [*7*]) than the Hessian of the distance function $r_k(.)$ in a manifold of constant sectional curvature $-k^2$, and

$$\nabla dr_k(u, u) = k \coth(kr_k) \| u^{\perp} \|^2 \geqslant k \| u^{\perp} \|^2 .$$

Continuing from the end of step 1,

$$\sum_{i=1}^{d} \nabla dr(X_s)(U_s(e_i), U_s(e_i))\lambda_i(s) \geqslant \lambda_2(s) \sum_{i=1}^{2} \nabla dr(X_s)(U_s(e_i)^{\perp}, U_s(e_i)^{\perp})$$

$$\geqslant \lambda_2(s)k \sum_{i=1}^{2} \| U_s(e_i)^{\perp} \|^2 \geqslant \lambda_2(s)k(3-2\sqrt{2})$$

by the lemma, using the fact that $U_s(e_i)$ is of unit length. Using (A1) this is

$$\geqslant \lambda_1(s)k(3 - 2\sqrt{2})/c_2$$

or in the notation of (A2).

$$\geqslant c_3\lambda_1(s).$$

On the other hand,

$$\sum_{i=1}^{d} (dr(U_s(e_i)))^2\lambda_i(s) \leqslant \lambda_1(s),$$

Since

$$\sum_{i=1}^{d} (dr(U_s(e_i)))^2 = \| \operatorname{grad} r \|^2 = 1.$$

The conclusion about formula (4.2) is that

$$\{ \dots \} \geqslant (c_3 - \varepsilon)\lambda_1(s) > (c_3 - \varepsilon)c_1 > 0 \tag{4.3}$$

by (A1), assuming $0 < \varepsilon < c_3$.

Step 3. Applying (A2).

Continuing from (4.1) for $0 < \varepsilon < c_3$

$$\mathbb{E}[e^{-\varepsilon r(X_{u(n)})} - e^{-\varepsilon r(X_{u(n-1)})} \mid \mathscr{F}_{u(n-1)}] \leqslant -\frac{1}{2}\varepsilon_1(c_3-\varepsilon)\mathbb{E}\left[\int_{u(n-1)}^{u(n)} e^{-\varepsilon r(X_s)}ds \mid \mathscr{F}_{u(n-1)}\right]$$

using (4.3) and the fact that the first integral on the right side of (4.1) is a martingale. By the definition of $u(n)$ in (3.5):

$$r(X_s) \leqslant r(X_{u(n-1)}) + 1, \qquad u(n-1) \leqslant s < u(n)$$

and hence

$$-e^{-\varepsilon r(X_s)} \leqslant -e^{-\varepsilon}e^{-\varepsilon r(X_{u(n-1)})}, \qquad u(n-1) \leqslant s < u(n).$$

If the constant c_4 is as in formula (3.4) we have:

$$\mathbb{E}[e^{-\varepsilon r(X_{u(n)})} - e^{-\varepsilon r(X_{u(n-1)})} \mid \mathscr{F}_{u(n-1)}] \leqslant -c_4 e^{-\varepsilon r(X_{u(n-1)})} \mathbb{E}[u(n) - u(n-1) \mid \mathscr{F}_{u(n-1)}]$$

where $\varepsilon = \varepsilon_0$ as in (3.4). (Note that $0 < \varepsilon_0 < c_3$). Hence

$$\mathbb{E}[e^{-\varepsilon r(X_{u(n)})} \mid \mathscr{F}_{u(n-1)}] \leqslant \{1 - c_4 \mathrm{E}[u(n) - u(n-1) \mid \mathscr{F}_{u(n-1)}]\} e^{-\varepsilon r(X_{u(n-1)})} \tag{4.4}$$

Refer back to the definition of $\mathrm{H}(n)$ in (3.6). Evidently

$$u(n) - u(n-1) \geqslant \delta 1_{\mathrm{H}(n)^c}.$$

By definition of δ, $c_4\delta = 1$; therefore

$$\begin{aligned} 1 - c_4 \mathbb{E}[u(n) - u(n-1) \mid \mathscr{F}_{u(n-1)}] &\leqslant 1 - c_4 \delta \mathbb{E}[1_{\mathrm{H}(n)^c} \mid \mathscr{F}_{u(n-1)}] \\ &= \mathrm{E}[1_{\mathrm{H}(n)} \mid \mathscr{F}_{u(n-1)}] \leqslant a(n). \end{aligned}$$

Therefore (4.4) says that

$$\mathbb{E}[e^{-\varepsilon r(X_{u(n)})} \mid \mathscr{F}_{u(n-1)}] \leqslant a(n) \mathbb{E}[e^{-\varepsilon r(X_{u(n-1)})} \mid \mathscr{F}_{u(n-1)}] = a(n) e^{-\varepsilon r(X_{u(n-1)})}.$$

By induction,

$$\mathrm{E}[e^{-\varepsilon r(X_{u(n)})}] \leqslant \prod_{j=1}^{n} a(j) e^{-\varepsilon r(X_0)}.$$

Therefore by assumption (A_2), equation (3.7),

$$\sum_{n=0}^{\infty} \mathbb{E}[e^{-\varepsilon r(X_{u(n)})}] < \infty \tag{4.5}$$

This proves that $r(X_{u(n)})$ goes to infinity almost surely as n tends to infinity. Therefore $r(X_t)$ tends to infinity as t tends to infinity, proving (i).

Step 4. Angular convergence.

For points x and y in M, let $\theta(x, y)$ denote the angle in the tangent space at m between the geodesics from m to x and from m to y respectively. Since we are assuming $\varepsilon \leqslant k$, geometric calculation shows that for some $c_5 > 0$,

$$\theta(x, y) \leqslant c_5 e^{-\varepsilon r(x)}, \qquad \text{provided} \quad d(x, y) < 1$$

(see Prat [*12*]). Consequently, if $u(n) \leqslant t \leqslant u(n+1)$,

$$\theta(X_{u(n)}, X_t) \leqslant c_5 e^{-\varepsilon r(X_{u(n)})},$$

and

$$\theta(X_{u(n)}, X_{u(m)}) \leqslant c_5 \sum_{j=n}^{m-1} e^{-\varepsilon r(X_{u(j)})}$$

which tends to zero as n, m tend to infinity, by (4.5). These two assertions prove that almost surely,

$$\theta(X_s, X_t) \to 0 \quad \text{as} \quad s, t \to \infty .$$

Together with the fact that $r(X_t)$ tends to infinity as t tends to infinity, this shows that there exists a random variable X_∞ with values in S^{d-1}, representing a limiting direction for (X_t), in the sense that

$\theta(X_t, X_\infty) \to 0$ as $t \to \infty$, almost surely.

§ 5. SOME ESTIMATES ON BROWNIAN MOTION IN R^m

We shall begin by deriving anew a classical estimate on m-dimensional Brownian motion. First we need some notation and a Lemma on 1-dimensional Brownian motion.

Notation. — If (Y_t) is a stochastic process taking non-negative real values, then define

$$Y_t^* = \sup_{0 \leqslant s \leqslant t} Y_s .$$

LEMMA 1. — Let (B_t) denote 1-dimensional Brownian motion, started at zero. Then for all $c > 0$

$$P(|B_t|^* < c) \geqslant (32t/\pi c^2)^{0.5}(1 - \exp(-c^2/2t)) \tag{5.1}$$

Proof. — Define

$$M_c = \inf\{ t : B_t = c \} .$$

Then

$$\begin{aligned} P(|B_t|^* \geqslant c) &= P(\{ M_c \leqslant t \} \cup \{ M_{-c} \leqslant t \}) \\ &= P(M_c \leqslant t) + P(M_{-c} \leqslant t) - P(\{ M_c \leqslant t \} \cap \{ M_{-c} \leqslant t \}) < 2P(M_c \leqslant t) \end{aligned}$$

using the symmetry of the distribution of Brownian motion about zero. The reflection principle (Ito and McKean [*16*, p. 26]) shows that

$$P(M_c \leqslant t) = 2 \int_c^\infty (2\pi t)^{-0.5} \exp(-u^2/2t) du .$$

Hence

$$\mathrm{P}(|\mathrm{B}_t|^* < c) \geqslant 1 - 4\int_c^\infty (2\pi t)^{-0.5} \exp(-u^2/2t)du$$

$$= \int_{-c}^{c} 2(2\pi t)^{-0.5} \exp(-u^2/2t)du \tag{5.2}$$

When u is in the interval $[0, c]$, $u^2 \leqslant uc$; therefore

$$\int_{-c}^{c} \exp(-u^2/2t)du \geqslant 2\int_0^c \exp(-uc/2t)du = \frac{4t}{c}(1-\exp(-c^2/2t)) \tag{5.3}$$

The combination of (5.2) and (5.3) give the result.

LEMMA 2. — Let m be a positive integer, and let $(\mathrm{W}_t) = (\mathrm{W}_t^1, \ldots, \mathrm{W}_t^m)$ denote Brownian motion in $\mathbb{R}^m$. Then

$$\mathrm{P}(|\mathrm{W}_t|^* < a) \geqslant (32t/\pi)^{m/2} a^{-m} e^{m\ln(m)/2}(1 - \exp(-a^2/2tm))^m \tag{5.4}$$

Proof.

$$\{|\mathrm{W}_t|^* < a\} \supseteq \bigcap_{j=1}^{m} \{|\mathrm{W}_t^j|^* < a/\sqrt{m}\},$$

because for every sample path in the set on the right,

$$|\mathrm{W}_t|^2 = \sum_{j=1}^{m} |\mathrm{W}_t^j|^2 \leqslant m(a/\sqrt{m})^2 = a^2.$$

By the independence of the co-ordinate processes,

$$\mathrm{P}(|\mathrm{W}_t|^* < a) \geqslant \prod_{j=1}^{m} \mathrm{P}(|\mathrm{W}_t^j|^* < a/\sqrt{m}).$$

By the previous Lemma, with $c = a/\sqrt{m}$, this is

$$\geqslant \{(32tm/\pi a^2)^{0.5}(1 - \exp(-a^2/2tm))\}^m$$
$$\geqslant (32t/\pi)^{m/2} a^{-m} e^{m\ln(m)/2}(1 - \exp(-a^2/2tm))^m$$

§ 6. ESTIMATES ON THE RADIAL PROCESS OF BROWNIAN MOTION ON M

PROPOSITION 2. — Suppose M is a d-dimensional Riemannian manifold with sectional curvatures $K(x)$ satisfying

$$-q^2 \leqslant K(x) \leqslant -k^2 < 0, \qquad x \text{ in } M$$

for constants k and q. Let (X_t) be a Brownian motion on M and let

$$R_t = d(X_0, X_t), \qquad R_t^* = \sup\{ R_s : 0 \leqslant s \leqslant t \}.$$

Then

$$P(R_t^* < a) \geqslant \left(\frac{32t}{\pi}\right)^{m/2} a^{-m} e^{m \ln(m)/2} \left(1 - \exp\left(\frac{-a^2}{2tm}\right)\right)^m \tag{6.1}$$

provided $m \geqslant d + (d-1)aq$.

In order to prove the proposition above, let M′ be a d-dimensional manifold of constant sectional curvature $-q^2$. Take and R^d-valued Brownian motion $(\overline{W}_t)$ and points x_0' in M′, x_0 in M, and solve the canonical stochastic dynamical systems of M′ and M to obtain Brownian motions (X_t') and (X_t) respectively, both driven by $(\overline{W}_t)$, with $X_0' = x_0'$ and $X_0 = x_0$. Define real-valued processes

$$R_t' = d(x_0', X_t'), \qquad R_t = d(x_0, X_t) \tag{6.2}$$

Let $r(y) = d(x_0, y)$ for y in M, and define

$$b(x) = \frac{1}{2}(d-1)q \coth(qx), \qquad x \text{ in } R. \tag{6.3}$$

By appling Ito's formula to the distance function $r(.)$ and to its counterpart in M′, Prat [*12*] shows that there exist a real-valued Brownian motions (B_t) and (B_t') on the same probability space such that

$$R_t = B_t + \int_0^t \frac{1}{2} \Delta_M r(X_s) ds \tag{6.4}$$

and

$$R_t' = B_t' + \int_0^t b(R_s') ds \tag{6.5}$$

In actual fact, the function $b(.)$ is half the Laplacian of the distance function in the manifold of constant curvature $-q^2$. Now define two more

real-valued processes to be the unique strong solutions, started at zero, to the stochastic integral equations:

$$S_t = B'_t + \int_0^t \frac{1}{2}(d-1)\left(q + \frac{1}{S_u}\right)du \tag{6.6}$$

$$Y_t^{(m)} = B'_t + \int_0^t \frac{1}{2}\left(\frac{m-1}{Y_s^{(m)}}\right)ds \tag{6.7}$$

where m is some positive integer $\geqslant 2$. It is well known—see Ikeda and Watanabe [*9*, p. 223]—that the law of $(Y_t^{(m)})$ is that of the distance of Brownian motion in $\mathbb{R}^m$ from the origin, known as the Bessel process of order m.

LEMMA 3. — *i*) $R'_t \leqslant S_t$ a.s., and $P(R_t < a) \geqslant P(R'_t < a)$ for all $a \geqslant 0$, $t \geqslant 0$. Assume that $m \geqslant d + (d-1)aq$, for some $a > 0$. Then

ii) On the set $\{ Y_t^{(m)*} \leqslant a \}$, $R'_s \leqslant Y_s^{(m)}$ a.s., for all $0 \leqslant s \leqslant t$.

iii) $P(R_t^* < a) \geqslant P(|W_t|^* < a)$

where (W_t) is Brownian motion in $\mathbb{R}^m$.

Remark. — The comparison of (R_t) and (R'_t) is due to Debiard, Gaveau and Mazet [*4*]. The idea of using (S_t) and $(Y_t^{(m)})$ was suggested to the author by S. R. S. Varadhan (Courant Institute).

Proof. — The Laplacian comparison theorem, found for example in Greene and Wu [*7*], shows that if $b(.)$ is the function defined in (6.3),

$$b(r(y)) \geqslant \frac{1}{2}\Delta_M r(y), \quad y \quad in \quad M.$$

It is a simple calculation to verify that

$$\frac{1}{2}(d-1)\left(q + \frac{1}{x}\right) = \frac{1}{2}(d-1)q\left(1 + \frac{1}{qx}\right) \geqslant b(x).$$

Part (*i*) of the Lemma now follows from the comparison theorem in Ikeda and Watanabe [*9*, p. 352]. The same theorem is used to prove part (*ii*), noting that

$$(m-1) \geqslant (d-1)(aq+1),$$

and

$$0 < y \leqslant a \Rightarrow (m-1) \geqslant (d-1)(yq+1) \Rightarrow \frac{m-1}{y} \geqslant \left(d-1\right)\left(q + \frac{1}{y}\right)$$

which proves $S_u \leqslant Y_u^{(m)}$ on $0 \leqslant u \leqslant t$, on the set $\{ Y_t^{(m)*} \leqslant a \}$. Combined with the result of (*i*), this gives part (*ii*).

As for part (*iii*), let

$$M_a^{(m)} = \inf \{ t : Y_t^{(m)} = a \}, \qquad N_a = \inf \{ t : R'_t = a \}.$$

Part (*ii*) implies that

$$s < M_a^{(m)} \Rightarrow R'_s \leqslant Y_s^{(m)} < a \Rightarrow s < N_a .$$

Hence

$$P(M_a^{(m)} > s) \leqslant P(N_a > s)$$

or equivalently

$$P(R_s^* < a) \geqslant P(R_s'^* < a) \geqslant P(Y_s^{(m)*} < a) = P(|W_s|^* < a).$$

From (*iii*) and Lemma 2 of § 5, Proposition 2 now follows.

§ 7. BROWNIAN MOTION ON A MANIFOLD WITH NEGATIVE CURVATURE GOING TO $-\infty$

Ichihara [*8*] has proved that for Riemannian manifolds M whose Ricci curvature at distance r along any minimal geodesic is bounded below by $-ar^2 - b$ (here a and b are some positive constants) the Brownian motion (X_t) on M is non-explosive, meaning that

$$P(X_t \in M) = 1, \qquad 0 \leqslant t < \infty .$$

It is well-known, and easy to prove, that if the sectional curvatures of M are bounded above by a negative constant, then the distance process

$$R_t = d(X_0, X_t)$$

tends to infinity almost surely as t tends to infinity. An open problem is to characterize those manifolds M of negative curvature on which Brownian motion has an angular limit; in other words, those for which X_∞ $(= \lim_{t \to \infty} X_t)$ exists in S^{d-1}. (See section § 3 for explanation.). A sufficient condition, given by Prat [*12*], is that the sectional curvatures of M are pinched between two negative constants. The following extension of Prat's result is included as an example of the application of Theorem A of § 3.

THEOREM B. — Suppose M is a d-dimensional Riemannian manifold containing some compact c_0 such that the sectional curvatures $K(x)$ at points x outside c_0 satisfy

$$-c^2 (\ln (r(x)))^{2-2h} \leqslant K(x) < -k^2 < 0 \tag{7.1}$$

for some $0 < h < 1$ and some constant c; here $r(x)$ is the distance of x from some fixed point x_0 in c_0. Then Brownian motion (X_t) on M has a limit almost surely on the sphere at infinity. In other words

$$X_\infty = \lim_{t \to \infty} X_t \text{ exists almost surely in } S^{d-1}.$$

COROLLARY. — A manifold with curvatures satisfying (7.1) admits non-constant bounded harmonic functions.

Proof of the Corollary. — Let h be any bounded, continuous function on S^{d-1}. Define for x in M

$$f(x) = \mathbb{E}_x[h(X_\infty)],$$

where $\mathbb{E}_x[\ldots]$ means that $X_0 = x$. Then f is a bounded harmonic function on M.

Proof of Theorem B. — For the sake of simplicity, let the Brownian motion (X_t) begin at $X_0 = x_0$, and define $r(x) = d(x_0, x)$ (the distance from x_0 to x in M). Notice that

$$u(n-1) \leqslant t \leqslant u(n) \Rightarrow r(X_t) \leqslant \sum_{j=0}^{n-2} d(X_{u(j)}, X_{u(j+1)}) + d(X_{u(n-1)}, X_{u(n)}).$$

From the definition of the stopping times $(u(n))$, it follows that $r(X_t) \leqslant n$ when $u(n-1) \leqslant t \leqslant u(n)$. Define for each n a real-valued process

$$R_s^n = d(X_{u(n-1)}, X_{u(n-1)+s}), \quad \text{stopped at } s = u(n) - u(n-1).$$

On the time interval $[0, u(n)]$, the Brownian motion (X_t) lives inside the ball of radius n in M; therefore it experiences sectional curvatures $K(.)$ bounded by

$$-q(n)^2 \leqslant K(x) \leqslant -k^2$$

where

$$q(n) = c(\ln(n))^{1-h} \tag{7.2}$$

Therefore the Corollary at the end of the previous section applies to the process (R_s^n), for each n. Recall the definition of $H(n)$ in § 3, assumption (A2), equation (3.6):

$$\begin{aligned} 1 - P(H(n)) &= P(u(n) = u(n-1) + \delta) = P(R_\delta^{n*} < 1) \\ &\geqslant \left(\frac{32\delta}{\pi}\right)^{m/2} e^{m \ln(m)/2}(1 - \exp(-1/2\delta m))^m \end{aligned} \tag{7.3}$$

where $m = m(n) = dq(n)$. Let $\gamma = (32\delta/\pi)^{1/2}$ and let $\lambda = \exp(-1/2\delta)$. It can be proved by calculus that for $0 < h < 1$

$$1 - \lambda^h \geqslant h(1 - \lambda).$$

Taking $h = 1/m$, it follows that

$$(1 - \lambda^{1/m})^m \geqslant e^{-m \ln(m) - m \ln(b)}, \qquad b = 1/(1 - \lambda).$$

Putting this into (7.3) gives:

$$1 - \mathrm{P}(\mathrm{H}(n)) \geqslant \exp(-m \ln(m)/2 + m \ln(\gamma) - m \ln(b)) \tag{7.4}$$

The fact that Brownian motion has independent increments implies that

$$\mathrm{P}(\mathrm{H}(n) \mid \mathscr{F}_{u(n-1)}) = \mathrm{P}(\mathrm{H}(n)).$$

Consider assumption (A2), equation (3.7) again; notice that

$$\prod_{n=1}^{\mathrm{N}} \left(1 - \frac{2}{n+3}\right) = \prod_{n=1}^{\mathrm{N}} \left(\frac{n+1}{n+3}\right) = \frac{2.3}{(\mathrm{N}+2)(\mathrm{N}+3)} \leqslant \frac{6}{\mathrm{N}^2}.$$

Hence if $a(n) = 1 - 2/(n+3)$, assumption (A2) holds. Hence it suffices to show that

$$\mathrm{P}(\mathrm{H}(n)) \leqslant 1 - \frac{2}{n+3},$$

or equivalently,

$$1 - \mathrm{P}(\mathrm{H}(n)) \geqslant \frac{2}{n+3}.$$

Using (7.4), it will be enough to verify that

$$m \ln(m)/2 - m(\ln(\gamma) - \ln(b)) \leqslant -\ln 2 - \ln(n+3) \tag{7.5}$$

for all sufficiently large n. Now

$$m = dq(n) = cd(\ln(n))^{1-h}$$

So

$$m \ln(m) = 0(\ln(n))^{1-h} \ln(\ln(n)).$$

Hence

$$\frac{m \ln(m)}{2 \ln(n+3)} = 0(\ln(\ln(n))/\ln(n))^h.$$

Moreover

$$\frac{m}{\ln(n+3)} = 0((\ln(n))^{-h}).$$

It follows that

$$\frac{m \ln (m)/2 + m (\ln (\gamma) - \ln (b))}{\ln (n + 3)} \to 0 \quad \text{as} \quad n \to \infty$$

and this verifies (7.5). The proof is complete.

ACKNOWLEDGMENTS

The author wishes to thank Paul Yang (U. S. C.) and Richard Durrett (U. C. L. A.) for their interest in this work.

REFERENCES

[1] R. W. R. DARLING, Martingales on manifolds and geometric Ito calculus. *Ph. D. Thesis*, University of Warwick, 1982.

[2] R. W. R. DARLING, Martingales in manifolds—definition, examples and béhaviour under maps. *Springer Lecture Notes in Math.*, t. **921**, 1982, p. 217-236.

[3] R. W. R. DARLING, Convergence of martingales on a Riemannian manifold. *Publ. R. I. M. S., Kyoto University*, t. **19**, 1983, p. 753-763.

[4] A. DEBIARD, B. GAVEAU and E. MAZET, Temps de sortie des boules normales et minoration locale de λ_1. *C. R. Acad. Sci. Paris*, t. **278 A**, 1974, p. 795-798.

[5] K. D. ELWORTHY, Stochastic differential equations on manifolds. *London Math. Soc. Lecture Notes*, t. **70**, Cambridge U. P., 1982.

[6] M. EMERY, *Convergence des martingales dans les variétés*. Actes du Colloque Laurent Schwartz (Astérisque), 1984, to appear.

[7] R. E. GREENE and H. WU, Function theory on manifolds which possess a pole. *Springer Lecture Notes in Math.*, t. **699**, 1979.

[8] K. ICHIHARA, Curvature, goedesics and the Brownian motion on a Riemannian manifold. II: explosion properties, *Nagoya Math. J.*, t. **87**, 1982, p. 115-125.

[9] N. IKEDA and S. WATANABE, *Stochastic differential equations and diffusion processes*. North Holland, 1981.

[10] P. A. MEYER, Géométrie stochastique sans larmes. Sém. de Probabilités XV, 1979/1980, *Springer Lecture Notes in Math.*, t. **850**, 1981, p. 44-102.

[11] P. A. MEYER, Le théorème de convergence des martingales dans les variétés riemanniennes. Sém. Prob. XVII, *Springer Lecture Notes in Math.*, t. **986**, 1983, p. 187-193.

[12] J.-J. PRAT, Étude asymptotique et convergence angulaire du mouvement brownien sur une variété à courbure négative. *C. R. A. S.*, t. **280**, ser. A, 1975, p. 1539-1542.

[13] D. W. STROOCK and S. R. S. VARADHAN, *Multidimensional diffusion processes*. Springer-Verlag, 1979.

[14] D. SULLIVAN, The Dirichlet problem at infinity for a negatively curved manifold. *J. Differential Geometry*, t. **18**, 1983, p. 723-732.

[15] W. A. ZHENG, Sur la convergence des martingales dans une variété riemannienne. *Zeit. fur Wahrsch.*, t. **63**, 1983, p. 511-515.

[16] K. ITO and H. P. MCKEAN, *Diffusion processes and their sample paths*. Springer, Berlin, 1965.

(Manuscrit reçu le 16 juillet 1984)