

# ANNALES DE L'I. H. P., SECTION C

EUGÈNE GUTKIN

## **Conservation laws for the nonlinear Schrödinger equation**

*Annales de l'I. H. P., section C*, tome 2, n° 1 (1985), p. 67-74

[http://www.numdam.org/item?id=AIHPC\\_1985\\_\\_2\\_1\\_67\\_0](http://www.numdam.org/item?id=AIHPC_1985__2_1_67_0)

© Gauthier-Villars, 1985, tous droits réservés.

L'accès aux archives de la revue « *Annales de l'I. H. P., section C* » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Conservation laws for the nonlinear Schrödinger equation

by

Eugène GUTKIN (\*)

Department of Mathematics, Columbia University, New York  
and MSRI, 2223 Fulton Street, Berkeley, CA 94720 (\*\*)

---

ABSTRACT. — We propose a method of calculating the operator densities  $\hat{h}_n$   $n = 0, 1, \dots$  of the conservation laws for the quantum nonlinear Schrödinger equation. It follows from the method that  $\hat{h}_n$  are polynomials in fields and their derivatives and in the coupling constant. The densities  $\hat{h}_n$   $n \leq 4$  are explicitly calculated. Comparison with the integral densities  $b_n$   $n = 0, 1, \dots$  for the classical nonlinear Schrödinger equation shows that the correspondence between  $\hat{h}_n$  and  $b_n$  breaks down after  $n = 3$ .

RÉSUMÉ. — On propose une méthode pour calculer les densités opératoires  $\hat{h}_n$   $n = 0, 1, \dots$  pour les intégrales de l'équation de Schrödinger non linéaire quantique. Il s'ensuit que les  $\hat{h}_n$  sont des fonctions polynomiales des champs, de leurs dérivées et de la constante de couplage. Les densités  $\hat{h}_n$ ,  $n \leq 4$ , sont calculées explicitement. En les comparant avec les densités intégrales  $b_n$   $n = 0, 1, \dots$  pour l'équation de Schrödinger non linéaire classique, on voit que la correspondance entre  $b_n$  et  $\hat{h}_n$  n'est plus valable pour  $n > 3$ .

---

### 1. INTRODUCTION

We consider the quantum nonlinear Schrödinger equation (NLSE) in  $1 + 1$  space-time dimensions

$$i\Psi_t = -\Psi_{xx} + 2c\Psi^\dagger\Psi^2. \quad (1.1)$$

---

(\*) Research supported in part by NSF Grant # MCS 8101739.

(\*\*) Present address: Mathematics, University of Southern California, Los Angeles, CA 90089, USA.

Its Hamiltonian

$$\hat{H}_2 = - \int dx (\Psi^\dagger \Psi_{xx} - c \Psi^\dagger \Psi^2) \quad (1.2)$$

is the second quantized form of the many body Hamiltonian

$$H_2^{(N)} = - \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} + c \sum_{i \neq j} \delta(x_i - x_j) \quad (1.3)$$

Hamiltonian (1.3) describes the interaction of  $N$  identical particles on the line via elastic collisions and  $c$  is the strength of interaction. The famous « Bethe Ansatz » [1] [2] exhibits the system of generalized eigenstates  $|\Psi_N(k_1, \dots, k_N)\rangle = |\Psi_N(\vec{k})\rangle$  of  $H_2^{(N)}$  which is complete if  $c \geq 0$ . We have

$$H_2^{(N)} |\Psi_N(\vec{k})\rangle = \left( \sum_{i=1}^N k_i^2 \right) |\Psi_N(\vec{k})\rangle. \quad (1.4)$$

Since Bethe Ansatz eigenstates depend on  $N$  quantum numbers  $k_1, \dots, k_N$  the Hamiltonian (1.3) must be completely integrable. This means that there are  $N$  independent operators  $H_n^{(N)}$   $n = 1, \dots, N$  such that

$$H_n^{(N)} |\Psi_N(\vec{k})\rangle = \left( \sum_{i=1}^N k_i^n \right) |\Psi_N(\vec{k})\rangle \quad (1.5)$$

$H_1^{(N)} = (-i) \sum_{i=1}^N \partial/\partial x_i$  is of course the total momentum and  $H_2^{(N)}$  is the

Hamiltonian (1.3). Existence of  $H_n^{(N)}$  should imply the infinite sequence of independent conservation laws  $\hat{H}_n$   $n = 1, 2, \dots$  for the NLSE given by their operator densities  $\hat{h}_n$

$$\hat{H}_n = \int dx \hat{h}_n(x). \quad (1.6)$$

We have

$$\hat{h}_1 = (-i) \Psi^\dagger \Psi_x \quad (1.7)$$

$$\hat{h}_2 = (-i)^2 (\Psi^\dagger \Psi_{xx} - c \Psi^\dagger \Psi^2). \quad (1.8)$$

Operators  $\hat{H}_n$  are completely characterized by the property that for any  $N$

$$\hat{H}_n |\Psi_N(\vec{k})\rangle = \left( \sum_{i=1}^N k_i^n \right) |\Psi_N(\vec{k})\rangle. \quad (1.9)$$

It is desirable to have explicit expressions for the operator densities  $\hat{h}_n$ . In this paper I suggest a method for calculating  $\hat{h}_n$  for any  $n$ . Using this method I calculate  $\hat{h}_3$  and  $\hat{h}_4$ . In section 4 I compare  $\hat{h}_n$  with the functional densities  $b_n$  of the integrals of motion for the classical NLSE

$$i\varphi_t = -\varphi_{xx} + 2c|\varphi|^2\varphi \tag{1.10}$$

Thacker [3] has obtained  $\hat{h}_3$  using a completely different approach. Kulish and Sklyanin [4] and Thacker [4] have integrated (1.1) using the quantum inverse scattering method. Their method however does not yield explicit formulas for  $\hat{h}_n$  in terms of the fields (\*).

### 2. N-PARTICLE SECTOR

In this section we fix  $N$  and omit the superscript  $N$  in formulas. The Hamiltonian  $H_2$  is equal to the Laplacean  $-\sum_{i=1}^N \partial^2/\partial x_i^2$  with the boundary conditions

$$(\partial/\partial x_j - \partial/\partial x_i)F = cF \tag{2.1}$$

on hyperplanes  $\{x_i - x_j = 0\}$   $i, j = 1, \dots, N$ .

Because of the symmetry of function  $F$  it suffices to restrict it to  $R_+^N = \{x_1 \leq x_2 \leq \dots \leq x_N\}$  and to impose boundary conditions

$$(\partial/\partial x_{k+1} - \partial/\partial x_k)F = cF \tag{2.2}$$

on hyperplanes  $x_k = x_{k+1}$   $k = 1, \dots, N - 1$ .

I will use the following fact. There is an operator  $P$  on symmetric functions in  $R^N$  that intertwines Laplacean with the Neumann boundary conditions

$$(\partial/\partial x_{k+1} - \partial/\partial x_k)F = 0 \tag{2.3}$$

and Laplacean with boundary conditions (2.2) for  $c \geq 0$ . The operator  $P$  constructed as follows. For any  $i \neq j$  let  $P_{ij}$  be given by

$$(P_{ij}f)(x_1, \dots, x_N) = \int_0^\infty dt e^{-ct} f(x_1, \dots, x_i - t, \dots, x_j + t, \dots, x_N). \tag{2.4}$$

Denote by  $S$  the operator from all functions  $f$  on  $R^N$  into symmetric functions on  $R^N$  obtained by restricting  $f$  to  $R_+^N$  and then extending it to  $R^N$  by symmetry. Then [5]

$$P = S \prod_{i < j} (1 - cP_{ij}). \tag{2.5}$$

---

(\*) *Added in proofs:* in a forthcoming paper I show that the formulas for integrals of the NLSE obtained in [4] via the quantum scattering method are false.

Denoting by  $\Delta_2$  the Laplacean with boundary conditions (2.3) we express the intertwining property of  $P$  by

$$H_2 P = P \Delta_2. \quad (2.6)$$

Let  $\Delta_n$  be given by  $(-i)^n \sum_{i=1}^N \partial^n / \partial x_i^n$  with « higher » Neumann boundary conditions

$$(\partial / \partial x_{k+1} - \partial / \partial x_k)^{2i+1} f = 0 \quad (2.7)$$

for  $i = 0, 1, \dots, [n/2] - 1$  on hyperplanes  $\{x_k = x_{k+1}\}$   $k = 1, \dots, N-1$ . Let  $H_n$  be defined from

$$P \Delta_n = H_n P \quad (2.8)$$

for  $n = 1, \dots$ . Since operators  $\Delta_n$  commute,  $H_n$  also commute. It follows from (2.5) that  $P$  takes boundary conditions (2.7) into boundary conditions

$$(\partial / \partial x_{k+1} - \partial / \partial x_k)^{2i+1} f = c(\partial / \partial x_{k+1} - \partial / \partial x_k)^{2i} f \quad (2.9)$$

So  $H_n$  is equal to  $(-i)^n \sum_{i=1}^N \partial^n / \partial x_i^n$  with boundary conditions (2.9) for

$i = 0, \dots, [n/2] - 1$ . It remains to obtain formulas for  $H_n$  similar to the formula (1.3) for  $H_2$ .

Let  $g(x_1, \dots, x_N)$  be an infinitely differentiable function and let  $f$  satisfy the boundary conditions (2.9). Then

$$\langle g | H_3 | f \rangle = (-i)^3 \int d^N x \bar{g} \left( \sum_{i=1}^N \frac{\partial^3}{\partial x_i^3} f \right). \quad (2.10)$$

Integrating by parts and taking (2.9) into account we get

$$\begin{aligned} \langle g | H_3 | f \rangle &= -(-i)^3 \int d^N x \sum_{i=1}^N \frac{\partial}{\partial x_i} \bar{g} \frac{\partial^2}{\partial x_i^2} f \\ &\quad - c(-i)^3 \sum_{i \neq j} \int d^N x \delta(x_i - x_j) \bar{g} \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right) f. \end{aligned} \quad (2.11)$$

Integrating by parts again

$$\begin{aligned} \langle g | H_3 | f \rangle &= (-i)^3 \int d^N x \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} \bar{g} \frac{\partial}{\partial x_i} f \\ &\quad + (-i)^3 c \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \frac{\partial}{\partial x_i} \bar{g} f \\ &\quad + (-i)^3 c \int d^N x \sum_{i \neq j} \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right) \delta(x_i - x_j) \bar{g} f. \end{aligned} \quad (2.12)$$

After one more integration by parts and obvious transformations (2.12) becomes

$$\begin{aligned} \langle g | H_3 | f \rangle &= i^3 \int d^N x \sum_{i=1}^N \frac{\partial^3}{\partial x_i^3} \bar{g} f \\ &\quad + (-i)^3 \frac{3}{2} c \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right) \bar{g} f \end{aligned} \quad (2.13)$$

which yields

$$H_3 = (-i)^3 \left( \sum_{i=1}^N \frac{\partial^3}{\partial x_i^3} - \frac{3}{2} c \sum_{i \neq j} \delta(x_i - x_j) \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right) \right). \quad (2.14)$$

For  $H_4$  we have

$$\langle g | H_4 | f \rangle = \int d^N x \bar{g} \sum_{i=1}^N \frac{\partial^4}{\partial x_i^4} f. \quad (2.15)$$

Integrating by parts the right hand side

$$\begin{aligned} & - \int d^N x \sum_{i=1}^N \frac{\partial}{\partial x_i} \bar{g} \frac{\partial^3}{\partial x_i^3} f \\ & \quad - c \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \bar{g} \left( \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial x_i \partial x_j} + \frac{\partial^2}{\partial x_j^2} \right) f. \end{aligned} \quad (2.16)$$

Integrating by parts the first term in (2.16) we get

$$\begin{aligned} & \int d^N x \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} \bar{g} \frac{\partial^2}{\partial x_i^2} f \\ & \quad + \frac{1}{2} c \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right) \bar{g} \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right) f. \end{aligned} \quad (2.17)$$

Integrating (2.17) by parts again and remembering the second term in (2.16) we have

$$\begin{aligned}
 \langle g | H_4 | f \rangle = & \int d^N x \sum_{i=1}^N \frac{\partial^4}{\partial x_i^4} \bar{g} f \\
 & - \frac{c}{2} \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \left( \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial x_j^2} \right) \bar{g} f \\
 & - \frac{c}{2} \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \left( \frac{\partial}{\partial x_i} + \frac{\partial}{\partial x_j} \right)^2 \bar{g} f \\
 & - c \int d^N x \sum_{i \neq j} \delta(x_i - x_j) \bar{g} \left( \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial x_i \partial x_j} + \frac{\partial^2}{\partial x_j^2} \right) f. \quad (2.18)
 \end{aligned}$$

The last term in (2.18) can again be integrated by parts yielding

$$\begin{aligned}
 - c \int d^N x \sum_{i \neq j} \left( \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial x_i^2 \partial x_j} + \frac{\partial^2}{\partial x_j^2} \right) \delta(x_i - x_j) \bar{g} f \\
 + 3c^2 \int d^N x \sum_{i \neq j \neq k} \delta(x_i - x_j) \delta(x_j - x_k) \bar{g} f. \quad (2.19)
 \end{aligned}$$

From (2.18) and (2.19) we have

$$\begin{aligned}
 H_4 = & \sum_{i=1}^N \frac{\partial^4}{\partial x_i^4} - c \sum_{i \neq j} \delta(x_i - x_j) \left( \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial x_i \partial x_j} + \frac{\partial^2}{\partial x_j^2} \right) \\
 & - c \sum_{i \neq j} \left( \frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial x_i \partial x_j} + \frac{\partial^2}{\partial x_j^2} \right) \delta(x_i - x_j) + 3c^2 \sum_{i \neq j \neq k} \delta(x_i - x_j) \delta(x_j - x_k). \quad (2.20)
 \end{aligned}$$

### 3. SECOND QUANTIZED FORM OF $H_n$

A standard calculation gives

$$\begin{aligned}
 \hat{H}_3 = & (-i)^3 \int dx \Psi^\dagger \Psi_{xxx} \\
 & - \frac{3}{2} c (-i)^3 \int dx \int dy \Psi^\dagger(x) \Psi^\dagger(y) \delta(x-y) \left( \frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right) \Psi(x) \Psi(y) \\
 = & (-i)^3 \int dx [\Psi^\dagger \Psi_{xxx} - 3c \Psi^{\dagger 2} \Psi \Psi_x]. \quad (3.1)
 \end{aligned}$$

Thus

$$\hat{h}_3(x) = (-i)^3(\Psi^\dagger\Psi_{xxx} - 3c\Psi^{\dagger 2}\Psi\Psi_x). \tag{3.2}$$

Also

$$\begin{aligned} \hat{H}_4 = & \int dx \Psi^\dagger\Psi_{xxxx} \\ & - c \int dx \int dy \Psi^\dagger(x)\Psi^\dagger(y)\delta(x-y)\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial x\partial y} + \frac{\partial^2}{\partial y^2}\right)\Psi(x)\Psi(y) \\ & - c \int dx \int dy \Psi^\dagger(x)\Psi^\dagger(y)\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial x\partial y} + \frac{\partial^2}{\partial y^2}\right)\delta(x-y)\Psi(x)\Psi(y) \\ & + 3c^2 \int dx \int dy \int dz \Psi^\dagger(x)\Psi^\dagger(y)\Psi^\dagger(z)\delta(x-y)\delta(y-z)\Psi(x)\Psi(y)\Psi(z). \end{aligned} \tag{3.3}$$

After obvious integrations by parts we have

$$\hat{H}_4 = \int dx [\Psi^\dagger\Psi_{xxx} - 2c\Psi^{\dagger 2}\Psi\Psi_{xx} - c\Psi^{\dagger 2}\Psi_x^2 - 2c\Psi^\dagger\Psi_{xx}^2 - c\Psi_x^{\dagger 2}\Psi^2 + 3c^2\Psi^{\dagger 3}\Psi^3]. \tag{3.4}$$

Thus

$$\hat{h}_4(x) = \Psi^\dagger\Psi_{xxxx} - 2c\Psi^{\dagger 2}\Psi\Psi_{xx} - c\Psi^{\dagger 2}\Psi_x^2 - 2c\Psi^\dagger\Psi_{xx}^2 - c\Psi_x^{\dagger 2}\Psi^2 + 3c^2\Psi^{\dagger 3}\Psi^3. \tag{3.5}$$

#### 4. COMPARISON WITH CLASSICAL INTEGRALS

The classical NLSE (or Zakharov-Shabat equation [6])

$$i\varphi_t = -\varphi_{xx} + 2c|\varphi|^2\varphi \tag{4.1}$$

is a completely integrable Hamiltonian system with infinitely many degrees of freedom [7]. In particular (4.1) has an infinite number of integrals of motion  $B_n(\bar{\varphi}, \varphi)$ . The functionals  $B_n$  are determined by the local densities  $b_n$

$$B_n(\bar{\varphi}, \varphi) = \int_{-\infty}^{\infty} dx b_n(\bar{\varphi}(x), \varphi(x)). \tag{4.2}$$

The densities  $b_n$  are found from the recurrence relation

$$b_{n+1} = \bar{\varphi} \frac{d}{dx} \left( \frac{b_n}{\bar{\varphi}} \right) - c \sum_{i+j=n-1} b_i b_j \tag{4.3}$$

and

$$b_0 = \bar{\varphi}\varphi. \tag{4.4}$$

From (4.3) and (4.4) we get

$$b_1 = \bar{\varphi}\varphi_x \tag{4.5}$$

$$b_2 = \bar{\varphi}\varphi_{xx} - c|\varphi|^4 \tag{4.6}$$



$$b_3 = \bar{\varphi}\varphi_{xxx} - 2c\bar{\varphi}^2(\varphi^2)_x - c\bar{\varphi}\varphi_x\varphi^2 \quad (4.7)$$

$$b_4 = \bar{\varphi}\varphi_{xxxx} - 2c\bar{\varphi}^2(\varphi^2)_{xx} - 2c\bar{\varphi}^2\varphi\varphi_{xx} - c\bar{\varphi}^2\varphi_x^2 - 3c\bar{\varphi}\varphi_x(\varphi^2)_x - c\bar{\varphi}\varphi_x\varphi^2 + 2c^2|\varphi|^6. \quad (4.8)$$

The local densities  $h$  and  $g$  that differ by a total derivative are equivalent

$h \simeq g$  because they define the same functional  $\int_{-\infty}^{\infty} dxh(x) = \int_{-\infty}^{\infty} dxg(x)$ . We have

$$b_3 \simeq \bar{\varphi}\varphi_{xxx} - \frac{3}{2}c\bar{\varphi}^2(\varphi^2)_x = b'_3 \quad (4.9)$$

$$b_4 \simeq \bar{\varphi}_{xx}\varphi_{xx} + 2c(\bar{\varphi}^2)_x(\varphi^2)_x + c\bar{\varphi}^2\varphi_x^2 + c\bar{\varphi}_x^2\varphi^2 + 2c^2|\varphi|^6 = b'_4. \quad (4.10)$$

We see from (3.2) that  $\hat{h}_3$  differs from  $b'_3$  by the factor  $(-i)^3$  only. On the other hand the difference between  $\hat{h}_4$  (3.5) and  $b_4$  is essential. Replacing  $\hat{h}_4$  by an equivalent operator density  $h'_4$  the closest that we can get to  $b'_4$  is

$$\hat{h}'_4 = \Psi_{xx}^\dagger\Psi_{xx} + 2c(\Psi^{\dagger 2})_x(\Psi^2)_x + c\Psi^{\dagger 2}\Psi_x^2 + c\Psi_x^{\dagger 2}\Psi^2 + 3c^2\Psi^{\dagger 3}\Psi^3. \quad (4.11)$$

The difference corresponds to  $c^2|\varphi|^6$  which is a nontrivial density.

## 5. CONCLUSION

The nonlinear Schrödinger equation (1.1) has an infinite sequence of conservation laws  $\hat{H}_n$  given by the operator densities  $\hat{h}_n(\Psi^\dagger(x), \Psi(x))$ . The densities  $\hat{h}_n$  can be found using the method of sections 2 and 3. It is clear from the method that  $\hat{h}_n$  are polynomials in the fields and their derivatives. Besides  $\hat{h}_n$  are polynomials in the coupling constant  $c$ . The degree of  $\hat{h}_n$  in  $c$  is  $[n/2]$ .

Correspondence between  $\hat{h}_n$  and the integral densities  $b_n$  of the classical NLSE (4.1) breaks down at  $n = 4$ .

## REFERENCES

- [1] E. LIEB and W. LINIGER, *Phys. Rev.*, t. **130**, 1963, p. 1605.
- [2] C. N. YANG, *Phys. Rev. Lett.*, t. **19**, 1967, p. 1312.
- [3] H. B. THACKER, *Phys. Rev.*, t. **D17**, 1978, p. 1031.
- [4] Articles in Integrable Quantum Field Theories, Proceedings, Tvarminne, Finland 1981, *Lecture Notes in Physics*, 151, Springer-Verlag, New York, 1982.
- [5] Another intertwining operator  $\tilde{P}$  was constructed in E. GUTKIN, *Duke Mat. J.*, t. **49**, 1, 1982 on the space of all functions not regarding the symmetry.
- [6] V. E. ZAKHAROV and A. B. SHABAT, *Zh. Eksp. Teor. Fiz.*, t. **61**, 1971, p. 118; *Sov. Phys., JETP*, t. **34**, 1972, p. 62.
- [7] V. E. ZAKHAROV and S. V. MANAKOV, *Teor. Mat. Fiz.*, t. **19**, 1974, p. 332. *Theor. Math. Phys.*, t. **19**, 1975, p. 551.

(Manuscrit reçu le 25 juin 1984)