ANNALES DE L'I. H. P., SECTION C

G. BUTAZZO

E. CAVAZZUTI

Limit problems in optimal control theory

Annales de l'I. H. P., section C, tome S6 (1989), p. 151-160

http://www.numdam.org/item?id=AIHPC_1989__S6__151_0

© Gauthier-Villars, 1989, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

LIMIT PROBLEMS IN OPTIMAL CONTROL THEORY

G. BUTTAZZO

Dipartimento di Matematica, Via Machiavelli, 35, 44100 Ferrara, Italy

E. CAVAZZUTI

Dipartimento di Matematica, Via Campi, 213/B, 41100 Modena, Italy

1. INTRODUCTION

In this paper we deal with sequences of optimal control problems of the form

$$(P_h) \quad \min \left\{ \int_0^1 f_h(t, y, u) dt : y' = g_h(t, y, u), y(0) = y_h^0 \right\}$$

where the state variable y belong to the Sobolev space $W^{1,1}(0,1;\mathbf{R}^n)$ and the control variable u is in $L^1(0,1;\mathbf{R}^m)$. We are interested in the asymptotic behaviour (as $h \rightarrow +\infty$) of the optimal pairs (u_h, y_h) of (P_h) ; more precisely, we shall construct a new problem (P_∞) such that

if (u_h, y_h) is an optimal pair of (P_h) and if (u_h, y_h) tends to (u_∞, y_∞) in the topology $wL^1(0,1;\mathbf{R}^m) \times L^\infty(0,1;\mathbf{R}^n)$, then (u_∞, y_∞) is an optimal pair for (P_∞) .

The basic tool for treating the asymptotic problem above is the Γ -convergence theory which has been already used very fruitfully for many variational problems (see for instance [1],[2],[4],[5],[8],[9])

Here we use a more sophisticated version of the usual Γ -limits, because we shall consider our problems (P_h) as minimization problems on the product space $U \times Y$ (U is the space of controls and Y the space of states) for the functionals

$$F_h(u, y) = \begin{cases} \int_0^1 f_h(t, y, u) dt & \text{if } y' = g_h(t, y, u), y(0) = y_h^0 \\ +\infty & \text{otherwise} \end{cases}$$

and the spaces U and Y will play a different role with respect to Γ -convergence.

In Section 2 we develop the abstract theory we shall need in the following; in Section 3 we show the applications to problems (P_h) above, and we give an example showing that in some situations the domain of problem (P_∞) is not given by a state equation $y' = g_\infty(t, y, u)$ but coincides with the entire product space $U \times Y$.

2. THE ABSTRACT FRAMEWORK

Let us denote by U and Y two topological spaces and let $F_h: U \times Y \rightarrow \overline{\mathbf{R}}$ be a sequence of functions; by $Z(+)$ we shall denote the "sup" operator and by $Z(-)$ the "inf" operator. For every $u \in U$ and $y \in Y$ we define

$$\Gamma_{\text{seq}}(N^\alpha, U^\beta, Y^\gamma) \lim_h F_h(u, y) = \bigwedge_{(u_h) \in S(u)} \bigvee_{(y_h) \in S(y)} \bigwedge_{k \in N} \bigvee_{h \geq k} F_h(u_h, y_h)$$

where α, β, γ are the signs $+$ or $-$, and $S(u)$ and $S(y)$ respectively denote the set of all sequences $u_h \rightarrow u$ in U and $y_h \rightarrow y$ in Y . For example we have

$$\Gamma_{\text{seq}}(N^+, U^-, Y^+) \lim_h F_h(u, y) = \inf_{u_h \rightarrow u} \sup_{y_h \rightarrow y} \limsup_{h \rightarrow \infty} F_h(u_h, y_h).$$

When a Γ -limit is independent of the sign $+$ or $-$ associated to one of the spaces N, U, Y this sign will be omitted. For example, if

$$\Gamma_{\text{seq}}(N^+, U^-, Y^+) \lim_h F_h(u, y) = \Gamma_{\text{seq}}(N^+, U^+, Y^+) \lim_h F_h(u, y),$$

then their common value will be indicated by $\Gamma_{\text{seq}}(N^+, U, Y^+) \lim_h F_h(u, y)$.

The following propositions are proved in [4].

PROPOSITION 2.1. *Let (u_h, y_h) be a minimum point for F_h , or simply a pair such that*

$$\lim_h F_h(u_h, y_h) = \lim_h \left[\inf_{U \times Y} F_h \right].$$

Assume that (u_h, y_h) converges to (u_∞, y_∞) in $U \times Y$ and that there exist

$$F_\infty = \Gamma_{\text{seq}}(N, U^-, Y^-) \lim_h F_h.$$

Then we have

- (i) (u_∞, y_∞) is a minimum point for F_∞ on $U \times Y$;
- (ii) $\lim_h \left[\inf_{U \times Y} F_h \right] = \min_{U \times Y} F_\infty$.

PROPOSITION 2.2. *Let $\{F_h\}$ and $\{G_h\}$ be two sequences of function from $U \times Y$ into $[0, +\infty]$, and let $(u, y) \in U \times Y$. Assume there exist*

$$\Gamma_{\text{seq}}(N, U^-, Y^-) \lim_h F_h(u, y) \quad \text{and} \quad \Gamma_{\text{seq}}(N, U, Y^-) \lim_h G_h(u, y).$$

Then we have

$$\Gamma_{\text{seq}}(N, U^-, Y^-) \lim_h [F_h + G_h](u, y) = \Gamma_{\text{seq}}(N, U^-, Y^-) \lim_h F_h(u, y) + \Gamma_{\text{seq}}(N, U, Y^-) \lim_h G_h(u, y).$$

In many applications, the introduction of a new auxiliary variable can be helpful; the following proposition shows the behaviour of Γ -limits with respect to this operation.

PROPOSITION 2.3. *Let $F_h: U \times Y \rightarrow \overline{\mathbb{R}}$ be a sequence of functions, let V be another topological space, and let $\Xi_h: U \times Y \rightarrow \wp(V)$ be a sequence of multimappings. Assume that the following compactness condition is satisfied:*

for every converging sequence (u_h, y_h) with $F_h(u_h, y_h)$ bounded, there exist a sequence $v_h \in \Xi_h(u_h, y_h)$ relatively compact in V .

Then setting

$$\Phi_h(u,v,y) = \begin{cases} F_h(u,y) & \text{if } v \in \Xi_h(u,y) \\ +\infty & \text{otherwise,} \end{cases}$$

we have for every $(u,y) \in U \times Y$

$$\begin{aligned} & \inf \left\{ \Gamma_{\text{seq}}(N^-, (U \times V)^-, Y^-) \lim_h \Phi_h(u,v,y) : v \in V \right\} \leq \\ & \leq \Gamma_{\text{seq}}(N^-, U^-, Y^-) \lim_h F_h(u,y) \leq \Gamma_{\text{seq}}(N^+, U^-, Y^-) \lim_h F_h(u,y) \leq \\ & \leq \inf \left\{ \Gamma_{\text{seq}}(N^+, (U \times V)^-, Y^-) \lim_h \Phi_h(u,v,y) : v \in V \right\} . \end{aligned}$$

Therefore, if for every $(u,v,y) \in U \times V \times Y$ there exists

$$\Gamma_{\text{seq}}(N, (U \times V)^-, Y^-) \lim_h \Phi_h(u,v,y) ,$$

we have

$$\Gamma_{\text{seq}}(N, U^-, Y^-) \lim_h F_h(u,y) = \inf \left\{ \Gamma_{\text{seq}}(N, (U \times V)^-, Y^-) \lim_h \Phi_h(u,v,y) : v \in V \right\} .$$

Proof. It is enough to repeat, with just some slight modifications, the proof of Proposition 2.4 of [3]. ■

In the following, if A is a set we denote by χ_A the function

$$\chi_A(x) = \begin{cases} 0 & \text{if } x \in A \\ +\infty & \text{otherwise.} \end{cases}$$

3. APPLICATIONS TO CONTROL PROBLEMS

Let k, m, n be positive integers and let $p > 1$. The space Y of states we consider is the Sobolev space $W^{1,1}(0,1; \mathbb{R}^n)$ endowed with the $L^\infty(0,1; \mathbb{R}^n)$ topology, and the space U of

controls is the space $L^p(0,1;\mathbb{R}^n)$ endowed with its weak topology (weak* if $p=+\infty$). The cost functions are of the form

$$(3.1) \quad J_h(u,y) = \int_0^1 f_h(t,y,u) dt$$

where $f_h:[0,1]\times\mathbb{R}^n\times\mathbb{R}^m\rightarrow[0,+\infty]$ are Borel functions. Finally, the state equations are

$$(3.2) \quad \begin{cases} y' \in a_h(t,y) + B_h(t,y) b_h(t,u) \\ y(0)=y_h^0 \end{cases}$$

where $a_h:[0,1]\times\mathbb{R}^n\rightarrow\mathbb{R}^n$ and $B_h:[0,1]\times\mathbb{R}^n\rightarrow\mathbb{R}^{nk}$ are Borel functions, and the multimappings $b_h:[0,1]\times\mathbb{R}^m\rightarrow\wp(\mathbb{R}^{nk})$ are Borel measurable (i.e. the sets $\{(t,u,v)\in[0,1]\times\mathbb{R}^n\times\mathbb{R}^k : v\in b_h(t,u)\}$ are Borel sets).

Then the control problems we are concerned are

$$(P_h) \quad \inf \{ J_h(u,y) : (u,y)\in\Lambda_h \}$$

or equivalently

$$(P_h) \quad \inf \{ F_h(u,y) : (u,y)\in U\times Y \}$$

where

$$(3.3) \quad \Lambda_h = \{ (u,y)\in U\times Y : y'\in a_h(t,y) + B_h(t,y) b_h(t,u), y(0)=y_h^0 \}$$

$$(3.4) \quad F_h = J_h + \chi_{\Lambda_h}.$$

We introduce now an auxiliary variable $v\in L^q(0,1;\mathbb{R}^k)$ with $q>1$ and define a new sequence of functionals by setting

$$(3.5) \quad \Phi_h(u,v,y) = \begin{cases} F_h(u,y) & \text{if } v\in b_h(t,u) \\ +\infty & \text{otherwise.} \end{cases}$$

In this way the problems (P_h) take the form

$$\inf \left\{ \int_0^1 [f_h(t,y,u) + \chi_{\{v\in b_h(t,u)\}}] dt : y'=a_h(t,y) + B_h(t,y) v, y(0)=y_h^0 \right\}.$$

In order to apply the abstract theory presented in Section 2 (more precisely Proposition 2.1), we have to calculate the $\Gamma_{\text{seq}}(N,U^-,Y^-)$ limit of the sequence F_h . To do this, we make

some hypotheses on f_h, a_h, B_h, b_h .

(3.6) For every $t \in [0,1], r \geq 0, y \in \mathbb{R}^n$ with $|y| \leq r$ we have (if $1/q + 1/q' = 1$)

$$\begin{aligned} |a_h(t,y)| &\leq M_h(t,r) & \text{with} & \quad \|M_h(\cdot, r)\|_{L^1(0,1)} \leq M(r) < +\infty \\ |B_h(t,y)| &\leq N_h(t,r) & \text{with} & \quad \|N_h(\cdot, r)\|_{L^{q'}(0,1)} \leq N(r) < +\infty. \end{aligned}$$

(3.7) For every $t \in [0,1], r \geq 0, y_1, y_2 \in \mathbb{R}^n$ with $|y_1|, |y_2| \leq r$ we have

$$\begin{aligned} |a_h(t, y_1) - a_h(t, y_2)| &\leq \alpha_h(t, r) |y_1 - y_2| & \text{with} & \quad \|\alpha_h(\cdot, r)\|_{L^1(0,1)} \leq \alpha(r) < +\infty \\ |B_h(t, y_1) - B_h(t, y_2)| &\leq \beta_h(t, r) |y_1 - y_2| & \text{with} & \quad \|\beta_h(\cdot, r)\|_{L^{q'}(0,1)} \leq \beta(r) < +\infty. \end{aligned}$$

(3.8) There exist $\lambda > 0$ and $a \in L^1(0,1)$ such that

$$\lambda(|u|^p + |v|^q) - a(t) \leq f_h(t, 0, u) + \chi_{\{v \in b_h(t, u)\}}$$

for every $t \in [0,1], u \in \mathbb{R}^m, v \in \mathbb{R}^k$. When $p = +\infty$ or $q = +\infty$ the quantities $|u|^p$ and $|v|^q$ in the left-hand side have to be substituted by $\chi_{\{u \in H\}}$ and $\chi_{\{v \in K\}}$ respectively, where $H \subset \mathbb{R}^m$ and $K \subset \mathbb{R}^k$ are bounded sets.

(3.9) For every $t \in [0,1], r \geq 0, u \in \mathbb{R}^m, y_1, y_2 \in \mathbb{R}^n$ with $|y_1|, |y_2| \leq r$ we have

$$f_h(t, y_1, u) \leq f_h(t, y_2, u) + \rho_r(t, |y_1 - y_2|) + \sigma_r(t, |y_1 - y_2|) |f_h(t, y_2, u)|^{(\alpha-1)/\alpha}$$

for a suitable $\alpha \in [1, +\infty]$ and functions $\rho_r(t, s), \sigma_r(t, s)$ from $[0,1] \times [0, +\infty[$ into $[0, +\infty[$ measurable in t , increasing and continuous in s , with $\rho_r(t, 0) = \sigma_r(t, 0) = 0$, and such that $z \rightarrow \rho_r(t, |z(t)|), z \rightarrow \sigma_r(t, |z(t)|)$ are continuous operators from Y into $L^1(0,1), L^\alpha(0,1)$ respectively.

(3.10) There exist $u_h \in L^p(0,1; \mathbb{R}^m)$ and $v_h \in L^q(0,1; \mathbb{R}^k)$ such that $v_h(t) \in b_h(t, u_h(t))$ for a.e. $t \in (0,1)$, and the sequence $f_h(t, 0, u_h(t))$ is weakly compact in $L^1(0,1)$.

LEMMA 3.1. *Under the previous assumptions, the following compactness condition is satisfied: for every converging sequence (u_h, y_h) with $F_h(u_h, y_h)$ bounded, there exists a sequence $\{v_h\}$ relatively compact in V such that for a.e. $t \in [0,1]$*

$$v_h(t) \in b_h(t, u_h(t)) \quad \text{and} \quad y'_h(t) = a_h(t, y_h(t)) + B_h(t, y_h(t)) v_h(t).$$

Proof. Let (u_h, y_h) be converging in $U \times Y$ with $F_h(u_h, y_h)$ bounded; then we have $(u_h, y_h) \in \Lambda_h$, so that we can find measurable functions $v_h(t)$ with

$$v_h(t) \in b_h(t, u_h(t)) \quad \text{and} \quad y'_h(t) = a_h(t, y_h(t)) + B_h(t, y_h(t)) v_h(t)$$

for a.e. $t \in [0, 1]$. It remains to prove that the sequence v_h is bounded in $L^q(0, 1; \mathbb{R}^k)$. Since y_h is uniformly bounded, by (3.8) and (3.9) we have for a suitable $r > 0$

$$\begin{aligned} \lambda(|u_h|^p + |v_h|^q) - a(t) &\leq f_h(t, 0, u_h) \leq \\ &\leq f_h(t, y_h, u_h) + \rho_r(t, r) + \sigma_r(t, r) \|f_h(t, y_h, u_h)\|^{(\alpha-1)/\alpha} \leq \\ &\leq c f_h(t, y_h, u_h) + \gamma(t) \end{aligned}$$

where $c > 0$ is a constant and $\gamma \in L^1(0, 1)$. Then, from the boundedness of $J_h(u_h, y_h)$ we get that v_h is bounded in $L^q(0, 1; \mathbb{R}^k)$. ■

By Lemma 3.1, Proposition 2.3 applies, so that we have reduced our problem to the characterization of the $\Gamma_{\text{seq}}(\mathbb{N}, (U \times Y)^-, Y^-)$ limit of the sequence $\Phi_h(u, v, y)$ defined in (3.5). Set now

$$\begin{aligned} \bar{f}_h(t, y, u, v) &= f_h(t, y, u) + \chi_{\{v \in b_h(t, u)\}} \\ \bar{J}_h(u, v, y) &= \int_0^1 \bar{f}_h(t, y, u, v) dt \\ \bar{\Lambda}_h &= \left\{ (u, v, y) \in U \times V \times Y : y' = a_h(t, y) + B_h(t, y) v \text{ a.e. on } [0, 1], y(0) = y_h^0 \right\}. \end{aligned}$$

Therefore

$$\Phi_h = \bar{J}_h + \chi_{\bar{\Lambda}_h}^-$$

and, by Proposition 2.2 we may split the $\Gamma_{\text{seq}}(\mathbb{N}, (U \times Y)^-, Y^-)$ limit of Φ_h into the sum

$$(3.11) \quad \Gamma_{\text{seq}}(\mathbb{N}, (U \times V)^-, Y) \lim_h \bar{J}_h + \Gamma_{\text{seq}}(\mathbb{N}, U \times V, Y^-) \lim_h \chi_{\bar{\Lambda}_h}^-.$$

The two terms in the sum above can be computed by using Lemma 3.1 and Theorem 3.4 of [4].

More precisely we have

PROPOSITION 3.2. Assume that (3.8),(3.9),(3.10) hold and that for every $y \in \mathbb{R}^n$, $\xi \in \mathbb{R}^n$, $\eta \in \mathbb{R}^k$

$$(3.12) \quad \bar{F}_h^*(\cdot, y, \xi, \eta) \rightarrow \varphi(\cdot, y, \xi, \eta) \quad \text{weakly in } L^1(0,1)$$

where $\bar{F}_h^*$ are the polar functions of $\bar{F}_h$ defined by

$$\bar{F}_h^*(t, y, \xi, \eta) = \sup \left\{ \xi u + \eta v - \bar{F}_h(t, y, u, v) : u \in \mathbb{R}^m, v \in \mathbb{R}^k \right\}.$$

Then, for every $(u, v, y) \in U \times V \times Y$ we have

$$\Gamma_{\text{seq}}(N, (U \times V)^-, Y) \lim_h \bar{J}_h(u, v, y) = \int_0^1 \varphi^*(t, y, u, v) dt$$

where φ^* is the polar function of φ .

PROPOSITION 3.3. Assume that (3.6),(3.7) hold and that

$$(3.13) \quad \text{for every } y \in \mathbb{R}^n \quad a_h(\cdot, y) \rightarrow a(\cdot, y) \quad \text{weakly in } L^1(0,1; \mathbb{R}^n);$$

$$(3.14) \quad \text{for every } y \in \mathbb{R}^n \quad B_h(\cdot, y) \rightarrow B(\cdot, y) \quad \text{strongly in } L^q(0,1; \mathbb{R}^{nk});$$

$$(3.15) \quad y_h^0 \rightarrow y^0 \quad \text{in } \mathbb{R}^n.$$

Then we have

$$\Gamma_{\text{seq}}(N, U \times V, Y^-) \lim_h \chi_{\Lambda_h}^- = \chi_{\bar{\Lambda}}^-$$

where

$$\bar{\Lambda} = \left\{ (u, v, y) \in U \times V \times Y : y' = a(t, y) + B(t, y) v, y(0) = y^0 \right\}.$$

Finally, we are in a position to compute the $\Gamma_{\text{seq}}(N, U^-, Y^-)$ limit of F_h . In fact, by Propositions 2.3, 3.2, and 3.3, and by (3.11) we get for every $(u, y) \in U \times Y$

$$\begin{aligned} \Gamma_{\text{seq}}(N, U^-, Y^-) \lim_h F_h(u, y) &= \\ &= \inf \left\{ \int_0^1 \varphi^*(t, y, u, v) dt : y' = a(t, y) + B(t, y) v, y(0) = y^0 \right\} = \end{aligned}$$

$$= \int_0^1 f(t, y, u, y') dt + \chi_{\{y(0)=y^0\}}$$

where the function f is defined by

$$f(t, y, u, w) = \inf \{ j(t, y, u, v) : w = a(t, y) + B(t, y) v \} .$$

We conclude with an example showing that in general the domain of the limit functional

$$F(u, y) = \int_0^1 f(t, y, u, y') dt + \chi_{\{y(0)=y^0\}}$$

is not given by a differential equation of the form $y'=g(t, y, u)$ but may be the whole space $U \times Y$.

EXAMPLE 3.4. Consider the sequence of optimal control problems

$$(P_h) \quad \min \left\{ \int_0^1 [u^2 + |y - y_0(t)|^2] dt : y' = a_h(t) y + b_h(t) u, y(0) = \xi \right\}$$

where u varies in $U = L^2(0, 1)$, y varies in $Y = W^{1,1}(0, 1)$, and $y_0 \in L^2(0, 1)$, $\xi \in \mathbf{R}$ are given.

About the functions a_h and b_h we assume that

$$\begin{cases} a_h \rightarrow a & \text{weakly in } L^1(0, 1) \\ b_h \rightarrow b & \text{weakly}^* \text{ in } L^\infty(0, 1) \\ b_h^2 \rightarrow \beta^2 & \text{weakly}^* \text{ in } L^\infty(0, 1) . \end{cases}$$

It is not difficult to check that all hypotheses (3.6),..., (3.10) and (3.12),..., (3.15) are satisfied, and after some standard calculations we find that the limit problem (P_∞) has the form

$$(P_\infty) \quad \min \left\{ \int_0^1 [u^2 + |y - y_0(t)|^2 + \frac{|y' - a(t) y - b(t) u|^2}{\beta^2(t) - b^2(t)}] dt : y(0) = \xi \right\} .$$

Note that it is $\beta^2(t) \geq b^2(t)$ for a.e. $t \in [0, 1]$, and

$$\beta^2 = b^2 \text{ a.e. on } [0, 1] \Leftrightarrow b_h \rightarrow b \text{ a.e. on } [0, 1] .$$

In this last situation, problem (P_∞) takes the usual form

$$(P_\infty) \quad \min \left\{ \int_0^1 [u^2 + |y - y_0(t)|^2] dt : y' = a(t) y + b(t) u, y(0) = \xi \right\} ,$$

but this does not arrive in the general case. Take for instance

$$b_h(t) = \sin(ht)$$

and we get $b=0$ and $\beta^2=1/2$, so that the limit problem is

$$\min \left\{ \int_0^1 [u^2 + |y - y_0(t)|^2 + 2|y' - a(t)y|^2] dt : y(0)=\xi \right\}.$$

REFERENCES

- [1] H.ATTOUCH: *Variational Convergence for Functions and Operators*. Appl.Math.Ser., Pitman, Boston (1984).
- [2] G.BUTTAZZO: *Su una definizione generale dei Γ -limiti*. Boll.Un.Mat.Ital., **14-B** (1977), 722-744.
- [3] G.BUTTAZZO: *Some relaxation problems in optimal control theory*. J.Math.Anal.Appl., **125** (1987), 272-287.
- [4] G.BUTTAZZO & G.DAL MASO: *Γ -convergence and optimal control problems*. J.Optim. Theory Appl., **38** (1982), 385-407.
- [5] E.CAVAZZUTI: *G-convergenze multiple, convergenze di punti di sella e di max-min*. Boll.Un.Mat.Ital., **1-B** (1982), 251-274.
- [6] F.H.CLARKE: *Admissible relaxation in variational and control problems*. J.Math.Anal. Appl., **51** (1975), 557-576.
- [7] F.H.CLARKE: *Optimization and Nonsmooth Analysis*. Wiley Interscience, New York (1983).
- [8] E.DE GIORGI: *Convergence problems for functionals and operators*. Proceedings of "Recent Methods in Nonlinear Analysis", Rome 1978, edited by E.De Giorgi & E. Magenes & U.Mosco, Pitagora, Bologna (1979), 131-188.
- [9] E.DE GIORGI & T.FRANZONI: *Su un tipo di convergenza variazionale*. Atti Accad.Naz. Lincei Rend.Cl.Sci.Fis.Mat.Natur., **58** (1975), 842-850.
- [10] T.ZOLEZZI: *On equiwellset minimum problems*. Appl.Math.Optim., **4** (1978), 209-223.