

# ANNALES DE L'I. H. P., SECTION C

MIN-CHUN HONG

## **On the Jäger-Kaul theorem concerning harmonic maps**

*Annales de l'I. H. P., section C*, tome 17, n° 1 (2000), p. 35-46

[http://www.numdam.org/item?id=AIHPC\\_2000\\_\\_17\\_1\\_35\\_0](http://www.numdam.org/item?id=AIHPC_2000__17_1_35_0)

© Gauthier-Villars, 2000, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## On the Jäger–Kaul theorem concerning harmonic maps

by

**Min-Chun HONG<sup>1</sup>**

School of Mathematical Sciences, The Australian National University,  
Canberra ACT 0200, Australia

Manuscript received 30 March 1998

---

**ABSTRACT.** – In 1983, Jäger and Kaul proved that the equator map  $u^*(x) = (\frac{x}{|x|}, 0) : B^n \rightarrow S^n$  is unstable for  $3 \leq n \leq 6$  and a minimizer for the energy functional  $E(u, B^n) = \int_{B^n} |\nabla u|^2 dx$  in the class  $H^{1,2}(B^n, S^n)$  with  $u = u^*$  on  $\partial B^n$  when  $n \geq 7$ . In this paper, we give a new and elementary proof of this Jäger–Kaul result. We also generalize the Jäger–Kaul result to the case of  $p$ -harmonic maps. © 2000 Éditions scientifiques et médicales Elsevier SAS

**RÉSUMÉ.** – En 1983, Jäger et Kaul ont démontrés que l'application équatorielle  $u^*(x) = (\frac{x}{|x|}, 0) : B^n \rightarrow S^n$  n'est pas stable si  $3 \leq n \leq 6$  et que c'est une minimisateur pour la fonctionnelle d'énergie  $E(u, B^n) = \int_{B^n} |\nabla u|^2 dx$  dans la classe  $H^{1,2}(B^n, S^n)$  avec  $u = u^*$  sur  $\partial B^n$  si  $n \geq 7$ . Nous donnons une preuve nouvelle, élémentaire de ce résultat de Jäger–Kaul. En plus nous généralisons le résultat de Jäger–Kaul au cas des applications  $p$ -harmoniques. © 2000 Éditions scientifiques et médicales Elsevier SAS

---

<sup>1</sup> Research of the author was supported by the Australian Research Council.

## 1. INTRODUCTION

Let  $(M, g)$  be a compact Riemannian manifold with (possibly empty) boundary  $\partial M$  and let  $(N, h)$  be another compact Riemannian manifold without boundary. Let  $u$  be a map from  $M$  to  $N$  which belongs to  $H^{1,2}(M, N)$ . We define the energy of  $u$  by

$$E(u, M) = \int_M |du|^2 dM, \quad (1.1)$$

where  $|du|$  denotes the Hilbert–Schmidt norm of the differential  $du(x)$ . The critical point of  $E$  is called “harmonic”. In a fundamental paper [4], Eells and Sampson established existence of smoothly harmonic maps from  $M$  to  $N$  assuming  $N$  has nonpositive section curvature. Let  $K \geq 0$  be an upper bound for the section curvature of  $N$  and  $B_\rho(q)$  the open geodesic ball in  $N$  with center  $q$  and radius  $\rho$ . Assuming essentially the size restriction

$$u(\partial M) \subset B_\rho(q), \quad \rho \leq \frac{\pi}{2\sqrt{K}}, \quad (1.2)$$

Hildebrandt et al. [10] showed existence of “small” smooth harmonic maps satisfying the condition (1.2) and also discovered that for  $n \geq 3$  the equator map  $u^* = (\frac{x}{|x|}, 0): B^n \rightarrow S^n$  is a weakly harmonic maps. The uniqueness of harmonic maps in [10] was later proved by Jäger and Kaul [11]. Many important contributions on the regularity of minimizing harmonic maps have been made since then. Schoen and Uhlenbeck [15, 16] obtained that the minimizer of  $E$  in  $H^{1,2}(M, N)$  is smooth except for a singular set, where the Hausdorff dimension of the singular set is less than or equal to  $n - 3$ . Meanwhile, Giaquinta and Giusti in [5,6] also proved this result for the case when the image lies in a coordinate chart (boundary regularity by Jost and Meier [13]). We would also like to mention Simon’s deep works on the structure of singularity of minimizing harmonic maps (e.g., [18]).

Let  $B^n$  be the unit ball in  $\mathbb{R}^n$  with boundary  $\partial B^n = S^{n-1}$ , where  $S^{n-1}$  is the unit sphere in  $\mathbb{R}^n$ .

A map  $u: B^n \rightarrow S^n$  is called “weakly harmonic” if  $u \in H^{1,2}(B^n, S^n)$  and it is a critical point of the energy  $\int_{B^n} |\nabla u|^2 dx$ , i.e.,

$$\int_{B^n} \nabla u \cdot \nabla \phi dx = \int_{B^n} |\nabla u|^2 u \cdot \phi dx$$

for all  $\phi \in H_0^{1,2}(B^n, \mathbb{R}^{n+1}) \cap L^\infty(B^n, \mathbb{R}^{n+1})$ .

In 1983, Jäger and Kaul [12] proved the following result:

**THEOREM (Jäger–Kaul).** –

- (i) When  $3 \leq n \leq 6$ , the equator map  $u^* = (\frac{x}{|x|}, 0)$  is unstable.
- (ii) When  $n \geq 7$ , the equator map  $u^*$  is a minimizer of the energy functional  $E(u, B^n) = \int_{B^n} |\nabla u|^2$  for all maps  $u \in H^{1,2}(B^n, S^n)$  with  $u = u^*$  on  $\partial B^n$ .

After this theorem, Giaquinta and Soucek [7] and Schoen and Uhlenbeck [17] proved that the Hausdorff dimension of the singular set of minimizing harmonic maps into a hemisphere is less than or equal to  $n - 7$ .

For any  $p \in \mathbb{R}$  with  $n > p \geq 2$ , we define the  $p$ -energy of maps in  $H^{1,p}(B^n, S^n)$  by

$$E_p(u, B^n) = \int_{B^n} |\nabla u|^p dx.$$

A map  $u \in H^{1,p}(B^n, S^n)$  is called “weakly  $p$ -harmonic” if  $u$  satisfies

$$\int_{B^n} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi dx = \int_{B^n} |\nabla u|^p u \cdot \phi dx$$

for all  $\phi \in H_0^{1,p}(B^n, \mathbb{R}^{n+1}) \cap L^\infty(B^n, \mathbb{R}^{n+1})$ . It is also easy to check that the equator map  $u^*$  is a weakly  $p$ -harmonic map from  $B^n$  to  $S^n$  for  $2 \leq p < n$ .

In this paper, we first generalize the Jäger–Kaul theorem to the  $p$ -energy by the following:

**THEOREM A.** – Assume that  $n > p \geq 2$ .

- (i) For  $3 \leq n < 2 + p + 2\sqrt{p}$ , the equator map  $u^* = (\frac{x}{|x|}, 0)$  is an unstable  $p$ -harmonic map from  $B^n$  to  $S^n$ .
- (ii) When  $n \geq 2 + p + 2\sqrt{p}$ , then  $u^*$  is a minimizer of the  $p$ -energy  $E_p$  in the class  $H^{1,p}(B^n, S^n)$  with boundary value  $u^*$  on  $\partial B^n$ . Moreover, if  $n \geq 2 + p + 2\sqrt{p}$ ,  $u^*$  is the unique minimizer.

We present a new proof of Theorem A(ii) and also point out that our proof is different from and much simpler than the original proof by Jäger and Kaul.

*Remark a.* – When  $p = 2$ ,  $n \geq 7 > 4 + 2\sqrt{2}$ . Thus the proof of Theorem A also extends the Helein theorem in [9, Theorem 3.1] for  $n \geq 9$

to the case when  $n \geq 7$ , i.e., for  $n \geq 7$ , we have

$$E(u) - E(u^*) \geq K_n \|u - u^*\|_{H^{1,2}(B^n)}^2,$$

where  $u$  is a map in  $H^{1,2}(B^n, S^n)$  which agree with  $u^*$  on  $\partial B^n$ , and  $K_n$  is a strict positive constant.

*Remark b.* – Consider the case of maps from  $B^n$  into  $S^{n-1}$ . Then  $u^* = \frac{x}{|x|}$  is a minimizer of  $E(u; B^n)$  for maps from  $B^n$  into  $S^{n-1}$ . This result has been proved by Brezis et al. [2] for  $n = 3$ , by Jäger and Kaul [12] for  $n \geq 7$  and by Lin [14] for  $n \geq 3$ . Moreover, Coron and Gulliver proved that  $u^* = \frac{x}{|x|}$  is a minimizer of the  $p$ -energy functional  $E_p(u; B^n)$  for maps from  $B^n$  into  $S^{n-1}$  for  $p \leq n - 1$ . Theorem A recovers the partial result of Coron and Gulliver [3] for  $n \geq 2 + p + 2\sqrt{p}$ . Perhaps, the uniqueness of the minimizer of  $p$ -energy functional of maps from  $B^n$  into  $S^{n-1}$  for  $n \geq 2 + p + 2\sqrt{p}$  is also a new result.

Let  $M = B^n$  again and let  $N$  be an ellipsoid of  $\mathbb{R}^{n+1}$ , i.e.,

$$N := \left\{ u = (v, z): |v|^2 + \frac{z^2}{a^2} = 1 \right\} \subset \mathbb{R}^{n+1},$$

where  $v \in \mathbb{R}^n$ ,  $z \in \mathbb{R}$ , and  $a > 0$  is a constant.

Baldes [1] in 1984 and Helein [8] in 1988 generalized the work of Jäger and Kaul when  $N$  is an ellipsoid:

THEOREM. –

- (i) (Baldes) When  $a^2 \geq 1$  and  $n \geq 7$ , the equator map  $u^*$  is the unique minimizer of the energy functional  $E$  in  $H^{1,2}(B^n; N)$  with boundary value  $(x, 0)$ .
- (ii) (Helein) If  $0 < a < 1$  and  $a^2 \geq 4(n-1)/(n-2)^2$ , the equator map  $u^*$  is the unique minimizer of the energy functional  $E$  in  $H^{1,2}(B^n; N)$  with boundary value  $(x, 0)$ .

We would point out that all proofs of Baldes [1] and Helein [8] are variants of the proof of Jäger and Kaul [12]. After the first version of this paper (CMA-Preprint, September 1996), the author was asked whether one could recover and generalize the results of Baldes and Helein using our proof of Theorem A. Here we generalize their results to  $p$ -harmonic maps with  $n > p \geq 2$  by the following:

**THEOREM B.** –

- (i) When  $a^2 \geq 1$  and  $n \geq 2 + p + 2\sqrt{p}$ , the equator map  $u^*$  is the unique minimizer of the  $p$ -energy functional  $E_p$  in  $H^{1,p}(B^n; N)$  with boundary value  $(x, 0)$ .
- (ii) If  $0 < a < 1$  and  $a \geq 4(n - 1)/(n - p)^2$ , the equator map  $u^*$  is the unique minimizer of the  $p$ -energy functional  $E_p$  in  $H^{1,p}(B^n; N)$  with boundary value  $(x, 0)$ .

**2. PROOF OF THEOREM A**

LEMMA 1. – For any  $p < n$ , we have

$$(n - p)^2 \int_{B^n} |\phi(x)|^2 \frac{1}{|x|^p} dx \leq 4 \int_{B^n} \frac{1}{|x|^{p-2}} \left| \frac{\partial \phi}{\partial r} \right|^2 dx$$

for all  $\phi \in H_0^{1,p} \cap L^\infty(B^n, \mathbb{R})$  with  $r = |x|$  where that equality occurs only if  $\phi = 0$ .

*Proof.* – The case of  $p = 2$  was proved in [1]. Integrating by parts and using Cauchy’s inequality, we have

$$\begin{aligned} \int_{B^n} |\phi|^2 \frac{1}{|x|^p} dx &= \int_{|\omega|=1} \int_0^1 \phi^2 r^{n-p-1} dr d\omega \\ &= \frac{2}{n - p} \int_{|\omega|=1} \int_0^1 \phi \frac{\partial \phi}{\partial r} r^{n-p} dr d\omega \\ &\leq \frac{1}{2} \int_{|\omega|=1} \int_0^1 \phi^2 r^{n-p-1} dr d\omega \\ &\quad + \frac{2}{(n - p)^2} \int_{|\omega|=1} \int_0^1 \left( \frac{\partial \phi}{\partial r} \right)^2 r^{n-p+1} dr d\omega \end{aligned}$$

for all  $\phi \in H_0^{1,p} \cap L^\infty(B^n, \mathbb{R})$ . The above inequality becomes equality iff

$$\phi = \frac{2}{(n - p)} \frac{\partial \phi}{\partial r},$$

this is possible only if  $\phi = 0$ . This proves our claim.  $\square$

*Remark c.* – Lemma 1 can be also proved in following way:

$$\begin{aligned} & \inf_{\phi \neq 0, \text{supp } \phi \subset \bar{B}^n \setminus \{0\}} \frac{\int_{B^n} |\nabla \phi|^2 r^{-(p-2)} dx}{\int_{B^n} r^{-p} |\phi|^2 dx} \\ &= \inf_{\tilde{\phi} \neq 0, \text{supp } \tilde{\phi} \subset (0,1]} \frac{\int_0^1 \left| \frac{\partial \tilde{\phi}}{\partial r} \right|^2 r^{n+1-p} dr}{\int_0^1 |\tilde{\phi}|^2 r^{n-1-p} dr} \\ &\geq \inf_{\tilde{\phi} \neq 0, \text{supp } \tilde{\phi} \subset (0,\infty)} \frac{\int_0^\infty \left| \frac{\partial \tilde{\phi}}{\partial r} \right|^2 r^{n+1-p} dr}{\int_0^\infty |\tilde{\phi}|^2 r^{n-1-p} dr} = \frac{(n-p)^2}{4}. \end{aligned}$$

This can be done by modifying a lemma taken from [17, Lemma 1.3].

## 2.1. Proof of Theorem A(ii)

Let  $u^* = (\frac{x}{|x|}, 0)$  be the “equator map” from  $B^n \rightarrow S^n$ . It is easy to see

$$|\nabla u^*|^2 = \frac{n-1}{r^2}.$$

Let  $w \in H^{1,2}(B^n, S^n)$  be any function with boundary value  $w = u^*$  on  $\partial B^n$ .

By Lemma 1, we obtain

$$\int_{B^n} |\nabla u^*|^{p-2} |\nabla \phi|^2 dx \geq \frac{(n-p)^2}{4(n-1)} \int_{B^n} |\nabla u^*|^p \phi^2 dx.$$

When  $n \geq 2 + p + \sqrt{4p}$ , we have

$$\frac{(n-p)^2}{4(n-1)} \geq 1.$$

When  $n \geq 2 + p + \sqrt{4p}$ , we get

$$\int_{B^n} |\nabla u^*|^{p-2} |\nabla(u^* - w)|^2 dx \geq \int_{B^n} |\nabla u^*|^p (u^* - w)^2 dx \quad (2.1)$$

for all  $w \in H^{1,p}(B^n, S^n)$  with  $w = u^*$  on  $\partial B^n$ . Moreover, we know

$$\int_{B^n} |\nabla u^*|^{p-2} |\nabla(u^* - w)|^2 dx$$

$$\begin{aligned}
 &= \int_{B^n} |\nabla u^*|^p \, dx - 2 \int_{B^n} |\nabla u^*|^{p-2} \nabla u^* \cdot \nabla w \, dx \\
 &\quad + \int_{B^n} |\nabla u^*|^{p-2} |\nabla w|^2 \, dx
 \end{aligned} \tag{2.2}$$

and

$$\int_{B^n} |\nabla u^*|^p (u^* - w)^2 \, dx = \int_{B^n} |\nabla u^*|^p (2 - 2u^* \cdot w) \, dx. \tag{2.3}$$

Notice that  $u^*$  is a weakly  $p$ -harmonic map, i.e.,

$$\int_{B^n} |\nabla u^*|^{p-2} \nabla u^* \cdot \nabla \phi \, dx = \int_{B^n} |\nabla u^*|^p u^* \cdot \phi \, dx$$

for all  $\phi \in H_0^{1,p}(B^n, \mathbb{R}^{n+1})$ . By taking  $\phi = u^* - w$ , we have

$$\int_{B^n} |\nabla u^*|^{p-2} \nabla u^* \cdot \nabla w \, dx = \int_{B^n} |\nabla u^*|^p u^* \cdot w \, dx. \tag{2.4}$$

From (2.1)–(2.4), we get for  $n \geq 2 + p + \sqrt{4p}$

$$\int_{B^n} |\nabla u^*|^{p-2} |\nabla w|^2 \, dx \geq \int_{B^n} |\nabla u^*|^p \, dx.$$

By the Hölder inequality, we have

$$\int_{B^n} |\nabla u^*|^{p-2} |\nabla w|^2 \, dx \leq \left( \int_{B^n} |\nabla u^*|^p \, dx \right)^{(p-2)/p} \left( \int_{B^n} |\nabla w|^p \, dx \right)^{2/p}.$$

Combing two above inequalities, we have

$$\int_{B^n} |\nabla u^*|^p \, dx \leq \int_{B^n} |\nabla w|^p \, dx$$

for all  $w \in H^{1,p}(B^n, S^n)$  with boundary value  $w = u^*$  on  $\partial B^n$  when  $n \geq 2 + p + 2\sqrt{p}$ .



From Lemma 1, we know that (2.1) become equality only if  $w = u^*$ . If  $n \geq 2 + p + 2\sqrt{p}$ , we have

$$\int_{B^n} |\nabla u^*|^p \, dx < \int_{B^n} |\nabla w|^p \, dx$$

for all  $w \in H^{1,p}(B^n, S^n)$  with boundary value  $w = u^*$  on  $\partial B^n$  and  $w \not\equiv u^*$ . It means that  $u^*$  is the unique minimizer for  $n \geq 2 + p + 2\sqrt{p}$ . This proves Theorem A(ii).

Assume that  $p = 2$ . When  $n \geq 7 > 4 + 2\sqrt{2}$ , one sees  $(n - 2)^2 / (4(n - 1)) > 1$ . By Lemma 1, we know

$$\int_{B^n} |\nabla \phi|^2 \, dx \geq \left(1 - \frac{4(n - 1)}{(n - 2)^2}\right) \int_{B^n} |\nabla \phi|^2 \, dx + \int_{B^n} |\nabla u^*|^2 \phi^2 \, dx$$

for all  $\phi \in H_0^{1,2}(B^n, \mathbb{R}^{n+1})$ . Taking  $\phi = w - u^*$ , we have

$$\int_{B^n} |\nabla w|^2 \, dx - \int_{B^n} |\nabla u^*|^2 \, dx \geq \left(1 - \frac{4(n - 1)}{(n - 2)^2}\right) \int_{B^n} |\nabla(w - u^*)|^2 \, dx,$$

where  $w \in H^{1,2}(B^n, S^n)$  agrees with  $u^*$  on  $\partial B^n$ . This proves our claim in Remark 1.

For  $t$  small we define  $u_t : B^n \rightarrow S^n$  by setting

$$u_t(x) = \frac{\left(\frac{x}{|x|}, t\phi(x)\right)}{(1 + t^2\phi^2)^{1/2}}$$

for a smooth function  $\phi$  on  $B^n$  vanishing near 0 and  $\partial B^n$ . A simple calculation gives

$$\frac{\partial \nabla u_t(x)}{\partial t} \Big|_{t=0} = (0, \dots, 0, \nabla \phi(x)), \quad \frac{\partial}{\partial t} |\nabla u_t|^2 \Big|_{t=0} = 0$$

and

$$\frac{\partial^2 \nabla u_t(x)}{\partial^2 t} \Big|_{t=0} = \phi^2 \left( \nabla \frac{x}{|x|}, 0 \right).$$

Now

$$\frac{1}{2} \frac{\partial^2}{\partial^2 t} |\nabla u_t|^2 \Big|_{t=0} = \left| \frac{\partial \nabla u_t(x)}{\partial t} \Big|_{t=0} \right|^2 + \left( \frac{\partial^2 \nabla u_t(x)}{\partial^2 t}, \nabla u_t \right) \Big|_{t=0}.$$

Then

$$\frac{d^2}{dt^2} E_p(u_t) \Big|_{t=0} = \int_{B^n} |\nabla u^*|^{p-2} [|\nabla \phi|^2 - \phi^2 |\nabla u^*|^2] dx.$$

It is easy to check that  $u^* = (\frac{x}{|x|}, 0)$  is a weakly  $p$ -harmonic map from  $B^n$  into  $S^n$ . If  $u^*$  is stable, we have

$$\int_{B^n} |\nabla u^*|^{p-2} [|\nabla \phi|^2 - \phi^2 |\nabla u^*|^2] dx \geq 0$$

for all smooth  $\phi$  vanishing near 0 and  $\partial B^n$ .

### 2.2. Proof of Theorem A(i)

Let us consider the following equation:

$$\begin{cases} \phi''(r) + \frac{n-p+1}{r} \phi'(r) + \frac{n-1-\varepsilon}{r^2} \phi(r) = 0, \\ \phi(r_0) = \phi(1) = 0. \end{cases} \tag{2.5}$$

for  $0 < r_0 < 1$ .

By setting  $\xi(t) = \phi(e^t)$ , Eq. (2.5) becomes

$$\xi''(t) + (n-p)\xi'(t) + (n-1-\varepsilon)\xi(t) = 0.$$

Let

$$v := v(\varepsilon) = \frac{1}{4} [n^2 - 2(p+2)n + p^2] + 1 + \varepsilon.$$

When  $3 \leq n < 2 + p + 2\sqrt{p}$ , there exists a small  $\varepsilon$  such that  $v < 0$  and we choose  $r_0$ :  $0 < r_0 < 1$  such that  $\sqrt{-v} \ln r_0$  is a multiple of  $2\pi$ . Then it is easily checked (see [12]) that the function

$$\phi(r) = \begin{cases} r^{(p-n)/2} \sin(\sqrt{-v} \cdot \ln r), & \text{for } r_0 < r \leq 1, \\ 0, & \text{for } r \leq r_0. \end{cases}$$

solves Eq. (2.5). This means that for  $3 \leq n < 2 + p + 2\sqrt{p}$ , there exists  $\varepsilon$  small and a non-zero  $\phi(r)$  on  $[r_0, 1]$ ,  $\phi(1) = \phi(r_0) = 0$ , such that

$$\int_0^1 r^{2-p} \left[ \phi'(r)^2 - \frac{(n-1)}{r^2} \phi^2 \right] r^{n-1} dr = - \int_0^1 \frac{\varepsilon}{r^2} r^{2-p} \phi^2 r^{n-1} dr < 0.$$

In other words, we see that  $u^*$  is unstable for  $3 \leq n < 2 + p + 2\sqrt{p}$ . This proves Theorem A(i).

From the proof of Theorem A, we have

**COROLLARY 2.** – Assume that  $u$  is a stable  $p$ -harmonic map from  $B^n$  into  $S^n$  and the values of  $u$  are on the equator  $(S^{n-1}, 0)$  of  $S^n$ . Then  $u$  is a local minimizer of the energy functional  $E_p$  in  $H^{1,p}(B^n, S^n)$ .

### 3. PROOF OF THEOREM B

In this section, let  $N$  be the ellipsoid of  $\mathbb{R}^{n+1}$  defined in Section 1 and suppose that  $p \in \mathbb{R}$  with  $n > p \geq 2$ . We define the  $p$ -energy of maps in  $H^{1,p}(B^n, N)$  by

$$E_p(v, z; B^n) = \int_{B^n} (|\nabla v|^2 + |\nabla z|^2)^{p/2} dx.$$

We write  $u = (v, z)$  with  $v \in \mathbb{R}^n, z \in \mathbb{R}$ . A map  $u = (v, z) \in H^{1,p}(B^n, N)$  is called “weakly  $p$ -harmonic” if  $u$  satisfies in the sense of distributions the following equations:

$$\begin{aligned} \operatorname{div}(|\nabla u|^{p-2} \nabla v) + |\nabla u|^{p-2} \lambda v &= 0, \\ \operatorname{div}(|\nabla u|^{p-2} \nabla z) + |\nabla u|^{p-2} \lambda \frac{z}{a^2} &= 0, \end{aligned}$$

where

$$\lambda = \left( |\nabla v|^2 + \frac{|\nabla z|^2}{a^2} \right) \frac{a^4}{a^4 v^2 + z^2}.$$

*Proof of Theorem B.* – Let  $u = (v, z)$  be any function in  $H^{1,p}(B^n, N)$  with boundary values  $(x, 0)$  on  $\partial B^n$ . Using Lemma 1 again, we have

$$\int_{B^n} \left| \nabla \frac{x}{|x|} \right|^{p-2} \left| \nabla \left( \frac{x}{|x|} - v \right) \right|^2 dx$$

$$\geq \frac{(n-p)^2}{4(n-1)} \int_{B^n} \left| \nabla \frac{x}{|x|} \right|^p \left| \frac{x}{|x|} - v \right|^2 dx \tag{3.1}$$

and

$$\int_{B^n} \left| \nabla \frac{x}{|x|} \right|^{p-2} |\nabla z|^2 dx \geq \frac{(n-p)^2 a^2}{4(n-1)} \int_{B^n} \left| \nabla \frac{x}{|x|} \right|^p \frac{|z|^2}{a^2} dx. \tag{3.2}$$

(i) Assume that  $a \geq 1$  and  $n \geq 2 + p + 2\sqrt{p}$ . Note the fact  $v^2 + \frac{z^2}{a^2} = 1$ . Thus using (3.1) and (3.2), the proof of Theorem A(ii) yields

$$\int_{B^n} \left| \nabla \frac{x}{|x|} \right|^{p-2} |\nabla u|^2 dx \geq \frac{(n-p)^2}{4(n-1)} \int_{B^n} \left| \nabla \frac{x}{|x|} \right|^p dx$$

for all  $u = (v, z)$  with same boundary values  $(x, 0)$ . The same argument in the proof of Theorem A(ii) gives that  $(\frac{x}{|x|}, 0)$  is the unique minimizer of  $E_p$  if  $n \geq 2 + p + 2\sqrt{p}$ .

(ii) Assume that  $0 < a < 1$  and  $a^2 \geq 4(n-1)/(n-p)^2$ . Thus using (3.1) and (3.2) again, we get

$$\int_{B^n} \left| \nabla \frac{x}{|x|} \right|^{p-2} |\nabla u|^2 dx > \frac{(n-p)^2 a^2}{4(n-1)} \int_{B^n} \left| \nabla \frac{x}{|x|} \right|^p dx$$

for all  $u = (v, z) \neq (\frac{x}{|x|}, 0)$  with same boundary values. The same argument in the proof of Theorem A(ii) gives that  $(\frac{x}{|x|}, 0)$  is the unique minimizer of  $E_p$ . This proves Theorem B.  $\square$

*Remark d.* – It is obvious from the proof of Theorem A(i) to see that  $(\frac{x}{|x|}, 0)$  is unstable for  $E_p$  when  $a^2 < 4(n-p)/(n-2)^2$ .

### REFERENCES

- [1] A. BALDES, Stability and uniqueness properties of the equator map from a ball into an ellipsoid, *Math. Z.* 185 (1984) 505–516.
- [2] H. BREZIS, J.-M. CORON and E.H. LIEB, Harmonic maps with defects, *Comm. Math. Phys.* 107 (1986) 649–705.
- [3] J.-M. CORON and R. GULLIVER, Minimizing  $p$ -harmonic maps into spheres, *J. Reine Angew. Math.* 401 (1989) 82–100.
- [4] J. EELLS and J.H. SAMPSON, Harmonic mappings of Riemannian manifolds, *Amer. J. Math.* 86 (1964) 109–160.

- [5] M. GIAQUINTA and E. GIUSTI, On the regularity of the minima of variational integrals, *Acta Math.* 148 (1982) 31–46.
- [6] M. GIAQUINTA and E. GIUSTI, The singular set of the minima of certain quadratic functionals, *Ann. Scuola Norm. Sup. Pisa* (4) 11 (1984) 45–55.
- [7] M. GIAQUINTA and J. SOUCEK, Harmonic maps into a hemisphere, *Ann. Scuola Norm. Sup. Pisa* (4) 12 (1985) 81–90.
- [8] F. HELEIN, Regularity and uniqueness of harmonic maps into an ellipsoid, *Manuscripta Math.* 60 (1988) 235–257.
- [9] F. HELEIN, Approximations of Sobolev maps between an open set and an Euclidean sphere, boundary data and singularities, *Math. Ann.* 285 (1989) 125–140.
- [10] S. HILDEBRANDT, H. KAUL and K.-O. WIDMAN, An existence theorem for harmonic mappings of Riemannian manifolds, *Acta Math.* 138 (1977) 1–16.
- [11] W. JÄGER and H. KAUL, Uniqueness and stability of harmonic maps and their Jacobi fields, *Manuscripta Math.* 28 (1979) 269–291.
- [12] W. JÄGER and H. KAUL, Rotationally symmetric harmonic maps from a ball into a sphere and the regularity problem for weak solutions of elliptic systems, *J. Reine Angew. Math.* 343 (1983) 146–161.
- [13] J. JOST and M. MEIER, Boundary regularity for minima of certain quadratic functionals, *Math. Ann.* 262 (1983) 549–561.
- [14] F.-H. LIN, Une remarque sur l'application  $x/|x|$ , *C. R. Acad. Sci. Paris* 305 (1987) 529–531.
- [15] R. SCHOEN and K. UHLENBECK, A regularity theory for harmonic maps, *J. Differential Geom.* 17 (1982) 307–335.
- [16] R. SCHOEN and K. UHLENBECK, Boundary theory and the Dirichlet problem for harmonic maps, *J. Differential Geom.* 18 (1983) 253–268.
- [17] R. SCHOEN and K. UHLENBECK, Regularity of minimizing harmonic maps into the sphere, *Invent. Math.* 78 (1984) 89–100.
- [18] L. SIMON, *Theorems on Regularity and Singularity of Energy Minimizing Maps*, Lecture Notes in Mathematics, ETH Zürich, Birkhäuser, Basel, 1996.