

ANN VERDOODT

Continued fractions for finite sums

Annales mathématiques Blaise Pascal, tome 1, n° 2 (1994), p. 71-84

http://www.numdam.org/item?id=AMBP_1994__1_2_71_0

© Annales mathématiques Blaise Pascal, 1994, tous droits réservés.

L'accès aux archives de la revue « Annales mathématiques Blaise Pascal » (<http://math.univ-bpclermont.fr/ambp/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

CONTINUED FRACTIONS FOR FINITE SUMS

Ann Verdoodt

Abstract

Our aim in this paper is to construct continued fractions for sums of the type $\sum_{i=0}^n b_i z^{c(i)}$ or $\sum_{i=0}^n b_i/z^{c(i)}$, where (b_n) is a sequence such that b_n is different from zero if n is different from zero, and $c(n)$ is an element of \mathbb{N} .

Résumé

Le but est de construire des fractions continues pour des sommes du type $\sum_{i=0}^n b_i z^{c(i)}$ or $\sum_{i=0}^n b_i/z^{c(i)}$, où (b_n) est une suite telle que b_n est différent de zéro pour n différent de zéro, et $c(n)$ est un élément de \mathbb{N} .

1. Introduction

$[a_0, a_1, a_2, \dots]$ denotes the continued fraction $a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$,

and $[a_0, a_1, \dots, a_n]$ denotes $a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots + \frac{1}{a_{n-1} + \frac{1}{a_n}}}}$.

The a_i 's are called the partial quotients (or simply the quotients), and $[a_0, a_1, \dots, a_n]$ is called a finite continued fraction .

Our aim in this paper is to construct continued fractions for sums of the type $\sum_{i=0}^n b_i z^{c(i)}$ or

$\sum_{i=0}^n b_i/z^{c(i)}$, where $c(i)$ is an element of \mathbb{N} .

In section 2 , we find continued fractions for finite sums of the type $\sum_{i=0}^n b_i z^i$ ($c(i) = i$)

or $\sum_{i=0}^n b_i z^{q^i}$ ($c(i) = q^i$), where (b_n) is a sequence such that b_n is different from zero if n is different from zero , and where q is a natural number different from zero and one .

Therefore , we start by giving a continued fraction for the sum $\sum_{i=0}^n b_i T^{3^i}$, where b_i is different from zero for all i different from zero (b_i is a constant in T) . This can be found in theorem 1 .

If we replace b_i by $b_i z^i$ in theorem 1 , and we put T equal to one , we find a continued

fraction for $\sum_{i=0}^n b_i z^i$ (theorem 2), and if we replace b_i by $b_i z^{q^i}$ in theorem 1 , and we put

T equal to one , we find a continued fraction for $\sum_{i=0}^n b_i z^{q^i}$ (theorem 3) (q is a natural number different from zero and one) .

In section 3 we find continued fractions for finite sums of the type $\sum_{i=0}^n \frac{b_i}{z^{c(i)}}$, for some sequences (b_n) and $(c(n))$, where $c(n)$ is a natural number .

In theorem 4 , we find a result for $c(i)$ equal to 2^i (for all i) .

Finally , in theorem 5 , we give a continued fraction for $\sum_{i=0}^v \frac{b_i}{z^{c(i)}}$, where $c(0)$ equals zero , and $c(n+1) - 2c(n) \geq 0$.

The results in this paper are extensions of results that can be found in [2] , [3] and [4] .

Acknowledgement : I thank professor Van Hamme for the help and the advice he gave me during the preparation of this paper .

2. Continued fractions for sums of the type $\sum_{i=0}^n b_i z^i$

All the proofs in sections 2 and 3 can be given with the aid of the following simple lemma :

Lemma

$$\text{Let i) } p_0 = a_0, \quad q_0 = 1, \quad p_1 = a_1 a_0 + 1, \quad q_1 = a_1, \\ p_n = a_n p_{n-1} + p_{n-2}, \quad q_n = a_n q_{n-1} + q_{n-2} \quad (n \geq 2),$$

then we have

$$\text{ii) } \frac{p_n}{q_n} = [a_0, a_1, \dots, a_n]$$

$$\text{iii) } p_n q_{n-1} - p_{n-1} q_n = (-1)^{n-1} \quad (n \geq 1)$$

$$\text{iv) } \frac{q_n}{q_{n-1}} = [a_n, a_{n-1}, \dots, a_1] \quad (n \geq 1)$$

These well-known results can e.g. be found in [1].

First we give a continued fraction for the sum $\sum_{i=0}^n b_i T^{3^i}$, where b_i is different from zero for all i different from zero (b_i is a constant in T):

Theorem 1

Let (b_n) be a sequence such that $b_n \neq 0$ for all $n > 0$.

Define a sequence (x_n) by putting $x_0 = [b_0 T]$, $x_1 = [b_0 T, b_1^{-1} T^{-3}]$, and if

$$x_n = [a_0, a_1, \dots, a_{2n-1}] \text{ then setting } x_{n+1} = [a_0, a_1, \dots, a_{2n-1}, -b_n^2/b_{n+1} T^{-3^n}, -a_{2n-1}, \dots, -a_1].$$

$$\text{Then } x_n = \sum_{i=0}^n b_i T^{3^i} \text{ for all } n \in \mathbb{N}.$$

Proof

For $n = 0$ the theorem clearly holds.

$$\text{If } n \text{ is at least one, we prove that } x_n = \sum_{i=0}^n b_i T^{3^i} \text{ and } q_{2n-1} = b_n^{-1} T^{-3^n}.$$

We prove this by induction. For $n = 1$ the assertion holds.

Suppose it holds for $1 \leq n \leq j$. We then prove the assertion for $n = j+1$.

$$x_{j+1} = [a_0, a_1, \dots, a_{2j+1-1}]$$

$$= [a_0, a_1, \dots, a_{2j-1}, a_{2j}, -[a_{2j-1}, \dots, a_1]] \quad (\text{using the definition of a continued fraction})$$

$$= \frac{-q_{2j-1} p_{2j} + q_{2j-2} p_{2j-1}}{-q_{2j-1} q_{2j} + q_{2j-2} q_{2j-1}} \quad (\text{by i), ii) and iv) of the lemma})$$

$$= \frac{-q_{2j-1} (a_{2j} p_{2j-1} + p_{2j-2}) + q_{2j-2} p_{2j-1}}{-q_{2j-1} (a_{2j} q_{2j-1} + q_{2j-2}) + q_{2j-2} q_{2j-1}} \quad (\text{by i) of the lemma})$$

$$\text{now we have } p_{2j-1} q_{2j-2} - p_{2j-2} q_{2j-1} = (-1)^{2j-2} = 1 \quad (\text{by iii) of the lemma})$$

$$= \frac{p_{2j-1}}{q_{2j-1}} - \frac{1}{a_{2j} (q_{2j-1})^2}$$

$$\text{now } a_{2j} (q_{2j-1})^2 = -T^{-3j} \frac{b_j^2}{b_{j+1}} (b_j^{-1} T^{-3j})^2 = -T^{-3j+1} b_{j+1}^{-1}$$

$$= [a_0, a_1, \dots, a_{2j-1}] + T^{3j+1} b_{j+1} = \sum_{i=0}^{j+1} b_i T^{3i} \quad (\text{by the induction hypothesis})$$

We still have to prove $q_{2j+1-1} = b_{j+1}^{-1} T^{-3(j+1)}$. Let k be at least one.

Then p_k and q_k are polynomials in $U = T^{-1}$. $\deg q_k > \deg q_{k-1}$, and the term with the highest degree in q_k is given by $a_k \cdot a_{k-1} \cdot \dots \cdot a_1$. This follows from i).

If r is a polynomial in U that divides p_k and q_k , then r must be a constant in U . This

immediately follows from iii). If r divides p_k and q_k , then r divides $(-1)^{k-1}$. So r must be a constant.

$$\text{Since } \sum_{i=0}^{j+1} b_i T^{3i} = [a_0, a_1, \dots, a_{2j+1-1}] = \frac{p_{2j+1-1}}{q_{2j+1-1}}, \text{ we have}$$

$$\frac{p_{2j+1-1}}{q_{2j+1-1}} = \sum_{i=0}^{j+1} \frac{b_i T^{3i} T^{-3j+1}}{T^{-3j+1}} = \sum_{i=0}^{j+1} \frac{b_i U^{3j+1-3i}}{U^{3j+1}} = \frac{b_{j+1} + \sum_{i=0}^j b_i U^{3j+1-3i}}{U^{3j+1}}$$

and we conclude that $q_{2j+1-1} = C U^{3j+1} = C T^{-3j+1}$ where C is a constant.

By the previous remark, we have that

$$q_{2j+1-1} = C T^{-3j+1} = C U^{3j+1} = a_1 \cdot a_2 \cdot \dots \cdot a_{2j+1-1}$$

$$= (-1)^{2j-1} (a_1 \cdot a_2 \cdot \dots \cdot a_{2j-1})^2 \cdot a_{2j} = -(q_{2j-1})^2 \cdot a_{2j}$$

$$(\text{by the induction hypothesis, since } q_{2j-1} = b_j^{-1} T^{-3j} = a_1 \cdot a_2 \cdot \dots \cdot a_{2j-1})$$

$$= - (b_j^{-1} T^{-3j})^2 \cdot (- T^{-3j} \frac{b_i^2}{b_{j+1}}) = \frac{T^{-3j+1}}{b_{j+1}} \quad \text{which we wanted to prove .}$$

We immediately have the following

Proposition

Let $x_0 = [a_0]$, $x_1 = [a_0, a_1]$ and if $x_n = [a_0, a_1, \dots, a_{2n-1}]$, then

$$x_{n+1} = [a_0, a_1, \dots, a_{2n-1}, a_{2n}, -a_{2n-1}, \dots, -a_1] .$$

If n is at least two, then the continued fraction of x_n consists only of the partial quotients

$a_{2n-1}, a_{2n-2}, -a_{2n-2}, \dots, a_1, -a_1$ and a_0 .

Then the distribution of the partial quotients for x_n is as follows ($n \geq 2$):

partial quotient

$$a_{2n-1} \quad a_{2n-2} \quad -a_{2n-2} \quad a_{2n-3} \quad -a_{2n-3} \quad \dots \quad a_{2i} \quad -a_{2i} \quad \dots \quad a_1 \quad -a_1 \quad a_0$$

number of occurrences

$$1 \quad 1 \quad 1 \quad 2 \quad 2 \quad \dots \quad 2^{n-i-2} \quad 2^{n-i-2} \quad \dots \quad 2^{n-2} \quad 2^{n-2} \quad 1$$

Proof

We give a proof by induction on n .

$$x_2 = [a_0, a_1, a_2, a_3] = [a_0, a_1, a_2, -a_1] , \text{ so the quotients } a_0, a_1, -a_1, a_2, \text{ occur once .}$$

So for n equal to 2 the assertion holds . Suppose it holds for $2 \leq n \leq j$. Then we prove it holds

for $n = j+1$. Since $x_{j+1} = [a_0, a_1, \dots, a_{2j+1-1}] = [a_0, a_1, \dots, a_{2j-1}, a_{2j}, -a_{2j-1}, \dots, -a_1]$, it is clear that the partial quotients a_{2j} and a_0 occur only once .

In the partial quotients a_1, \dots, a_{2j-1} we have

partial quotient

$$a_{2j-1} \quad a_{2j-2} \quad -a_{2j-2} \quad a_{2j-3} \quad -a_{2j-3} \quad \dots \quad a_{2i} \quad -a_{2i} \quad \dots \quad a_1 \quad -a_1$$

number of occurrences

$$1 \quad 1 \quad 1 \quad 2 \quad 2 \quad \dots \quad 2^{j-i-2} \quad 2^{j-i-2} \quad \dots \quad 2^{j-2} \quad 2^{j-2}$$

so in the partial quotients $-a_1, \dots, -a_{2j-1}$ we have

partial quotient

$$-a_{2j-1} \quad a_{2j-2} \quad -a_{2j-2} \quad a_{2j-3} \quad -a_{2j-3} \quad \dots \quad a_{2i} \quad -a_{2i} \quad \dots \quad a_1 \quad -a_1$$

number of occurrences

$$1 \quad 1 \quad 1 \quad 2 \quad 2 \quad \dots \quad 2^{j-i-2} \quad 2^{j-i-2} \quad \dots \quad 2^{j-2} \quad 2^{j-2}$$

This proves the proposition .

Using theorem 1, we immediately have the following :

Theorem 2

Let (b_n) be a sequence such that b_n is different from zero for all n different from zero .

Define a sequence (x_n) by putting $x_0 = [b_0]$, $x_1 = [b_0, b_1^{-1}z^{-1}]$ and if $x_n = [a_0, a_1, \dots, a_{2n-1}]$

then setting $x_{n+1} = [a_0, a_1, \dots, a_{2n-1}, -b_n^2 / b_{n+1} z^{n-1}, -a_{2n-1}, \dots, -a_1]$,

then $x_n = \sum_{i=0}^n b_i z^i$ for all $n \in \mathbb{N}$.

Proof

Replace b_i by $b_i z^i$ in theorem 1, and put T equal to one .

Some examples

1) Let $x_n = \sum_{i=0}^n x^i$ (i.e. $b_i = 1$ for all i) . Then $a_0 = 1$, $a_1 = x^{-1}$ and $a_{2n} = -x^{n-1}$ ($n \geq 1$)

2) Let $x_n = \sum_{i=0}^n \frac{x^i}{i!}$ (i.e. $\lim_{n \rightarrow \infty} x_n = e^x$) .

Then $a_0 = 1$, $a_1 = x^{-1}$ and $a_{2n} = -\frac{n+1}{n!} x^{n-1}$ ($n \geq 1$)

3) Let $x_n = \sum_{i=0}^n \frac{(-1)^i x^{2i}}{(2i)!}$ (i.e. $\lim_{n \rightarrow \infty} x_n = \cos x$) .

Then $a_0 = 1$, $a_1 = -2x^{-2}$ and $a_{2n} = (-1)^n \frac{(2n+2)(2n+1)}{(2n)!} x^{2n-2}$ ($n \geq 1$)

4) Let $x_n = \sum_{i=0}^n \frac{(-1)^i x^{2i+1}}{(2i+1)!}$ (i.e. $\lim_{n \rightarrow \infty} x_n = \sin x$) .

Then $a_0 = x$, $a_1 = -6x^{-3}$ and $a_{2n} = (-1)^n \frac{(2n+3)(2n+2)}{(2n+1)!} x^{2n-1}$ ($n \geq 1$)

In an analogous way as in the previous theorem , we have

Theorem 3

Let (b_n) be a sequence such that b_n is different from zero for all n different from zero, and let q be a natural number different from zero and one.

Define a sequence (x_n) by putting $x_0 = [b_0 z]$, $x_1 = [b_0 z, b_1^{-1} z^{-q}]$ and if $x_n = [a_0, a_1, \dots, a_{2n-1}]$ then setting $x_{n+1} = [a_0, a_1, \dots, a_{2n-1}, -b_n^2/b_{n+1} z^{-q(q-2)}, -a_{2n-1}, \dots, -a_1]$.

Then $x_n = \sum_{i=0}^n b_i z^{qi}$ for all $n \in \mathbb{N}$.

Proof

Replace b_i by $b_i z^{qi}$ in theorem 1, and put T equal to one.

An Example

In [4] we find the following :

Let F_q be the finite field of cardinality q . Let $A = F_q[X]$, $K = F_q(X)$, $K_\infty = F_q((1/X))$

and let Ω be the completion of an algebraic closure of K_∞ . Then A, K, K_∞, Ω are well-known analogues of $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ respectively.

Let $[i] = X^{qi} - X$ (the symbol $[i]$ does not have the same meaning as in $x_0 = [a_0]$). This is just the product of monic irreducible elements of A of degree dividing i .

Let $D_0 = 1, D_i = [i] D_{i-1}^{q-1}$ if $i > 0$. This is the product of monic elements of A of degree i .

Let us introduce the following function : $e(Y) = \sum_{i=0}^{\infty} \frac{Y^{qi}}{D_i}$ ($Y \in \Omega$).

Then Thakur gives the following theorem :

Define a sequence x_n by setting $x_1 = [0, Y^{-q} D_1]$ and if $x_n = [a_0, a_1, \dots, a_{2n-1}]$ then setting

$x_{n+1} = [a_0, a_1, \dots, a_{2n-1}, -Y^{-q(q-2)} D_{n+1}/D_n^2, -a_{2n-1}, \dots, -a_1]$, then $x_n = \sum_{i=1}^n \frac{Y^{qi}}{D_i}$ for all $n \in \mathbb{N}$.

In particular, $e(Y) = Y + \lim_{n \rightarrow \infty} x_n$.

If we put $b_i = D_i^{-1}$ if $i > 0$, and $b_0 = 0$ in theorem 3, then we find the result of Thakur.

3. Continued fractions for sums of the type $\sum_{i=0}^n \frac{b_i}{z^{c(i)}}$

In this section, b_i is a constant in z , and $c(i)$ is a natural number. Our first theorem in this

section gives the continued fraction for the sum $\sum_{i=0}^n \frac{b_i}{z^{2^i}}$ (i.e. $c(i) = 2^i$ for all i):

Theorem 4

Let (b_n) be a sequence such that b_n is different from zero for all n . A continued fraction for

the sum $\sum_{i=0}^n \frac{b_i}{z^{2^i}}$ can be given as follows:

Put $x_0 = [0, z/b_0]$, $x_1 = [0, \frac{z}{b_0} - \frac{b_1}{b_0^2}, \frac{b_0^3 z}{b_1^2} + \frac{b_0^2}{b_1}]$ and if $x_k = [a_0, a_1, \dots, a_{2k}]$ then setting

$x_{k+1} = [a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}, \gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1}, a_{2k+2}, \dots, a_{2k+1}]$ where $\gamma_{k+1} = b_{k+1} \frac{(b_0)^{2^{k+1}}}{(b_1)^{2^{k+1}}}$,

$a_{2k+i} = \gamma_{k+1}^2 a_{2k-i+1}$ if i is even, and $a_{2k+i} = \gamma_{k+1}^{-2} a_{2k-i+1}$ if i is odd ($2 \leq i \leq 2^k$),

then $x_k = \sum_{i=0}^k \frac{b_i}{z^{2^i}}$ for all $k \in \mathbb{N}$.

Proof

If we have $x_n = [a_0, a_1, \dots, a_{2^n}] = \frac{p_{2^n}}{q_{2^n}}$, we show by induction that x_n equals $\sum_{i=0}^n \frac{b_i}{z^{2^i}}$, and

that q_{2^n} equals $z^{2^n} \frac{b_0^{2^n}}{b_1^{2^n}}$. For $n = 0, 1$ this follows by an easy calculation.

Suppose the assertion holds for $0 \leq n \leq k$. Then we show it holds for $n = k+1$.

The first part of the proof, i.e. showing that $x_{k+1} = \sum_{i=0}^{k+1} \frac{b_i}{z^{2^i}}$ is analogous to the first part of the proof of [2], theorem 1.

$$x_{k+1} = [a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}, \gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1}, a_{2k+2}, \dots, a_{2k+1}]$$

$$= [a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}, \gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1}, \gamma_{k+1}^2 [a_{2k-1}, a_{2k-2}, a_{2k-3}, \dots, a_2, a_1]]$$

(using the definition of a continued fraction)

Now if $[a_0, a_1, \dots, a_{2k}] = \frac{p_{2k}}{q_{2k}}$, then $[a_0, a_1, \dots, a_{2k-1}] = \frac{p_{2k-1}}{q_{2k-1}}$ and so

$$[a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}] = \frac{(a_{2k} + \gamma_{k+1})p_{2k-1} + p_{2k-2}}{(a_{2k} + \gamma_{k+1})q_{2k-1} + q_{2k-2}} = \frac{p_{2k} + \gamma_{k+1}p_{2k-1}}{q_{2k} + \gamma_{k+1}q_{2k-1}}$$

(by i) and ii) of the lemma)

$$\text{Then } [a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}, \gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1}] = \frac{(\gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1})(p_{2k} + \gamma_{k+1}p_{2k-1}) + p_{2k-1}}{(\gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1})(q_{2k} + \gamma_{k+1}q_{2k-1}) + q_{2k-1}}$$

(by i) and ii) of the lemma)

And so

$$[a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}, \gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1}, \gamma_{k+1}^2 [a_{2k-1}, a_{2k-2}, a_{2k-3}, \dots, a_2, a_1]]$$

$$= \frac{a_{2k} q_{2k-1} p_{2k} + \gamma_{k+1} a_{2k} q_{2k-1} p_{2k-1} - \gamma_{k+1} q_{2k-1} p_{2k} + q_{2k-2} p_{2k} + \gamma_{k+1} q_{2k-2} p_{2k-1}}{a_{2k} q_{2k-1} q_{2k} + \gamma_{k+1} a_{2k} q_{2k-1} q_{2k-1} - \gamma_{k+1} q_{2k-1} q_{2k} + q_{2k-2} q_{2k} + \gamma_{k+1} q_{2k-2} q_{2k-1}}$$

(by iv) of the lemma)

If we use the following equalities

$$(p_n - p_{n-2})q_{n-1} = a_n p_{n-1} q_{n-1}$$

$$(q_n - q_{n-2})p_n = a_n p_n q_{n-1}$$

$$(q_n - q_{n-2})q_n = a_n q_n q_{n-1}$$

$$(q_n - q_{n-2})q_{n-1} = a_n q_{n-1}^2 \quad (\text{ by i) of the lemma })$$

then we find that the numerator equals $q_{2k} p_{2k} + \gamma_{k+1}$ (by iii) of the lemma) and the denominator equals $(q_{2k})^2$.

So we conclude

$$x_{k+1} = \frac{p_{2k}}{q_{2k}} + \frac{\gamma_{k+1}}{(q_{2k})^2} = \sum_{i=0}^k \frac{b_i}{z^{2i}} + \frac{(b_1)^{2k+1}}{z^{2k+1}(b_0)^{2k+1}} b_{k+1} \frac{(b_0)^{2k+1}}{(b_1)^{2k+1}} = \sum_{i=0}^{k+1} \frac{b_i}{z^{2i}}$$

$$\text{We still have to show } q_{2k+1} = z^{2k+1} \frac{(b_0)^{2k+1}}{(b_1)^{2k+1}}.$$

In the same way as in the proof of theorem 1, we find that $q_{2k+1} = C z^{2k+1}$ where C is a constant .

Let α_i be the coefficient of z in a_i .

Then for C , the coefficient of z^{2k+1} in q_{2k+1} . we have

$$C = \alpha_1 \alpha_2 \dots \alpha_{2k-1} \alpha_{2k} (\gamma_{k+1}^{-2} \alpha_{2k}) (\gamma_{k+1}^2 \alpha_{2k-1}) (\gamma_{k+1}^{-2} \alpha_{2k-2}) (\gamma_{k+1}^2 \alpha_{2k-3}) \dots (\gamma_{k+1}^2 \alpha_1)$$

$$= (\alpha_1 \alpha_2 \dots \alpha_{2k-1} \alpha_{2k})^2 = (\text{coefficient of } z^{2k} \text{ in } q_{2k})^2 = \left(\frac{(b_0)^{2k}}{(b_1)^{2k}} \right)^2 = \frac{(b_0)^{2k+1}}{(b_1)^{2k+1}}$$

and we conclude $q_{2k+1} = z^{2k+1} \frac{(b_0)^{2k+1}}{(b_1)^{2k+1}}$. This finishes the proof .

Some examples

1) If we put b_i equal to one for all i , and z is an integer at least 3, then we find theorem 1 of [2]:

$$\text{Let } B(u,v) = \sum_{i=0}^v \frac{1}{u^{2i}} = \frac{1}{u} + \frac{1}{u^2} + \frac{1}{u^4} + \dots + \frac{1}{u^{2v}} \quad (u \geq 3, u \text{ an integer})$$

Then $B(u,0) = [0,u]$, $B(u,1) = [0,u-1,u+1]$, and if $B(u,v) = [a_0, a_1, \dots, a_n] = \frac{p_n}{q_n}$

then $B(u,v+1) = [a_0, a_1, \dots, a_{n-1}, a_n+1, a_n-1, a_{n-1}, a_{n-2}, \dots, a_2, a_1]$.

2) Put $b_i = \lambda^i$. Then we have $x_0 = [0, u]$, $x_1 = [0, u - \lambda, \frac{u}{\lambda^2} + \frac{1}{\lambda}]$ and if $x_k = [a_0, a_1, \dots, a_{2k}]$,

then $x_{k+1} = [a_0, a_1, \dots, a_{2k-1}, a_{2k} + \gamma_{k+1}, \gamma_{k+1}^{-2} a_{2k} - \gamma_{k+1}^{-1}, a_{2k+2}, \dots, a_{2k+1}]$, where $\gamma_{k+1} = \lambda^{k+1-2k+1}$,

$a_{2k+i} = \gamma_{k+1}^2 a_{2k-i+1}$ if i is even, and $a_{2k+i} = \gamma_{k+1}^{-2} a_{2k-i+1}$ if i is odd ($2 \leq i \leq 2k$),

$$\text{then } x_k = \sum_{i=0}^k \frac{\lambda^i}{u^{2i}} \text{ for all } k \in \mathbb{N}.$$

For some some sequences (b_n) and $(c(n))$, we can give a continued fraction for the sum

$$\sum_{i=0}^v \frac{b_i}{z^{c(i)}} \text{ as follows:}$$

Theorem 5

Let (b_n) be a sequence such that $b_n \neq 0$ for all n , and $b_0 \neq 0, 1, -1$, and $1/2$, and let $(c(n))$

be a sequence such that $c(0) = 0$, and $c(n+1) - 2c(n) \geq 0$.

$$\text{Put } x_0 = [-b_0^2, \frac{1}{b_0} - 1, \frac{1}{b_0} + 1] = [a_0, a_1, a_2] = \frac{p_2}{q_2} = \frac{p(0)}{q(0)},$$

$$\text{and if } x_v = [a_0, a_1, \dots, a_n] = \frac{p_n}{q_n} = \frac{p(v)}{q(v)},$$

then setting $x_{v+1} = [a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1, 1, a_n - 1, a_{n-1}, \dots, a_2, a_1]$,

where $d(v) = c(v+1) - 2c(v)$, $\alpha_v = \frac{b_v^2}{b_{v+1}}$ if $v \geq 1$ and $\alpha_0 = \frac{b_0^4}{b_1}$,

$$\text{then } x_v = \sum_{i=0}^v \frac{b_i}{z^{c(i)}} \text{ for all } v \text{ in } \mathbb{N}, \text{ and } q(v) = \frac{z^{c(v)}}{b_v} \text{ if } v \geq 1, q(0) = \frac{1}{(b_0)^2}$$

Remarks

- 1) The special form of b_0 , $x_0 = b_0 = [-b_0^2, \frac{1}{b_0} - 1, \frac{1}{b_0} + 1] = [a_0, a_1, a_2]$ is needed since in the expression $[a_0, a_1, \dots, a_n] = \frac{p_n}{q_n}$ the integer n must be even .
- 2) The value of n is $n = 2^{v+1} + 2^v + 2$ (this can be easily seen by induction)
- 3) The only partial quotients that appear are $-b_0^2, \frac{1}{b_0} - 1, \frac{1}{b_0} + 1, \frac{1}{b_0}, \frac{1}{b_0} - 2, \alpha_v z^{d(v)} - 1$, and 1 ,
so b_0 must be different from $0, 1, -1$, and $1/2$.

Proof

For v equal to $0, 1$ or 2 we find this result by an easy computation .

We prove the theorem by induction on v .

Suppose we have $x_v = \sum_{i=0}^v \frac{b_i}{z^{c(i)}} = [a_0, a_1, \dots, a_n] = \frac{p_n}{q_n} = \frac{p_{(v)}}{q_{(v)}}$ with $q_{(v)} = \frac{z^{c(v)}}{b_v}$

Then we show that $x_{v+1} = [a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1, 1, a_n - 1, a_{n-1}, \dots, a_2, a_1] = \sum_{i=0}^{v+1} \frac{b_i}{z^{c(i)}}$

with $q_{(v+1)} = \frac{z^{c(v+1)}}{b_{v+1}}$.

The first part of the proof , i.e. showing that $x_{v+1} = \sum_{i=0}^{v+1} \frac{b_i}{z^{c(i)}}$, is analogous to the first part of the proof of the theorem in [3] .

Now , by repeated use of i) an ii) of the lemma , we have

$$[a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1] = \frac{(\alpha_v z^{d(v)} - 1)p_n + p_{n-1}}{(\alpha_v z^{d(v)} - 1)q_n + q_{n-1}} ;$$

$$[a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1, 1] = \frac{\alpha_v z^{d(v)} p_n + p_{n-1}}{\alpha_v z^{d(v)} q_n + q_{n-1}} ;$$

$$[a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1, 1, a_n - 1] = \frac{a_n \alpha_v z^{d(v)} p_n + a_n p_{n-1} - p_n}{a_n \alpha_v z^{d(v)} q_n + a_n q_{n-1} - q_n}$$

$$x_{v+1} = [a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1, 1, a_n - 1, a_{n-1}, \dots, a_1]$$

$$= [a_0, a_1, \dots, a_n, \alpha_v z^{d(v)} - 1, 1, a_n - 1, [a_{n-1}, \dots, a_1]]$$

(using the definition of a continued fraction)

$$= \frac{a_n q_{n-1} \alpha_v z^{d(v)} p_n + q_{n-2} \alpha_v z^{d(v)} p_n + a_n q_{n-1} p_{n-1} - q_{n-1} p_n + q_{n-2} p_{n-1}}{a_n q_{n-1} \alpha_v z^{d(v)} q_n + q_{n-2} \alpha_v z^{d(v)} q_n + a_n (q_{n-1})^2 - q_{n-1} q_n + q_{n-2} q_{n-1}}$$

(by i) , ii) and iv) of the lemma)

$$= \frac{p_n}{q_n} + \frac{1}{(q_n)^2 \alpha_v z^{d(v)}} \quad \text{(by i) and iii) of the lemma since n is even)}$$

$$\begin{aligned} \text{So } x_{v+1} &= \frac{p_n}{q_n} + \frac{1}{(q_n)^2 \alpha_v z^{d(v)}} = \sum_{i=0}^v \frac{b_i}{z^{c(i)}} + \frac{(b_v)^2 b_{v+1}}{z^{2c(v)} (b_v)^2 z^{d(v)}} \quad \text{since } q_n = q_{(v)} = \frac{z^{c(v)}}{b_v}, \alpha_v = \frac{(b_v)^2}{b_{v+1}} \\ &= \sum_{i=0}^{v+1} \frac{b_i}{z^{c(i)}} \end{aligned}$$

We still have to prove $q_{(v+1)} = q_{2n+2} = \frac{z^{c(v+1)}}{b_{v+1}}$, and since $\frac{z^{c(v+1)}}{b_{v+1}} = (q_n)^2 \alpha_v z^{d(v)}$, it suffices to prove that $q_{2n+2} = (q_n)^2 \alpha_v z^{d(v)}$.

We can not use the same trick here as in the proofs of theorems 1 and 4, since we do not necessarily have $\deg q_{k+1} > \deg q_k$ (q_k as a polynomial in z)

We already know that $q_{n+1} = (\alpha_v z^{d(v)} - 1)q_n + q_{n-1}$, $q_{n+2} = \alpha_v z^{d(v)} q_n + q_{n-1}$

Repeated use of i) of the lemma gives

$$q_{n+3} = q_{(n+2)+1} = a_n \alpha_v z^{d(v)} q_n + a_n q_{n-1} - q_n = r_1 \alpha_v z^{d(v)} q_n - q_{n-2} \quad \text{(where we put } a_n = r_1 \text{)}$$

$$q_{n+4} = q_{(n+2)+2} = (a_{n-1} a_n + 1) \alpha_v z^{d(v)} q_n - a_{n-1} q_{n-2} + q_{n-1} = r_2 \alpha_v z^{d(v)} q_n + q_{n-3}$$

(where we put $a_{n-1} a_n + 1 = r_2$)

$$\begin{aligned} q_{n+5} &= q_{(n+2)+3} = (a_{n-2}(a_{n-1} a_n + 1) + a_n) \alpha_v z^{d(v)} q_n + a_{n-2} q_{n-3} - q_{n-2} \\ &= r_3 \alpha_v z^{d(v)} q_n - q_{n-4} \quad \text{(where we put } a_{n-2}(a_{n-1} a_n + 1) + a_n = r_3 \text{)} \end{aligned}$$

etc...

Continuing this way, we find

$$q_{(n+2)+k} = r_k \alpha_v z^{d(v)} q_n + (-1)^k q_{n-(k+1)}, \quad q_{(n+2)+k+1} = r_{k+1} \alpha_v z^{d(v)} q_n + (-1)^{k+1} q_{n-(k+2)}$$

$$\begin{aligned} \text{Then } q_{(n+2)+k+2} &= (a_{n-(k+1)} r_{k+1} + r_k) \alpha_v z^{d(v)} q_n + (-1)^{k+1} a_{n-(k+1)} q_{n-k-2} + (-1)^k q_{n-k-1} \\ &= r_{k+2} \alpha_v z^{d(v)} q_n + (-1)^{k+2} q_{n-(k+3)} \end{aligned}$$

and finally we have $q_{2n} = q_{(n+2)+n-2} = r_{n-2} \alpha_v z^{d(v)} q_n + q_{n-(n-1)}$

$$q_{2n+1} = q_{(n+2)+n-1} = r_{n-1} \alpha_v z^{d(v)} q_n - q_{n-n} \quad (\text{we remark that } n \text{ is even})$$

$$\text{and so } q_{2n+2} = q_{(n+2)+n} = r_n \alpha_v z^{d(v)} q_n - a_1 q_0 + q_1 = r_n \alpha_v z^{d(v)} q_n$$

So if we want to show that $q_{2n+2} = (q_n)^2 \alpha_v z^{d(v)}$, we must show that r_n equals q_n .

For the sequence (r_n) we have $r_0 = 1$, $r_1 = a_n$, $r_2 = a_{n-1} a_n + 1 = a_{n-1} r_1 + r_0$,

$$r_3 = a_{n-2} (a_{n-1} a_n + 1) + a_n = a_{n-2} r_2 + r_1, \text{ and continuing this way we find } r_{k+2} = a_{n-(k+1)} r_{k+1} + r_k.$$

From this it follows that $[1, a_n, \dots, a_1] = [1, c_1, \dots, c_n] = \frac{t_n}{r_n}$ (we put $a_i = c_{n+1-i}$)

$$\text{with } t_0 = c_0, \quad r_0 = 1, \quad t_1 = c_1 c_0 + 1, \quad r_1 = c_1,$$

$$t_n = c_n t_{n-1} + t_{n-2}, \quad r_n = c_n r_{n-1} + r_{n-2} \quad (n \geq 2),$$

Now n can be written as $n = 2k+2$ (see remark 2 following theorem 5) and so

$$[a_0, a_1, \dots, a_n] = [a_0, a_1, \dots, a_k, \alpha_{v-1} z^{d(v-1)} - 1, 1, a_k - 1, a_{k-1}, \dots, a_1] = \frac{p_n}{q_n}$$

$$\text{and then } [1, a_1, \dots, a_k, \alpha_{v-1} z^{d(v-1)} - 1, 1, a_k - 1, a_{k-1}, \dots, a_1] = [1, a_1, \dots, a_n] = \frac{p'_n}{q_n}$$

where the q_i ($0 \leq i \leq n$) stay the same since q_i does not depend on a_0 .

$$\text{So } [1, a_1, \dots, a_{k-1}, a_k - 1, 1, \alpha_{v-1} z^{d(v-1)} - 1, a_k, a_{k-1}, \dots, a_1] = [1, a_n, \dots, a_1] = \frac{t_n}{r_n}$$

and we conclude $q_i = r_i$ for $0 \leq i \leq k-1$.

We have to show $q_n = r_n$. Now (by repeated use of i) of the lemma)

$$q_k = a_k q_{k-1} + q_{k-2}, \quad r_k = q_k - q_{k-1};$$

$$q_{k+1} = \alpha_{v-1} z^{d(v-1)} q_k - q_k + q_{k-1}, \quad r_{k+1} = q_k;$$

$$q_{k+2} = \alpha_{v-1} z^{d(v-1)} q_k + q_{k-1}, \quad r_{k+2} = \alpha_{v-1} z^{d(v-1)} q_k - q_{k-1};$$

$$\begin{aligned} q_{k+3} &= q_{(k+2)+1} = \alpha_{v-1} z^{d(v-1)} a_k q_k + a_k q_{k-1} - q_k = a_k \alpha_{v-1} z^{d(v-1)} q_k - q_{k-2} \\ &= R_1 \alpha_{v-1} z^{d(v-1)} q_k - q_{k-2}, \text{ where we put } a_k = R_1, \end{aligned}$$

$$r_{k+3} = r_{(k+2)+1} = a_k \alpha_{v-1} z^{d(v-1)} q_k + q_{k-2} = R_1 \alpha_{v-1} z^{d(v-1)} q_k + q_{k-2};$$

$$\begin{aligned} q_{k+4} &= q_{(k+2)+2} = (a_{k-1} a_k + 1) \alpha_{v-1} z^{d(v-1)} q_k - a_{k-1} q_{k-2} + q_{k-1} \\ &= (a_{k-1} a_k + 1) \alpha_{v-1} z^{d(v-1)} q_k + q_{k-3} \end{aligned}$$

$$= R_2 \alpha_{v-1} z^{d(v-1)} q_k + q_{k-3} \text{ where we put } (a_{k-1} a_k + 1) = R_2,$$

$$r_{k+4} = r_{(k+2)+2} = (a_{k-1}a_k+1)\alpha_{v-1}z^{d(v-1)}q_k + a_{k-1}q_{k-2} - q_{k-1}$$

$$= (a_{k-1}a_k+1)\alpha_{v-1}z^{d(v-1)}q_k - q_{k-3} = R_2\alpha_{v-1}z^{d(v-1)}q_k - q_{k-3}$$

....

If we continue this way, we find $q_{(k+2)+i} = R_i\alpha_{v-1}z^{d(v-1)}q_k + (-1)^i q_{k-(i+1)}$, and

$$r_{(k+2)+i} = R_i\alpha_{v-1}z^{d(v-1)}q_k - (-1)^i q_{k-(i+1)} \quad (0 \leq i \leq k, R_0 = 1), \text{ and so we have}$$

$$q_{2k} = q_{(k+2)+k-2} = R_{k-2}\alpha_{v-1}z^{d(v-1)}q_k + q_{k-(k-1)}, \quad q_{2k+1} = q_{(k+2)+k-1} = R_{k-1}\alpha_{v-1}z^{d(v-1)}q_k - q_{k-k} \quad (\text{we}$$

remark that k is even) and thus $q_{2k+2} = q_{(k+2)+k} = R_k\alpha_{v-1}z^{d(v-1)}q_k - a_1q_0 + q_1 = R_k\alpha_{v-1}z^{d(v-1)}q_k$,

$$\text{and } r_{2k} = r_{(k+2)+k-2} = R_{k-2}\alpha_{v-1}z^{d(v-1)}q_k - q_{k-(k-1)}, \quad r_{2k+1} = r_{(k+2)+k-1} = R_{k-1}\alpha_{v-1}z^{d(v-1)}q_k + q_{k-k} \quad \text{and}$$

$$\text{thus } r_{2k+2} = r_{(k+2)+k} = R_k\alpha_{v-1}z^{d(v-1)}q_k + a_1q_0 - q_1 = R_k\alpha_{v-1}z^{d(v-1)}q_k,$$

So we conclude that $q_{2k+2} = q_n$ equals $r_{2k+2} = r_n$. This finishes the proof.

The case b_i equal to one, where z is an integer at least two, is studied by Shallit ([3]):

Let $(c(k))$ be a sequence of positive integers such that $c(v+1) \geq 2c(v)$ for all $v \geq v'$, where v' is a non-negative integer. Let $d(v) = c(v+1) - 2c(v)$. Define $S(u, v)$ as follows:

$$S(u, v) = \sum_{i=0}^v u^{-c(i)}, \text{ where } u \text{ is an integer, } u \geq 2. \text{ Then Shallit proved the following theorem:}$$

Suppose $v \geq v'$. If $S(u, v) = [a_0, a_1, \dots, a_n]$ and n is even, then

$$S(u, v+1) = [a_0, a_1, \dots, a_n, u^{d(v)-1}, 1, a_{n-1}, a_{n-1}, a_{n-2}, \dots, a_2, a_1].$$

References

- [1] G. H. Hardy and E. M. Wright, An Introduction to the Theory of Numbers, Oxford University Press, 1979.
- [2] J. O. Shallit, Simple Continued Fractions for Some Irrational Numbers, Journal of Number Theory, vol. 11 (1979), p. 209-217.
- [3] J. O. Shallit, Simple Continued Fractions for Some Irrational Numbers II, Journal of Number Theory, vol. 14 (1982), p. 228-231.
- [4] D. S. Thakur, Continued Fraction for the Exponential for $F_q[t]$, Journal of Number Theory, vol. 41 (1992), p. 150-155.

Ann VERDOODT

Vrije Universiteit Brussel,
Faculty of Applied Sciences
Pleinlaan 2, B - 1050 Brussels
Belgium