ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

RAGHAVAN NARASIMHAN

On holomorphic functions of polynomial growth in a bounded domain

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3e série, tome 21, nº 2 (1967), p. 161-166

<http://www.numdam.org/item?id=ASNSP_1967_3_21_2_161_0>

© Scuola Normale Superiore, Pisa, 1967, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (http://www.sns.it/it/edizioni/riviste/annaliscienze/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

ON HOLOMORPHIC FUNCTIONS OF POLYNOMIAL GROWTH IN A BOUNDED DOMAIN

by RAGHAVAN NARASIMHAN (*)

Let Ω be a bounded open set in $\mathbb{C}^n$. For $z \in \Omega$, let

$$d(z) = d(z, \Omega)$$

denote the distance of the point z from the boundary $\partial\Omega$ of Ω.

DEFINITION. *A holomorphic function f in Ω is said to be of polynomial growth if there exist constants $M, \varrho > 0$ such that, for $z \in \Omega$ we have*

$$|f(z)| \leq M d(z)^{-\varrho};$$

M and ϱ may depend on f.

The object of this note is to prove a theorem which is preliminary to certain questions on the existence of sections of coherent analytic sheaves on Stein spaces satisfying conditions of growth similar to the one occurring in the definition above. The result is the following.

THEOREM. *Let g be holomorphic in $\overline{\Omega}$* (i. e. *in a neighborhood of $\overline{\Omega}$ and let f be a function of polynomial growth in Ω. If (g does not vanish identically on any non-empty open set in Ω and) f/g is holomorphic in Ω, then f/g has polynomial growth.*

PROOF. Let $h = f/g$. It is sufficient to prove that for any $a \in \partial\Omega$ there is a neighborhood U and constants $M, \varrho > 0$ with

$$|h(z)| \leq M d(z)^{-\varrho} \quad \text{for} \quad z \in \Omega \cap U.$$

We may suppose that $a = 0$. Further, it is sufficient to prove our inequality after a linear change of coordinates in $\mathbb{C}^n$. Thus, by the Weierstrass preparation theorem, we may make a linear change of coordinates such that

---

Pervenuto alla Redazione il 12 Ottobre 1966.

(*) Supported by Air Force Grant AF-AFOSR-1071-66.

g assumes the form

$$g(z) = \omega(z) P(z); P(z) = z_n^p + \sum_{\nu=1}^{p} a_\nu(z_1, \dots, z_{n-1}) z_n^{p-\nu}, a_\nu(0, \dots, 0) = 0,$$

here ω and P are holomorphic on the closure $\bar{U}$ of a neighborhood U of 0 in $\mathbb{C}^n$ and ω has no zeros on $\bar{U}$. Further, we may suppose that $U = U' \times U''$, $U' \subset \mathbb{C}^{n-1}$, $U'' \subset \mathbb{C}$ and that, if $z' \in \bar{U}'$, $z' = (z_1, \dots, z_{n-1})$, and $P(z'; t) = 0$, $t \in \mathbb{C}$, then $t \in U''$. [The notation $P(z'; t)$, $z' \in U'$, is self-explanatory].

We have only to obtain an estimate of the form

$$\left| \frac{f}{P}(z) \right| = | \omega(z) h(z) | \leq M d(z)^{-\varrho} \quad \text{for} \quad z \in \Omega \cap U.$$

Let V be a neighborhood of $\bar{\Omega}$ in which g is holomorphic; and let $A = \{z \in V \mid g(z) = 0\}$. We may clearly suppose that $0 \in A$ and that $U \subset V$. For $z = (z', z_n) \in U' \times U'' = U$, we define

$$\delta_A(z) = \min_\nu | z_n - z_n^{(\nu)} |$$

where $z_n^{(\nu)}$, $\nu = 1, \dots, p$ are the zeros of the polynomial $P(z'; t)$ in t.

LEMMA 1. For $z \in U$, we have

$$| P(z) | \geq \delta_A(z)^p.$$

PROOF OF LEMMA 1. Given $z' \in U'$, let $\lambda_1, \dots, \lambda_p$ be the zeros of $P(z'; t)$. Then

$$| P(z) | = | (z_n - \lambda_1) \dots (z_n - \lambda_p) | \geq \delta_A(z)^p,$$

since, by definition, we have $| z_n - \lambda_j | \geq \delta_A(z)$ for each j.

Let $0 < \varepsilon \leq 1/2$ and let

$$\Delta_1 = \{z \in \Omega \cap U \mid \varepsilon d(z) \leq \delta_A(z)\},$$

$$\Delta_2 = \{z \in \Omega \cap U \mid \varepsilon d(z) > \delta_A(z)\}.$$

Clearly Δ_2 is a neighborhood of $A \cap U \cap \Omega$.

Since f has polynomial growth, there are constants $M, \varrho > 0$ with

$$| f(z) | \leq M d(z)^{-\varrho}, z \in \Omega.$$

Hence, for $z \in \Delta_1$, we have

$$\left| \frac{f}{P}(z) \right| \leq Md(z)^{-\varrho} \, | P(z) |^{-1} \leq Md(z)^{-\varrho} \, \delta_A(z)^{-p} \leq M\varepsilon^{-p} \, d(z)^{-\varrho-p}$$

by Lemma 1 and the definition of Δ_1.

To deal with Δ_2, we proceed as follows. For given $\zeta' = (\zeta_1, \dots, \zeta_{n-1}) \in U'$, let L_ζ, denote the complex line

$$z_1 = \zeta_1, \dots, z_{n-1} = \zeta_{n-1}.$$

Clearly $L_{\zeta'} \cap A$ contains at most p points for any $\zeta' \in U'$.

LEMMA 2. $\Delta_2 \cap L_{\zeta'}$ is relatively compact in $\Omega \cap L_{\zeta'}$ (considered as open sets in the plane $L_{\zeta'}$).

PROOF OF LEMMA 2. Suppose that $\{z^{(k)}\}$, $(k = 1, 2, \dots)$ is a sequence of points of $\Delta_2 \cap L_{\zeta'}$ converging to a point $z^{(0)} \in \partial\Omega$. Since $\delta_A(z^{(0)}) = \lim_{k \to \infty} \delta_A(z^{(k)}) \leq$ $\leq \lim_{k \to \infty} \varepsilon d(z^{(k)}) = 0$, $z^{(0)} \in A$. If then W is a small neighborhood of $z^{(0)}$ not containing any other point of $L_{\zeta'} \cap A$, we see that for $z \in W \cap L_{\zeta'}$ we have

$$\delta_A(z) = | z_n - z_n^{(0)} | = | z - z^{(0)} | \geq d(z, \Omega) = d(z)$$

since $z^{(0)} \in \partial\Omega$. In particular, if k is large enough,

$$d(z^{(k)}) \leq \delta_A(z^{(k)}) < \varepsilon d(z^{(k)})$$

by definition of Δ_2, which is absurd. Hence no sequence of points of $\Delta_2 \cap L_{\zeta'}$ can converge to a point of $\partial\Omega$, which proves Lemma 2.

LEMMA 3. If $\zeta' \in U'$, then, for any points x_1, x_2 lying in the same connected component of $\Delta_2 \cap L_{\zeta'}$, we have

$$\varepsilon^{2p} \, d(x_1) \leq d(x_2) \leq \varepsilon^{-2p} \, d(x_1).$$

PROOF OF LEMMA 3. Let $z^{(1)}, \dots, z^{(m)}$, $m \leq p$, be the points in $\Delta_2 \cap L_{\zeta'}$ and let $D^{(j)} = \{z \in L_{\zeta'} \cap \Omega \mid | z_n - z_n^{(j)} | < \varepsilon d(z)\}$ $(j = 1, \dots, m)$. If $z \in D^{(j)}$, we have

$$d(z) \leq d(z^{(j)}) + | z - z^{(j)} | = d(z^{(j)}) + | z_n - z_n^{(j)} | < d(z^{(j)}) + \varepsilon d(z),$$

so that

$$d(z) < (1 - \varepsilon)^{-1} \, d(z^{(j)}) < \varepsilon^{-1} \, d(z^{(j)}).$$

On the other hand

$$d(z) \geq d(z^{(j)}) - |z - z^{(j)}| > d(z^{(j)}) - \varepsilon\, d(z),$$

so that

$$d(z^{(j)}) < (1+\varepsilon)\, d(z) < \varepsilon^{-1}\, d(z).$$

Hence, if x, y are points of $D^{(j)}$, we have

$$\varepsilon^2\, d(x) < d(y) < \varepsilon^{-2}\, d(x).$$

Since, by Lemma 2, $\Delta_2 \cap L_{\zeta'} \subset D^{(1)} \cup \ldots \cup D^{(m)}$, if D is any connected component of $\Delta_2 \cap L_{\zeta'}$ and $x_1, x_2 \in D$, there exist indices $j_1, \ldots, j_r$, $r \leq m \leq p$, and points $y_1, \ldots, y_{r-1}$ with $y_\nu \in D^{(j_\nu)} \cap D^{(j_{\nu+1})}$ such that

$$x_1,\ x_2 \in D^{(j_1)} \cup \ldots \cup D^{(j_r)}.$$

If now $x_1, x_2 \in D$, we may suppose that $x_1 \in D^{(j_1)}$ and $x_2 \in D^{(j_s)}$, $s \leq r$. Then, by what we have seen above, if $s \geq 2$,

$$d(x_1) \leq \varepsilon^{-2}\, d(y_1) \leq \varepsilon^{-4}\, d(y_2) \leq \ldots \leq \varepsilon^{-2(s-1)}\, d(y_{s-1}) \leq \varepsilon^{-2s}\, d(x_2),$$

so that $d(x_1) \leq \varepsilon^{-2p}\, d(x_2)$, and this is true also if $s = 1$. In the same way, $d(x_2) \leq \varepsilon^{-2p}\, d(x_1)$, and this proves Lemma 3.

We now proceed to the proof of our theorem.

If Δ is any open subset of $L_{\zeta'}$ for $\zeta' \in U'$, we denote by $\partial\Delta$ its boundary in $L_{\zeta'}$. We claim that if ε is sufficiently small, then for all ζ' near $0 \in U'$, $\partial(\Delta_2 \cap L_{\zeta'}) \subset \Delta_1$. Because of Lemma 2, it is sufficient to show that $\partial(\Delta_2 \cap L_{\zeta'})$ does not meet $\zeta' \times \partial U''$. Now, if ζ' is restricted to a relatively compact subset U_1' of U', then $\zeta' \times \partial U''$ has a distance from $L_{\zeta'} \cap A$ which is bounded below. Hence $\delta_A(z)$ is bounded below on $(U_1' \times \partial U'') \cap \Omega$; since $d(z)$ is bounded above, if ε is small enough, no point of $\zeta' \times \partial U''$, $\zeta' \in U_1'$, can have points of $\Delta_2 \cap L_{\zeta'}$ arbitrarily close to it. Hence, for $\zeta' \in U_1'$, if ε is small, $\partial(\Delta_2 \cap L_{\zeta'}) \subset \Delta_1$.

If $z \in \Delta_2 \cap (U_1' \times U'')$, let $\zeta' = (z_1, \ldots, z_{n-1})$ and let D be the connected component of $\Delta_2 \cap L_{\zeta'}$ containing z. Since $\dfrac{f}{P}$ is holomorphic in D, we obtain, by the maximum principle,

$$\left| \frac{f}{P}(z) \right| \leq \sup_{x \in \partial D} \left| \frac{f}{P}(x) \right|,$$

Since $\partial D \subset \partial(\Delta_2 \cap L_{\zeta'}) \subset \Delta_1$, Lemma 1 gives, for $x \in \partial D$,

$$\left|\frac{f}{P}(x)\right| \leq M\, d(x)^{-\varrho}\, \delta_A(x)^{-p} \leq \frac{M}{\varepsilon^p}\, d(x)^{-\varrho-p} \quad (\text{since} \quad x \in \Delta_1)$$

$$\leq \frac{M}{\varepsilon^p}\, \varepsilon^{-2p(\varrho+p)}\, d(z)^{-\varrho-p} \quad (\text{by Lemma 3}).$$

Hence, for $z \in \Delta_2 \cap (U_1' \times U'')$,

$$\left|\frac{f}{P}(z)\right| \leq \sup_{x \in \partial D} \left|\frac{f}{P}(z)\right| \leq M_1\, d(z)^{-\varrho_1}, \quad M_1 = M\, \varepsilon^{-p-2p(\varrho+p)}, \quad \varrho_1 = \varrho + p.$$

Since for $z \in \Delta_1$, we have

$$\left|\frac{f}{P}(z)\right| \leq M\, \varepsilon^{-p}\, d(z)^{-\varrho-p} \leq M_1\, d(z)^{-\varrho_1}$$

we have

$$\left|\frac{f}{P}(z)\right| \leq M_1\, d(z)^{-\varrho_1} \quad \text{for} \quad z \in \Omega \cap (U_1' \times U'').$$

As already remarked, this proves the theorem.

REMARK 1. It is easily seen that Theorem 1 when $f \equiv 1$ (case when g has no zeros in Ω but may have zeros in $\bar{\Omega}$) is equivalent to the following statement:

Let g be holomorphic in an open set $U \subset \mathbb{C}^n$. Let $A = \{z \in U \mid g(z) = 0\}$ and $d(z, A)$ be the distance of $z \in \Omega$ from A. Then, for any compact set $K \subset U$, there exist constants c, $\varrho > 0$ with

$$|g(z)| \geq c\, d(z, A)^{\varrho} \quad \text{for} \quad z \in K.$$

This is a very special case of a famous inequality of Lojasiewicz which asserts the validity of a similar inequality for all *real* analytic (and even *semi-analytic*) functions in open sets in $\mathbb{R}^n$; see e. g. S. Lojasiewicz: *Ensembles sémi-analytiques*; Inst. Hautes Etudes Sc. Paris (1965).

REMARK 2. The assumption that g be holomorphic in $\bar{\Omega}$ is essential. For example, if $g(z) = \exp(-1/z)$ in $\Omega = \{z \in \mathbb{C} \mid \operatorname{Re} z > 0,\ |z| < 1\}$, we have $|g(z)| < 1$, $g(z) \neq 0$ in Ω, but $1/g(z)$ tends to infinity like $\exp(1/d(z))$ when $z \to 0$ along the real axis.

If $\Omega = \left\{ z \in \mathbb{C} \mid |\operatorname{Im} z| < \frac{\pi}{2} |z|^2, \ |z| < 1 \right\}$ and $g(z) = \exp\left(-\exp\left(\frac{1}{z}\right)\right)$, again $|g(z)| < 1$ in Ω. This time $1/g(z) \longrightarrow \infty$ like $\exp\{\exp(c/d(z))\}$ for a suitable $c > 0$ when $z \longrightarrow 0$ along the real axis.

It is possible to give examples where the growth is even faster.

REMARK 3. We have actually shown that there are constants $C, \alpha > 0$ depending only on g such that if $|f(z)| \leq M\, d(z)^{-\varrho}$ and f/g is holomorphic in Ω, then $\left| \frac{f}{g}(z) \right| \leq MC^{\varrho+1}\, d(z)^{-\varrho-\alpha}$. Somewhat better inequalities can be obtained, but this requires more complicated reasoning.

The Institute for Advanced Study, Princeton