

ANNALI DELLA  
SCUOLA NORMALE SUPERIORE DI PISA  
*Classe di Scienze*

D. D. J. HACON

**Manifolds of the homotopy type of a bouquet of spheres**

*Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3<sup>e</sup> série*, tome 24,  
n° 4 (1970), p. 703-715

[http://www.numdam.org/item?id=ASNSP\\_1970\\_3\\_24\\_4\\_703\\_0](http://www.numdam.org/item?id=ASNSP_1970_3_24_4_703_0)

© Scuola Normale Superiore, Pisa, 1970, tous droits réservés.

L'accès aux archives de la revue « *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

# MANIFOLDS OF THE HOMOTOPY TYPE OF A BOUQUET OF SPHERES

By D. D. J. HACON

## 1. Introduction.

This note is concerned with manifolds homotopy equivalent to  $\vee S_i$  (a bouquet of spheres of varying dimensions). In this connection a useful concept is that of *thickening* which is a homotopy generalization of the idea of regular neighbourhood. The reader is referred to [7] for definitions in the general case. Here we shall be concerned with the specific problem of describing the set of thickenings of  $\vee S_i$  and it will be convenient to adopt a definition of thickening that differs slightly from that to be found in [6] (see § 2).

In [3] Haefliger classified thickenings of (simply-connected) bouquets of spheres subject to certain dimensional restrictions. Our purpose here is to improve on Haefliger's result in the piecewise-linear case and deal with the non-simply-connected case, reducing it to the problem of classifying concordance classes of embeddings of a number of solid tori in a certain manifold, as follows.

Denote by  $P^n$  the solid  $n$ -pretzel i. e. an  $n$ -ball with a finite number of 1-handles attached orientably. Then the classification of thickenings is reduced to the classification of concordance classes of embeddings of the disjoint union of solid tori in  $\partial P$ , which is a simpler question. For instance, if the bouquet in question is simply-connected then  $\partial P$  is a sphere and the problem is now to classify concordance classes of embeddings of solid tori in a sphere. See [3]. If, on the other hand, the bouquet consists of a circle and a sphere we need to look at knots of a solid torus in  $S^1 \times S^q$  ( $q$  being  $\dim P - 2$ ).

Suppose  $f: \vee S_i \rightarrow W$  is a homotopy equivalence (and hence a simple homotopy equivalence since the Whitehead group of a free group is trivial). If  $f$  is homotopic to a piecewise linear embedding  $g: \vee S_i \rightarrow W$  proceed as follows. Take a regular neighbourhood  $N$  of  $g \vee S_i$  in  $\text{Int } W$ , the interior of

$W$  (if  $g \vee S_i$  meets  $\partial W$  isotop it into  $\text{Int } W$ ).  $N$  is homeomorphic to  $W$ , for by the  $s$ -cobordism theorem [1],  $W - \text{Int } N$  is homeomorphic to  $\partial N \times [0, 1]$ . We thus obtain a handlebody decomposition of  $W$  suffixed by the cell structure of  $\vee S_i$ .

In general, however, there exist homotopy equivalences  $f: \vee S_i \rightarrow W$  which are *not* homotopic to an embedding and consequently the above procedure cannot be followed. But a theorem of Stallings [5] allows us to factor  $f: \vee S_i \rightarrow W$  up to homotopy through a simple homotopy equivalence  $f': \vee S_i \rightarrow N$  where  $N$  is a  $p_N + 1$ -dimensional polyhedron in  $W$ . We seek a simple description of  $N$  in terms of  $\vee S_i$  which will (as in the case when  $f$  is an embedding) provide a handlebody decomposition of  $W$  suffixed by the cell structure of  $\vee S_i$ . In fact it will be shown (§ 3) that, if  $f: \vee S_i \rightarrow W$  is a homotopy equivalence,  $W$  may be expressed as  $P$  plus handles of index two or more and that handles of sufficiently large index are attached disjointly i. e. after a certain point in the construction of  $W$  the order in which handles are subsequently attached is immaterial.

## 2. The main theorem.

Throughout we restrict ourselves to the piecewise linear ( $PL$ ) category [7].

Write  $\cup S_i$  for the disjoint union  $S_1 \cup, \dots, \cup S_N$  of spheres  $S_1, \dots, S_N$  of dimensions  $p_1, \dots, p_N$  subject to the condition  $1 \leq p_1 \leq, \dots, \leq p_N$ . Let  $* = (*, \dots, *)$  be a point of  $S_1 \times, \dots, \times S_N$ . Then  $\vee S_i = S_1 \vee, \dots, \vee S_N$  is the subpolyhedron

$$(S_1 \times \{*\} \times, \dots, \times \{*\}) \cup, \dots, \cup (\{*\} \times, \dots, \times \{*\} \times S_N) \text{ of } S_1 \times, \dots, \times S_N.$$

Let  $\pi: \cup S_i \rightarrow \vee S_i$  be the obvious identification map. If we write  $\vee S_i$  in the form  $S^1 \vee, \dots, \vee S^1 \vee S_1 \vee, \dots, \vee S_N$  it is understood that  $p_i \geq 2$ .

Now let  $M$  be a compact, connected, oriented manifold with nonempty boundary  $\partial M$  and such that

$$(1) \quad \dim M \geq \text{Max}(6, \dim \cup S_i + 3)$$

$$(2) \quad \partial M \subset M \text{ induces an isomorphism of fundamental groups.}$$

We will be considering pairs  $(M, f)$ ,  $M$  as above and  $f: \cup S_i \rightarrow M$  homotopic to  $g \circ \pi$  where  $g: \vee S_i \rightarrow M$  is a homotopy equivalence. We call such a pair  $(M, f)$  a *thickening*.

REMARKS (1). Since  $M$  is assumed connected any map  $f: \cup S_i \rightarrow M$  factors up to homotopy through  $\pi: \cup S_i \rightarrow \vee S_i$ .

(2) Suppose  $f: \cup S_i \rightarrow M$  factors up to homotopy through a homotopy equivalence  $g: \vee S_i \rightarrow M$ . Let  $g': \vee S_i \rightarrow M$  be any other homotopy factorization. Then  $g'$  is also a homotopy equivalence.


(3) If  $S_1, \dots, S_N$  are all circles then all thickenings  $(M, f)$  are equivalent in the sense below.

$(M, f)$  and  $(M', f')$  are said to be *equivalent* if  $M \cong M'$  are homeomorphic and the homeomorphism  $H$  can be chosen to preserve all the data i. e..

(a)  $H$  is orientation-preserving

(b) the diagram

homotopy commutes.



REMARK. In the simply-connected case ( $p_1 \geq 2$ ) these definitions coincide with those of Haefliger [3].

Let  $P$  be the manifold defined in the introduction. That is a ball plus 1-handles. Let  $\dim P = q$  where  $q \geq p_N + 3$  and  $p_1 \geq 2$ . Any inessential orientation-preserving embedding  $h: \bigcup_1^N \partial \Delta^{p_i} \times \Delta^{q-p_i} \rightarrow \partial P$  determines an oriented manifold of the homotopy type of a bouquet of spheres. The manifold is  $P$  plus handles  $\Delta^{p_i} \times \Delta^{q-p_i}$  attached by means of the given embedding. Since the embedding  $h$  was assumed inessential we obtain a well-defined homotopy class of maps  $f: \cup S_i \rightarrow P + \text{handles}$ . It is easily seen that this  $f$  is in fact a thickening.

REMARK. The restriction that  $h$  be inessential is only a restriction if  $p_1 = 2$ , since  $\pi_i(P) = 0$  for  $i > 1$ .

Suppose  $h, k$  are concordant embeddings. It is an immediate consequence of the concordance extension theorem [4] that the two thickenings defined by  $h$  and  $k$  are equivalent.

Thus there is defined a function  $\Phi$  from the set of concordance classes of embeddings of solid tori in  $\partial P$  to the set of equivalence classes of thickenings of  $\vee S_i$ .

Here is the main result of this note.

THEOREM. If, for the bouquet  $S^1 \vee \dots \vee S^1 \vee S_1 \vee \dots \vee S_N$ ,  $\dim P = q \geq \text{Max}(6, p_N + 3)$  then

- (1)  $\Phi$  is surjective if  $2p_N - q + 1 < p_1$
- (2)  $\Phi$  is injective if  $2p_N - q + 2 < p_1$ .

The proof is deferred to section 3.

### 3. The factorization lemma.

In this section it will be shown that any thickening  $f: \cup S_i \rightarrow W$  factors up to homotopy through a homotopy equivalence  $g: \vee S_i \rightarrow N$  where  $N$  is a subpolyhedron of  $\text{Int } W$  with special properties. But first some notation. Write  $X$  for  $\vee S_i$  and filter  $X$  by  $* = X_0 \subset, \dots, \subset X_N = X$  where  $X_i = S_i \vee, \dots, \vee S_i$  and  $*$  is the base point of  $X$ . Let  $f_k = f|S_k$ . Finally, denote by  $\Sigma f$  the singular set of  $f$  i.e. the closure of the set  $\{x \in X | f^{-1}fx \neq x\}$ .

LEMMA 1. (factorization lemma) Let  $f: X \rightarrow \text{Int } W$  be a thickening (or more accurately let  $f \circ \pi$  be one). Suppose that  $f$  is nondegenerate and that for  $k = 1, \dots, N$ ,  $\dim \Sigma f \cap S_k \leq p_k + p_N - q$ . Then there exist polyhedra  $Y_0 \subset Y_1 \subset, \dots, \subset Y_N$  in  $\text{Int } W$  such that

- (1)  $f|X_k \rightarrow Y_k \cup fS_k$  is a homotopy equivalence
- (2)  $Y_{k+1}$  collapses to  $Y_k \cup fS_k$
- (3)  $\dim Y_k \cap fS_{k+i} \leq p_k + p_N - q$  ( $i > 0$ )
- (4)  $\Sigma f \cap S_k \subset f_k^{-1}Y_k$  and the latter collapses to  $*$ .

REMARK 1. If the first few  $S_i$  of  $X$  have small enough dimension then  $f$  embeds them (by the general position hypotheses on  $f$ ) and the first few  $Y_i$  are defined by  $Y_i = fX_{i-1}$  satisfying conditions (1), ..., (4).

REMARK 2. Lemma 1 yields a minimal handlebody decomposition for  $W$  as follows. Define inductively handlebodies  $H_k$  in  $\text{Int } W$  ( $k = 1, \dots, N$ ) by first triangulating  $f: X \rightarrow W$  so that  $Y_0, \dots, Y_N$  appear as subcomplexes (to be denoted by the same symbols). Take the barycentric second-derived subdivisions of  $X, W$ .  $f$  remains simplicial being non-degenerate. Define  $H_k = N^2(Y_k \cup fS_k; W)$ , the simplicial neighbourhood of  $Y_k \cup fS_k$  in the second-derived subdivision of  $W$ . Then  $H_{k+1}$  is  $H_k$  plus a handle. For  $(Y_{k+1} \cup fS_{k+1}) - \text{Int } N^2(Y_{k+1}; W) = fS_{k+1} - \text{Int } N^2(Y_{k+1}; W)$  and the latter is a  $p_{k+1}$ -disk in  $\text{Int } W$  that meets  $N^2(Y_{k+1}; W)$  in its boundary. This follows from condition (4).

And  $H_{k+1} - \text{Int } N^2(Y_{k+1}; W)$  is a ball meeting  $N^2(Y_{k+1}; W)$  in  $N^2(fS_{k+1}; W) \cap \partial N^2(Y_{k+1}; W)$  which is a solid torus. By regular neighbo-

uhoods theory,  $H_{k+1}$  is homeomorphic to

$$N^2(Y_{k+1}; W) \cup_{\psi} \Delta^{2k+1} \times \Delta^{q-2k+1}$$

where  $\psi: \partial\Delta \times \Delta \rightarrow \partial N^2(Y_{k+1}; W)$  is an embedding. But by condition (2)  $Y_{k+1}$  collapses to  $Y_k \cup fS_k$  and so  $N^2(Y_{k+1}; W)$  and  $H_k = N^2(Y_k \cup fS_k; W)$  are homeomorphic. Therefore  $H_{k+1}$  is  $H_k$  plus a handle and we obtain a handlebody decomposition of  $H_N$  (and hence of  $W$ ) suffixed by the cell structure of  $X$ . Furthermore, (1) implies that the thickening  $f: X \rightarrow W$  is filtered by a series of thickenings  $f|X_k \rightarrow N^2(Y_k \cup fS_k; W)$ .

As observed in the introduction, it is possible (under certain dimensional restrictions) to find a handlebody decomposition of  $W$  in which the handles are attached independently of one another after a certain stage. To show this we need a modification of the factorization lemma.

LEMMA 2. Let  $f$  satisfy the hypothesis of lemma 1. Suppose, in addition, that  $p_{k+1} \geq 2p_N - q + 2$  for some  $k (1 \leq k < N)$ , and that  $Y_0, \dots, Y_k$  have been found satisfying conditions (1) through (4) of Lemma 1. Then there exists a polyhedron  $Y$  in  $\text{Int } W$  such that

- (a)  $Y$  collapses to  $Y_k \cup fS_k$
- (b)  $f: X \rightarrow Y \cup fX$  is a homotopy equivalence
- (c)  $\Sigma f \subset f^{-1}Y$
- (d)  $f_{k+i}^{-1}Y$  collapses to  $*$  for all  $i > 0$ .

REMARK. As before we have a handlebody decomposition of  $W$ . Triangulate  $f: X \rightarrow W$  so that  $Y_0, \dots, Y_k, Y$  are subcomplexes of  $W$ . Define

$$H_j = N^2(Y_j \cup fS_j; W) \quad (0 \leq j \leq k)$$

and

$$H = N^2(Y \cup fX; W).$$

Then the handles  $N^2(fS_j; W) - \text{Int } N^2(Y; W)$  are attached independently to  $N^2(Y; W)$  i.e.  $N^2(fS_j; W) \cap \partial N^2(Y; W)$  are disjoint solid tori ( $k+1 \leq j \leq N$ ). For, by (c) of Lemma 2,  $f$  embeds

$$\begin{aligned} f^{-1} \left\{ \bigcup_{j=k+1}^N fS_j - \text{Int } N^2(Y; W) \right\} &\text{ and } S_j - f^{-1} \text{Int } N^2(Y; W) \\ &= S_j - \text{Int } N^2(f_j^{-1}Y; W) \\ &= S_j \text{ minus the interior of a ball, by (d)} \\ &= a \text{ ball, } (j = k+1, \dots, N); \end{aligned}$$

and the ball  $fS_j - \text{Int } N^2(Y; W)$  meets  $N^2(Y; W)$  in its boundary only, by (c). Thus  $N^2(Y; W) \cup fX$  is  $N^2(Y; W)$  plus balls  $fS_j - \text{Int } N^2(Y; W)$  attached disjointly to  $\partial N^2(Y; W)$ . This completes the proof that  $H$  is  $H_k$  plus disjointly attached handles.

To prove Lemma 1 and 2 we will need some general position and engulfing lemmas.

**DEFINITION.** If  $Y_0, Y, Z$  are polyhedra in the manifold  $M$  and  $Y_0 \subset Y$ , then  $Y - Y_0$  is said to be in *general position with respect to  $Z$*  if  $\dim(Y - Y_0) \cap Z \leq \dim Y - Y_0 + \dim Z - \dim M$ .

**DEFINITION.** If  $Y$  is a polyhedron and  $M$  a manifold, a map  $f: Y \rightarrow M$  is in *general position* if

- (1)  $f$  is non-degenerate
- (2)  $\dim \Sigma f < 2 \dim Y - \dim M$ .

**COROLLARY TO THEOREM 15 [7].** If  $Y_0, Y, A_1, \dots, A_n$  are polyhedra in a manifold  $M$  with  $Y_0 \subset Y$  and  $Y - Y_0 \subset \text{int } M$ , then there exists a homeomorphism  $h: M \rightarrow M$  such that

- (1)  $h|_{Y_0 \cup \partial M} = \text{Identity}$
- (2)  $h(Y - Y_0)$  is in general position with respect to  $A_1, \dots, A_n$ .

**PROOF.** By induction on  $\dim A_1 \cup \dots \cup A_n$ .

**COROLLARY TO THEOREM 18 [7].** Let  $f: Y \rightarrow \text{Int } M$  be a map and  $Y_0$  a subpolyhedron of  $Y$ . Suppose  $f|_{Y_0}$  is in general position. Then  $f$  is homotopic to  $g$ , a map in general position, by an arbitrarily small homotopy that keeps  $Y_0$  fixed.

**LEMMA 3.** If  $f: X \rightarrow M$ ,  $X$  a sphere-bouquet,  $M$  a manifold, then  $f$  is homotopic to  $g: X \rightarrow \text{Int } M$  where  $g$  is in general position and  $\dim \Sigma g \cap S_k \leq \dim S_k + \dim X - \dim M$  ( $k = 1, \dots, N$ ).

**PROOF.** First homotop  $fX$  into  $\text{Int } M$  and then use induction on  $N$ , the number of spheres in the bouquet. If  $N = 1$  apply the second corollary above. If not, the inductive step is proved by homotoping  $f|_{S_N}$  into general position keeping  $f^*$  fixed and then applying the first corollary to minimize the dimension of  $fS_N \cap fX_{N-1}$  by putting  $f(S_N - *)$  into general position with respect to  $fX_{N-1}$  keeping  $f^*$  fixed.

To state the engulfing lemmas we need

DEFINITION. A subpolyhedron  $C$  of a manifold  $M$  is called a  $k$ -spine of  $M$  if the pair  $M, C$  is  $k$ -connected.

DEFINITION. A polyhedron is called  $t$ -collapsible if it can be collapsed to a polyhedron of dimension not greater than  $t$ . The following lemma is a special case of Theorem 21 [7].

LEMMA 4 (Zeeman). Let  $C$  be an  $m$ -3-collapsible  $k$ -spine of the manifold  $M$  ( $\dim M$  being  $m$ ),  $Y$  a polyhedron in  $M$  and

$$\dim Y \cap \partial M < \dim Y \leq k \leq m - 3.$$

Then  $Y$  may be engulfed from  $C$  relative to  $\partial M$  i. e. there exists  $C^+$  in  $M$  such that  $C \cup Y \subset C^+$ ,  $(C \cup Y) \cap \partial M = C^+ \cap \partial M$ ,  $C^+$  collapses to  $C$ , and  $\dim C^+ - C \leq \dim Y + 1$ .

ADDENDUM TO LEMMA 4. Suppose that  $A_1, \dots, A_n$  are polyhedra in  $M$ . By the corollary to Theorem 15 we may insist that  $C^+ - (C \cap Y)$  be in general position with respect to  $A_1, \dots, A_n$ .

LEMMA 5. Let  $C$  be an  $m$ -3-collapsible  $k$ -spine of  $M$  and  $D$  a  $q$ -3-collapsible  $k + 1$ -spine of  $Q$  and let  $f: M, C \rightarrow Q, D$  be non-degenerate and proper (i.e.  $f^{-1} \partial Q = \partial f^{-1} Q$ ). Suppose that  $\dim (f^{-1} D - C) = x \leq k \leq m - 3 \leq q - 6$  and that  $\partial M \cap (f^{-1} D - C)$  is empty.

Then there exist polyhedra  $C^+ \subset M, D^+ \subset Q$  such that

- (A)  $C^+ = f^{-1} D^+$  (i.e.  $\dim f^{-1} D^+ - C^+ < 0$ )
- (B)  $C^+ \cap \partial M = C \cap \partial M; D^+ \cap \partial Q = D \cap \partial Q$
- (C)  $C^+$  collapses to  $C; D^+$  collapses to  $D$
- (D)  $\dim C^+ - C \leq x + 1; \dim D^+ - D \leq x + 2$ .

If, further,  $A_1, \dots, A_n \subset Q$  are polyhedra in general position with respect to  $fM$ , then  $C^+, D^+$  may be chosen to satisfy (A), ..., (D) and the extra condition

- (E)  $D^+ - D$  is in general position with respect to  $A_1, \dots, A_n$ .

PROOF. The proof resembles that of Lemma 63 [7]. We will define inductively polyhedra  $C_i \subset M, D_i \subset Q$  such that

- (a)  $fC_i \subset D_i$  and  $\dim f^{-1} D_i - C_i \leq x - i$ .
- (b)  $C_i$  collapses to  $C; D_i$  collapses to  $D$ .



- (c)  $C_i \cap \partial M = C \cap \partial M$ ;  $D_i \cap \partial Q = D \cap \partial Q$ .  
 (d)  $\dim C_i - C_{i-1} \leq x + 2 - i$ ;  $\dim D_i - D_{i-1} \leq x + 3 - i$ .  
 (e)  $D_i - D_{i-1}$  is in general position with respect to  $A_1, \dots, A_n$ .

The induction starts with  $C_i = C$ ,  $D_i = D$  ( $i \leq 0$ ) and finishes with  $i = x + 1$  because then  $\dim f^{-1} D_i - C_i < 0$ . Condition (E) will be satisfied because  $D^+ - D = \bigcup_{i \geq 0} (D_{i+1} - D_i)$  and each  $D_{i+1} - D_i$  is in general position with respect to  $A_1, \dots, A_n$ .

*The inductive step ( $i \geq 0$ ).*

Assume that  $C_j, D_j$  have been chosen satisfying (a), ..., (e) for  $j \leq i$ .

By (a)  $\dim f^{-1} D_i - C_i \leq x - i$ .

By (b)  $C_i$  is an  $m - 3$ -collapsible  $k$ -spine of  $M$  (since  $C$  is).

So by Lemma 4 there exists  $C_{i+1} \subset M$  such that  $C_{i+1}$  collapses to  $C_i$ ,  $f^{-1} D_i \subset C_{i+1}$ ,  $\partial M \cap C_{i+1} = \partial M \cap f^{-1} D_i$ ,  $\dim C_{i+1} - C_i \leq x + 1 - i$  and  $C_{i+1} - f^{-1} D_i$  is in general position with respect to  $f^{-1} A_1, \dots, f^{-1} A_n$ . This implies that  $\dim f C_{i+1} - D_i \leq \dim f(C_{i+1} - C_i) \leq x + 1 - i$ ; also that  $\partial M \cap C_{i+1} = \partial M \cap C_i \cup \partial M \cap f^{-1} D_i$ . But

$$\begin{aligned}
 \partial M \cap f^{-1} D_i &= \\
 &= f^{-1}(\partial Q \cap D_i) && (f \text{ is proper}) \\
 &= f^{-1}(\partial Q \cap D) && (\text{by (c)}) \\
 &= \partial M \cap f^{-1} D \\
 &= \partial M \cap C && (\text{by initial hypothesis}).
 \end{aligned}$$

By (b)  $D_i$  is a  $q - 3$ -collapsible  $k + 1$ -spine of  $Q$ . So by Lemma 4, there exists  $D_{i+1} \subset Q$  such that  $D_{i+1}$  collapses to  $D_i$ ,  $f C_{i+1} \subset D_{i+1}$ ,  $\dim D_{i+1} - D_i \leq x + 2 - i$ ,  $\partial Q \cap D_{i+1} = \partial Q \cap (D_i \cup f C_{i+1})$  and  $D_{i+1} - (D_i \cup f C_{i+1})$  is in general position with respect to  $fM, A_1, \dots, A_n$ . This implies that  $\dim f^{-1} D_{i+1} - C_{i+1} \leq \dim f f^{-1} D_{i+1} - f C_{i+1} = \dim fM \cap (D_{i+1} - f C_{i+1}) = \dim fM \cap (D_{i+1} - (f C_{i+1} \cup D_i)) \leq x + 2 - i - 3$ . Also we have that  $\partial Q \cap D_{i+1} = \partial Q \cap D_i \cup \partial Q \cap f C_{i+1}$ . But  $\partial Q \cap f C_{i+1} = f(\partial M \cap C_{i+1}) = f(\partial M \cap C) \subset \subset \partial Q \cap D$ . So  $\partial Q \cap D_{i+1} = \partial Q \cap D$ . We have thus defined  $C_{i+1}, D_{i+1}$  satisfying (a), ..., (d). The  $\Delta$  also satisfies (e); for,  $D_{i+1} - (f C_{i+1} \cup D_i)$  is in general position with respect to  $A_1, \dots, A_n$ ; and  $f C_{i+1} - D_i = f(C_{i+1} - f^{-1} D_i)$  is in general position with respect to  $A_1, \dots, A_n$  since  $A_1, \dots, A_n$  are (by hypothesis) in general position with respect to  $fM$  and  $C_{i+1} - f^{-1} D_i$  was chosen

to be in general position with respect to  $f^{-1}A_1, \dots, f^{-1}A_n$ . This completes the proof of the inductive step and hence of lemma 5.

PROOF OF LEMMA 1. Let us write  $Z_k = Y_k \cup fS_k$ . Construct  $Y_k$  (and hence  $Z_k$ ) inductively starting with  $Y_0 = Z_0 = fX_0 = f*$ . Suppose that we have found  $Y_0, \dots, Y_k$  satisfying conditions (1), ..., (4). By lemma 4 and the fact that  $\dim \Sigma f \cap S_{k+1} \leq p_{k+1} - 3$  there exists  $C_{k+1}$  in  $S_{k+1}$  such that  $\Sigma f \cap S_{k+1} \subset C_{k+1}$ ,  $C_{k+1}$  collapses to  $*$ , and  $\dim C_{k+1} \leq 1 + p_N + p_{k+1} - q$ . Now  $Z_k$  is a  $p_{k+1} - 1$ -spine of  $\text{Int } W$  and  $1 + p_N + p_{k+1} - q \leq p_{k+1} - 2$ . Therefore by lemma 4 there exists  $D_{k+1}$  in  $\text{Int } W$  such that  $fC_{k+1} \subset D_{k+1}$ ,  $D_{k+1}$  collapses to  $Z_k$ ,  $\dim D_{k+1} - Z_k \leq 2 + p_N + p_{k+1} - q$  and  $D_{k+1} - (Z_k \cup fC_{k+1})$  is in general position with respect to  $fS_{k+1}, \dots, fS_N$ . This and condition (3) imply that for  $i > 1$

$$\begin{aligned} & p_N + p_{k+1} - q \geq \\ & \geq \dim fS_{k+i} \cap [D_{k+1} - (Z_k \cup fC_{k+1}) \cup Z_k \cup fC_{k+1}] \\ & \geq \dim fS_{k+i} \cap D_{k+1}. \end{aligned}$$

Now  $f_{k+1} : S_{k+1}, C_{k+1} \rightarrow W$ ,  $D_{k+1}, C_{k+1}$  is a  $p_{k+1} - 2$ -spine of  $S_{k+1}$ ,  $D_{k+1}$  is a  $p_{k+1} - 1$ -spine of  $\text{Int } W$  and  $\dim f_{k+1}^{-1} D_{k+1} - C_{k+1} \leq p_N + p_{k+1} - q$ . So, by lemma 5, there exists  $Y_{k+1}$  in  $\text{Int } W$  such that  $Y_{k+1}$  collapses to  $D_{k+1}$ ,  $f_{k+1}^{-1} Y_{k+1}$  collapses to  $*$  and  $\dim fS_{k+i} \cap (Y_{k+1} - D_{k+1}) \leq p_N + p_{k+1} - q$  ( $i > 1$ ). It follows that  $\dim fS_{k+i} \cap Y_{k+1} \leq p_N + p_{k+1} - q$  ( $i > 1$ ). Thus  $Y_{k+1}$  is defined and satisfies (2) (3) and (4).

The proof of the induction step will be complete once it has been shown that  $f|X_{k+1} \rightarrow Z_{k+1}$  is a homotopy equivalence. First triangulate  $f : X \rightarrow W$  and pass to the barycentric second derived triangulations of  $X, W$ .  $f$  remains simplicial.

We showed that  $f_{k+1}^{-1} Y_{k+1}$  collapsed to  $*$ . Thus  $N^2(f_{k+1}^{-1} Y_{k+1}; S_{k+1}) = f_{k+1}^{-1} N^2(Y_{k+1}; W)$  is a ball.

Further,  $\Sigma f \cap S_{k+1} \subset f_{k+1}^{-1} Y_{k+1}$  and so  $f_{k+1}$  maps

$$S_{k+1} - \text{Int } N^2(f_{k+1}^{-1} Y_{k+1}; S_{k+1})$$

homeomorphically onto  $Z_{k+1} - \text{Int } N^2(Y_{k+1}; W)$ .

To prove that  $f|X_{k+1} \rightarrow Z_{k+1}$  is a homotopy equivalence, we show that

(\*)  $f|X_{k+1} \rightarrow N^2(Y_{k+1}; W) \cup fS_{k+1}$  is a homotopy equivalence.

(\*\*)  $N^2(Y_{k+1}; W) \cup fS_{k+1}$  collapses to  $Y_{k+1} \cup fS_{k+1}$ .

Composing (\*\*) with (\*), we obtain a homotopy equivalence:

$$X_{k+1} \xrightarrow{f|} N^2(Y_{k+1}; W) \cup fS_{k+1} \supset Y_{k+1} \cup fS_{k+1}.$$

PROOF OF (\*).  $f$  maps the pair  $X_{k+1}, X_k \cup N^2(f_{k+1}^{-1} Y_{k+1}; S_{k+1})$  into the pair  $fS_{k+1} \cup N^2(Y_{k+1}; W), N^2(Y_{k+1}; W)$ .

$f|X_k \rightarrow N^2(Y_{k+1}; W)$  is a homotopy equivalence because  $f|X_k \rightarrow Z_k$  is one and  $N^2(Y_{k+1}; W)$  collapses to  $Z_k$  via  $Y_{k+1}$ .

Let us write  $U(\ )$  for «universal cover of». All spaces to which  $U(\ )$  is applied will have isomorphic fundamental groups for [by Remark (1) following Lemma 1]  $p_{k+1}$  may be assumed to be greater than one. Therefore the map  $f|X_{k+1}$  induces homology excision isomorphisms between  $H_*(U(X_{k+1}), U(X_k \cup N^2(f_{k+1}^{-1} Y_{k+1}; S_{k+1})))$  and

$$H_*(U(fS_{k+1} \cup N^2(Y_{k+1}; W)), U(N^2(Y_{k+1}; W))).$$

So, by the 5-Lemma and Whitehead's theorem, the map  $fX_{k+1} \rightarrow N^2(Y_{k+1}; W) \cup fS_{k+1}$  induces isomorphisms of homotopy groups in all dimensions and is thus a homotopy equivalence.

PROOF OF (\*\*).  $N^2(Y_{k+1}; W) \cup fS_{k+1}$  collapses to  $Y_{k+1} \cup fS_{k+1}$  because we may factor the collapse from  $N^2(Y_{k+1}; W)$  to  $Y_{k+1}$  through  $Y_{k+1} \cup U N^2(Y_{k+1} \cap fS_{k+1}; fS_{k+1})$ . This proves (\*\*) and completes the proof of Lemma 1.

PROOF OF LEMMA 2. Suppose polyhedra  $Y_0, \dots, Y_k$  have been found satisfying conditions (1), ..., (4) of Lemma 1. Recall that for  $i > 0$   $\dim \Sigma f \cap S_{k+i} \leq p_N + p_{k+i} - q$  and  $\dim f_{k+i}^{-1} Z_k \leq p_N + p_k - q$ . So by Lemma 4 there exists  $C$  in  $S_{k+1} \cup \dots \cup S_N$  such that  $(\Sigma f \cup f^{-1} Z_k) \cap (S_{k+1} \cup \dots \cup S_N) \subset C$ ,  $C$  collapses to  $*$  and  $\dim C \cap S_{k+i} \leq p_N + p_{k+i} - q + 1$ . Now  $Z_k$  is a  $p_{k+1} - 1$ -spine of  $\text{Int } W$  and by hypothesis  $1 + 2p_N - q \leq p_{k+1} - 1$ . Therefore by Lemma 4 there exists  $D$  such that  $fC \subset D$ ,  $D$  collapses to  $Z_k$   $\dim D - Z_k \leq 2 + 2p_N - q$  and  $D - (fC \cup Z_k)$  is in general position with respect to  $fS_{k+1}, \dots, fS_N$ .

Then

$$\begin{aligned} \dim f_{k+i}^{-1} D - (C \cap S_{k+i}) &= \dim f S_{k+i} \cap (d - (fC \cup Z_k)) \\ &\leq p_{k+i} + 2 + 2p_N - q - q \\ &\leq p_{k+1} - 3. \end{aligned}$$

Now  $C \cap S_{k+i}$  is a (collapsible)  $p_{k+1} - 2$ -spine of  $S_{k+i}$  and  $D$  is a  $p_{k+1} -$

— 1-spine of  $\text{Int } W$  and so there exists  $Y$  in  $\text{Int } W$  such that  $Y$  collapses to  $Z_k$ ,  $\Sigma f \subset f^{-1} Y$ , and  $f_{k+i}^{-1} Y$  collapses to  $*$  ( $i > 0$ ). The proof of lemma 2 is completed by showing that (as in lemma 1)  $f: X \rightarrow Y \cup f X$  is a homotopy equivalence.

It remains to prove the theorem of § 2.

**PROOF OF THEOREM.** (1) Surjectivity of  $\Phi$ . If  $f: X \rightarrow W$  is a thickening, homotop  $f$  into general position in the sense of lemma 3 and use lemma 2 to obtain a manifold  $W_0$  in  $\text{Int } W$  such that  $f X \subset \text{Int } W$  and  $f: X \rightarrow W_0$  is a thickening representing an element in the image of  $\Phi$ . See Remark after lemma 2. The  $S$  cobordism theorem provides us with an equivalence between the thickenings  $f: X \rightarrow W_0$  and  $f: X \rightarrow W$  and so  $\Phi$  is surjective.

(2) Injectivity of  $\Phi$ . Consider the special case  $X = S^1 \vee, \dots, \vee S^1 \vee S^p$ ; the proof for more spheres is similar.

Let  $*$  be the barycenter of the simplex  $\Delta$ . Let  $h_0, h_1$  be two embeddings of the solid torus  $\partial \Delta \times \Delta$  in  $\partial P$  ( $\dim \partial \Delta = p - 1$  and  $\dim \partial \Delta \times \Delta = \dim \partial P$ ). Let the handlebody corresponding to  $h_i$  be  $H(h_i) = P \cup_{h_i} \Delta \times \Delta$  ( $i = 0, 1$ ). Let  $\delta_i: \Delta \times \Delta \rightarrow H(h_i)$  and  $p_i: P \rightarrow H(h_i)$  be the associated embeddings (thus  $p_i^{-1} \delta_i = h_i$  i. e.  $\forall x \in \partial \Delta \times \Delta, \delta_i x = p_i h_i x$ ). Suppose that  $h_0, h_1$  determine equivalent thickenings (the equivalence being a homeomorphism  $G: H(h_1) \rightarrow H(h_0)$ ). Then a relative version of the proof of surjectivity shows that there exist embeddings

$$\alpha: \Delta \times \Delta \times [0, 1] \rightarrow H(h_0) \times [0, 1]$$

$$\beta: P \times [0, 1] \rightarrow H(h_0) \times [0, 1] \text{ such that}$$

$$\begin{aligned} 1) \quad & \alpha(x, 0) = (\delta_0 x, 0) \\ & \alpha(x, 1) = (G \delta_1 x, 1) \end{aligned}$$

$$\beta(x, 0) = (p_0 x, 0)$$

and

$$\beta P \times \{1\} = G p_1 P \times \{1\}$$

$$2) \quad \alpha^{-1} \text{Im } \beta = \partial \Delta \times \Delta \times [0, 1].$$

Thus we have a concordance  $\alpha^{-1} \circ \beta | \partial \Delta \times \Delta \times [0, 1] \rightarrow \partial P \times [0, 1]$  between  $h_0$  and  $\lambda \circ h_1$  where  $\lambda: P \rightarrow P$  is a self equivalence (i. e. an orientation — preserving homeomorphism homotopic to the identity). We need to show that  $\lambda \circ h_1$  and  $h_1$  are concordant.

First we choose  $\lambda$  of a special type. Let  $I^1 = [-1, +1]$  and  $I^k = I^1 \times \dots \times I^1 \subset \mathbb{R}^k$ . Then if  $q \geq 3$  we take  $P^q = P^3 \times I^{q-3}$ .  $P^3 = B \cup H$  is the union of a 3-ball  $B$  and disjointly-attached 1-handles.

Let  $C$  be the union of the set of cores of these handles. Then the reader may verify the following.

**PROPOSITION.** Any self-equivalence  $\lambda: P^3 \times I^k \rightarrow P^3 \times I^k$  is concordant to one of the form  $\mu \times \text{Id}$ , where  $\mu|_{B \cup C} = \text{Id}$ . As for  $h_1$ , we may clearly assume that  $\text{Im } h_1$  lies in  $\text{Int } P^q \times \{-1\} \subset \partial(P^q \times I^1)$ . It will suffice then to prove the following.

**LEMMA.** If  $\lambda: P^q \rightarrow P^q$  is a self-equivalence and  $\Sigma^p \subset \text{Int } P$  a sphere ( $p \leq q - 3$ ) then  $\lambda$  is concordant to  $\lambda'$  where  $\lambda'$  fixes (pointwise) a neighbourhood of  $\Sigma$  in  $P$ .

**PROOF.** By the proposition above choose  $\lambda = \mu \times \text{Id}$ , with  $\mu|_{B \cup C} = \text{Id}$ . Thus  $\lambda|_{B \times I^{q-3} \cup C \times I^{q-3}} = \text{Id}$ . The result of [2] is easily generalized to show that  $\Sigma$  can be compressed (by an ambient isotopy) into  $B \times I^{q-3} \cup C \times I^{q-3}$  [the intersection of  $\Sigma$  with  $C \times I^{q-3}$  being a set of disjoint cylinders (= homeomorphs of  $S^{p-1} \times [0, 1]$ ). Thus  $\lambda$  fixes  $\Sigma$ . It remains to show that after an isotopy  $\lambda$  fixes not only  $\Sigma$  but some neighbourhood of  $\Sigma$  in  $P$ .

Let  $\tilde{P}$  be the universal cover of  $P$  with covering projection  $\pi: \tilde{P} \rightarrow P$ . Since  $\Sigma$  is inessential in  $P$  choose a connected component  $\tilde{\Sigma}$  of  $\pi^{-1}\Sigma$ ; thus  $\Sigma, \tilde{\Sigma}$  are homeomorphic via  $\pi$ . Furthermore, in a neighbourhood of  $\tilde{\Sigma}$ ,  $\pi$  is (1-1). Let  $\tilde{\lambda}: \tilde{P} \rightarrow \tilde{P}$  be the lift of  $\lambda$  that fixes  $\tilde{\Sigma}$  pointwise i. e.  $\pi \circ \tilde{\lambda} = \lambda \circ \pi$  and  $\tilde{\lambda}|_{\tilde{\Sigma}} = \text{Id}$ .

Since  $\lambda|_{B \times I^{q-3}} = \text{Id}$  there is a  $q$ -ball  $R$  in  $\tilde{P}$  with  $\tilde{\lambda}R = R$  and  $\tilde{\Sigma} \subset \text{Int } R$ .

It follows from Lemma 59 of [7] that  $\tilde{\lambda}$  is isotopic (fixing  $\tilde{\Sigma}$  to  $A: \tilde{P} \rightarrow \tilde{P}$  that fixes  $R$  pointwise. Projecting down by  $\pi$  we see that there is an ambient isotopy of  $P$  that takes  $\lambda$  to  $\lambda'$  where  $\lambda'$  is the inclusion in a neighbourhood of  $\Sigma$ . This completes the proof of the lemma and hence of the theorem.

REFERENCES

- [1] D. BARDEN; *Thesis*, Cambridge.
- [2] D. D. J. HACON: *Knotted spheres in Tori*, Q. J. M. Oxford (2) 20 (1969), 431-445.
- [3] A. HAEFLIGER: *Enlacements de sphères en codimension supérieure à deux*. Comm. Math. Helv. 41 (1966) 51-72.
- [4] J. F. P. HUDSON and W. B. R. LICKORISH: *Extending PL concordances*, to appear.
- [5] J. STALLINGS: *The embedding of homotopy types into manifolds*, to appear.
- [6] C. T. C. WALL: *Classification problems in differential topology, IV*, Topology 5 (1966) 73-94.
- [7] E. C. ZEEMAN: *Seminar on Combinatorial Topology I. H. E. S.* (1963), revised 1966).