

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

SAMUEL I. GOLDBERG

Invariant submanifolds of codimension 2 of almost contact manifolds

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3^e série, tome 25, n° 3 (1971), p. 377-388

http://www.numdam.org/item?id=ASNSP_1971_3_25_3_377_0

© Scuola Normale Superiore, Pisa, 1971, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

INVARIANT SUBMANIFOLDS OF CODIMENSION 2 OF ALMOST CONTACT MANIFOLDS

SAMUEL I. GOLDBERG ⁽¹⁾

1. Introduction.

In his dissertation, Smyth [8] classified the complex hypersurfaces M of the simply connected complex space forms \tilde{M} under the conditions that in the induced metric they are complete Einstein spaces. M is then a totally geodesic submanifold, or else the holomorphic sectional curvature of \tilde{M} is positive and M is a complex hypersphere. A local analogue for odd dimensional manifolds was subsequently obtained by Yano and Ishihara [9]. They proved that if M is an invariant submanifold of codimension 2 of a normal contact Riemannian manifold \tilde{M} of constant sectional curvature and if in the induced metric M is an Einstein space, then M is a totally geodesic submanifold of \tilde{M} . Observe that the exceptional part of Smyth's result does not occur, that is positive curvature yields the same result in all cases.

Consider either a $(2n + 1)$ -dimensional normal contact Riemannian manifold or a cosymplectic space and let M be an invariant submanifold immersed as an orientable hypersurface (M, j) of a hypersurface (P, i) along which the fundamental vector field of \tilde{M} is tangent. Then, if the induced f -structure on P (of rank $2n - 2$) is normal, or, if the unit normal field of $j(M)$, with respect to the induced Riemannian metric, is a Killing vector field, M is a totally geodesic submanifold of \tilde{M} . This is an odd dimensional analogue of a result on complex hypersurfaces of Kaehler manifolds obtained in [3].

As in [3], no assumption on the metric structure of \tilde{M} is made. Indeed, it is not assumed that the ambient space is a space form or that the submanifold is an Einstein space.

Pervenuto alla Redazione il 20 Luglio 1970.

⁽¹⁾ Research partially supported by the National Science Foundation.

2. Hypersurfaces of almost contact manifolds.

Let \tilde{M} be an almost contact metric manifold of dimension $2n + 1$, $n \geq 2$, with fundamental affine collineation $\tilde{\varphi}$, fundamental vector field \tilde{E} , compatible metric \tilde{g} and contact form $\tilde{\eta}$, where

$$\tilde{\eta} = \tilde{g}(\tilde{E}, \cdot).$$

Let \tilde{N} be the field of unit normals to $i(P)$ with respect to \tilde{g} . Consider a $2n$ -dimensional hypersurface P immersed in \tilde{M} with immersion $i: P \rightarrow \tilde{M}$ having the property

(T): For each $p \in P$, the vector $\tilde{E}_{i(p)}$ belongs to the tangent hyperplane of $i(P)$.

Then,

$$(2.1) \quad \tilde{\varphi} i_* X = i_* fX + \alpha(X) \tilde{N},$$

$$(2.2) \quad \tilde{\varphi} \tilde{E} = 0,$$

$$(2.3) \quad \tilde{\eta}(\tilde{N}) = 0,$$

where f and α are tensor fields on P of types (1,1) and (0,1), respectively, i_* is the induced tangent map and $X \in \mathcal{X}(P)$ — the module of C^∞ vector fields on P . Since i is a regular map, there is a vector field E' on P such that

$$(2.4) \quad \tilde{E} = i_* E'.$$

Hence, by (2.1) and (2.2), $fE' = 0$ and $\alpha(E') = 0$. Putting $\eta' = i^* \tilde{\eta}$, we have

$$(2.5) \quad \eta'(E') = 1.$$

Since $\tilde{\varphi} \tilde{N}$ is orthogonal to \tilde{N} with respect to \tilde{g} , it is tangent to the hypersurface, so there is a vector field A on P such that

$$(2.6) \quad \tilde{\varphi} \tilde{N} = -i_* A.$$

Applying $\tilde{\varphi}$ to both sides of (2.1) gives $f^2 X = -X + \eta'(X) E' + \alpha(X) A$ and $\alpha(fX) = 0$.

Applying $\tilde{\varphi}$ to both sides of (2.6) yields $fA = 0$ and $\alpha(A) = 1$. Summarizing, we have the following result established in [5].

PROPOSITION 1. *Let P be a $2n$ -dimensional hypersurface immersed in the almost contact manifold \tilde{M} with immersion i . Then, there exist tensor fields f, E', η', A and α on P satisfying the relations*

$$(2.7) \quad f^2 = -I + \eta' \otimes E' + \alpha \otimes A,$$

$$(2.8) \quad \eta' \circ f = 0, \quad \alpha \circ f = 0,$$

$$(2.9) \quad fE' = 0, \quad fA = 0,$$

$$(2.10) \quad \eta'(E') = 1, \quad \eta'(A) = 0,$$

$$(2.11) \quad \alpha(E') = 0, \quad \alpha(A) = 1,$$

where I is the identity transformation of P_p , that is the induced structure on P is a globally framed f -structure of rank $2n - 2$.

Let \tilde{V} be the Riemannian connection of (\tilde{M}, \tilde{g}) and let D be the induced connection on P , that is, the Riemannian connection of $G = i^*\tilde{g}$. Then, the equations of Gauss and Weingarten are

$$(2.12) \quad \tilde{V}_{i_*X} i_* Y = i_* D_X Y + h(X, Y) \tilde{N}$$

and

$$(2.13) \quad \tilde{V}_{i_*X} \tilde{N} = -i_* HX,$$

respectively, where h and H are the second fundamental tensors of the immersion of types (0,2) and (1,1), respectively and

$$h(X, Y) = G(HX, Y).$$

If the structure on \tilde{M} is *normal*, that is, if the almost complex structure \tilde{J} on $\tilde{M} \times R$ defined by

$$\tilde{J}\left(\tilde{X}, \varrho \frac{d}{dt}\right) = \left(\tilde{\varphi}\tilde{X} - \varrho\tilde{E}, \tilde{\eta}(\tilde{X}) \frac{d}{dt}\right),$$

where ϱ is a C^∞ real valued function and \tilde{X} is a C^∞ vector field on \tilde{M} , gives rise to a complex structure on $\tilde{M} \times R$, then the tensor field $[\tilde{\varphi}, \tilde{\varphi}] + d\tilde{\eta} \otimes \tilde{E}$ (of type (1,2)) vanishes, where $[\tilde{\varphi}, \tilde{\varphi}](\tilde{X}, \tilde{Y}) = [\tilde{\varphi}\tilde{X}, \tilde{\varphi}\tilde{Y}] - \tilde{\varphi}[\tilde{\varphi}\tilde{X}, \tilde{Y}] - \tilde{\varphi}[\tilde{X}, \tilde{\varphi}\tilde{Y}] + \tilde{\varphi}^2[\tilde{X}, \tilde{Y}]$, $\tilde{X}, \tilde{Y} \in \mathcal{X}(\tilde{M})$.

An almost contact metric structure is called *quasi-Sasakian* if it is normal and its fundamental form $\tilde{\Phi}$ is closed, where $\tilde{\Phi}(\tilde{X}, \tilde{Y}) = \tilde{g}(\tilde{\varphi}\tilde{X}, \tilde{Y})$. Thus, Sasakian ($\tilde{\Phi} = d\eta$) and cosymplectic ($d\tilde{\eta} = 0$) manifolds are quasi-Sasakian [1].

The hypersurface P carries an almost hermitian structure. To see this, we set

$$(2.14) \quad J = f + \eta' \otimes A - \alpha \otimes E'.$$

Then, from (2.7) (2.11), it is seen that J is an almost complex structure, that is $J^2 = -I$. From (2.1), (2.4) and (2.6), we see that

$$\eta' = G(E', \cdot), \quad \alpha = G(A, \cdot).$$

The fields E' and A are therefore orthonormal by (2.10) and (2.11). Observe that

$$JE' = A.$$

By (2.1), since $\tilde{\varphi}$ is skew symmetric with respect to \tilde{g} , f is skew symmetric with respect to G . We put $F(X, Y) = G(fX, Y)$, that is $F = i^* \tilde{\Phi}$. Then, from (2.14),

$$G(JX, Y) = F(X, Y) + \eta'(X)\alpha(Y) - \alpha(X)\eta'(Y),$$

from which J is skew symmetric with respect to G .

Putting $\Omega(X, Y) = G(JX, Y)$, we obtain

$$(2.15) \quad \Omega = F + 2\eta' \wedge \alpha.$$

Observe that if \tilde{M} is quasi-Sasakian, then the 2 form F is closed.

If the ambient space is cosymplectic, η' is also closed. The following result was obtained in [5].

PROPOSITION 2. *If the ambient space is cosymplectic,*

$$(D_X f) Y = \alpha(Y)HX - h(X, Y)A,$$

$$D_X E' = 0, \quad D_X A = fHX,$$

$$D_X \eta' = 0, \quad (D_X \alpha)(Y) = -h(X, fY),$$

$$h(X, E') = \eta'(HX) = 0, \quad h(X, A) = \alpha(HX).$$

When the vector bundle over P , with fibre the vector space spanned by E' and A at each point of P , is endowed with an affine connection γ ,

it admits an almost complex structure \widehat{J} . If \widehat{J} is integrable, the globally framed f -structure is *normal* [7]. By defining γ in such a way that E' and A are parallel fields, it has zero curvature. The f structure is then normal if $[f, f] + d\eta' \otimes E' + d\alpha \otimes A$ vanishes. The following result was also obtained in [5].

PROPOSITION 3. *If the ambient space is cosymplectic, then a necessary and sufficient condition for the induced globally framed f -structure on P to be normal is that*

$$fH - Hf = \alpha \otimes D_A A.$$

3. Hypersurfaces of almost complex manifolds.

Let M be an immersed orientable hypersurface of P . We denote by j the immersion and by N the field of unit normals to $j(M)$ with respect to G (with orientation determined by P). Let ∇ be the Riemannian connection of (M, g) , $g = j^* G$. Then,

$$(3.1) \quad D_{j_* x} j_* y = j_* \nabla_x y + k(x, y) N$$

and

$$(3.2) \quad D_{j_* x} N = -j_* Kx,$$

where k and K are the second fundamental tensors of the immersion j , of types (0,2) and (1,1), respectively, and $x, y \in \mathcal{X}(M)$. We set

$$(3.3) \quad \eta(x) = G(Jj_* x, N)$$

and

$$(3.4) \quad \Phi(x, y) = G(Jj_* x, j_* y).$$

Then, Φ is a 2-form on M . If E is the contravariant form of η with respect to g , then it is a vector field on M satisfying

$$(3.5) \quad JN = -j_* E.$$

An endomorphism φ of $\mathcal{X}(M)$ is defined by the relation

$$(3.6) \quad \Phi(x, y) = g(\varphi x, y).$$

Thus, Φ being a 2-form, φ is skew symmetric with respect to g . Moreover, by (3.3),

$$(3.7) \quad Jj_* x = j_* \varphi x + \eta(x) N.$$

It follows that

$$(3.8) \quad \varphi^2 = -I + \eta \otimes E,$$

where I is the identity transformation field of M_m , $m \in M$. In addition, (3.3) and (3.7) yield

$$\eta(\varphi x) = 0,$$

which is equivalent to

$$\varphi E = 0$$

by the skew symmetry of φ . Consequently, M is an almost contact manifold [2].

4. Invariant submanifolds of codimension 2 of a cosymplectic space.

In the sequel, M is an *invariant submanifold* of \tilde{M} , that is

$$\tilde{\varphi} \iota_* x = \iota_* \varphi x, \quad \iota = i \circ j,$$

namely, at each point of M , the tangent space is invariant under the action of $\tilde{\varphi}$. Then, by means of (2.1), (2.4), (2.6), (2.14) and (3.7),

$$\eta(x)N = \tilde{\eta}(\iota_* x)A$$

and

$$j^* \alpha = 0.$$

Putting $x = E$, we obtain $N = A$. For, by (3.5), since $JE' = A$,

$$E' = \pm j_* E.$$

Hence, $N = \tilde{\eta}(\iota_* E)A = \eta'(j_* E)A = A$, by choosing $\eta'(j_* E) = 1$, since N and A are each of length 1. Thus, *if M is an invariant submanifold of an almost contact manifold with immersion ι , the vector field A coincides with the normal field N and $j^* \alpha = 0$.*

PROPOSITION 4. *If \tilde{M} is a cosymplectic manifold, then M is also cosymplectic.*

PROOF. Since $\eta = i^* \tilde{\eta}$, $(V_x \eta)(y) + \eta(V_x y) = (\tilde{V}_{\iota_* x} \tilde{\eta})(\iota_* y) + \tilde{\eta}(\tilde{V}_{\iota_* x} \iota_* y) = \tilde{\eta}(\tilde{V}_{\iota_* x} \iota_* y)$. For, in a cosymplectic manifold the covariant derivative of

the contact form is zero. From (2.12) and (3.1), we obtain

$$\tilde{V}_{\iota_* x} \iota_* y = \iota_* V_x y - k(x, y) \tilde{\varphi} \tilde{N} + h'(x, y) \tilde{N},$$

where $h'(x, y) = h(j_* x, j_* y)$. Applying (2.2) and (2.3), we get $V_x \eta = 0$ (see also § 5).

Defining the (1,1) tensor field H' by $h'(x, y) = g(H' x, y)$, we get

$$Hj_* x = j_* H' x - \omega(x) N$$

for some 1-form ω on M .

PROPOSITION 5. *Let M be an invariant submanifold of the cosymplectic space \tilde{M} with the immersion ι . Then,*

$$K = -\varphi H' = H' \varphi.$$

PROOF. We differentiate the function $\alpha(j_* y)$ in the direction x , then apply Proposition 2, formulae (2.14) and (3.7), and observe that $j^* \alpha = 0$:

$$\begin{aligned} x(\alpha(j_* y)) &= (D_{j_* x} \alpha)(j_* y) + \alpha(D_{j_* x} j_* y) \\ &= -h(j_* x, fj_* y) + k(x, y) \\ &= -h(j_* x, Jj_* y - \eta'(j_* y) A) + k(x, y) \\ &= -h(j_* x, j_* \varphi y) + k(x, y) \\ &= -G(Hj_* x, j_* \varphi y) + k(x, y) \\ &= -G(j_* H' x, j_* \varphi y) + k(x, y) \\ &= -g(H' x, \varphi y) + g(Kx, y), \end{aligned}$$

so

$$g(Kx, y) = -g(\varphi H' x, y)$$

and

$$g(x, Ky) = g(x, H' \varphi y).$$

COROLLARY. *Under the conditions in the proposition,*

$$K^2 = H'^2$$

and

$$\text{trace } H' = \text{trace } K = 0,$$

so M is a minimal submanifold.

PROOF. By the proposition, $K^2 = -H' \varphi^2 H' = H'^2 - (\eta \circ H') \otimes H' E$. But, by Proposition 2, $h(X, E') = 0$, so since $G(HX, E') = G(X, HE) = G(X, Hj_* E)$, we get $G(j_* x, Hj_* E) = G(j_* x, j_* H' E) = g(x, H' E) = 0$. That M is a minimal submanifold is a consequence of the fact that the second fundamental tensors are symmetric and Φ is skew symmetric.

THEOREM 1. *Let M be an invariant submanifold of a cosymplectic manifold \tilde{M} . If M is immersed in \tilde{M} as an orientable hypersurface of a hypersurface with the property (T), and if the field of unit normals N on P is a Killing vector field, then M is a totally geodesic submanifold of \tilde{M} .*

PROOF. By Proposition 2, $h(X, fY) + h(Y, fX) = 0$ which is equivalent to the statement that H commutes with f . Applying this to the vector field $j_* x$, we get

$$H' \varphi = \varphi H', \quad \omega \circ \varphi = 0.$$

For,

$$\begin{aligned} Hfj_* x &= H\{Jj_* x - \eta'(j_* x)A\} \\ &= H\{j_* \varphi x + \eta(x)N\} - \eta(x)HA \\ &= Hj_* \varphi x \\ &= j_* H' \varphi x - \omega(\varphi x)N, \end{aligned}$$

and

$$\begin{aligned} fHj_* x &= f\{j_* H' x - \omega(x)N\} \\ &= fj_* H' x \\ &= Jj_* H' x - \eta'(j_* H' x)A \\ &= j_* \varphi H' x + \eta(H' x)N - \eta(H' x)A \\ &= j_* \varphi H' x. \end{aligned}$$

Applying Proposition 5, $K = 0$ and $H' = 0$, the latter being due to the Corollary to Proposition 5.

COROLLARY. *Under the conditions in the theorem, the hypersurface P is a Kaehler manifold.*

PROOF. Since H and f commute $D_A A = fHA = HfA = 0$. Applying Proposition 3, the induced globally framed structure on P is normal. Hence, J is integrable (see [6]). By (2.15), P is Kaehlerian if η' and α are closed. That η' is closed is immediate since $\tilde{\eta}$ is closed. That α is closed is a consequence of the fact that A is a parallel field. To see this, we express any C^∞ vector field on P as $j_* X + \mu N$ for some $x \in \mathcal{X}(M)$ and C^∞ function μ on P , and show that $(D_{j_* x} \alpha)(j_* y)$, $(D_{j_* x} \alpha)(A)$, $(D_{j_* x} \alpha)(E')$, $(D_N \alpha)(j_* y)$, $(D_N \alpha)(E')$, and $(D_N \alpha)(A)$ vanish. That this is the case follows from Proposition 2, the vanishing of H' and the fact that $j^* \alpha$ is zero.

If the induced almost complex structure tensor J is integrable there exists an affine connection D on P such that $DJ = 0$. If this is the Riemannian connection induced by \tilde{g} , then the geometrical condition on N may be replaced by the condition that J be integrable. For, then by [4], Proposition 20, K and φ commute.

5. Invariant submanifolds of codimension 2 of a Sasakian space.

Theorem 1 has an analogue for normal contact metric spaces, that is for *Sasakian manifolds*. To this end, we state the appropriate analogue of Proposition 2 (see [5]).

PROPOSITION 6. *Let \tilde{M} be a Sasakian manifold. Then, the relations*

$$(D_X f) Y = -G(X, Y) E' + \eta'(Y) X + \alpha(Y) HX - h(X, Y) A,$$

$$D_X E' = fX, \quad D_X A = fHX,$$

$$(D_X \eta')(Y) = F(X, Y), \quad (D_X \alpha)(Y) = -h(X, fY),$$

$$h(X, E') = \eta'(HX) = \alpha(X), \quad h(X, A) = \alpha(HX)$$

hold on P .

Observe that E' is a killing vector field.

REMARK. We have shown (Proposition 4) that an invariant submanifold of a cosymplectic manifold with the immersion ι is also a cosymplectic manifold. A more general statement can be made, namely, *an invariant submanifold of a quasi-Sasakian manifold with the immersion ι is a quasi-Sasakian manifold*. To see this, observe that $\Phi = i^* \tilde{\Phi}$, since $\Phi = j^* \Omega$, $i^* \tilde{\Phi} = \Omega - 2\eta' \wedge \alpha$ and j^* is a ring homomorphism. Moreover, the condition $[\tilde{\varphi}, \tilde{\varphi}] + d\tilde{\eta} \otimes \tilde{E} = 0$ implies $[\varphi, \varphi] + d\eta \otimes E = 0$. However, Theorem 1 and

Theorem 2 (below) do not extend to quasi-Sasakian spaces in general. The key statement required is that if A is a Killing vector field, then H' and φ commute. Observe also that

$$(\tilde{V}_{i_*x} \tilde{\varphi}) \tilde{N} = i_*(\varphi H' x + Kx).$$

This is an identity if the ambient space is either cosymplectic or Sasakian (see Proposition 8 for the latter) since $\tilde{V}_{\tilde{x}} \tilde{\varphi}$ vanishes in the former case, and although this is not so for normal contact manifolds $(\tilde{V}_{i_*x} \tilde{\varphi}) \tilde{N} = \tilde{\eta}(\tilde{N}) i_* X - \tilde{g}(i_* X, \tilde{N}) \tilde{E} = 0$ by virtue of (2.3).

For quasi-Sasakian manifolds of different rank, $K \neq -\varphi H'$ unless the immersion is further restricted (see [1], Proposition 5.1).

PROPOSITION 7. *If \tilde{M} is a Sasakian manifold, then M is also a Sasakian manifold.*

PROOF. The structure tensors of \tilde{M} are related by

$$(\tilde{V}_{\tilde{x}} \tilde{\varphi}) \tilde{Y} = \tilde{\eta}(\tilde{Y}) \tilde{X} - \tilde{g}(\tilde{X}, \tilde{Y}) \tilde{E}.$$

Hence, $(\tilde{V}_{i_*x} \tilde{\varphi})(i_*y) = i_*\{\eta(y)x - g(x, y)E\}$. Since M is invariant $\tilde{V}_{i_*x}(\tilde{\varphi}i_*y) = \tilde{V}_{i_*x}(i_*\varphi y) = i_*\{(V_x \varphi)y + \varphi V_x y\} - k(x, \varphi y)\tilde{\varphi}\tilde{N} + h'(x, \varphi y)\tilde{N}$. But $\tilde{V}_{i_*x}(\tilde{\varphi}i_*y) = (\tilde{V}_{i_*x} \tilde{\varphi})(i_*y) + \tilde{\varphi}\{i_*V_x y - k(x, y)\tilde{\varphi}\tilde{N} + h'(x, y)\tilde{N}\} = i_*\{\eta(y)x - g(x, y)E + \varphi V_x y\} + k(x, y)\tilde{N} + h'(x, y)\tilde{\varphi}\tilde{N}$. Thus,

$$(V_x \varphi)y = \eta(y)x - g(x, y)E$$

which says that M is a Sasakian manifold.

Observe that the above proof also yields the formulae

$$k(x, \varphi y) = -h'(x, y)$$

and

$$k(x, y) = h'(x, \varphi y).$$

Hence,

PROPOSITION 8. *Let M be an invariant submanifold of the Sasakian manifold \tilde{M} with the immersion i . Then,*

$$K = H'\varphi, \quad H' = \varphi K.$$

COROLLARY. *Under the conditions in the proposition,*

$$K^2 = H'^2$$

and

$$\text{trace } H' = \text{trace } K = 0,$$

so M is a minimal submanifold.

The proof of Theorem 2 below parallels that of Theorem 1, Propositions 2 and 5 being replaced by Propositions 6 and 8, respectively. The following fact is also required.

LEMMA. *If the ambient space is a normal contact manifold, then $H'E$ vanishes.*

PROOF. By Proposition 6, $h(j_*x, E') = 0$ since $j^*\alpha = 0$. The remainder of the proof may be found in the proof of the Corollary to Proposition 5.

THEOREM 2. *Let M be an invariant hypersurface of a Sasakian manifold \tilde{M} . If M is immersed in \tilde{M} as an orientable hypersurface of a hypersurface with the property (T), and if the field of unit normals N on P is a Killing vector field, then M is a totally geodesic submanifold of \tilde{M} .*

COROLLARY. *The hypersurface P is a non-Kaehlerian hermitian manifold. J is integrable by Theorem 10 of [5] and Theorem 1 of [6].*

That P is not Kaehlerian is a consequence of the fact that η' is not closed. For, by Proposition 6, if η' were closed, then F' would vanish and this is not possible.

BIBLIOGRAPHY

- [1] D. E. BLAIR, *The theory of quasi-Sasakian structures*, *J. of Differential Geometry*, 1 (1967), 331-345.
- [2] S. I. GOLDBERG, *Totally geodesic hypersurfaces of Kaehler manifolds*, *Pacific J. Math.*, 27 (1968), 275-281.
- [3] S. I. GOLDBERG, *On immersions of complex hypersurfaces of Kaehler manifolds*, in *Revue Roumaine de Math. Pures et Appl.*, XV, No. 9 (1970), 1407-1413.
- [4] S. I. GOLDBERG and K. YANO, *Affine hypersurfaces of complex spaces*, *J. London Math. Soc.*, 2 (1970), 241-250.
- [5] S. I. GOLDBERG and K. YANO, *Globally framed f -manifolds, to appear in Illinois, J. Math.*
- [6] S. I. GOLDBERG and K. YANO, *On normal globally framed f -manifolds*, *Tohoku Math. J.*, 22 (1970), 362-370.
- [7] S. ISHIHARA, *Normal structure f satisfying $f^3 + f = 0$* , *Kōdai Math. Sem. Rep.*, 18 (1966), 36-47.
- [8] B. SMYTH, *Differential geometry of complex hypersurfaces*, *Ann. of Math.*, 85 (1967), 247-266.
- [9] K. YANO and S. ISHIHARA, *On a problem of Nomizu-Smyth on a normal contact Riemannian manifold*, *J. of Differential Geometry*, 3 (1969), 45-58.

*University of Illinois
Urbana, Illinois*