

ANNALI DELLA SCUOLA NORMALE SUPERIORE DI PISA *Classe di Scienze*

ROBERT KAUFMAN

Approximation of smooth functions and covering properties of sets

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 2, n° 3 (1975), p. 479-481

[<http://www.numdam.org/item?id=ASNSP_1975_4_2_3_479_0>](http://www.numdam.org/item?id=ASNSP_1975_4_2_3_479_0)

© Scuola Normale Superiore, Pisa, 1975, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Approximation of Smooth Functions and Covering Properties of Sets.

ROBERT KAUFMAN (*)

1. — Let Q be a closed cube in Euclidean space E^{n+1} ($n \geq 1$); using various spaces of differentiable functions on Q , one obtains corresponding classes of massive or negligible sets $F \subseteq Q$. The space $C^1(Q)$ is well known; for a number $1 < \alpha < 2$ we define the space $C^\alpha(Q)$ to contain functions whose partial derivatives satisfy a uniform Lipschitz condition with exponent $\alpha - 1$. A closed set $F \subseteq Q$ is called N_1 if $f(F)$ has linear Lebesgue measure 0 for all f in $C^1(Q)$ except at most a set of first category. Let $C^{\alpha n}$ be the Banach space of mappings into E^n , whose co-ordinates are of class C^α .

THEOREM 1. There is a closed N_1 -set F in Q , and an open set U in $C^{\alpha n}(Q)$, such that $f(F)$ contains a ball in E^n , for every f in U .

Let us say that a subset S in C^{1n} has *uniform rank n* if all tangent (Jacobian) mappings $J(f, x)$ ($f \in S, x \in Q$) transform the unit ball in E^{n+1} onto the unit ball in E^n (or a larger set). If a ball $B(r, x_0)$ is contained in Q , then $f(B) \supseteq B(r, f(x_0))$; this can be seen by a variant of the Cauchy-Peano method in ordinary differential equations [1, pp. 1-7].

THEOREM 1'. Let Q_0 be a compact set interior to Q and S a bounded subset of the space $C^{\alpha n}$, of uniform rank n . Then there is an N_1 -set $F \subseteq Q$, such that $f(F) \supseteq f(Q_0)$ for all f in S .

2. — Let T be a bounded subset of the Banach space $C^\alpha[0, 1]$, defined similarly to $C^\alpha(Q)$. For small numbers $r > 0$, T is contained in $\exp [Ar^{-1/\alpha}]$ ball of radius r in the *uniform metric*—a theorem of Kolmogorov [3, p. 153]. It is essential that the domain of the functions be a linear set, but the same bound holds for bounded subsets of $C^{\alpha n}[0, 1]$.

(*) University of Illinois - Urbana, Ill.
Pervenuto alla Redazione il 23 Gennaio 1974.

3. — Let K be a compact subset of Q , and let W be a neighborhood of K , interior to Q . For some $\varepsilon > 0$ every point of K has distance $> 2\varepsilon$ from the boundary of W . Let T be the set of all level curves $\Gamma(t)$, $0 \leq t \leq \varepsilon$, of functions f in S , meeting K . We choose arc-length as the parameter of each curve Γ , that is, $\|\Gamma'(t)\| = 1$. This condition on Γ , together with the boundedness of S in $C^{\alpha n}(Q)$ and the uniform rank n of S , implies that the set T is bounded in $C^{\alpha, n+1}[0, \varepsilon]$. Therefore Kolmogorov's estimate is valid for small $r > 0$.

The curves $\Gamma(t)$ have length ε and have equicontinuous tangent vectors $\Gamma'(t)$, so their diameters exceed some $c_1 > 0$. Let rA be the set of vectors $(ru_1, \dots, ru_{n+1})$, where each u_i is an integer. Since a curve Γ has diameter $> c_1$, some co-ordinate, say x_1 , increases $> c_1 n^{-1}$ along Γ ; when r is small x_1 then assumes $> c_2 r^{-1}$ values ru along Γ . Taking a point $(ru_1^0, x_2, \dots, x_{n+1})$ on Γ , we observe that some element $(ru_1^0, ru_2, \dots, ru_{n+1})$ of rA has distance $< nr$ from Γ . Thus at least $c_2 r^{-1}$ elements λ of rA have distance $< nr$ from the curve Γ . For small r , all these elements λ belong to Q .

Now we form a random selection A^* from the set $rA \cap Q$. To define the distribution of this selection, we fix once and for all a number τ in the interval $0 < \tau < 1 - \alpha^{-1}$. Then we select or reject the elements of $rA \cap Q$ independently of each other, with the probability of each selection exactly r^τ . The probability that A^* contains none of the $c_2 r^{-1}$ elements λ , found above, is $< \exp - c_2 r^{\tau-1}$ (because $1 - y < \exp - y$ for $y > 0$).

By Kolmogorov's estimate, we can select at most $\exp Ar^{-1/\alpha}$ curves $\Gamma_1 \in T$, so that every curve Γ is within r of some Γ_1 in the uniform metric on $[0, \varepsilon]$. The probability that *every* curve Γ_1 has distance $< nr$ from *some* element of A^* , exceeds $1 - \exp c_2 r^{\tau-1} \exp Ar^{-1/\alpha} \rightarrow 1$, because $\tau < 1 - \alpha^{-1}$. But then the same is true for all curves Γ , and a distance $(n+1)r$.

Suppose now that $x \in K$ and $f \in S$. Then, considering the level curve Γ of f through x , we see that $\|f(\lambda) - f(x)\| < c_3 r$ for some element λ of A^* . If r is small enough, then the ball $B(\lambda, c_3 r)$ —of center λ and radius $c_3 r$ —is contained in W and $f(x) \in f(B)$.

4. — To construct N_1 -sets by the random method, we require another property of A^* . We choose in succession an integer $k > 1$ and a number π in $(0, 1)$ so that $k\tau + (\pi - 1)k(n+1) > n+1$. Consider the event $M: A^*$ contains some k distinct elements $\lambda_1, \dots, \lambda_k$ with all distances $\|\lambda_i - \lambda_j\| \leq 3r^\pi$. To estimate $P(M)$ we bound the number J of the k -tuples: $rA \cap Q$ has $\ll r^{-(n+1)}$ elements; and each ball of radius $3r^\pi$ contains $\ll r^{(\pi-1)(n+1)}$ elements of rA . Hence $J \ll r^{-a}$, where $a = -(n+1) + (\pi-1)(n+1)(k-1)$. But $P(M) \leq J r^{k\tau} \rightarrow 0$ because $k\tau + a > 0$.

For small r we can choose A^* to have the covering property found in 3, while avoiding the event M . We write W_1 for the union of balls $B(\lambda, 2c_3r)$, $\lambda \in A^*$, and V_1 for $\cup B(\lambda, c_3r)$. Then $f(V_1) \supseteq f(K)$ for all f in S , and $W_1 \subseteq W$ for small r .

Now we can repeat this process, using V_1^- for K and $W_1 \cap W$ in place of W . Then we find a small r_2 , and corresponding sets V_2 and W_2 so that $f(V_2) \supseteq f(V_1) \supseteq f(K)$, etc. Moreover, c_3 and π are uncharged in the successive applications of the basic construction. The set $F = \bigcap_1^\infty W_m^-$ then has the property $f(F) \supseteq f(K)$, and we prove finally that F is an N_1 -set.

5. - To each g in $C^1(Q)$, and each $\varepsilon > 0$, we construct g_1 in $C^1(Q)$ so that $\|g - g_1\| < \varepsilon$ in $C^1(Q)$ and $g_1(F)$ has measure $< \varepsilon$. This shows that the elements of C^1 , transforming F onto a null set, are a dense G_δ . We begin with a partition of the centers λ_q , that is, the elements of A^* , corresponding to a small value of the radius r . Let Y_1 be a maximal selection of centers λ_q , having distances at least r^π ; let Y_2 be a maximal selection from the remaining centers, etc. If λ belongs to Y_k , then $\|\lambda - \lambda_q\| < r^\pi$ for $k-1$ centers $\lambda_q \neq \lambda$. But then we have k centers with distances $< 2r^\pi$ a contradiction. Therefore $Y_1 \cup \dots \cup Y_{k-1}$ exhausts A^* .

Let $s^{k+1} = r^{1-\pi}$, and observe that every real number has distance $< rs^{-k}$ from some multiple urs^{-k} of rs^{-k} . Hence we can define h_1 in $C^1(Q)$ so that each number $g(\lambda) + h_1(\lambda)$, with λ in Y_1 , is a multiple of rs^{-k} . In view of the distance r^π between the members of Y_1 , we can take h_1 to have norm $s^{-k}r(1+r^{-\pi})$, as in [2]. Then we construct h_2 so that each number $(g+h_1+h_2)\lambda$, with λ in Y_2 , is a multiple of rs^{1-k} . The norm of h_2 is again $\ll s^{-k}r^{1-\pi}$, and moreover $|h_2| < rs^{1-k}$. By this process we construct $g_1 = g + h_1 + \dots + h_{k-1}$, and $\|g - g_1\|$ is small because $s^{-k}r^{1-\pi} \rightarrow 0$. Moreover, $|g + h_1 + \dots + h_j - g_1| \ll \ll rs^{j-k}(1 \leq j < k)$, so $|g_1(\lambda) - urs^{j-k-1}| \ll rs^{j-k}$ for each λ in Y_j . When r is small, the partial derivatives of g_1 are bounded by some $B = B(g)$; thus $B(\lambda, c_3r)$ is mapped inside a ball of radius $\ll r + rs^{j-k} \leq 2rs^{j-k}$, centered at urs^{j-k-1} . Since the set $g_1(Q)$ remains within some finite interval, the union $\cup B(\lambda, c_3r)$ is mapped onto a set of measure $\ll s$, and this completes the proof.

REFERENCES

- [1] E. A. CODDINGTON - N. LEVINSON, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
- [2] R. KAUFMAN, *Metric properties of some planar sets*, Colloq. Math., **23** (1971), pp. 117-120.
- [3] G. G. LORENTZ, *Approximation of Functions*, Holt, New York, 1966.