ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

JULIO E. BOUILLET

Dirichlet problem for parabolic equations with continuous coefficients

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 3, n° 3 (1976), p. 405-441

<http://www.numdam.org/item?id=ASNSP_1976_4_3_3_405_0>

© Scuola Normale Superiore, Pisa, 1976, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (http://www.sns.it/it/edizioni/riviste/annaliscienze/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Dirichlet Problem for Parabolic Equations with Continuous Coefficients. (*)

JULIO E. BOUILLET (**)

Summary. – *L^p-boundary value problems for strongly parabolic operators such that the coefficients of the highest order derivatives are only uniformly continuous have been studied by V. A. Solonnikov* [10]. *In the case of a cylindre $\Omega\times(0, T)$, Ω a smooth spatial domain, and initial data zero, Solonnikov assumes the Dirichlet data to belong to a trace space. More precisely, if $L \equiv \sum_{|\alpha|\leqslant 2b} a_\alpha(P, t)D_p^\alpha - D_t$ is the strongly parabolic operator, w^k, $k = 0, 1, \ldots, b-1$, the Dirichlet data, then the problem*

$$Lu = 0 \quad in \quad \Omega\times(0, T), \qquad D_N^k u = w^k \quad at \quad \partial\Omega\times(0, T), \qquad u = 0 \quad on \quad \Omega \quad for \quad t = 0,$$

D_N indicating normal derivative to $\partial\Omega$, admits a unique solution in the space of functions whose spatial derivatives up to order $2b$, and the time derivative, belong in $L^p(\Omega\times(0, T))$, $p>1$. Solonnikov observed that this implies that w^k must have spatial derivatives up to order $2b-1-k$, and a «fractional» time derivative of order $(2b-1-k)/2k$ in $L^p(\partial\Omega\times(0, T))$. Moreover, a spatial derivative of order $2b-1-k$ of w^k will have a «fractional» derivative in the spatial direction of order $1/p'$ and in the time direction of order $1/2bp'$. With this information let us denote, for right now, the space of w^k by $\overset{\circ}{L}{}^p_{(2b-1-k+1/p'),\,(2b-1-k+1/p')/2b}(\partial\Omega\times(0, T))$; in the present work we find a class of existence and uniqueness to the problem above with the assumption that $w^k \in \overset{\circ}{L}{}^p_{(b-1-k+\varepsilon),\,(b-1-k+\varepsilon)/2b}(\partial\Omega\times(0, T))$. Here $\varepsilon<1$ is an arbitrary but fixed positive number. This means that we have reduced the smoothness requirements on the data w^k by at least b derivatives in the space direction, and $b/2b = \frac{1}{2}$ in the time direction. In a subsequent paper we shall discuss the non-zero initial value case. Outline: the definitions and notation appear in I, §§ 1-5, *for the case of a half-space, and in* VIII, § 24 *for the bounded domain Ω. The problem in a half-space is treated in* I-VI, *using certain surface and volume potentials* (III-IV). *Using the half-space results, we obtain an elliptic a priori estimate* (VII) *in the half-space. The problem in a general domain is studied in* VIII *for the parabolic case (a priori parabolic estimate: Theorem,* § 25; *existence and uniqueness theorems: Theorem* 1 *and Theorem* 2, § 27). *In* IX *an a priori estimate for the strongly elliptic case is derived.* *This work is part of a modified version of our Dissertation, under the direction of Professor Eugene B. Fabes. We wish to thank Professor Fabes for many invaluable talks and advise.*

(*) This work has been partially supported by grants NSF GP15832 and AFOSR 883-67 at the University of Minnesota, U.S.A.

(**) Departamento de Ciencias Exactas, Universidad Nacional de Salta, Salta (FGMB), Argentina.

Pervenuto alla Redazione il 4 Dicembre 1974 ed in forma definitiva il 5 Gennaio 1976.

I. – Definitions and notations for the problem in the half-space.

§ 1. – Points in R^n will be denoted by the letters x, z, w, while y, v, η, t, r, s will denote real numbers, the last three referring to time. $R_+^{n+1} = R^n x(0, \infty)$ will be the spatial domain (half-space), with points denoted by (x, y), (z, v). The differential operators will be defined for functions in the « cylinder » $R_+^{n+1} x(0, T)$, whose points are (x, y, t), t denoting boundary of this « cylinder » is $S_T = R^n x(0, T)$.

The following notations are standard: $f*g$ for the convolution of the functions f and g, $\mathcal{F}(f)(\cdot)$, occasionally also $\hat{f}(\cdot)$, the Fourier transform $\int f(\xi) \exp[i\langle \xi, \cdot \rangle] d\xi$, where $\langle \cdot, \cdot \rangle$ denotes scalar product of vectors. We will write, e.g. $\mathcal{F}_z(f)$ to specify the transformation in the variable z.

With $x = (x_1, \ldots, x_n)$ and $\alpha = (\alpha_1, \ldots, \alpha_n)$, α_i nonnegative integers, we set

$$x^\alpha = x_1^{\alpha_1} \ldots x_n^{\alpha_n}, \qquad |\alpha| = \alpha_1 + \ldots + \alpha_n,$$

$$D_x^\alpha = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}} = D_{x_1}^{\alpha_1} \ldots D_{x_n}^{\alpha_n}, \qquad \Delta = \Delta_x = \sum_{i=1}^{n} D_{x_i}^2 .$$

when there is no confusion we will use this notation to include $\alpha = (\alpha_1, \ldots, \alpha_n, \alpha_{n+1})$, and write $D_{x,y}^\alpha$ for $D_x^{(\alpha_1, \ldots, \alpha_n)} D_y^{\alpha_{n+1}}$. This will only apply to space variables.

We will denote by $X_D(\cdot)$ the characteristic function of the set D.

§ 2. – DEFINITION. A parabolic singular integral operator is an operator of the form (cf. [3], [4])

$$Kf(x, t = \alpha(x, t) f(x, t) + \lim_{\varepsilon \to 0^+} \int_0^{t-\varepsilon} \int_{R^n} k(x, t; x - z, t - s) f(z, s)\, dz\, ds ,$$

the limit in $L^p(R^n x(0, T))$, where $a(x, t)$ is a bounded measurable function on $R^n x(0, T)$, and the variable kernel $k(x, t; z, s)$ is defined for $t \in (0, T)$, $s > 0$ as

$$k(x, t; z, s) = \frac{\Omega(x, t; z/s^{1/2b})}{s^{1+n/2b}},$$

where

(1) $\Omega(x, t; \cdot) \in \mathcal{S}(R^n) =$ space of rapidly decreasing functions.

(2) $\int_{R_n} \Omega(x, t; w)\, dw = 0,$

(3) $\sup_{(y,s) \in S_T} \left(\int_{R^n} |w^\alpha D_w^\beta \Omega(y, s; w)|^2\, dw \right)^{\frac{1}{2}} < \infty$ (see, e.g. [6]).

If in the above definition $a(x, t)$ is a constant and $k(x, t; z, s) = k(z, s)$ is independent of (x, t), the operator Kf will be said to be of convolution type.

We define the symbol of K to be the function

$$\sigma(K)(x, t; z, s) = a(x, t) + \int_0^\infty \frac{\mathcal{F}_w(\Omega)(x, t; zr^{1/2b})}{r} \exp[itr]\, dr .$$

K is known to be a continuous mapping of $L^p(R^n x(0, T))$ for $1 < p < \infty$. Its properties (cf. [2], [3], [4]) will be assumed here.

Following [3], [5] we will denote by $\mathfrak{J}_p(S_T)$, $1 < p < \infty$, the class of operators $J: L^p(S_T) \to L^p(S_T)$ satisfying for any $a \geqslant 0$

(i) $JX_{(a,\infty)} = X_{(a,\infty)} JX_{(a,\infty)}$,

(ii) $\|X_{(a,a+\varepsilon)} J(X_{(a,a+\varepsilon)} f)\|_{L^p(S_T)} \leqslant \omega(\varepsilon) \cdot \|X_{(a,a+\varepsilon)} f\|_{L^p(S_T)}$,

where $\omega(\varepsilon) \to 0$ as $\varepsilon \to 0$, uniformly in $a \geqslant 0$. We set $\mathfrak{J}(S_T) = \bigcap_p \mathfrak{J}_p(S_T)$. $\mathfrak{J}_p(R_+^{n+1} x(0, T))$, $\mathfrak{J}(R_+^{n+1} x(0, T))$ are defined the same way.

§ 3. – We introduce the fundamental solution of the operator $(-1)^b \Delta_x^b + D_t$ on S_T, defined by

$$F(x, t) = \mathcal{F}_z(\exp[-|z|^{2b} t])(x) \quad \text{for } t > 0,$$
$$= 0 \quad \text{for } t \leqslant 0 .$$

DEFINITION. For β real > 0,

$$(\Lambda^{-\beta} f)(x, t) = (\Lambda^{-\beta} * f)(x, t) = \frac{1}{\Gamma(\beta/2b)} t^{\beta/2b - 1} F(x, t) * f(x, t) =$$
$$= \frac{1}{t^{1+(n-\beta)/2b}} \Omega_\beta(x/t^{1/2b}) * f(x, t), \qquad \Omega_\beta(\cdot) \in \mathcal{S} .$$

For $0 < \beta \leqslant 2b$, $\Lambda^{-\beta}(x, t)$ is a tempered distribution on R^{n+1}, whose Fourier transform is $(|x|^{2b} - it)^{-\beta/2b}$. Also if $f \in L^p(S_T)$, $D_x^\alpha \Lambda^{-\beta} f \in L^p(S_T)$ for $|\alpha| \leqslant [\beta]$ = largest integer $\leqslant \beta$, and $D_t \Lambda^{-2b+\beta} f \in L^p(S_T)$. We set $\Lambda^0 f = f$ and proceed to define Λ^β.

DEFINITION. For $0 < \beta \leqslant 2b$,

$$\Lambda^\beta f = \left((-1)^b \Delta_x^b + D_t\right) \Lambda^{-2b+\beta} f$$

Λ^β is well defined on rapidly decreasing functions f, which vanish for $t \leqslant 0$. As tempered distributions, $\mathcal{F}(\Lambda^\beta f) = (|x|^{2b} - it)^{\beta/2b} \cdot \hat{f}$. For β integral, $(-1)^b \Delta^b \Lambda^{-2b+\beta} f$ can also be written [3] as $\sum_{|\alpha|=\beta} K^\alpha D_x^\alpha f$, with K^α a parabolic singular integral operator with symbol

$$\sigma(K_\alpha) = i^{2b-\beta} \frac{P^\alpha(x)}{(|x|^{2b} - it)^{1-\beta/2b}},$$

$P_\alpha(x)$ defined by $|x|^{2b} = \sum_{|\alpha|=\beta} P_\alpha(x) x^\alpha$: considering $(-1)^b \Delta^b \Lambda^{-2b+\beta}$ as a tempered distribution on R^{n+1} this decomposition is clear if we recall that

$$\mathcal{F}(\Lambda^{-2b+\beta}) = \frac{1}{(|x|^{2b} - it)^{1-\beta/2b}}.$$

If $f = f(z, \eta, s; x, y, v, t)$, z, η, s, y, and v being parameters, we introduce the notation $\Lambda^\beta f(z, \eta, s; x, y, v, t)$ to mean $[\Lambda^\beta f(z, \eta, s; \cdot, y, v, \cdot)](x, t)$.

§ 4. – For $\delta \geqslant 0$, $L^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ is the space of functions whose derivatives $D^\alpha_{x,y} u$, for $|\alpha| \leqslant 2b$, and $D_t u$ in the sense of distributions are given by functions belonging to $L^p(R^n \times (\delta, \infty) \times (0, T))$. $L^p_{2b,1}$ is a Banach space with the norm

$$\|u\|_{L^p_{2b,1}(R^n \times (\delta,\infty) \times (0,T))} = \sum_{|\alpha| \leqslant 2b} \|D^\alpha_{x,y} u\|_{L^p(R^n \times (\delta,\infty) \times (0,T))} + \|D_t u\|_{L^p(R^n \times (\delta,\infty) \times (0,T))}.$$

For $\delta = 0$ we write $L^p_{2b,1}(R^{n+1}_+ \times (0, T))$.

$\mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ denotes the space of functions $u \in L^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ which are limits in $L^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ of functions $\in C_0^\infty(R^{n+1} \times (0, \infty))$.

§ 5. – A linear differential operator

$$L \equiv \sum_{|\alpha| \leqslant 2b} a_\alpha(x, y, t) D^\alpha_{x,y} - D_t$$

is said to be parabolic in the sense of Petrovski if

$$\operatorname{Re}\Big(\sum_{|\alpha|=2b} a_\alpha(x, y, t)(i\xi)^\alpha \Big) < -\pi |\xi|^{2b}, \quad \text{for } 0 \neq \xi \in R^{n+1},$$

$\pi > 0$ independent of (x, y, t) in $R^{n+1}_+ \times (0, T)$. Each $a_\alpha(x, y, t)$ is assumed to be measurable, bounded, and for $|\alpha| = 2b$, uniformly continuous in $R^{n+1}_+ \times (0, T)$.

We will introduce a distance function: for given $\gamma, 0<\gamma<1$, and $1<p<\infty$, let γ_p be a number such that

$$1-1/p \leqslant \gamma_p < \frac{2b}{2b-\gamma}(1-1/p).$$

We define $d_p(y, t) = \min(y, t^{\gamma_p/2b})$.

Throughout this work C will denote a constant, not necessarily the same at each occurrence. The connection between C and other parameters (eg. parameter of parabolicity, dimension, etc.) will be made explicit when relevant.

We will also let $\psi(r)$ denote any function of the form: constant$\cdot$ $\cdot \exp[-\text{constant}\cdot r]$, the constants and r being real and positive. When related to a solution of the operator L, these constants will depend only on the parameter of parabolicity π and on the $\max_{|\alpha|\leqslant 2b} \sup |a_\alpha(x, y, t)|$.

II. – The parametrix.

§ 6. – We will construct a kernel for a generalized volume potential following [5]. We consider first a differential operator with constant coefficients

$$L_0 \equiv \sum_{|\alpha|=2b} a_\alpha D_{xy}^\alpha - D_t .$$

Let $F(x, y, t) = \mathcal{F}_\xi\left(\exp \sum_{|\alpha|=2b} a_\alpha (i\xi)^\alpha t\right)(x, y)$, $\xi \in R^{n+1}$, be the fundamental solution of L_0. We construct a function $G_0(x, y, v, t)$, $y, v > 0$, satisfying as a function of (x, y, t),

(i) $G_0(\cdot, \cdot, v, \cdot) \in \mathring{L}^p_{2b,1}\left(R^n \times (\delta, \infty) \times (0, T)\right)$ for every $\delta > 0$, $1<p<\infty$,

(ii) $L_0 G_0(x, y, v, t) = 0$ for $y > 0$,

(iii) for $k = 0, \ldots, b-1$,

$$\lim_{y\to 0^+} \Lambda^{-k}(D_y^k G_0)(x, y, v, t) = [\Lambda^{-k} D_y^k F(\cdot, -v, \cdot)](x, t),$$

the limit taken in $L^p(S_T)$.

It is known [3], that G_0 can be written

$$G_0(x, y, v, t) = \sum_{k,j=0}^{b-1} \int_0^t \int_{R^n} \Lambda^{2b-1-k} D_y^k F(x-z, y, t-s) \cdot \cdot T_{k,j}[\Lambda^{-1}(D_y^j F)(\cdot, -v, \cdot)(x, t)](z, s)\, dz ds ,$$

where $(T_{k,j})$ is a $b \times b$ matrix of parabolic singular integral operators (of convolution type).

Considered as a function of x, v, t for $y > 0$, $G_0(x, y, v, t)$ is also a solution of the boundary value problem

(i') $G_0(\cdot, y \cdot, \cdot) \in \overset{\circ}{L}{}^{p}_{2b,1}\big(R^n \times (\delta, \infty) \times (0, T)\big)$ for every $\delta > 0$, $1 < p < \infty$, $y > 0$,

(ii') $\Big(\sum_{|\alpha| \leqslant 2b} a_\alpha D_x^{\alpha_1 \dots \alpha_n} (-D_v)^{\alpha_{n+1}} - D_t\Big) G_0(x, y, v, t) = 0$ for $v > 0$,

(iii') for $l = 0, \dots, b-1$,

$$\lim_{v \to 0^+} \text{in } L^p(S_T) \Lambda^{-l} D_v^l G_0(x, y, v, t) = (-1)^l \Lambda^{-l} D_y^l F(x, y, t).$$

Since for $y > 0$ $\lim_{v \to 0^+} D_v^l G_0$ exists in $L^p(S_T)$, (iii') implies that for $l = 0, \dots, b-1$,

$$D_v^l G_0(x, y, 0, t) = (-1)^l D_y^l F(x, y, t) \quad \text{for every } y > 0.$$

We now introduce the function

$$K(x, y, v, t) = F(x, y - v, t) - G_0(x, y, v, t).$$

For $l = 0, \dots, b-1$,

$$D_v^l K(x, 0, v, t) = 0, \quad v > 0, \quad \text{and} \quad D_y^l K(x, y, 0, t) = 0, \quad y > 0$$

§ 7. – The proof of the following theorem is long and technical in nature, and will not be included here.

THEOREM. For $y > 0$, $v > 0$, $y \neq v$, and $B > 0$

(i) $$|D_x^\alpha D_y^j D_v^h \Lambda^B F(x, y - v, t)| \leqq \frac{\psi\big(|x|/t^{1/2b}\big)\psi\big(|y - v|/t^{1/2b}\big)}{t^{(n+1+|\alpha|+j+h+B)/2b}},$$

(ii) $$|D_x^\alpha D_y^j D_v^h \Lambda^B G_0(x, y, v, t) \leqq \frac{\psi\big(|x|/t^{1/2b}\big)\psi\big(y/t^{1/2b}\big)\psi\big(v/t^{1/2b}\big)}{t^{(n+1+|\alpha|+j+h+B)/2b}},$$

(iii) $$|D_x^\alpha D_y^j D_v^h \Lambda_*^B K(x, y, v, t)| \leqq \frac{\psi\big(|x|/t^{1/2b}\big)\psi\big(|y - v|/t^{1/2b}\big)}{t^{(n+1+|\alpha|+j+h+B)/2b}}.$$

Clearly, (iii) follows from (i) and (ii).

We shall consider the operator with constant coefficients L_{0zrs},

$$(L_{0zrs} u)(x, y, t) \equiv \sum_{|\alpha| = 2b} a_\alpha(z, r, s) D_{xy}^\alpha u(x, y, t) - D_t u(x, y, t).$$

Let $F(z, r, s; x, y, t)$, $G_0(z, r, s; x, y, v, t)$, and $K(z, r, s; x, y, v, t)$ denote the fundamental solution F, and the functions G_0 and K (introduced in § 6) which are associated with the operator L_{0zrs}, $z \in R_n$, r, s, and v being real parameters. Clearly, these functions are solutions of the equation $L_{0zrs}u = 0$.

III. – Estimates for some surface potentials.

§ 8. – For $\Phi(x, t) \in L^p(S_T)$, $1 < p < \infty$, $j = 0, 1, \ldots, b-1$, $0 < \mu < 1$, we introduce the following potentials

$$T_{\mu j}\Phi)(x, y, t) = \int_0^t \int_{R^n} \Lambda^{b-j-\mu} D_y^j F(z, 0, s; x - z, y, t - s)\, \Phi(z, s)\, dz\, ds\,.$$

LEMMA 1. For $\Phi \in L^p(S_T)$, $1 < p < \infty$,

(i) $\|D_{x,y}^{\alpha} \Lambda^{\mu} T_{\mu j}\Phi(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C \cdot T^{(b-1-|\alpha|)/2b} \|\Phi\|^{L^p(S_T)}$ for $|\alpha| < b-1$,

(ii) $\|\Lambda^{b-1+\mu-k} D_y^k T_{\mu j}\Phi(\cdot, y, \cdot)\|_{L^p(S_T)} + \sum_{|\alpha|=b-1} \|D_{x,y}^{\alpha} \Lambda^{\mu} T_{\mu j}\Phi(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C \cdot \|\Phi\|_{L^p(S_T)}$,

(iii) $y^{|\alpha|-(b-1)} \|D_{x,y}^{\alpha} \Lambda^{\mu} T_{\mu j}\Phi(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C \cdot \|\Phi\|_{L^p(S_T)}$ for $|\alpha| > b-1$.

(iv) For $|\alpha| \leqslant b-1$,

$$\|D_{x,y}^{\alpha} \Lambda^{\mu} T_{\mu j}\Phi\|_{L^p(R_+^{n+1} \times (0,T))} \leqslant C \cdot T^{(b-1-|\alpha|+1/p)/2b} \cdot \|\Phi\|_{L^p(S_T)}.$$

PROOF. By the estimates on F, § 7, and Young's inequality in $dx\,dt$, we have

$$\|D_{xy}^{\alpha} \Lambda^{\mu} T_{\mu_j}\Phi(\cdot, y, \cdot)\|_{(L^p S_T)} \leqslant C \cdot \left\{ y^{b-1-|\alpha|} \int_0^{T/y^{2b}} \frac{\Psi(1/s^{1/2b})}{s^{(b+1+|\alpha|)/2b}}\, ds \right\} \|\Phi\|_{L^p S_T)}.$$

When $|\alpha| < b-1$ the expression in brackets is bounded by $C \cdot T^{(b-1-|\alpha|)/2b}$. When $|\alpha| > b-1$, the integral in ds is finite and bounded independently of y. In both case (i) and case (iii), the constants C depend on $|\alpha|$, the parameter of parabolicity π, and the $\max_{|\alpha|=2b} \sup |a_{\alpha}(x, y, t)|$.

The discussion above hints that in the case (ii), for $|\alpha| = b-1$ we will find the singularity of a parabolic singular operator. In fact, (ii) is a consequence of the theory of parabolic singular interals with variable kernel (see [3]). (iv) is obtained by taking L^p-norms in dy on both sides of the estimate above.

The proof of the following Lemma is straightforward

LEMMA 2. If Φ e $L^p(S_T)$, $1<p<\infty$, and $\gamma-1/p<\mu$, then

$$\|y^{b+1-\gamma}L(T_{\mu_j}\Phi)(x,y,t)\|_{L^p(R_+^{n+1}\times(0,T))}\leqslant\omega(T)\cdot\|\Phi\|_{L^p(S_T)},$$

where $\omega(T)\to 0$ as $T\to 0^+$.

IV. – Estimates for the volume potential.

§ 9. – We shall study a volume potential whose kernel is the function $K(z,\eta,s;x,y,v,t)$ (cf. § 6).

DEFINITION. For $f\in L^p\big(R_+^{n+1}\times(0,T)\big)$, $1<p<\infty$,

$$V_f(x,y,t)=\int_0^t\iint_{R_+^{n+1}}K(z,v,s;x-z,y,v,t-s)f(z,y,s)\,dz\,dv\,ds\,.$$

THEOREM. If $f\in L^p\big(R_+^{n+1}\times(0,T)\big)$, $1<p<\infty$, $0\leqslant\beta<\gamma-1/p$, $\gamma<1$, then

(i) for $k=0,\dots,b-1$, $\|\Lambda^{b-1+\beta-k}D_y^kV_f(\cdot,y,\cdot)\|_{L^p(S_T)}\to 0$ as $y\to 0^+$;

(ii) $$\sum_{k=0}^{b-1}\|\Lambda^{b-1-k+\beta}D_y^kV_f(\cdot,y,\cdot)\|_{L^p(S_T)}+\sum_{|\alpha|\leqslant b-1}\|D_{x,y}^\alpha\Lambda^\beta V_f(\cdot,y,\cdot)\|_{L^p(S_T)}+$$
$$+\sum_{b\leqslant|\alpha|\leqslant 2b-1}\|d_y(y,\cdot)^{|\alpha|-(b-1)}D_{x,y}^\alpha\Lambda^\beta V_f(\cdot,y,\cdot)\|_{L^p(S_T)}\leqslant$$
$$\leqslant C\cdot T^{\gamma''/2b}\cdot\|d_y(\cdot,\cdot)^{b+1-\gamma}f(\cdot,\cdot,\cdot)\|_{L^p(R_+^{n+1}\times(0,T)},$$

the constant depending on π, $\max_{|\alpha|=2b}\sup|a_\alpha|$, and p, and $\gamma''>0$ depending on γ,γ_p, b and a number $\gamma'<\gamma-\beta-1/p$.

$d_y(y,t)=\min(y,t^{\gamma_p/2b})$ is the distance function introduced in § 5.

PROOF. We recall that $D_v^lK(z,\eta,s;x,0,v,t)=0$ for $l\leqslant b-1$ (§ 6). We may therefore write, for $0\leqslant l\leqslant b-1$

$$K(z,\eta,s;x,y,v,t)=\frac{y^{l+1}}{l!}\int_0^1(1-\lambda)^lD_y^{l+1}K(z,\eta,s;x,\lambda y,v,t)\,d\lambda\,.$$

A similar remark applies to D_v^lK.

Part (i) is a direct consequence of the techniques used below, applied to

$$\Lambda^{b-1+\beta-k}D_y^kV_f=\frac{y^{b-k}}{(b-k-1)!}\int_0^t\iint_{R_+^{n+1}}\int_0^1\Lambda^{b-1+\beta-k}D_y^b\cdot$$

$$\cdot K(z,v,s;x-z,\lambda y,v,t-s)(1-\lambda)^{b-k-1}\,d\lambda f(z,v,s)\,dz\,dv\,ds\,.$$

PROOF OF (ii). We prove this in three Lemmas. In Lemmas A and B (§ 10) we prove the estimate for $d_p = y$. In Lemma C (§ 11) we show the estimate for $d_p = t^{\gamma_p/2b}$. For the general case we set

$$f(x, y, t) = X_{(y>t^{\gamma_p/2b})} \cdot f(x, y, t) + X_{(y \geqslant t^{\gamma_p/2b})} \cdot f(x, y, t) = f_1 + f_2$$

and apply to each potential V_{f_1}, V_{f_2} the corresponding estimate.

§ 10. – LEMMA A, (i). For $k = 0, \ldots, b-1$, $\beta + \gamma' < \gamma - 1/p$, $\gamma' > 0$,

$$\|\Lambda^{b-1+\beta-k} D_y^k V_f(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C \cdot T^{\gamma'/2b} \cdot \min(1, y^{\gamma-\beta-\gamma'-1/p}) \cdot \cdot \|v^{b+1-\gamma} f(z, v, s)\|_{L^p(R_+^{n+1} \times (0,T))} .$$

(ii) For $b-1 < |\alpha| \leqslant 2b-1$,

$$y^{|\alpha|-(b-1)} \|D_{x,y}^\alpha \Lambda^\beta V_f(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C \cdot T^{\gamma'/2b} \cdot \min(1, y^{\gamma-\beta-\gamma'-1/p}) \cdot \cdot \|v^{b+1-\gamma} f(z, v, s)\|_{L^p(R_+^{n+1} \times (0,T))} .$$

PROOF. Clearly,

$$\Lambda^{b-1+\beta-k} D_y^k V_f(x, y, t)| \leqslant \Big(\int_0^{y/2} + \int_{y/2}^{\infty} \Big) \int_0^t \int_{R^n} |\Lambda^{b-1+\beta-k} D_y^k \cdot K(z, v, s; x-z, y, v, t-s) \cdot f(z, v, s)| \, dz \, dv \, ds .$$

We estimate first the term $\int_0^{y/2}$. It can be written

$$\int_0^{y/2} \int_0^t \int_{R^n} \left| \Lambda^{b-1+\beta-k} D_y^k \int_0^1 \frac{(1-\lambda)^b}{(b-1)!} D_v^b K(z, v, s; x-z, y, \lambda v, t-s) d\lambda \cdot v^b \cdot f(z, v, s) \right| dz \, dv \, ds .$$

Applying the estimates for K (§ 7) and Young's inequality, and observing that $|y - \lambda v| \geqslant y/2$, we get

$$\left\| \int_0^{y/2} \right\|_{L^p(S_T)} \leqslant C \int_0^{y/2} \int_0^T \frac{\psi(y/s^{1/2b})}{s^{1+\beta/2b}} \left(\frac{T}{s} \right)^{\gamma'/2b} ds \cdot v^{b+1-\gamma} \|f(\cdot, v, \cdot)\|_{L^p(S_T)} \frac{dv}{v^{1-\gamma}} ,$$

where we have introduced the factor $(T/s)^{\gamma'/2b} > 1$, $\beta + \gamma' \leqslant \gamma - 1/p$. Applying now Hölder inequality in dv we obtain the desired estimate for $\int_0^{y/2}$, with right-hand side

$$C \cdot T^{\gamma'/2b} \cdot y^{\gamma-\beta-\gamma'-1/p} \cdot \int_0^{\infty} \frac{\psi(1/s^{1/2b})}{s^{1+(\beta+\gamma')/2b}} ds \cdot \|v^{b+1-\gamma} f(z, v, s)\|_{(L^p(R_+^{n+1} \times (0,T))} .$$

A similar argument gives the estimate (ii) for the corresponding $\int_0^{y/2}$.

For the term $\int_{y/2}^{\infty}$ in (i), we observe that

$$D_y^k K = \int_0^1 (1-\mu)^{k-1} v^k D_v^k \int_0^1 (1-\lambda)^{b-k-1} D_y^b K(\ldots; x-z, \lambda y, \mu v, t-s)\, d\lambda\, d\mu .$$

Therefore,

$$\left\| \int_{y/2}^{\infty} \right\|_{L^p(S_T)} \leqslant C \cdot y^{b-k} \int_{y/2}^{\infty} dv \int_0^1 d\mu \int_0^1 d\lambda \int_0^T \frac{\psi(|\lambda y - \mu v|/s^{1/2b})}{s^{1+\beta/2b}} \left(\frac{T}{s}\right)^{\gamma'/2b} ds \cdot$$
$$\cdot \frac{v^k v^{b-k+1-\gamma}}{v^{b-k+1-\gamma}} \cdot \|f(\cdot, v, \cdot)\|_{(L^p S_T)} .$$

Applying Hölder inequality in $dv\, d\mu\, d\lambda$ we get

$$\left\| \int_{y/2}^{\infty} \right\|_{L^p(S_T)} \leqslant C \cdot T^{\gamma'/2b} \cdot y^{b-k} \cdot \left(\int_{\gamma/2}^{\infty} \int_0^1 \int_0^1 \frac{dv\, d\lambda\, d\mu}{\{|\lambda y - \mu v|^{\beta+\gamma'} v^{b-k+1-\gamma}\}^{p/(p-1)}} \right)^{1-1/p} \cdot$$
$$\cdot \|v^{b+1-\gamma} f(z, v, s)\|_{L^p(R_+^{n+1} \times (0,T))}$$
$$\leqslant C \cdot T^{\gamma'/2b} \frac{y^{b-k} y^{1-1/p}}{y^{\beta+\gamma'+b-k+1-\gamma}} \cdot \left\{ \int_{\frac{1}{2}}^{\infty} \int_0^1 \int_0^1 \frac{dv\, d\lambda\, d\mu}{[|\lambda - \mu v|^{\beta+\gamma'} v^{b-k+1-\gamma}]^{p/(p-1)}} \right\}^{1-1/p} \cdot$$
$$\cdot \|v^{b+1-\gamma} f(z, v, s)\| .$$

For the term $\int_{y/2}^{\infty}$ in (ii) we have

$$\left\| \int_{y/2}^{\infty} \right\|_{L^p(S_T)} \leqslant C \int_{y/2}^{\infty} \int_0^1 v^{2b-1-|\alpha|} \int_0^T \frac{\psi(|y - \lambda v|/s^{1/2b})}{s^{1+\beta/2b}} \left(\frac{T}{s}\right)^{\gamma'/2b} ds \cdot \frac{v^{b+1-\gamma} \|f(\cdot, v, \cdot)\|^{L^p(S_T)} dv\, d\lambda}{v^{b+1-\gamma}}$$
$$\leqslant \frac{C \cdot T^{\gamma'/2b} y^{1-1/p}}{y^{|\alpha|-b} y^{2-\gamma+\beta+\gamma'}} \left\{ \int_{\frac{1}{2}}^{\infty} \int_1^0 \frac{dv\, d\lambda}{[|1 - \lambda v|^{\beta+\gamma'} v^{2-\gamma}]^{p/p-1}} \right\}^{1-1/p} \cdot$$
$$\cdot \|v^{b+1-\gamma} f(z, v, s)\|_{L^p(R_+^{n+1} \times (0,T))} .$$

For $|a| = 2b-1$, the integral $\int_0^1 d\lambda$ will not be present.

Observe that setting $\gamma' = \gamma - \beta - 1/p$ we have an estimate independent of $y^{\gamma-\beta-\gamma'-1/p}$. The proof is complete.

LEMMA *B*. For $|\alpha| \leqslant b-1$,

$$\|D^{\alpha}_{x,y} \Lambda^{\beta} V_f(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C \cdot T^{\gamma' 2/b} \cdot \min(1, y^{\gamma-\beta-\gamma'-1/p}) \cdot \cdot \|v^{b+1-\gamma} f(z, v, s)\|_{L^p(R^{n+1}_+ \times (0,T))} .$$

PROOF. Set $D^{\alpha}_{x,y} \Lambda^{\beta} = D^k_y D^{\delta}_x \Lambda^{\beta} = (D^{\delta}_x \Lambda^{-(b-1-k)}) D^k_y \Lambda^{b-1+\beta-k}$, $|\delta| + k = |\alpha| \leqslant b-1$. The result follows from Lemma *A* and the fact that $D^{\delta}_x \Lambda^{-(b-1-k)}$ is an $L^p(S_T)$ operator.

§ 11. – LEMMA *C*. For $|\alpha| \leqslant b-1$, γ_p as in § 5, and

$$\gamma_p < (2b-\beta-1/p)/(2b-\gamma),$$
$$\|D^{\alpha}_{x,y} \Lambda^{\beta} V_f(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C_1(T) \cdot \|s^{(\gamma_p/2b)(b+1-\gamma)} f(z, v, s)\|_{L^p(R^{n+1}_+ \times (0,T))},$$

and for $b \leqslant |\alpha| \leqslant 2b-1$,

$$\|t^{(\gamma_p/2b)(|\alpha|-(b-1))} D^{\alpha}_{xy} \Lambda^{\beta} V_f(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant C_2(T) \cdot \|s^{(\gamma_p/2b)(b+1-\gamma)} f\|_{L^p(R^{n+1}_+ \times (0,T))} .$$

PROOF. Case $|\alpha| \leqslant b-1$.

$$\|D^{\alpha}_{xy} \Lambda^{\beta} V_f(\cdot, y, \cdot)\|_{L^p(R^n)} \leqslant C \int_0^t \frac{1}{(t-s)^{(1+|\alpha|+\beta)/2b}} \int_0^{\infty} \Psi(|y-v|/(t-s)^{1/2b}) \cdot \|f(\cdot, v, s)\| \, dv ds .$$

We apply Hölder's inequality in dv and observe that

$$\Big(\int_0^{\infty} \Psi^{p'}(|y-v|/(t-s)^{1/2b}) \, dv \Big)^{1-1/p} \leqslant C \cdot (t-s)^{(1-1/p)/2b} .$$

Hence

$$\|D^{\alpha}_{xy} \Lambda^{\beta} V_f(\cdot, y, t)\|_{L^p(R^n)} \leqslant C \int_0^{\infty} \frac{X_{(t,0)}(s) (T/t)^{1-\{|\alpha|+\beta+1/p+\gamma_p(b+1-\gamma)\}/2b}}{(t-s)^{(|\alpha|+\beta+1/p)/2b} s^{(\gamma_p/2b)(b+1-\gamma)}} \cdot \cdot \big(s^{(\gamma_p/2b)(b+1-\gamma)} \|f[\cdot, \cdot, s)\|_{L^p(R^{n+1}_+)} \big) \, ds .$$

The power of (T/t) is positive, provided $\gamma_p < (b+1-(\beta+1/p))/(b+1-\gamma)$. Now the power of $(t-s)$ is clearly integrable, and $(\gamma_p/2b)(b+1-\gamma) + 1/p < 1$ due to the choice of γ_p. Hence by Hardy-Schur's Lemma (cf. [8], [9]), we obtain the desired inequality.

CASE $|\alpha| \geqslant b$. We proceed as in previous case, introducing now the factor $t^{(\gamma_p/2b)(|\alpha|-(b-1))}$, to obtain

$$(*) \quad t^{(\gamma_p/2b)(|\alpha|-(b-1))} \|D^{\alpha}_{xy} A^{\beta} V_f(\cdot, y, t)\|_{L^n(R^n)} \leqslant$$

$$\leqslant C \int_0^{\infty} \frac{X_{(0,t)}(s)(T/t)^{1-\{|\alpha|+\beta+1/p+\gamma_p(2b-\gamma-|\alpha|)\}/2b} t^{(\gamma_p/2b)(|\alpha|-(b-1))}}{(t-s)^{(|\alpha|+\beta+1/p)/2b} s^{(\gamma_p/2b)(b+1-\gamma)}} \cdot$$

$$\cdot \left(s^{(\gamma_p/2b)(b+1-\gamma)} \|f(\cdot, \cdot, s)\|_{L^p(R^{n+1}_+)}\right) ds .$$

We show that the power of (T/t) is positive by showing that the quantity in brackets is $< 2b$. The ratio $\big((2b-\beta-1/p)-|\alpha|\big)/\big((2b-\gamma)-|\alpha|\big)$ is an increasing function of $|\alpha|$; therefore its minimum value is attained at $|\alpha| = 0$: $1 < (2b-\beta-1/p)/(2b-\gamma)$. As the function $(r-\beta-1/p)/(r-\gamma)$ is decreasing, the condition $\gamma_p < (2b-\beta-1/p)/(2b-\gamma)$ also implies $\gamma_p < (b+1-\beta-1/p)/(b+1-\gamma)$ (see Case $|\alpha| \leqslant b-1$). It is clear that for $1 < p < \infty$ and $0 < \gamma < 1$, a number γ_p satisfying

$$1-1/p \leqslant \gamma_p < \frac{2b}{2b-\gamma}(1-1/p) \quad \text{and} \quad \gamma_p < \frac{2b-\beta-1/p}{2b-\gamma}$$

will suit to our requirements for all $|\alpha| \leqslant 2b-1$.

Returning to the estimate, we have $\|t^{(\gamma_p/2b)(|\alpha|-(b-1))} D^{\alpha}_{xy} A^{\beta} V_f(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant$ $\leqslant L^p$-norm in t of the integral in the right hand side of $(*)$. The result follows now from Hardy's Lemma.

V. – The operator J.

§ 12. – We shall study a commutator for the operator

$$(Jf)(x, y, t) = \lim_{\varepsilon \to 0^+} \int_0^{t-\varepsilon} \iint_{R^{n+1}_+} (L_{xyt} K)(z, v, s; x-z, y, v, t-s) f(z, v, s)\, dz\, dv\, ds ,$$

Consider

$$[y^{b+1-\gamma} J(f) - J\big(v^{b+1-\gamma} f(z, v, s)\big)](x, y, t) =$$

$$= \sum_{|\alpha|=2b} T^0_{\alpha}\big([a_{\alpha}(x, y, t) - a_{\alpha}(z, v, s)] f(z, v, s)\big)(x, y, t) +$$

$$+ \sum_{|\alpha|<2b} a_{\alpha}(x, y, t)\, T^0_{\alpha}\big(f(z, v, s)\big)(x, y, t) ,$$

where we have set

$$T_\alpha^0(f)(x, y, t) = \int_0^t \iint_{R_+^{n+1}} (y^{b+1-\gamma} - v^{b+1-\gamma}) D_{x,y}^\alpha K(z, v, s; x-z, y, v, t-s) \cdot \cdot f(z, v, s)\, dz\, dv\, ds .$$

We also set

$$T_\alpha^1(f)(x, y, t) = \int_0^t \iint_{R_+^{n+1}} (t^{(\gamma_p/2b)(b+1-\gamma)} - s^{(\gamma_p/2b)(b+1-\gamma)})(D_{x,y}^\alpha K) f(z, v, s)\, dz\, dv\, ds ,$$

T_α^1 being the corresponding operators for the commutator

$$[t^{(\gamma_p/2b)(b+1-\gamma)} J(f) - J(s^{(\gamma_p/2b)(b+1-\gamma)} f(z, v, s))](x, y, t) .$$

LEMMA. Let $f \in L^p(R_+^{n+1} \times (0, T))$. For $0 \leqslant |\alpha| \leqslant 2b$, $1/p < \gamma < 1$, and $1 < p < \infty$,

(i) $\|T_\alpha^0 f\|_{L^p(R_+^{n+1} \times (0,T))} \leqslant C \cdot T^{1-(|\alpha|/2b)} \|y^{b+1-\gamma} f(x, y, t)\|_{L^p(R_+^{n+1} \times (0,T))}$,

(ii) $\|T_\alpha^1 f\|_{L^p(R_+^{n+1} \times (0,T))} \leqslant C \cdot T^{1-(|\alpha|/2b)} \|t^{(\gamma_p/2b)(b+1-\gamma)} f(x, y, t)\|_{L(^p R_+^{n+1} \times (0,T))}$.

PROOF OF (i). We set as usual

$$|(T_\alpha^0 f)(x, y, t)| \leqslant \Big(\int_0^{y/2} + \int_{y/2}^{\infty} \Big) \int_0^t \int_{R^n} |y^{b+1-\gamma} - v^{b+1-\gamma}| \cdot |D_{x,y}^\alpha K| \cdot |f|\, dz\, dv\, ds .$$

To estimate the first term, we use the known properties of K together with Young's inequality in the variables z and s, and the fact that $v \leqslant y/2$ to obtain

$$\Big\| \int_0^{y/2} \Big\|_{L^p(S_T)} \leqslant C \cdot T^{1-(|\alpha|/2b)} \int_0^\infty \frac{X_{(0,y/2)}(v)}{y^\gamma \cdot v^{1-\gamma}} \big(v^{b+1-\gamma} \|f(\cdot, v, \cdot)\|_{L^p(S_T)}\big)\, dv .$$

The desired inequality follows from this one, applying Hardy's lemma ([8], [9]) and recalling that $\gamma > 1/p$.

For the second term, we apply the known estimate directly to K to get

$$\Big\| \int_{y/2}^{2y} \Big\|_{L^p(S_T)} \leqslant C \cdot T^{1-(|\alpha|/2b)} \int_0^\infty \frac{X_{(y/2,2y)}(v) |y^{b+1-\gamma} - v^{b+1-\gamma}|}{v^{b+1-\gamma} |y - v|} \cdot \big(v^{b+1-\gamma} \|f(\cdot, v, \cdot)\|_{L^p(S_T)}\big)\, dv .$$

We now apply Hardy's lemma to obtain the desired inequality.

PROOF OF (ii). It is clear that

$$|T^1_\alpha(f)(x, y, t)| \leqslant \int_0^t \iiint_{R^{n+1}_u} |t^{(\gamma_p/2b)(b+1-\gamma)} - s^{(\gamma_p/2b)(b+1-\gamma)}||D^\alpha_{x,y} K||f|\,dz\,dv\,ds\,.$$

By applying Young's inequality in the variables z, v to the known estimates for K, and observing that $(t-s)^{1-(|\alpha|/2b)} \leqslant T^{1-(|\alpha|/2b)}$, it is easy to see that

$$\|T^1_\alpha f\|_{L^p(R^{n+1}_+ \times (0,T))} \leqslant C \cdot T^{1-(|\alpha|/2b)} \Big\| \int_0^\infty k(t, s) \Big(s^{(\gamma_p/2b)(b+1-\gamma)} \|f(\cdot, \cdot, s)\|_{L^p(R^{n+1})} \Big)\, ds \Big\|_{L^p(0,\infty)},$$

where

$$k(t, s) = \frac{X_{(0,t)}(s)}{s^{(\gamma_p/2b)(b+1-\gamma)}} \, \frac{t^{(\gamma_p/2b)(b+1-\gamma)} - s^{(\gamma_p/2b)(b+1-\gamma)}}{t-s}\,.$$

Due to the conditions on γ_p and the integrability of the second factor $k(t, s)$ satisfies the hypotheses in Hardy's lemma, from which part (ii) follows. This completes the proof of the lemma.

§ 13. – LEMMA. For $f \in L^p(R^{n+1}_+ \times (0, T))$, $1/p < \gamma < 1$, $1 < p < \infty$,

(i) $\|y^{b+1-\gamma} J(f) - J(v^{b+1-\gamma} f)\|_{L^p(R^{n+1}_+ \times (0,T))} \leqslant \omega(T) \|y^{b+1-\gamma} f\|_{L^p(R^{n+1}_+ \times (0,T))}$

(ii) $\|t^{(\gamma_p/2b)(b+1-\gamma)} J(f) - J(s^{(\gamma_p/2b)(b+1-\gamma)} f)\|_{L^p(R^{n+1}_+ \times (0,T))} \leqslant$

$$\leqslant \omega(T) \|t^{(\gamma_p/2b)(b+1-\gamma)} f(x, y, t)\|_{L^p(R^{n+1}_+ \times (0,T))},$$

with $\omega(T) \to 0$ as $T \to 0^+$, ω depending on the moduli of continuity of the a_α, $|\alpha| = 2b$.

PROOF. Let $\varphi \in C_0^\infty(R^{n+2})$, $\varphi \geqslant 0$, with $\iiint_{R^{n+2}} \varphi = 1$ and support φ contained in the set $|x|^2 + y^2 + t^2 < 1$. For $|\alpha| = 2b$, we extend $a_\alpha(x, y, t)$ to all R^{n+2}, preserving uniform continuity and define, for $\lambda > 0$,

$$a^\lambda_\alpha(x, y, t) = (1/\lambda^{n+2}) \iiint_{R^{n+2}} a_\alpha(w, u, r)\, \varphi\left(\frac{x-w}{\lambda}\,,\ \frac{y-u}{\lambda}\,,\ \frac{t-r}{\lambda} \right) dw\,du\,dr\,.$$

Then $a^\lambda_\alpha \in C^\infty(R^{n+2})$, $|a_\alpha(x, y, t) - a^\lambda_\alpha(x, y, t)| \leqslant \omega(\lambda) =$ maximum of moduli of continuity of a_α, $|\alpha| = 2b$, $|Da_\alpha(x, y, t)| \leqslant (C/\lambda)\omega(\lambda)$, D being any derivative, and therefore $|a^\lambda_\alpha(x, y, t) - a^\lambda_\alpha(z, v, s)| \leqslant (C/\lambda)\omega(\lambda)(|x-z| + |y-v| + |t-s|)$.

According to the decomposition and the estimates in previous §, it will be sufficient to consider the case $|\alpha| = 2b$, $j = 0, 1$. We have

$$T_\alpha^j([a_\alpha(x, y, t) - a_\alpha(z, v, s)]f) = T_\alpha^j([a_\alpha(x, y, t) - a_\alpha^\lambda(x, y, t)]f) + \\ + T_\alpha^j([a_\alpha^\lambda(z, v, s) - a_\alpha(z, v, s)]f) + T_\alpha^j([a_\alpha^\lambda(x, y, t) - a_\alpha^\lambda(z, v, s)]f).$$

The first two terms are in absolute value less than or equal to $\omega(\lambda)$ times (bounds for $|T_\alpha^j f|$ in proof (i) and (ii) of lemma, § 12). For the third, we set $|y^{b+1-\gamma} - v^{b+1-\gamma}| = E^0$, $|t^{(\gamma_p/2b)(b+1-\gamma)} - s^{(\gamma_p/2b)(b+1-\gamma)}| = E^1$ and ovserve that it is in absolute value

$$\leqslant \int_0^t \int_0^\infty \int_{R^n} |D_{x\,y}^\alpha K| \cdot E^j \cdot (C/\lambda)\omega(\lambda)(|x - z| + |y - v| + |t - s|) \cdot |f(z, v, s)|\, dz\, dv\, ds .$$

The usual procedure applies to each term above. For the third, $t - s < T$, so we set $\lambda = T^{1/2b}$ and obtain with a new $\omega(T)$ that tends to zero as $T \to 0+$, the same type of bound for $|T_\alpha^j f|$ used in Lemma, § 12. The proofs of § 12 apply again.

§ 14. – LEMMA. Let $f \in L^p(R_+^{n+1} \times (0, T))$. Then the operator (Jf) maps $L^p(R_+^{n+1} \times (0, T))$ continuously into itself, and belongs to $\mathfrak{Z}(R_+^{n+1} \times (0, T))$ (cf. § 2).

PROOF. Condition (i), § 2 is clear. To prove condition (ii), we set $J = J_1 - J_2$, where

$$J_1 f(x, y, t) = \lim_{\varepsilon \to 0^+} \text{ in } L^p \int_0^{t-\varepsilon} \iint_{R_+^{n+1}} L_{xyt} F(z, v, s; x - z, y - v, t - s) f(z, v, s)\, dz\, dv\, ds ,$$

$$J_2 f(x, y, t) = \int_0^t \iint_{R_+^{n+1}} L_{xyt} G_0(z, v, s; x - z, y, v, t - s) f(z, v, s)\, dz\, dv\, ds .$$

J_1 is known to belong in $\mathfrak{Z}$ (cf. [5]). Furthermore,

$$J_1 = \sum_{|\alpha| = 2b} (a_\alpha \overline{K}_\alpha - \overline{K}_\alpha a_\alpha) + \sum_{|\alpha| < 2b} a_\alpha J_\alpha ,$$

$\overline{K}_\alpha$ being a variable kernel, and the J_α, L^p-operators in $\mathfrak{Z}$. For J_2, we see

that it can be decomposed in two sums, setting

$$N_\alpha(f) = \int_0^t \iint_{R_+^{n+1}} D_{x,y}^\alpha G_0(z, v, s; x-y, v, t-s) f(z, v, s)\, dz\, dv\, ds$$

$$J_2 f = \sum_{|\alpha|=2b} N_\alpha([a_\alpha(x, y, t) - a_\alpha(z, v, s)]f) + \sum_{|\alpha|<2b} a_\alpha N_\alpha(f).$$

Consider first the case $|\alpha| = 2b$. Replacing f by $X_{(a,a+\varepsilon)}(s) f(z, v, s)$ se see that each term in the first sum above can be written as

$$N_\alpha([a_\alpha(x, y\ \ t) - a_\alpha^\lambda(x\ \ y, t)] X_{(a,a+\varepsilon)} f) + N_\alpha([a_\alpha^\lambda(z, v, s) - a_\alpha(z, v, s)] X f) + \\ + N_\alpha([a_\alpha^\lambda(x, y, t) - a_\alpha^\lambda(z, v, s)] X_{(a,a+\varepsilon)} f),$$

where each term is in absolute value

$$\leqslant \omega(\varepsilon^{1/2b}) \int_0^t \iint_{R_+^{n+1}} (\text{bounds for } |D_{x,y}^\alpha G_0| \text{ in } \S\,7) \cdot X_{(a,a+\varepsilon)}(s) |f(z\ \ v\ \ s)|\, dz\, dv\, ds,$$

if $\lambda = \varepsilon^{1/2b}$. (We observe that for $a \leqslant t \leqslant a+\varepsilon$ and $a \leqslant s \leqslant t$, $t-s \leqslant \varepsilon$ and $X_{(0,\varepsilon)}(t-s) = 1$).

By Young's inequality in $dx\,dt$, the $L^p(R^n \times (a, a+\varepsilon))$-norm of the above expression is bounded by

$$C\omega(\varepsilon^{1/2b}) \int_0^\infty \frac{1}{|y+v|} \int_0^{\varepsilon/|y+v|^{2b}} \frac{\Psi(1/s^{1/2b})}{s^{1+1/2b}}\, ds \|f(\cdot, v, \cdot)\|_{L^p(S_{(a,a+\epsilon)})} dv.$$

(Here we have set $\Psi(y/s^{1/2b})\Psi(v/s^{1/2b}) \leqslant \Psi(|y+v|/s^{1/2b})$, the Ψ being of exponential type (cf. § 5). The integral in ds is bounded independently of y, v).

We now apply Hardy's lemma to obtain the inequality

$$\sum_{|\alpha|=2b} \|N_\alpha([a_\alpha(x, y, t) - a_\alpha(z, v, s)] X_{(a,a+\varepsilon)} f)\|_{L^p(R_+^{n+1} \times (a,a+\varepsilon))} \leqslant \\ \leqslant C\omega(\varepsilon^{1/2b}) \|f\|_{L^p(R_+^{n+1} \times (a,a+\varepsilon))}.$$

For $|\alpha| < 2b$, using the fact that $\Psi(y)\Psi(v) \leqslant \Psi(|y-v|)$, $y, v > 0$, and applying the known estimates for G_0 and the remarks on $X_{(0,\varepsilon)}(t-s)$, we easily get

$$\|N_\alpha(X_{(a,a+\varepsilon)} f)\|_{L^p(R_+^{n+1} \times (a,a+\varepsilon))} \leqslant C\varepsilon^{1-(|\alpha|/2b)} \|f\|_{L^p(R_+^{n+1} \times (a,a+\varepsilon))},$$

C depending on $\max_{|\alpha|<2b} \sup |a_\alpha|$, completing the proof of the lemma.

REMARK. We have essentially shown that

$$L_{xvt}\Big(\int_0^t \iint_{R_+^{n+1}} G_0(z, v, s; x-z, y, v, t-s) f(z, v, s)\, dz\, dv\, ds\Big) = J_2 f$$

belongs to $L^p\big(R_+^{n+1}\times(0, T)\big)$ provided $f \in L^p\big(R_+^{n+1}\times(0, T)\big)$, and its norm is $\leqslant C\|f\|_{L^p(R_+^{n+1}\times(0,T))}$. We observe that the boundary term in the computation of the D_t is zero, due to the estimates for G_0 (§ 7); we also see that

$$\int_0^t \iint_{R_+^{n+1}} G_0(z, v, s; x-z, y, v, t-s) f(z, v, s)\, dz\, dv\, ds \in \mathring{L}^p_{2b,1}\big(R_+^{n+1}\times(0, T)\big)\,.$$

Estimate for J. As a consequence of the results above, we have the following

LEMMA. For $f \in L^p\big(R_+^{n+1}\times(0, T)\big)$, $1/p<\gamma<1$, $1<p<\infty$,

$$\|d_y(y, t)^{b+1-\gamma} Jf(x, y, t)\|_{L(^pR_+^{n+1}\times(0,T))} \leqslant \omega(T)\|d_y(y, t)^{b+1-\gamma} f(x, y, t)\|_{L^p(R_+^{n+1}\times(0,T))}\,,$$

where $\omega \to 0$ as $T \to 0^+$.

§ 15. – The operator J having small norm as an operator on $L^p\big(R_+^{n+1}\times(a, a+\varepsilon)\big)$ for ε suitably small, it follows that $I-J$ is invertible over $L^p\big(R_+^{n+1}\times(0, T)\big)$. In fact, choosing m large enough,

$$\|J(X_{(a,a+(T/m))} g\|_{L^p(R_+^{n+1}\times(a,a+(T/m)))} \leqslant (\tfrac{1}{2})\|g\|_{L^p(R_+^{n+1}\times(a,a+(T/m)))})\,,$$

provided $a+(T/m)\leqslant T$. Let g_1 be a function with support in $R_+^{n+1}\times(0, T/m)$ such that $(I-J)g_1 = f$ on $R_+^{n+1}\times(0, T/m)$, and in general let g_k have support in $R_+^{n+1}\times\big((k-1)T/m, k(T/m)\big)$, and satisfy $(I-J)\,g_k = f - \sum_{h=1}^{k-1}(I-J)g_h$ on $R_+^{n+1}\times\big((k-1)T/m, k(T/m)\big)$. Set $g = \sum_{k=1}^{n} g_k$: $g \in L^p\big(R_+^{n+1}\times(0, T)$, being a sum of functions in that space, and $(I-J)g = f$.

With the construction above it is easy to prove the following

LEMMA. If $f \in L^p\big(R_+^{n+1}\times(0, T)\big)$, $1<p<\infty$, $1/p<\gamma<1$, then

$$\|d_y(y, t)^{b+1-\gamma}(I-J)^{-1} f\|_{L^p(R_+^{n+1}\times(0,T))} \leqslant C\|d_y(y, t)^{b+1-\gamma} f\|_{L^p(R_+^{n+1}\times(0,T))}\,.$$

($d_y(y, t)$ can be replaced by y or $t^{(\gamma_p/2b)}$).

REMARK. If the function f and the coefficients of L_{xyt} are differentiable, so is the function $(I-J)^{-1}f=g$. In fact, the only possible points of discontinuity of g or its derivatives with respect to time are those in the partition kT/m, $k\leqslant m$. To see that g is smooth at those points, we only have to construct $(I-J)^{-1}f$ with a different partition. The uniqueness of $(I-J)^{-1}f$ shows the differentiability of g at the points kT/m.

The differentiability of $(I-J)^{-1}=\sum J^k$ for small t can be seen from the fact that $DJ(f)=J^{\sim}(f)+J(Df)$, D denoting any derivative, and $J^{\sim}$ being a $\mathfrak{Z}$-operator that depends on the derivatives of the coefficients. From this identity we derive the recursion inequality

$$\|DJ^n(f)\|\leqslant\omega(r)\big(\|J^{n-1}(f)\|+\|DJ^{n-1}(f)\|\big)\,,\qquad \omega\to 0 \text{ as } r\to 0^+\,,$$

which shows convergence in norm of the series $\sum DJ^n(f)$ for r sufficiently small.

VI. – The main results in the half-space.

§ 16. – THEOREM. Let $f\in L^p\big(R_+^{n+1}\times(0,T)\big)$ for some p, $1<p<\infty$. Set

$$\begin{aligned}V(x,y,t)&=-\int_0^t\iint_{R_+^{n+1}}K(z,v,s;x-z,y,v,t-s)(I-J)^{-1}f(z,v,s)\,dz\,dv\,ds\\&=-V_{(I-J)^{-1}f}(x,y,t)\,.\end{aligned}$$

Then for $0\leqslant\beta<\gamma-1/p$,

(i) $V(x,y,t)\in\overset{\circ}{L}{}^p_{2b,1}\big(R_+^{n+1}\times(0,T)\big)$, and $LV=f$ for $y>0$,

(ii) for $k=0,\dots,b-1$,

$$\lim_{y\to 0^+}\text{ in } L^p(S_T)\Lambda^{b-1+\beta-k}D_y^kV(\cdot,y,\cdot)=0\,.$$

(iii) $$\sum_{k=0}^{b-1}\|\Lambda^{b-1-k+\beta}D_y^kV(\cdot,y,\cdot)\|_{L^p(S_T)}$$
$$+\sum_{|\alpha|\leqslant b-1}\|D_{x,y}^{\alpha}\Lambda^{\beta}V(\cdot,y,\cdot)\|_{L^p(S_T)}$$
$$+\sum_{b\leqslant|\alpha|\leqslant 2b-1}\|d_a(y,\cdot)^{|\alpha|-(b-1)}D_{x,y}^{a}\Lambda^{\beta}V(\cdot,y,\cdot)\|_{L^p(S_T)}\leqslant C(T)\|d_p^{b+1-\gamma}f\|_{L^p}\,,$$

where $C(T)\to 0$ as $T\to 0^+$.

(A similar estimate holds with d_p replaced by y, and with d_p replaced by $t^{\gamma p/2b}$).

PROOF (i). Recall that

$$K(z, v, x; y, v, t) = F(z, v, s; x, y - v, t) - G_0(z, v, s; x, y, v, t),$$

and that $(I - J)^{-1} f \in L^p(R_+^{n+1} \times (0, T))$. Set $V = - V_1 + V_2$, where

$$V_1(x, y, t) = \int_0^t \iint_{R_+^{n+1}} F(z, v, s; x - z, y - v, t - s)(I - J)^{-1} f(z, v, s)\, dz\, dv\, ds,$$

$$V_2(x, y, t) = \int_0^t \iint_{R_+^{n+1}} G_0(z, v, s; x - z, y, v, t - s)(I - J)^{-1} f(z, v, s)\, dz\, dv\, ds.$$

We showed (cf. Remark to Lemma, § 14) that $V_2 \in \mathring{L}^p_{2b,1}(R_+^{n+1} \times (0, T))$. It is known [2] that $V_1 \in \mathring{L}^p_{2b,1}(R_+^{n+1} \times (0, T))$, hence $V \in \mathring{L}^p_{2b,1}(R_+^{n+1} \times (0, T))$. Also

$$LV_1 = - (I - J)^{-1} f + J_1((I - J)^{-1} f),$$

and by same Remark, $LV_2 = J_2((I - J)^{-1} f)$. Therefore

$$\begin{aligned} LV &= (I - J)^{-1} f - (J_1 - J_2)(I - J)^{-1} f \\ &= (I - J)^{-1} f - J(I - J)^{-1} f = f. \end{aligned}$$

Part (ii) is an immediate consequence of Theorem § 9, (i), and the fact that $(I - J)^{-1} f$ belongs in $L^p(R_+^{n+1} \times (0, T))$.

Part (iii) follows from the estimates for the volume potential (Theorem, § 9, cf. lemmas A, B, C, §§ 10-11), and from estimate in Lemma, § 15.

§ 17. – For $j = 0, \ldots, b - 1$, we introduce the functions

$$u_j(x, y, t) = (T_{\mu j} \Phi)(x, y, t) + \\ + \int_0^t \iint_{R_+^{n+1}} K(z, v, s; x - z, y, v, t - s)(I - J)^{-1}(L T_{\mu j} \Phi)(z, v, s)\, dz\, dv\, ds \quad (*).$$

(cf. definitions in § 6(K), § 8($T_{\mu j}$), and in § 9 (volume potential)).

THEOREM. If $\Phi \in L^p(S_T)$, then for $j = 0, \ldots, b - 1$,

(i) $u_j(x, y, t) \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$,

(ii) $L\mu_j = 0$ in $R_+^{n+1} \times (0, T)$.

PROOF. Is included in §§ 17-19. We introduce the following notation. For $|\alpha| = 2b$, let $a_\alpha^\lambda(z, v, s)$ denote the regularization of the coefficient $a_\alpha(z, v, s)$ in L(cf. § 13). We will denote by $F^\lambda(z, v, s; x, y, t)$ the $\mathcal{F}_\xi\Big[\exp \sum_{|\alpha|=2b} a_\alpha^\lambda(z, v, s)(i\xi)^\alpha t\Big](x, y)$, by $T_{\mu j}^\lambda \Phi$ the surface potential

$$(T_{\mu j}^\lambda \Phi)(x, y, t) = \int_0^t \int_{R^n} \Lambda^{b-j-\mu} D_y^j F^\lambda(z, 0, s; x - z, y, t - s)\, \Phi(z, s)\, dz\, ds\,,$$

and by $u_j^\lambda(x, y, t)$ the corresponding functions (*).

If $\Phi \in C_0^\infty\big(R^n \times (0, \infty)\big)$, it can be seen that $u_j^\lambda(x, y, t) \in \mathring{L}_{2b,1}^p\big(R_+^{n+1} \times (0, T)\big)$, and that $Lu_j^\lambda = 0$ for $y > 0$. The second statement will follow from the theorem in § 16; the first is a consequence of the same theorem in § 16 and of the fact that $(I - J)^{-1}$ is a L^p-mapping, if we observe that $T_{\mu j}^\lambda \Phi \in \mathring{L}_{2b,1}^p\big(R_+^{n+1} \times (0, T)\big)$.

When the coefficients a_α, $|\alpha| = 2b$, are only bounded and uniformly continuous, and $\Phi \in C_0^\infty\big(R^n \times (0, \infty)\big)$, we will show that the expression in (*), § 17, belongs in $\mathring{L}_{2b,1}^p\big(R_E \times (\delta, \infty) \times (0, T)\big)$ for every $\delta > 0$. We first state two lemmas.

LEMMA 1. Let $\lambda, \lambda' > 0$. For $j = 0, \ldots, b - 1$,

$$\|(T_{\mu j}^\lambda - T_{\mu j}^{\lambda'})\, \Phi\|_{L^p(R_+^{n+1} \times (0,T))} \leqslant$$

$$\leqslant C \cdot T^{(\mu + 1/p)/2b} \max_{|\alpha|=2b} \|a_\alpha^\lambda(z, 0, s) - a_\alpha^{\lambda'}(z, 0, s)\|_{L^\infty(S_T)} \|\Phi\|_{L^p(S_T)}\,.$$

LEMMA 2. If $\Phi \in L^p(S_T)$, $1 < p < \infty$, $\gamma - 1/p < \mu$, then

$$\|y^{b+1-\gamma} L(T_{\mu j}^\lambda - T_{\mu j}^{\lambda'})\, \Phi(x, y, t)\|_{L^p(R_+^{n+1} \times (0,T))} \leqslant C\omega(T) \max_{|\alpha|=2b} \|a_\alpha^\lambda - a_\alpha^{\lambda'}\|_{L^\infty(S_T)} \|\Phi\|_{L^p(S_T)}\,.$$

These lemmas are L^p versions of similar results in Pogorzelski [12], and can be proved by similar arguments.

If now $\theta(y) \in C^\infty(0, \infty)$, $\theta = 0$ for $y < \delta$, $\theta = 1$ for $y \geqslant 2\delta$, and we set $\theta^{(l)}(y) = D_y^l \theta(y)$, we have the

COROLLARY. For $|\alpha| + l \leqslant 2b$,

$$\|\theta^{(l)}(y)\, D_{xy}^\alpha (T_{\mu j}^\lambda - T_{\mu j}^{\lambda'}) \Phi(x, y, t)\|_{L^p(R_+^{n+1} \times (0,T))} \leqslant C_\delta \omega(T) \max_{|\alpha|2=2b} \|a_\alpha^\lambda - a_\alpha^{\lambda'}\|_{L^\infty(S_T)} \|\Phi\|_{L^p(S_T)}\,.$$

We now continue with the proof of the Theorem, § 17.

§ 18. – We recall that for $\Phi \in C_0^\infty(R^n \times (0, \infty))$, $u_j^\lambda(x, y, t) = (T_{\mu j}^\lambda \Phi) + V_{(\Phi)}^\lambda$ belongs to $\mathring{L}^p_{2b,1}(R_+^{n+1} \times (0, T))$ (we have set $V_{(\Phi)}^\lambda = V_{(I-J)^{-1}(LT_j^\lambda \Phi)}$).

The product $\theta(y)(u_j^\lambda - u_j^{\lambda'})$ vanishes near $y = 0$ and $t = 0$. It is known [4] that

$$\|\theta(u_j^\lambda - u_j^{\lambda'})\|_{L^p_{2b,1}(R_+^{n+1}\times(0,T))} \leqslant C \cdot \|L(\theta(u_j^\lambda - u_j^{\lambda'}))\|_{L^p(R_+^{n+1}\times(0,T))} \leqslant$$

$$\leqslant C \sum_{|\alpha|+l\leqslant 2b} \|\theta^{(l)} D_{x,y}^\alpha (T_{\mu j}^\lambda - T_{\mu j}^{\lambda'})\Phi\|_{L^p(R_+^{n+1}\times(0,T))} +$$

$$+ C\|\theta L(T_{\mu j}^\lambda - T_{\mu j}^{\lambda'})\Phi\|_{L^p(R_+^{n+1}\times(0,T))} +$$

$$+ C\|\sum_{l>0} \theta^{(l)}(y) D_{x,y}^\alpha V_{(\Phi)}^{\lambda,\lambda'}(x, y, t)\|_{L^p(R_+^{n+1}\times(0,T))},$$

where the meaning of $V_{(\Phi)}^{\lambda,\lambda'}(x, y, t)$ is clear.

Now

$$\|\sum_{l>0} \theta^{(l)} D_{x,y}^\alpha V_{(\Phi)}^{\lambda,\lambda'}\|_{L^p(R_+^{n+1}\times(0,T))}$$

$$\leqslant \sum_{l>0} \|\theta^{(l)}(y)\| D_{x,y}^\alpha V_{(\Phi)}^{\lambda,\lambda'}(\cdot, y, \cdot)\|_{L^p(S_T)}\|_{L^p(0,\infty)}$$

$$\leqslant \sum_{l>0} \left\|\theta^{(l)}(y) \cdot \frac{1}{\delta^{|\alpha|}} \cdot y^{|\alpha|}\right\| D_{x,y}^\alpha V_{(\Phi)}^{\lambda,\lambda'}(\cdot, y, \cdot)\|_{L^p(S_T)}\|_{L^p(0,\infty)}$$

$$\leqslant C(T) \sum_{l>0} \left\|\frac{\theta(l)}{\delta^{|\alpha|}}\right\|_{L^\infty(0,\infty)} \|y^{b+1-\gamma}(I-J)^{-1} L(T_{\mu j}^\lambda - T_{\mu j}^{\lambda'})\Phi(x, y, t)\|_{(L^p(R_+^{n+1}\times(0,T))}$$

$$\leqslant C_\delta(T)\|y^{b+1-\gamma} L(T_{\mu j}^\lambda - T_{\mu j}^{\lambda'})\Phi(x, y, t)\|_{L^p(R_+^{n+1}\times(0,T))}.$$

(We have used the estimates for the volume potential (§ 9, and lemmas A and B, § 10) with $\beta = 0$, and the Lemma in § 15)

By Lemmas 1 and 2, § 17, we conclude that

$$\theta(u_j^\lambda - u_j^{\lambda'})\|_{L^p_{2b,1}(R_+^{n+1}\times(0,T))} \leqslant C(T), \qquad \max_{|\alpha|=2b} \|a_\alpha^\lambda - a_\alpha^{\lambda'}\|_{L^\infty(S_T)} \cdot \|\Phi\|_{L^p(S_T)}.$$

This shows that $\{u_j^\lambda\}_\lambda$ is a Cauchy sequence in $\mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$. Let $\bar{u}_j \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$ be its limit. By Lemmas 1 and 2, with $T_{\mu j}^{\lambda'}$ replaced by $T_{\mu j}$, it follows that the potentials $T_{\mu j}^\lambda \Phi$ and $V_{(\Phi)}^\lambda$ converge to $T_{\mu j}\Phi$,

$$V_{(\Phi)} = \int_0^t \iint_{R_+^{n+1}} K(z, v, s; x - z, y, v, t - s)(I - J)^{-1}(LT_{\mu j}\Phi)(z, v, s)\, dz\, dv\, ds.$$

Hence $\bar{u}_j$ admits the representation (*) for $\Phi \in C_0^\infty(R^n \times (0, \infty))$, i.e.,

$$\bar{u}_j(x, y, t) = (T_{\mu j}\Phi)(x, y, t) +$$

$$+ \int_0^t \iint_{R_+^{n+1}} K(z, v, s; x - z, y, v, t - s)(I - J)^{-1}(LT^{\mu j}\Phi)(z, v, s)\, dz\, dv\, ds\,, \quad (*)$$

and $\bar{u}_j \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$.

Clearly $L\bar{u}_j = 0$ for $y > 0$.

§ 19. – The same representation holds for $\Phi \in L^p(S_T)$. To see this, select $\Phi^\lambda \in C_0^\infty(R^n \times (0, \infty))$, $\|\Phi - \Phi^\lambda\|_{L^p(S_T)} \to 0$ as $\lambda \to 0$, and let $\bar{u}_j^\lambda$ be the function (*) above with Φ^λ in place of Φ. Again let $\theta(y) \in C^\infty(0, \infty)$, $\theta(y) = 0$ for $y < \delta$, $\theta(y) = 1$ for $y > 2\delta$. We have

$$\begin{aligned}\|\theta(\bar{u}_j - \bar{u}_j^\lambda)\|_{L^p_{2b,1}(R_+^{n+1} \times (0,T))} &\leqslant \|L[\theta(\bar{u}_j - \bar{u}_j^\lambda)]\|_{L^p(R_+^{n+1} \times (0,T))} \\ &\leqslant C \sum_{(|\alpha| + l \leqslant 2b} \|\theta^{(l)} D_{\mu j}^\alpha T_{\mu j}(\Phi - \Phi)\|_{L^p(R_+^{n+1} \times (0,T)} + \\ &+ \|\theta L T_{\mu j}(\Phi - \Phi^\lambda)\|_{L^p(R_+^{n+1} \times (0,T))} + \\ &+ C \| \sum_{l > 0} \theta^{(l)}(y) D_{x,y}^\alpha V_{(\Phi - \Phi^\lambda)(x,y,t)} \|_{L^p(R_+^{n+1} \times (0,T))}\end{aligned}$$

where $V_{(\Phi - \Phi)^\lambda}$ denotes the volume potential in (*) corresponding to $\Phi - \Phi^\lambda$.

A slight modification of the arguments in § 18 shows that these terms are majorized by $C\|\Phi - \Phi^\lambda\|_{L^p(S_T)}$: Therefore $\bar{u}_j^\lambda \to \bar{\bar{u}}_j$ in $\mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$. Also, by Lemmas 1 and 2, § 8, both

$$\|T_{\mu j}(\Phi - \Phi^\lambda)\|_{L^p(R_+^{n+1} \times (0,T))} \quad \text{and} \quad \|y^{b+1-\gamma} L T_{\mu j}(\Phi - \Phi^\lambda)\|_{L^p(R_+^{n+1} \times (0,T))}$$

tend to zero as $\lambda \to 0$. This shows that

$$\bar{\bar{u}}_j \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T)) \quad \text{for every } \delta > 0\,,$$

and admits the representation (*) with $\Phi \in L^p(S_T)$. From this we see that $\bar{\bar{u}}_j \equiv u_j$ and $Lu_j = 0$ for $y > 0$.

This concludes the proof of Theorem, § 17.

§ 20. – It is known (cf. [3]) that

$$\Lambda^{b-1+\mu-k} D_y^k T_{\mu j}\Phi(x, y, t)$$

$$= \int_0^t \int_{R^n} \Lambda^{2b-1-(k+j)} D_y^{k+j} F(z, 0, s; x - z, y, v, t - s)\, \Phi(z, s)\, dz\, ds$$

converges in $L^p(S_T)$ as $y \to 0^+$, to a limit of the form $(S_{k,j} + J_{k,j})\Phi$, where $(S_{k,j})$ is a matrix of parabolic singular integral operators whose matrix of symbols admits an inverse $[\sigma(S_{k,j})(z, s; x, t)]^{-1} = (\sigma(S^*_{k,j})(z, s; x, t))$ provided $(x, t) \neq (0, 0)$, $S^*_{k,j}$ being a parabolic singular integral operator such that $(S_{kj}) \cdot (S^*_{kj}) = I + (J^0_{kj})$, with $J^0_{kj} \in \mathfrak{J}(S_T)$ (cf. [4]). $J_{k,j}$ belongs to $\mathfrak{J}(S_T)$, and is a limit as $y \to 0$ of a series of commutators in the variables (x, y, t) which belong to $\mathfrak{J}(S_T)$ uniformly in y [3].

From the estimates for the volume potential (lemma A, § 10) and the estimates for $y^{b+1-\gamma}(I - J)^{-1}(\cdot)$ and $y^{b+1-\gamma} L T_{\mu j} \Phi$ (§ 15 and Lemma 2, § 8), it follows that for $\beta < \mu$, with $\beta < \gamma - 1/p < \mu$,

$$\Lambda^{b-1+\beta-k} D_y^k \int_0^t \iint_{R_+^{1+u}} K(z, v, s; x - z, y, v, t - s)(I - J)^{-1}(L T_{\mu j} \Phi)(z, v, s)\, dz\, dv\, ds \to 0$$

in $L^p(S_T)$ as $y \to 0^+$.

Also, for $f \in L^p(R_+^{n+1} \times (0, T))$ (Theorem § 16, (ii)),

$$\Lambda^{b-1+\beta-k} D_y^k \int_0^t \iint_{R_+^{n+1}} K(z, v, s; x - z, y, v, t - s)(I - J)^{-1} f(z, v, s)\, dz\, dv\, ds \to 0$$

in $L^p(S_T)$ as $y \to 0^+$.

We observe now that $(J_{kj}) \cdot (S^*_{kj})$ (the dot indicates matrix multiplication) is a matrix of $\mathfrak{J}$-operators, due to the fact that if $J \in \mathfrak{J}(S_T)$, and S is a singular integral operator on S_T, $JS \in \mathfrak{J}(S_T)$.

Therefore $I + [(J^0_{kj}) + (J_{kj}) \cdot (S^*_{kj})]$ has an inverse.

Set

$$(\mathfrak{U}_{kj}) = (S^*_{kj}) \cdot [I + (J^0_{kj}) + (J_{kj}) \cdot (S^*_{kj})]^{-1}.$$

THEOREM. Let there be given $f \in L^p(R_+^{n+1} \times (0, T))$, and for $k = 0, \ldots, b-1$, functions $w^k \in L^p(S_T)$ such that $\Lambda^{b-1-k+\mu} w^k \in L^p(S_T)$ for some μ, $0 < \mu < 1$.

Then there exists a function $u \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$ such that

(i) $Lu = f$ in $R_+^{n+1} \times (0, T)$),

(ii) $\|\Lambda^{b-1-k+\beta}(D_y^k u(\cdot, y, \cdot) - w^k(\cdot, \cdot))\|_{L^p(S_T)} \to 0$ as $y \to 0^+$, for every β, $0 \leqslant \beta < \mu$,

(iii) For $0 \leqslant \beta < \gamma - 1/p < \mu$,

$$\sum_{|\alpha| \leqslant b-1} \|D^{\alpha}_{x,y} \Lambda^{\beta} u(\cdot, y, \cdot)\|_{L^p(S_T)} +$$

$$+ \sum_{k=0}^{b-1} \|\Lambda^{b-1+\beta-k} D^k_y u(\cdot, y, \cdot)\|_{L^p(S_T)} +$$

$$+ \sum_{b \leqslant |\alpha| \leqslant 2b-1} \|d_p(y, \cdot)^{|\alpha|-(b-1)} D^{\alpha}_{x,y} \Lambda^{\beta} u(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant$$

$$\leqslant C \sum_{k=0}^{b-1} \|\Lambda^{b-1+\mu-k} w^k\|_{L^p(S_T)} + C(T) \|d_p^{b+1-\gamma_r} f\|_{L^p(R^{n+1}_+ \times (0,T))}$$

(d_p can be replaced with y or $t^{\gamma_p/2b}$).

PROOF. We set

$$g_{\mu l}(x, y, t) = \sum_{j=0}^{b-1} (\mathfrak{V}_{lj} \Lambda^{b-1+\mu-j} w)(x, y, t)$$

and define

$$u(x, y, t) = \sum_{l=0}^{b-1} (T_{\mu l} g_{\mu l})(x, y, t) +$$

$$+ \sum_{l=0}^{b-1} \int_0^t \iiint_{R^{n+1}_+} K(z, v, s; x-z, y, v, t-s) \cdot (I-J)^{-1} (L T_{\mu l} g_{\mu l})(z, v, s)\, dz\, dv\, ds -$$

$$- \int_0^t \iiint_{R^{n+1}_+} K(z, v, s; x-z, y, v, t-s) (I-J)^{-1} f(z, v, s)\, dz\, dv\, ds\,.$$

The fact that $u \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$, and Part (i) follow from the theorems, §§ 16 and 17. To prove (ii), it is enough to consider the terms $\sum_{l=0}^{b-1} (T_{\mu l} g_{\mu l})(x, y, t)$ and to recall the definition of $g_{\mu l}$ and the fact that $\Lambda^{\beta-\mu}$ is an integrable function.

§ 21. – THEOREM. Suppose $u \in \mathring{L}^p_{2b,1}(R^{n+1}_+ \times (0, T))$, $1 < p < \infty$. Then $u(x; y, t)$ admits the representation $u = u_1 + u_2$, where

$$u_1(x, y, t) =$$

$$= \sum_{l=0}^{b-1} \{(T_{\mu l} g_{\mu l})(x, y, t) + \int_0^t \iiint_{R^{n+1}_+} K(z, v, s; x-z, y, v, t-s)(I-J)^{-1} (L T_{\mu l} g)_{\mu l} dz\, dv\, ds\},$$

$$u_2(x, y, t) =$$

$$= -\int_0^t \iint_{R_+^{n+1}} K(z, v, s; x-z, y, v, t-s)(I-J)^{-1}(Lu)(z, v, s)\, dz\, dv\, ds\,,$$

with

$$g_{\mu l}(x, y, t) = \sum_{j=0}^{b-1} \big(\mathfrak{V}_{lj} \Lambda^{b-1+\mu-j} D_y^j u(\cdot, 0, \cdot)\big)(x, y, t)\,, \qquad \text{(cf. § 20)}\,.$$

Furthermore, for $0 \leqslant \beta < \gamma - 1/p < \mu$,

$$\sum_{k=}^{b-1} \|\Lambda^{b-1+\beta-k} D_y^k u(\cdot, y, \cdot)\|_{L^p(S_T)} + \sum_{|\alpha| \leqslant b-1} \|D_{xy}^\alpha \Lambda^\beta u(\cdot, y, \cdot)\|_{L(S_T)} +$$

$$+ \sum_{b \leqslant |\alpha| \leqslant 2b-1} \|d_p(y, \cdot)^{|\alpha|-(b-1)} D_{x,y}^\alpha \Lambda^\beta u(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant$$

$$\leqslant C \sum_{=0}^{b-1} \|\Lambda^{b-1+\mu-k} D_y^k u(\cdot, 0, \cdot)\|_{L^p(S_T)} + C(T) \cdot \|d_p^{b+1-\gamma}(Lu)\|_{L^p(R_+^{n+1} \times (0,T))}\,,$$

where $C(T) \to 0$ as $T \to 0$, d_p can be replaced by y or $t^{\gamma_p/2b}$, and C, $C(T)$ depend only on the parameter of parabolicity π and on the $\max\limits_{|\alpha|=2b} \sup |a_\alpha|$.

Proof. We assume first $u \in C_0^\infty\big(R_+^{n+1} \times (0, \infty)\big)$, and let $u^\lambda = u_1^\lambda + u_2^\lambda$, u_1^λ and u_2^λ being the terms in the decomposition above, with

$$T_{\mu l}^\lambda,\ g_{\mu l}^\lambda,\ \mathfrak{V}_{lj}^\lambda = \big((S_{lj*}^\lambda) \cdot [I + (J_{lj}^{0\lambda}) + (J_{lj}^\lambda) \cdot (S_{lj*}^\lambda)]^{-1}\big)_{lj}$$

being the expressions corresponding to $F^\lambda(z, 0, s; x, y, t)$ (cf. §§ 13 and 20).

We observe that

$$g_{\mu l}^\lambda(x, y, t) = \sum_{j=0}^{b-1} \{\mathfrak{V}_{lj}^\lambda \Lambda^{b-1+\mu-j} D_y u(\cdot, 0, \cdot)\}(x, y, t)$$

vanishes near $t = 0$ and has all derivatives in $L^p\big(R_+^{n+1} \times (0, T)\big)$.

By an argument similar to that in § 17, we see that

$$T_{\mu l}^\lambda g_{\mu l}^\lambda \in \mathring{L}_{2b,1}^p\big(R_+^{n+1} \times (0, T)\big)\,.$$

Hence $u_1^\lambda \in \mathring{L}_{2b,1}^p\big(R_+^{n+1} \times (0, T)\big)$. u_2^λ belongs to $\mathring{L}_{2b,1}^p\big(R_+^{n+1} \times (0, T)\big)$ by Theorem, § 16 ($Lu \in L^p$).

Therefore $u^\lambda \in \mathring{L}_{2b,1}^p\big(R_+^{n+1} \times (0, T)\big)$, and it is clear that $Lu^\lambda \equiv Lu$ for $y > 0$.

Also $\Lambda^{b-1+\mu-k} D_y^k u^\lambda$ tends to $\Lambda^{b-1+\mu-k} D_y^k u(\cdot, 0, \cdot)$ in $L^p(S^T)$ as $y \to 0^+$. On the other hand, $D_y^k u^\lambda$ and $D_y^k u$ converge in $L^p(S_T)$ as $y \to 0^+$, being func-

tions in $\mathring{L}^p_{2b,1}(R^{n+1}_+ \times (0, T))$. Hence we must have $D^k_y u^\lambda(\cdot, 0, \cdot) = D^k_y u(\cdot, 0, \cdot)$, and we see that $u - u^\lambda$ satisfies a homogeneous initial-boundary value problem with homogeneous data. By [3] it follows that

$$u \equiv u^\lambda \equiv u^\lambda_1 + u^\lambda_2 .$$

Now as $\lambda \to 0$, $g^\lambda_{\mu l} \to g_{\mu l}$ in $L^p(S_T)$ (it would be enough to show that $S^{\lambda^*} \to S^*$, $J^\lambda \to J$ as operators on $L^p(S_T) \times \ldots \times L^p(S_T)$, $(b-1)$ times; we observe that

$$\|J^\lambda_{lh}(X_{(a,a+\varepsilon)} f)\|_{L^p(R^n \times (a,a+\varepsilon))} \leqslant \omega(\varepsilon) \|f\|_{L^p(R^n \times (a,a+\varepsilon))} ,$$

ω in dependent of a and λ, cf. definition of J_{lh} in [3]).

Thus we have the representation $u = u_1 + u_2$ for $u \in C^\infty_0(R^{n+1} \times (0, \infty))$. Any function in $\mathring{L}^p_{2b,1}(R^{n+1}_+ \times (0, T))$ being a limit of $C^\infty_0(R^{n+1} \times (0, \infty))$-functions in the $L^p_{2b,1}$ sense, the general result follows by a density argument, recalling the obvious convergence in L^p of sequences like $(g_{\mu l})_\nu$ and Lu_ν, to their corresponding expressions for $u \in L^p_{2b,1}$, and the fact that

$$\|y^{b+1-\gamma} L T_{\mu l}((g_{\mu l})_\nu - g_{\mu l})\|_{L^p(R^n \times (0,T))} \to 0 .$$

The estimate is an immediate consequence.

§ 22. – Using the representation above, we can prove the uniqueness of the solution to the problem (cf. Theorem, § 20): $u \in L^p_{2b,1}(R^n \times (\delta, \infty) \times \times (0, T))$ for every $\delta > 0$,

$$Lu = f \in L^p(R^{n+1}_+ \times (0, T)) \quad \text{for } y > 0 ,$$

$$\|\Lambda^{b-1+\beta-k}(D^k_y u(\cdot, y, \cdot) - w^k(\cdot, \cdot))\|_{L^p(S_T)} \to 0 \quad \text{as } y \to 0^+$$

for every β, $0 \leqslant \beta < \mu < 1$, and $k = 0, \ldots, b-1$.

Here $\Lambda^{b-1+\mu-k} w^k \in L^p(S_T)$.

THEOREM. If

(i) $u(x, y, t) \in \mathring{L}^p_{2b,1}(R^n \times (\delta, \infty) \times (0, T))$ for every $\delta > 0$,

(ii) for $k = 0, \ldots, b-1$, and some μ, $0 < \mu < 1$,

$$\|\Lambda^{b-1+\mu-k} D^k_y u(\cdot, y, \cdot)\|_{L^p(S_T)} \to 0 \quad \text{as } y \to 0^+ ,$$

and

(iii) $Lu = 0$ for $y > 0$,

Then $u \equiv 0$.

PROOF. We observe that for $\delta > 0$,

$$u(x, y + \delta, t) \in \mathring{L}^p_{2b,1}\big(R^{n+1}_+ \times (0, T)\big)$$

is a solution of

$$L^\delta g \equiv \sum_{|\alpha| \leqslant 2b} a_\alpha(x, y + \delta, t) D^\alpha_{x,y} g + D_t g = 0 \quad \text{for } y > 0 .$$

By the estimate of previous theorem, with $\beta = 0$, we have

$$\|u(\cdot, y + \delta, \cdot)\|_{L^p(S_T)} \leqslant C \sum_{k=0}^{b-1} \|\Lambda^{b-1+\mu-k} D^k_y u(\cdot, \delta, \cdot)\|_{L^p(S_T)} .$$

Hence,

$$\|u(\cdot, y, \cdot)\|_{L^p(S_T)} \leqslant \varlimsup_{\delta \to 0^+} \|u(\cdot, y + \delta, \cdot)\|_{L^p(S_T)} = 0 \quad \text{for } y > 0 ,$$

proving the theorem (we recall that the constants in the representation theorem depend only on π and $\max\limits_{|\alpha|=2b} \sup |a_\alpha|$).

VII. – An elliptic estimate.

§ 23. – Let now

$$\mathcal{E} \equiv \sum_{|\alpha| \leqslant 2b} a_\alpha(x, y) D^\alpha_{x,y}$$

be an operator in R^{n+1}_+, strongly elliptic in the sense that

$$\operatorname{Re}\Big(\sum_{|\alpha|=2b} a_\alpha(x, y)(i\xi)^\alpha \Big) < -\pi \cdot |\xi^{2b} ,$$

$\pi > 0$ and independent of (x, y), $\xi \in R^{n+1}$, $\xi \neq 0$. We assume each $a_\alpha(x, y)$ to be bounded and measurable, and for $|\alpha| = 2b$, uniformly continuous in $\overline{R^{n+1}_+}$.

For $1 < p < \infty$ we define $\bar{d}_p(y) = \min(y, T^{\gamma_p/2b})$, T being a constant.

For B any real number, we will denote by $G_{-B}(x)$ the Bessel potential, defined by

$$\mathcal{F}(G_{-B}f)(x) = (1+|x|^2)^{B/2}\cdot\mathcal{F}(f)(x)\,.$$

Now set $L \equiv \mathcal{E} - D_t$ and assume $u(x,y)\in L^p_{2b}(R^{n+1}_+)$.
Clearly $t\cdot u(x,y)\in \mathring{L}^p_{2b,1}(R^{n+1}_+\times(0,T))$ and we have (cf. § 21)

$$\begin{aligned}\sum_{k=0}^{b-1}\|D_y^k\Lambda^{b-1+\beta-k}(s\cdot u)(\cdot,y,\cdot)\|_{L^p(S_T)} &+ \sum_{|\alpha|\leqslant b-1}\|\Lambda^\beta(s\cdot D^\alpha_{x,y}u)(\cdot,y,\cdot)\|_{L^p(S_T)} + \\ &+ \sum_{b\leqslant|\alpha|\leqslant 2b-1}\|(d_p(y,\cdot))^{|\alpha|-(b-1)}\Lambda^\beta(s\cdot D^\alpha_{x,y}u)(\cdot,y,\cdot)\|_{L^p(S_T)} \\ &\leqslant C\sum_{k=0}^{b-1}\|D_y^k\Lambda^{b-1+\mu-k}(s\cdot u)(\cdot,0,\cdot)\|_{L^p(S_T)} + \\ &+ C(T)\|(d_p(y,t))^{b+1-\gamma}u(x,y)\|_{L^p(R^{n+1}_+\times(0,T))} + \\ &+ C(T)\|(d_p(y,t))^{b+1-\gamma}tLu(x,y)\|_{L^p(R^{n+1}_+\times(0,T))}\quad(*)\,,\end{aligned}$$

where

$$0\leqslant\beta<\gamma<1/p<\mu\,,\qquad d_p(y,t)=\min(y,t^{\gamma_p/2b})\ \ (\S\,5)\,,$$

and $C(T)\to 0$ as $T\to 0^+$.

THEOREM. If $\mathcal{E}$ is a strongly elliptic operator in R^{n+1}_+, and $(x,y)\in L^p_{2b}(R^{n+1}_+)$, then, with $0\leqslant\beta<\gamma-1/p<\mu<1$,

$$\begin{aligned}\sum_{k=0}^{b-1}\|G_{-(b-1+\beta-k)}(D_y^k u)(\cdot,y)\|_{L^p(R^n)} &+ \sum_{|\alpha|\leqslant b-1}\|G_{-\beta}(D^\alpha_{x,y}u)(\cdot,y\|_{L^p(R^n)} + \\ &+ \sum_{b\leqslant|\alpha|\leqslant 2b-1}(\bar d_p(y))^{|\alpha|-(b-1)}\|G_{-\beta}(D^{x,y}_\alpha u)(\cdot,y)\|_{L^p(R^n)}\leqslant \\ &\leqslant C\sum_{k=0}^{b-1}\|G_{-(b-1+\beta-k)}(D_y^k u)(\cdot,0)\|_{L^p(R^n)} + C\|(\bar d_p(y))^{b+1-\gamma}\cdot \\ &\cdot\mathcal{E}u(x,y)\|_{L^p(R^{n+1}_+\times(0,T))} + C\|u\|_{L^p(R^{n+1}_+\times(0,T))}\,.\end{aligned}$$

PROOF. Consider first the following inequalities (see [5] for the first two)

$$\text{(i)}\quad \frac{1}{C}(\bar d(y))^{|\alpha|}T^{1/p}\leqslant\left(\int_\tau^T(d(y,t))^{|\alpha|p}\,dt\right)^{1/p}\leqslant C(\bar d_p(y))^{|\alpha|}\,T^{1/p}$$

where $\tau = 0, T/2$, and $|\alpha|$ can be replaced with $b + 1 - \gamma$;

(ii) $\|\Lambda^{-k} f\|_{L^p(S_T)} \leqslant C \cdot \|G_k f\|_{L^p(R^n)}$;

(iii) $\|(d_p(y,t))^{b+1-\gamma} \cdot t \cdot \mathcal{E}u(x,y)\|_{L^p(R_+^{n+1}\times(0,T))} \leqslant CT^{1+1/p} \|(\bar{d}_p(y))^{b+1-\gamma} \cdot \mathcal{E}u(x,y)\|_{L^p(R_+^{n+1})}$;

(iv) $\|(d_p(y,t))^{b+1-\gamma} u(x,y)\|_{L^p(R_+^{n+1}\times(0,T))} \leqslant CT^{(\gamma_p/2b)(b+1-\gamma)+1/p} \cdot \|u(x,y)\|_{L^p(R_+^{n+1})}$;

(v) for $0 < B < 2b$, $\|\Lambda^\beta (sf(z))\|_{L^p(S_T)} \leqslant C \|G_{-B} f\|_{L^p(R^n)}$.

(i)-(v) show that the right hand side of the parabolic estimate (*) is majorized by the right hand side of the elliptic one. It is clear that the proof will be completed by the following result, whose proof follows the lines of [5]. Appendix

LEMMA. Let $f(x) \in C_0^\infty(R^n)$. Then for $1 < p < \infty$, $0 \leqslant B \leqslant 2b$,

$$\|G_{-B} f\|_{L^p(R^n)} \leqslant C_T \|\Lambda^B (s \cdot f(z))\|_{L^p(R^n \times (T/2,T))}.$$

REMARK. If u is assumed to have support contained in $\{(x,y): |x|^2 + y^2 \leqslant r^2, y \geqslant 0\}$, the term $\|u\|_{L^p(R_+^{n+1})}$ in the estimate may be replaced by $\|u\|_{L^1(R_+^{n+1})}$, with a change in the constants.

VIII. – The main results in a general domain.

§ 24. – We now consider the Initial-Dirichlet boundary-value problem with initial data zero, for a parabolic equation where the cylinder is defined as the product of a domain in R^{n+1} with the time interval $(0, T)$. P shall denote a point inside that domain, $D^\alpha = D_P^\alpha$ a spatial derivative of order $|\alpha|$, Q a point on the boundary. The differential operator, L, is assumed to satisfy Petrovski's condition and the conditions on the coefficients stated in § 5 with (x, y, t) replaced by (P, t).

From here on, Ω will denote a bounded, smooth domain in R^{n+1}. By this we mean

(i) There exists a finite number of functions f_i having continuous and bounded derivatives up to order $2b + 2$, and each mapping the disc $\{(x,0): |x|^2 < r_i^2, r_i > 0\} \subset R^{n+1}$ into $\partial\Omega$ in a $1-1$ manner, such that every point $Q \in \partial\Omega$ can be written $Q = f_i(x, 0)$, $|x|^2 < r_i^2$, for some i;

(ii) If N_Q denotes the unit inner normal to $\partial\Omega$ at Q, and $\Omega_\delta = \{P \in \Omega: \operatorname{dist}(P, \partial\Omega) > \delta, \delta \geqslant 0\}$, then there is a number $\delta_0 > 0$ such that for each i

the function

$$(x, y) \to f_i(x, y) = Q + y \cdot N_Q, \qquad Q = f_i(x, 0) \in \partial\Omega$$

maps the set $\{(x, y): |x|^2 < r_i^2,\ 0 \leqslant y \leqslant 4\delta_0\}$ into $\bar{\Omega} - \Omega_{4\delta_0}$ in a $1-1$ manner. We assume further that every point $P \in \bar{\Omega} - \Omega_{4\delta_0}$ can be uniquely written as $P = Q + y \cdot N_Q$, $Q \in \partial\Omega$.

We consider the finite covering of the set $\bar{\Omega} - \Omega_{\delta_0}$ by the sets U_i, image of

$$\{(x, y): |x|^2 < r_i^2,\ 2\delta_0 > y > \varkappa,\ \varkappa \text{ a suitable number } < 0\}$$

under $f_i(x, y) = f_i(x, 0) + y \cdot N_{f_i(x,0)}$. We let $\{\varphi_i\}$ denote a fixed C_0^∞ partition of unity subordinate to $\{U_i\}$, and we denote by $\{\zeta_i\}$ a family of functions such that $\zeta_i \in C_0^\infty(U_i)$ and $\zeta_i \equiv 1$ in a neighborhood of the support of φ_i. We point out here that when $0 < \delta < \delta_0$, for the domain Ω_δ we can associate the sets $U_{\delta i}$, and the families $\{\varphi_{\delta i}\}$, $\{\zeta_{\delta i}\}$ obtained from the ones above by the transformation $P \to P + \delta N_Q$, where $P = Q + yN_Q \in U_i$. It follows that the derivatives of $\varphi_{\delta i}$, $\zeta_{\delta i}$ can be bounded uniformly on δ, $0 < \delta < \delta_0$.

If $u(P)$ is a function defined in $\bar{\Omega}$ we will set, for simplicity, $u_i^{\sim}(x, y) = u \circ f_i(x, y)$. For the functions φ_i, ζ_i, we will set, e.g., $(\varphi_i) = \varphi_i^{\sim}$ when there is no confusion. We will also write $u_i^{\sim}(x, y, t)$ for $u(f_i(x, y), t)$. We observe that $(D_{N_Q} u_i^{\sim})(x, y) = D_y(u_i^{\sim})(x, y)$.

The L^p-norm of a function $u(Q, t)$ defined on $\partial\Omega \times (0, T)$ is equivalent to $\sum_i \|\varphi_i u\|$, or $\sum_i \|\zeta_i u\|$, which are computed as integrals over $R^n \times (0, T) = S_T$.

We introduce now the operators Λ on $\partial\Omega$.

DEFINITION. If $u(Q, t) \in L^p(\partial\Omega \times (0, T))$, $1 < p < \infty$, and $0 \leqslant \beta \leqslant 2b$,

$$(\Lambda^{-\beta} u)(Q, t) = \sum_i \zeta_i(Q) \Lambda^{-\beta}[\varphi_i^{\sim} \cdot u_i^{\sim}](fi^{-1}(Q), t)\,,$$

where $\Lambda^{-\beta}[\cdot]$ is the operator already defined on S_T.

DEFINITION. For functions $u(Q, t) \in C^\infty(\partial\Omega \times (0, T))$ which are identically zero for t near zero,

$$(\Lambda^{\beta} u)(Q, t) = \sum_i \zeta_i(Q) \Lambda^{\beta}[\varphi_i^{\sim} \cdot u_i^{\sim}](fi^{-1}(Q), t)\,.$$

It can be shown that $\Lambda^\beta \Lambda^{-\beta}$ is not in general the identity on C^∞ functions that vanish near $t = 0$, but it is extendible to an invertible operator on $L^p(\partial\Omega \times (0, T))$, $1 < p < \infty$.

The Bessel potentials are defined in a similar way.

DEFINITION. If $u(Q) \in C^\infty(\partial\Omega)$,

$$(G_{-\beta}u)(Q) = \sum_i \zeta_i(Q)\, G_{-\beta}[\varphi_i^\sim \cdot u_i^\sim](fi^{-1}(Q), t)\,,$$

where $\mathcal{F}_x(G_{-\beta}[\varphi_i^\sim \cdot u_i^\sim]) = (|x|^2+1)^{b/2} \cdot \mathcal{F}_x[\varphi_i^\sim \cdot u_i^\sim]$, $x \in R^n$.

In the next paragraphs we follow the method used in [5], article (4.1), for the corresponding problem with initial data zero.

§ 25. – THEOREM. If $u \in \mathring{L}^p_{2b,1}(\Omega \times (0, T))$, $1 < p < \infty$, $0 \leqslant \beta < \mu < 1$, and $Lu = 0$ in $\Omega \times (0, T)$, then

$$\sup_{y > \delta_0} \Big\{ \sum_{|\alpha| \leqslant 2b-1} y^{[|\alpha|-(b-1)]} \| D^\alpha (\Lambda^\beta u)(Q + yN_Q, t) \|_{L^p(\partial\Omega \times (0,T))} + \\ + \sum_{k=0}^{b-1} \| \Lambda^{b-1+\beta-k} [D^k_{N_Q} u](Q + yN_Q, t) \|_{L^p(\partial\Omega \times (0,T))} \Big\} \\ \leqslant C \cdot \sum_{k=0}^{b-1} \| \Lambda^{b-1+\mu-k} [D^k_{N_Q} u](Q, t) \|_{L^p(\partial\Omega \times (0,T))}\,.$$

(Here and in the following, the expression $y^{[|\alpha|-(b-1)]}$ is meant to be replaced by 1 for $|\alpha| \leqslant b-1$).

PROOF. We shall sketch the proof of this result, which proceeds first for small T (this §), and then in the general case (next §).

By applying the definition of Λ^β (§ 24), dropping continuous functions of compact support, and introducing a new constant C_{δ_0}, we can see that the left hand side $(LHS^\beta_T(u; \Omega))$ of the estimate above is less than or equal to

$$\|u\|^\sim \equiv C_{\delta_0} \sum_i \sup_{y < 2\delta_0} \Big\{ \sum_{|\alpha| \leqslant 2b-1} y^{[|\alpha|-(b-1)]} \| D^\alpha_{xy} \Lambda^\beta (\varphi_i^\sim \cdot u_i^\sim)(\cdot, y, \cdot) \|_{L^p(S_T)} + \\ + \sum_{k=0}^{b-1} \| \Lambda^{b-1+\beta-k} D^k_y (\varphi_i^\sim \cdot u_i^\sim)(\cdot, y, \cdot) \|_{L^p(S_T)} + \\ + \sum_{k=1}^{b-1} \sum_{m=1}^{k} \| \Lambda^{b-1+\beta-k} (D^m_y \varphi_i^\sim \cdot D^{k-m}_y u_i^\sim)(\cdot, y, \cdot) \|_{L^p(S_T)} \Big\}\,.$$

We now

(i) Define a parabolic operator $L_i^\sim$ on $R^{n+1}_+ \times (0, T)$, with coefficients bounded and measurable, and those of the leading terms, uniformly continuous in $\overline{R^{n+1}_+} \times [0, T]$, that satisfies

$$L_i^\sim (u_i^\sim)(x, y, t) = L(u)(f_i(x, y), t) \quad \text{for } f_i(x, y) \in U_i\,;$$

(ii) Apply the estimates for $L_i^{\sim}$ (§ 20) to the first two terms above;

(iii) Omit somewhat lengthy considerations to obtain, with $0<\delta<\delta_0$ and $C(T)\to 0$ as $T\to 0^+$

$$
\begin{aligned}
LHS_T^{\beta}(u;\Omega)\leqslant\|u\|^{\sim}\leqslant C\sum_{k=0}^{b-1}\|\Lambda^{b-1+\mu-k}(D_{N_Q}^k u)(Q,t)\|_{L^p(\partial\Omega\times(0,T))}+\\
+C_{\delta,\delta_0}C(T)\sup_{\delta<y<2\delta_0}\sum_{|\alpha|\leqslant 2b-1}\|(D^\alpha u)(Q+yN_Q,t)\|_{L^p(\partial\Omega\times(0,T))}+\\
+C_{\delta_0}C(T)\delta^{1-\gamma+1/p}\sup_{y<\delta}\sum_{|\alpha|=\leqslant 2b-1}y^{[|\alpha|-(b-1)]}\|(D^\alpha u)(Q+yN_Q,t)\|_{L^p(\partial\Omega\times(0,T))}+\\
+C_{\delta_0}\sum_i\sup_{y<2\delta_0}\sum_{k\geqslant m\geqslant 0}\|\Lambda^{-1+\mu-\beta}\Lambda^{b-1+\beta-(k-m)}(D_y^m\varphi_i^{\sim}D_y^{k-m}u_i^{\sim})(\cdot,y,\cdot)\|_{L^p(S_T)};
\end{aligned}
$$

(iv) Take δ small enough so we can move the third term in (iii) over to the left hand sides. We modify the constants (introducing C_δ in front of $\|u\|^{\sim}$) and fix this δ from now on. And finally

(v) CLAIM. There exists a $T_0>0$ such that the estimate in the Theorem holds for $T\leqslant T_0$.

Clearly, the second term in (iii) (modified as in (iv)) can be bounded with

$$C_{\delta,\delta_0}C(T)\,LHS_T^{\beta}(u;\Omega)\leqslant C_\delta C_{\delta,\delta_0}C(T)\cdot\|u\|^{\sim}.$$

Also, the fourth term in (iii) can be shown to be bounded by

$$C_{\beta,\mu}(T)\,C_{\delta,\delta_0}C_\delta\|u\|^{\sim},\qquad\text{with } C_{\beta,\mu}(T)\to 0 \text{ as } T\to 0^+.$$

Therefore, if T is selected so that

$$C_{\delta_0}C_{\beta,\mu}(T)+C_{\delta,\delta_0}C(T)<1,$$

then

$$(\#)\qquad LHS_T^{\beta}(u;\Omega)\leqslant C_{\delta,\delta_0,\beta,\mu}\sum_{k=1}^{b-1}\|\Lambda^{b-1+\mu-k}(D_{N_Q}^k u)(Q,t)\|_{L^p(\partial\Omega\times(0,T))},$$

i.e. there exists a $T_0>0$, depending on δ, δ_0, β, μ such that the theorem is true for $\partial\Omega\times(0,T)$, $T\leqslant T_0$.

REMARK. For $T\leqslant T_0$, (#) is also true for all the domains $\Omega_{\delta_0^1}$, $0\leqslant\delta_0^1<\delta_0$, with the same constant $C_{\delta,\delta_0,\beta,\mu}$ (which, we recall, depends on the families $\{U_i\}$, $\{\varphi_i\}$, $\{\zeta_i\}$, cf. definitions in § 24).

§ 26. – To prove the theorem for any T, we rewrite the estimates in (iii) with third term deleted, see (iv), as

$$LHS_T^{\beta}(u;\Omega) \leqslant C_{\delta,\delta_0}\Big\{\sum_{k=0}^{b-1}\|\Lambda^{b-1+\mu-k}(D_{N_Q}^k u)\|_{L^p(\partial\Omega\times(0,T))} + \|u\|^{\approx}\Big\},$$

where we have set

$$\|u\|^{\approx} \equiv \sup_{\delta<y<2\delta_0}\sum_{|\alpha|\leqslant 2b-1}\|(D^{\alpha}u)(Q+yN_Q,t)\|_{L^p(\partial\Omega\times(0,T))} +$$

$$+\sum_i \sup_{y<2\delta_0}\sum_{k\geqslant m\geqslant 0}^{b-1}\|\Lambda^{-1+\mu-\beta}\Lambda^{b-1+\beta-(k-m)}(D_y^m\varphi_i^{\sim}D_y^{k-m}u_i^{\sim})(\cdot,y,\cdot)\|_{L^p(S_T)},$$

and observe that the desired result is a consequence of the following lemma, whose proof is also omitted

LEMMA – Given $\varepsilon>0$, there exists a constant C_ε such that for all $u\in \mathring{L}^p_{2b,1}(\Omega\times(0,T))$ that satisfy $Lu=0$ we have

$$\|u\|^{\approx} \leqslant \varepsilon LHS_T^{\beta}(u;\Omega) + C_\varepsilon\sum_{k=0}^{b-1}\|\Lambda^{b-1+\mu-k}(D_{N_Q}^k u)\|_{L^p(\partial\Omega\times(0,T))}.$$

REMARK. The a priori estimate in Theorem, § 25, holds for all domains $\Omega_{\delta_0^1}$, with $\delta_0^1 \ll \delta_0$. The constant in the theorem is independent of δ_0^1 (cf. also Remark to § 25).

§ 27. – We are now in a position to prove the main results.

THEOREM 1. Let $w^k\in L^p(\partial\Omega\times(0,T))$, $k=0,\ldots,b-1$, be such that $\Lambda^{b-1+\mu-k}w^k\in L^p(\partial\Omega\times(0,T))$ for some μ, $0<\mu<1$. Then there exists a solution to the problem

(i) $u(P,t)\in \mathring{L}^p_{2b,1}(\Omega^*\times(0,T))$ for every subdomain Ω^* such that $\bar{\Omega}^*\subset\Omega$,

(ii) $Lu=0$ in $\Omega\times(0,T)$, and

(iii) for every β, $0\leqslant\beta<\mu$,

$$\lim_{y\to 0^+}\|\Lambda^{b-1+\beta-k}[D_{N_Q}^k u(Q+yN_Q,t)-w^k(Q,t)]\|_{L^p(\partial\Omega\times(0,T))}=0.$$

PROOF. Let $w_j^k\in C_0^{\infty}(\partial\Omega\times(0,\infty))$, such that $w_j^k\to w^k$ and $\Lambda^{b-1+\mu-k}w_j^k\to$ $\to\Lambda^{b-1+\mu-k}w^k$, in $L^p(\partial\Omega\times(0,T))$ as $j\to\infty$, $0\leqslant 1\leqslant b-1$. It is known [10] that we can find $u_j(P,t)\in \mathring{L}^p_{2b,1}(\Omega\times(0,T))$ satisfying (ii) and (iii) with w^k replaced by w^k.

Now if $\theta(P) \in C_0^\infty(\Omega)$, $\theta \equiv 1$ in Ω_δ, then

$$\|\theta u_j\|_{L^p_{2b,1}(\Omega\times(0,T))} \leqslant C_\delta \sum_{\substack{|\beta|>0 \\ |\beta|+|\alpha|\leqslant 2b}} \|(D^\beta\theta)(D^\alpha u_j)\|_{L^p(\Omega\times(0,T))}\,.$$

This, together with the theorem, § 25, imply that if $\bar{\Omega}^* \subset \Omega$,

$$\|u_j - u_l\|_{L^p_{2b,1}(\Omega^*\times(0,T))} \leqslant C_{\Omega^*} \sum_{k=0}^{b-1} \|\Lambda^{b-1+\mu-k}(w_j^k - w_l^k)\|_{L^p(\partial\Omega\times(0,T))}\,.$$

Hence $\{u_i\}$ is a Cauchy sequence in $L^p_{2b,1}(\Omega^*\times(0,\,T))$, for every subdomain Ω^* with $\bar{\Omega}^* \subset \Omega$. Let $u(P,\,t)$ denote the limit of $\{u_j\}$. Clearly $u(P,\,t)$ satisfies (i) and (ii).

To prove (iii) we observe that

$$\sup_{y<\delta_0} \sum_{k=0}^{b-1} \|\Lambda^{b-1+\beta-k} D^k_{N_Q}(u_j - u_l)(Q + yN_Q,\,t)\|_{L^p(\partial\Omega\times(0,T))} \leqslant$$
$$\leqslant C \sum_{k=0}^{b-1} \|\Lambda^{b-1+\mu-k}[w_j^k - w_l^k]\|_{L^p(\partial\Omega\times(0,T))}\,,$$

which implies

$$\sup_{y<\delta_0} \sum_{k=0}^{b-1} \|\Lambda^{b-1+\beta-k} D^k_{N_Q}(u_j - u)(Q + yN_Q,\,t)\|_{L^p(\partial\Omega\times(0,T))} \leqslant$$
$$\leqslant C\cdot \lim_{l\to\infty} \sum_{k=0}^{b-1} \|\Lambda^{b-1+\mu-k}[w_j^k - w_l^k]\|_{L^p(\partial\Omega\times(0,T))} \to 0 \qquad \text{as } j\to\infty\,.$$

Now we observe that

$$\Lambda^{b-1+\beta-k} D^k_{N_Q} u = \Lambda^{b-1+\beta-k} D^k_{N_Q}(u - u_j) + \Lambda^{b-1+\beta-k}[D^k_{N_Q} u_j]\,,$$

and that

$$\lim_{y\to 0^+} \|\Lambda^{b-1+\beta-k}[D^k_{N_Q}(u - u_j)]\|_{L^p(\partial\Omega\times(\,,T))} \leqslant \sup \|\Lambda^{b-1+\beta-k} D^k_{N_Q}(u - u_j)\|_{L^p(\partial\Omega\times(0,T))}\,,$$

and we see that

$$\lim_{y\to 0^+} \Lambda^{b-1+\beta-k}[D^k_{N_Q} u] = \lim_{j\to\infty} \Lambda^{b-1+\beta-k} w_j^k = \Lambda^{b-1+\beta-k} w^k \qquad \text{in } L^p\big(\partial\Omega\times(0,\,T)\big)\,.$$

THEOREM 2. Let $u(P,\,t)$ be any function such that

(i) $u(P,\,t)$ belongs in $\mathring{L}^p_{2b,1}\big(\Omega^*\times(0,\,T)\big)$ for every subdomain Ω^* such that $\bar{\Omega}^* \subset \Omega$,

(ii) $Lu \equiv 0$ in $\Omega \times (0, T)$,

(iii) for $k = 0, \ldots, b-1$, and for some μ, $0 < \mu < 1$,

$$\varliminf_{y \to 0^+} \|\Lambda^{b-1+\mu-k}[D^k_{N_Q} u](Q + yN_Q, t)\|_{L^p(\partial\Omega \times (0,\chi T))} < \infty .$$

Then, for $0 \leqslant \beta < \mu$,

$$\sup_{y<\delta_0} \Big\{ \sum_{|\alpha| \leqslant 2b-1} y^{[|\alpha|-(b-1)]} \|D^\alpha(\Lambda^\beta u)(Q + yN_Q, t)\|_{L^p(\partial\Omega\times(0,T))} + \\ + \sum_{k=0}^{b-1} \|\Lambda^{b-1+\beta-k}[D^k_{N_Q} u](Q + yN_Q, t)\|_{L^p(\partial\Omega\times(0,T))} \Big\} \leqslant \\ \leqslant C \cdot \sum_{k=0}^{b-1} \varliminf_{y\to 0^+} \|\Lambda^{b-1+\mu-k}[D^k_{N_Q} u](Q + yN_Q, t)\|_{L^p(\partial\Omega\times(0,T))} .$$

COROLLARY. The solution to problem (i)-(iii) in Theorem 1 is unique.

PROOF OF THEOREM 2. We recall that $(\cdot)^{[|\alpha|-(b-1)]} \equiv 1$ for $|\alpha| \leqslant b-1$ (§ 25). We consider any fixed y, $0 < y < \delta_0$, we take $0 < \delta_0^1 \ll y$, and study $\Omega_{\delta_0^1} \times (0, T)$; clearly $u \in \mathring{L}^p_{2b,1}(\Omega_{\delta_0^1} \times (0, T))$ and $Lu = 0$. By the a priori estimate in Theorem, § 25 (cf. also the Remark to § 26), and with $0 \leqslant \beta < \mu$,

$$\sum_{|\alpha| \leqslant 2b-1} (y - \delta_0^1)^{[|\alpha|-(b-1)]} \|D^\alpha(\Lambda^\beta u)(Q + yN_Q, t)\|_{L^p(\partial\Omega\times(0,T))} + \\ + \sum_{k=0}^{b-1} \|\Lambda^{b-1+\beta-k}[D^k_{N_Q} u](Q + yN_Q, t)\|_{L^p(\partial\Omega\times(0,T))} \leqslant \\ \leqslant C_{\beta,\mu,\delta_0} \cdot \sum_{k=0}^{b-1} \|\Lambda^{b-1+\mu-k}[D^k_{N_Q} u](Q + \delta_0^1 N_Q, t)\|_{L^p(\partial\Omega\times(0,T))} .$$

The theorem follows by taking $\varliminf$ on both sides and recalling that C_{β,μ,δ_0} does not depend on δ_0^1 (cf. loc. cit.).

IX. – The elliptic estimate.

§ 28. – As in § 23, we let $\mathcal{E} \equiv \sum_{|\alpha| \leqslant 2b} a_\alpha(P) D^\alpha$ be a strongly elliptic operator on Ω, that is, $\mathcal{E} - D_t$ is parabolic in the sense of Petrovski. Again $\pi > 0$ will denote the parameter of parabolicity of $\mathcal{E} - D_t$ and the a-'s are assumed to be bounded and measurable, and for $|\alpha| = 2b$, uniformly continuous in $\bar{\Omega}$. $G_{-\beta}$ shall denote the Bessel potentials defined in § 24.

As done in §§ 25-27, the expression $(\cdot)^{[|\alpha|-(b-1)]}$ is to be replaced by 1 for $|\alpha| \leqslant b-1$.

THEOREM. If $u \in L^p_{2b}(\Omega)$ and $\mathcal{E}u = 0$, then for $0 \leqslant \beta < \mu$,

$$\sup_{y<\delta_0}\Big\{ \sum_{|\alpha|\leqslant 2b-1} y^{[|\alpha|-(b-1)]} \|D_\alpha[G_{-\beta}u](Q+yN_Q)\|_{L^p(\partial\Omega)} + \\ + \sum_{k=0}^{b-1} \|G_{-(b-1+\beta-k)}[D^k_{N_Q}u](Q+yN_Q)\|_{L^p(\partial\Omega)}\Big\} \\ \leqslant C\Big\{ \sum_{k=0}^{b-1} \|G_{-(b-1+\mu-k)}[D^k_{N_Q}u](Q)\|_{L^p(\partial\Omega)} + \|u\|_{L^1(\Omega)}\Big\}.$$

The proof of this result follows lines analogous to those in the parabolic estimate of § 25. We shall only sketch them.

By application of the definitions of the Bessel potentials to the left hand side $(LHS^\beta(u))$ of the inequality, a bound is obtained to whose terms the estimates of § 23 and Remark, § 23 apply with elliptic operators $\mathcal{E}_i^{\sim}$ defined by

$$\mathcal{E}_i^{\sim}(u_i^{\sim})(x,y) = \mathcal{E}u(f_i(x,y)) \quad \text{for } f_i(x,y) \in U_i\,.$$

Support considerations on the C_0^∞ functions in the definition of $G_{-\beta}$ (§ 24) lead to the estimate

$$LHS^\beta(u) \leqslant C\Big(\sum_{k=0}^{b-1} \|G_{-(b-1+\mu-k)}(D^k_{N_Q}u)(Q)\|_{L^p(\partial\Omega)} + \|u\|_{L^1(R^{n+1}_+)}\Big) + \\ + \|u\|^{\approx} + C_{\delta_0}\delta^{1-\gamma+1/p} \sup_{y<\delta} \sum_{|\alpha|\leqslant 2b-1} y^{[|\alpha|-(b-1)]} \|D^\alpha u(Q+yN_Q)\|_{L^p(\partial\Omega)} \\ + C_{\delta,\delta_0} \sup_{\delta<\gamma<2\delta_0} \sum_{|\alpha|\leqslant 2b-1} \|D^\alpha u(Q+yN_Q)\|_{L^p(\partial\Omega)},$$

where we have set

$$\|u\|^{\approx} \equiv C_{\delta_0} \sum_i \sup_{y<2\delta_0} \sum_{k\geqslant m\geqslant 0}^{b-1} \|G_{1-\mu+\beta} G_{-(b-1+\beta-(k-m))}(D_y^m \varphi_i^{\sim} D_y^{k-m} u_i^{\sim})(y)\|_{L^p(R^n)}.$$

Fixing δ small enough, the third term in the estimate above can be moved over to the left hand side. The last term can be shown to be $\leqslant \varepsilon \cdot C\|\mathcal{E}(\varphi u)\|_{L^p(\Omega)} + C_\varepsilon \|u\|_{L^1(\Omega)}$, where $\varphi \in C_0^\infty(\Omega)$, $\varphi \equiv 1$ on $\Omega' \supset\supset \Omega_\delta$, by using a trace theorem [11], and an estimate in [1], already needed for Remark, § 23.

But $\|\mathcal{E}(\varphi u)\|_{L^p(\Omega)} \leqslant C_{\delta\varphi} LHS^\beta(u)$, so choosing ε small enough we even-

tually get

$$LHS^{\beta}(u) \leqslant C\Big\{ \sum_{k=0}^{b-1} \|G_{-(b-1+\mu-k)}(D_{N_Q}^k u)\|_{L^p(\partial\Omega)} + \|u\|_{L^1(\Omega)} + \|u\|^{\approx} \Big\}.$$

The proof can be completed by proving the following

LEMMA. To every $\varepsilon' > 0$ there is a constant $C_{\varepsilon'}$ such that every $u \in L^p_{2b}(\Omega)$ with $\mathcal{E}u = 0$ satisfies

$$\|u\|^{\approx} \leqslant \varepsilon' LHS^{\beta}(u) + C_{\varepsilon'} \|u\|_{L^1(\Omega)}.$$

REFERENCES

[1] S. AGMON - A. DOUGLIS - L. NIRENBERG, *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions, I*, Communications on Pure and Applied Mathematics, **12**, no. 4 (1959).

[2] E. B. FABES, *Singular integrals and partial differential equations of parabolic type*, Studia Mathematica, **28** (1966).

[3] E. B. FABES - M. JODEIT Jr., *L^p-boundary value problems for parabolic equations*, Bulletin A.M.S., **74**, no. 6 (1968). Also *Singular integrals and boundary value problems for parabolic equations in a half-space*, University of Minnesota.

[4] E. B. FABES - N. M. RIVIÈRE, *Systems of parabolic equations with uniformly continuous coefficients*, Journal d'Analyse Mathématique, **17** (1966).

[5] E. B. FABES - N. M. RIVIÈRE, *L^p-estimates near the boundary for solutions of the Dirichlet problem*, Annali della Scuola Normale Superiore, Pisa, **24**, fasc. III (1970).

[6] E. B. FABES, *Singular integrals and their applications to the Cauchy and Cauchy-Dirichlet problems for parabolic equations*, Notes from a course given at the Istituto Matematico, Università di Ferrara, Italia, 1970.

[7] A. FRIEDMAN, *Partial Differential Equations of Parabolic Type*, Prentice-Hall, Inc., Englewood Cliffs, N. J., 1964.

[8] G. H. HARDY - J. E. LITTLEWOOD - G. PÖLYA, *Inequalities*, Cambridge at the University Press, 1952.

[9] E. M. STEIN, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.

[10] V. A. SOLONNIKOV, *On boundary value problems for linear parabolic systems of differential equations of a general type* (Russian), Trudy Matematicheskogo Instituta Akademii Nauk (Steklov), **83**, Leningrad, 1965.

[11] E. GAGLIARDO, *Proprietà di alcune classi di funzioni di più variabili*, Ricerche di Matematica, **7** (1958).

[12] W. POGORZELSKI, *Étude de la matrice des solutions fondamentales du système parabolique d'équations aux dérivées partielles*, Ricerche di Matematica, **7** (2) (1958).