

ANNALI DELLA  
SCUOLA NORMALE SUPERIORE DI PISA  
*Classe di Scienze*

JANOS PINTZ

SAVERIO SALERNO

**On the comparative theory of primes**

*Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4<sup>e</sup> série*, tome 11,  
n° 2 (1984), p. 245-260

[http://www.numdam.org/item?id=ASNSP\\_1984\\_4\\_11\\_2\\_245\\_0](http://www.numdam.org/item?id=ASNSP_1984_4_11_2_245_0)

© Scuola Normale Superiore, Pisa, 1984, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## On the Comparative Theory of Primes.

JANOS PINTZ - SAVERIO SALERNO (\*)

1. Knapowski and Turàn investigated in a series of papers ([3]) sign changes of the functions (in the case of  $i = 1, 2, 4$ )

$$(1.1) \quad \left\{ \begin{array}{l} \Delta_1(x, q, l_1, l_2) = \sum_{p \leq x} \varepsilon(p, q, l_1, l_2) \\ \Delta_2(x, q, l_1, l_2) = \sum_{n \leq x} \varepsilon(n, q, l_1, l_2) \frac{\Lambda(n)}{\log n} \\ \Delta_3(x, q, l_1, l_2) = \sum_{p \leq x} \varepsilon(p, q, l_1, l_2) \log p \\ \Delta_4(x, q, l_1, l_2) = \sum_{n \leq x} \varepsilon(n, q, l_1, l_2) \Lambda(n) \end{array} \right.$$

where

$$(1.2) \quad \Lambda(n) = \begin{cases} \log p & \text{if } n = p^n \\ 0 & \text{otherwise} \end{cases}$$

$$\varepsilon(n) = \varepsilon_1(n) - \varepsilon_2(n), \quad \varepsilon_i(n) = \begin{cases} 1 & \text{if } n \equiv l_i(q) \\ 0 & \text{otherwise} \end{cases}$$

and (what we shall always assume without mentioning)

$$(1.3) \quad (l_1, q) = (l_2, q) = 1, \quad l_1 \not\equiv l_2 \pmod{q}.$$

For a general modulus  $q$  Knapowski and Turàn needed always the so called Haselgrove condition (H), that the  $L$ -functions have no real non-trivial zeros or, in an explicit formulation, they assumed the existence of

(\*) This work was written whilst the first named author was « visiting professor » at the University of Salerno with the grant of C.N.R.

The author want to express their gratitude to C.N.R. for making this collaboration possible.

Pervenuto alla redazione il 6 Giugno 1983.

a number  $A(q)$ ,  $0 < A(q) < 1$ , such that (for  $s = \sigma + it$ )

$$(1.4) \quad L(s, \chi, q) \neq 0 \quad \text{for} \quad |t| \leq A(q).$$

It is easy to see that in the case of the existence of a real zero, being on the right from all complex zeros of all  $L(s, \chi, q)$  functions, the functions  $\Delta_i(x)$  are of constant sign for suitable pairs  $l_1, l_2$  for  $x > x_0$ .

Thus, at the present stage of analytic number theory, a hypothesis of type (1.4) is necessary in all investigations.

We note that (1.4) has been verified by Spira [6] for all  $q < 25$ . The other assumption, the so called finite Riemann-Piltz conjecture (FR-P),

$$(1.5) \quad L(s, \chi, q) \neq 0 \quad \text{for} \quad \sigma \neq \frac{1}{2}, \quad |t| \leq D = cq^{10},$$

used in many investigations of Knapowski and Turàn, is of more technical nature. The aim of Knapowski and Turàn was to achieve effective results, i.e. explicitly dependent only on  $q$ ,  $A(q)$  (and in given cases on  $D$ ). In the case of  $l_1 = 1$  or  $l_2 = 1$  they were able to find, using only Haselgrove condition, infinitely many sign changes of  $\Delta_i(x)$  for  $1 \leq i < 4$ . (Although they did not treat the case  $i = 3$ , we mention the corresponding results for  $i = 3$  too, if it is possible to obtain it, using the method applied by them for the other cases). Also, they could give a lower estimation of the number  $V_i(Y, q, l_1, l_2)$  of sign changes of  $\Delta_i(x)$  in the interval  $(2, Y)$  and an explicit upper bound for the first sign change (cf. parts I-III of [3]).

For general  $l_1$  and  $l_2$ , besides (H), they had to assume also the finite Riemann-Piltz conjecture and even this led to results only for  $i = 2$  and  $4$  (with other numerical estimates, naturally than in the earlier mentioned case). In the case  $i = 1$  and  $3$  they needed the additional assumption that  $l_1$  and  $l_2$  would be both quadratic residues or both non-residues, besides (FR-P) and (H), (cf. parts V and VI of [3]). In the case of  $i = 4$  they succeeded in showing for general  $l_1$  and  $l_2$  the above mentioned results in a quantitatively much weaker but (as all the earlier mentioned results) also effective form, without supposing (FR-P), i.e. only assuming (H) (cf. part VII of [3]).

We note that the most important open problem of the comparative prime number theory is to assure infinitely many sign changes of  $\Delta_1(x, q, l_1, l_2)$  for all pairs  $l_1, l_2$  with (1.3). This problem seems to be hopeless at present, even supposing the (infinite) Riemann-Piltz conjecture, besides (H) naturally. (One has nearly the same difficulties for  $\Delta_3(x)$ , but this is not so important from an arithmetic point of view).

In this work (assuming (H) and (FR-P) in a slightly weaker form) we shall treat the case  $i = 4$ , for general  $l_1, l_2$ , (and the case  $i = 3$  if  $l_1$  and  $l_2$  have the same quadratic character) improving earlier results of Knapowski and Turàn, which we summarize now (in some cases with slight changes) as follows:

We shall use the notations  $\exp_{v+1}(x) = \exp(\exp_v(x))$ ,  $\exp_1(x) = \exp(x)$ ,  $\log_{v+1} x = \log \log_v x$ ,  $\log_1 x = \log x$ .

**THEOREM A.** Assume (H) and (FR-P) (cf. (1.4)-(1,5)). Then for

$$(1.6) \quad Y > \max \left\{ \exp_2(cq^{20}), \exp_2\left(\frac{c}{A^3(q)}\right) \right\}$$

one has for  $i = 2, 4$

$$(1.7) \quad \max_{Y^{1/2} \leq x \leq Y} \Delta_i(x, q, l_1, l_2) > \sqrt{Y} \exp\left(-44 \frac{\log Y \log_3 Y}{\log_2 Y}\right)$$

for all pairs  $l_1$  and  $l_2$  with (1.3), where, as always in the following, the generic symbol  $c$  replaces an explicitly calculable positive absolute constant, which might have different values at various appearances.

Since the opposite inequality clearly holds, by changing the role of  $l_1$  and  $l_2$ , one obtains the following

**COROLLARY.** On the above conditions, for  $i = 2, 4$  one has for the number of sign changes of  $\Delta_i(x)$  the lower estimate

$$(1.8) \quad V_i(Y, q, l_1, l_2) > \log_2 Y.$$

**THEOREM B.** Assume (H) (cf. (1.4)). Then for

$$(1.9) \quad Y > \max \left( \exp_2(q^c), \exp_2\left(\frac{c}{A^3(q)}\right) \right)$$

all functions  $\Delta_4(x, q, l_1, l_2)$  with (1.3) change their sign in the interval

$$(1.10) \quad [\log^2 Y/4, Y].$$

For the proofs of Theorem A and B see parts V and VII of [3], respectively.

In the present work we shall show (roughly speaking) that in the case of  $i = 4$  (1.7) can be improved to

$$(1.11) \quad \max_{x \in [Y \exp(-c(q) \log^{\dagger} Y), Y]} \Delta_4(x) > \sqrt{Y} \exp(-c(q) \log^{\dagger} Y)$$

thereby obtaining

$$(1.12) \quad V_4(Y) > c(q) \log^{\dagger} Y .$$

Furthermore, we get for the first sign change of  $\Delta_4(x)$  the upper bound

$$(1.13) \quad Y_0 = \exp\left(\frac{q^4}{A(q)}\right),$$

which improves (1.6). (In the above formulas we neglected some  $\log q$ ,  $\log(1/A(q))$  and  $\log \log Y$  factors).

Due to the lower estimate (1.11) by the strong form of the prime number theorem of arithmetic progressions, we obtain the same results in the case of  $i = 3$ , if  $l_1$  and  $l_2$  are both quadratic residues or both quadratic non-residues. (In Knapowski-Turàn's proof this needs relatively many additional efforts, as can be seen from part VI of [3]).

Finally we remark that by ineffective methods (essentially due to Landau [5, § 197], Grosswald [2] and Anderson-Stark [1]) one obtains that if for a non-trivial zero  $\rho_0 = \beta_0 + i\gamma_0$  of an  $L(s, \chi, q)$ , function one has  $\gamma_0 \neq 0$ ,  $\beta_0 > \frac{1}{2}$  and

$$(1.14) \quad a(\rho_0) = \sum_{\substack{\chi(\bmod q) \\ L(\rho_0, \chi) = 0}} (\bar{\chi}(l_2) - \bar{\chi}(l_1)) m_{\chi}(\rho_0) \neq 0$$

(where  $m_{\chi}(\rho_0)$  denotes the multiplicity of  $\rho_0$  as a zero of  $L(s, \chi)$ ), then for every  $l_1, l_2$  with (1.3)

$$(1.15) \quad \overline{\lim}_{x \rightarrow \infty} \frac{\Delta_4(x)}{x^{\beta_0}} > \left| \frac{a(\rho_0)}{\rho_0} \right|, \quad \underline{\lim}_{x \rightarrow \infty} \frac{\Delta_4(x)}{x^{\beta_0}} < - \left| \frac{a(\rho_0)}{\rho_0} \right|.$$

This is valid for  $\Delta_3(x)$  too, if  $l_1$  and  $l_2$  have the same quadratic character, or in case of  $\beta_0 > \frac{1}{2}$  for every pair  $l_1, l_2$  with (1.3). However, this theorem does not yield any localisation of sign changes or lower estimation of  $V_i(Y, q, l_1, l_2)$ .

We further note that the method of Knapowski and Turàn, (also in the present refined form) does not furnish better lower estimates than (essentially)  $\sqrt{Y}$ , even assuming the existence of a zero  $\rho_0$  with  $\beta_0 > \frac{1}{2}$ ,  $\gamma_0 \neq 0$  and  $a(\rho_0) \neq 0$ .

**2.** In our results, we shall always assume that the Haselgrove condition (H) and the finite Riemann-Piltz conjecture (FR-P) hold, the second

one up to a level  $D$  with

$$(2.1) \quad D = c_0 q^2 \log^6 q$$

where  $c_0$  is a sufficiently large positive absolute constant.

Here and in the sequel, we shall denote by  $c_i, i = 0, 1, \dots$  explicitly calculable positive absolute constants; moreover, the generic symbol  $c$ , and the signs  $\gg, \ll, 0$ , replace such constants;  $\exp(x) = e^x, \log_2 Y = \log \log Y$ .

We shall also assume without any further mention the trivial condition (1.3) on  $l_1, l_2$ .

Our results are the following:

**THEOREM 1.** *Assume (H), (FR-P) and let be such that*

$$(2.2) \quad Y > Y_0 = \exp\left(c \frac{D^5}{A(q)} \log \frac{1}{A(q)}\right)$$

*Then, there exists  $x$  with*

$$(2.3) \quad x \in \left[ Y \exp\left(-\frac{Cq}{\sqrt{A(q)}} (\log Y)^{\frac{1}{2}} (\log_2 Y)^{\frac{3}{2}}\right), Y \right]$$

*such that*

$$(2.4) \quad \Delta_4(x, q, l_1, l_2) > \sqrt{Y} \exp\left(-\frac{Cq}{\sqrt{A(q)}} (\log Y)^{\frac{1}{2}} (\log_2 Y)^{\frac{3}{2}}\right).$$

**COROLLARY 1.** *Under the assumptions of Theorem 1, for  $Y$  verifying (2.2) we have at least one sign change of  $\Delta_4(x)$  for  $x$  belonging to the interval (2.3).*

*Hence, we obtain*

$$(2.5) \quad V_4(Y, q, l_1, l_2) \gg \frac{\sqrt{A(q)} \log^{\frac{1}{2}} Y}{q \log^{\frac{3}{2}} Y} \quad \text{for } Y > \exp\left(cD^5 A^{-1}(q) \log \frac{1}{A(q)}\right).$$

**THEOREM 2.** *Assume (H), (FR-P) and let*

$$(2.6) \quad Y > \exp\left(\frac{cD^2}{A(q)} \log^3 \frac{D}{A(q)}\right).$$

*Then, there exists  $x$  with*

$$(2.7) \quad x \in \left[ Y \exp\left(-\frac{c \log^{\frac{3}{2}} Y \log_2 Y}{(DA(q))^{\frac{1}{2}}}\right), Y \right]$$

*such that*

$$(2.8) \quad \Delta_4(x) > \sqrt{x} \exp\left(-\frac{cq^{8/3} \log^4 q}{A^{\frac{1}{2}}(q)} (\log Y)^{\frac{1}{2}} (\log_2 Y)^2\right).$$

Using the fact that the error term given by the Prime Number Theorem for arithmetic progression (see for instance Prachar [5, pp. 297-298]) is smaller than the lower bounds (2.4), (2.7), we obtain

**COROLLARY 2.** *The statements of Theorem 1, Corollary 1 and Theorem 2 continue to be true without any change also for  $\Delta_3(x)$ , if  $l_1$  and  $l_2$  are both quadratic non-residues or if they are both quadratic residues (mod  $q$ ).*

Actually one can assure (2.2)-(2.4) also for  $i = 3$  if  $l_1$  is a quadratic non-residue and  $l_2$  a residue. However, in this case we cannot guarantee the opposite inequality, which follows in the earlier cases just by changing the role of  $l_1$  and  $l_2$ .

**3. We introduce**

$$(3.1) \quad F(s) = \sum_{n=1}^{\infty} \frac{\varepsilon(n) \Lambda(n)}{n}.$$

We have

$$(3.2) \quad F(s) = \frac{1}{\varphi(q)} \sum_{\chi(\text{mod } q)} (\bar{\chi}(l_2) - \bar{\chi}(l_1)) \frac{L'}{L}(s, \chi) = \frac{f_{-1}}{s} + f_0 + \dots$$

where (see [5], 7.4.21 and 7.4.46)

$$(3.3) \quad |f_{-1}| = \left| \frac{1}{\varphi(q)} \sum_{\substack{\chi(\text{mod } q) \\ \chi(-1)=1}} (\bar{\chi}(l_2) - \bar{\chi}(l_1)) \right| \leq 2,$$

$$(3.4) \quad f_0 = O(\log^2 q)$$

if there exists no Siegel-zero, as assured by (H).

Furthermore, let

$$(3.5) \quad a_\varrho = \frac{1}{\varphi(q)} \sum_{\substack{\chi(\text{mod } q) \\ L(\varrho, \chi)=0}} (\bar{\chi}(l_2) - \bar{\chi}(l_1)) m(\varrho) = \text{Res}_{s=\varrho} F(s)$$

where  $m_\chi(\varrho)$  denotes the multiplicity of  $\varrho$  as a zero of  $L(s, \chi)$ .

**LEMMA 1.** *For  $\exp[-3\mu] \leq K \leq \mu/9$ ,  $\mu \geq \log q$ , we have*

$$(3.6) \quad \frac{1}{2\sqrt{\pi K}} \int_{\exp(\mu+3\sqrt{\mu K})}^{\exp(\mu-3\sqrt{\mu K})} \frac{\Delta_4(x)}{x} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx = \sum_{\varrho} \frac{a_\varrho}{\varrho} \exp(K\varrho^2 + \mu\varrho) + \mu f_{-1} + f_0 + O(\exp(-\mu/4))$$

where the sum is performed on the non-trivial zeros  $\rho$  ( $0 < \text{Re } \rho < 1$ ) of  $L$ -functions (mod  $q$ ).

PROOF. We use the following integral formula

$$(3.7) \quad \frac{1}{2i\pi} \int_{(2)} \exp(Ks^2 + \alpha s) ds = \frac{1}{2\sqrt{\pi K}} \exp\left(-\frac{\alpha^2}{4K}\right), \quad K \in \mathbf{R}^+, \quad \alpha \in \mathbf{C}$$

and the fact that, by partial summation

$$(3.8) \quad F(s) = \sum_{n=1}^{\infty} \frac{\varepsilon(n)A(n)}{n^s} = s \int_1^{\infty} \frac{A_4(x)}{x^{s+1}} dx.$$

Then, we have

$$(3.9) \quad \begin{aligned} \frac{1}{2\sqrt{\pi K}} \int_1^{\infty} \frac{A_4(x)}{x} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx \\ = \frac{1}{2\pi i} \int_1^{\infty} \frac{A_4(x)}{x} \int_{(2)} \exp[Ks^2 + (\mu - \log x)s] ds dx \\ = \frac{1}{2\pi i} \int_{(2)} \frac{F(s)}{s} \exp[Ks^2 + \mu s] ds. \end{aligned}$$

Since the following well-known estimate holds:

$$(3.10) \quad \frac{L'}{L}(s, \chi) \ll \log(q(|t| + 2)) \ll (\log q) \log(|t| + 2) \quad \text{for } s = -\frac{1}{2} + it$$

we obtain

$$(3.11) \quad \begin{aligned} R = \frac{1}{2\pi i} \int_{(-\frac{1}{2})} \frac{F(s)}{s} \exp(Ks^2 + \mu s) ds \ll (\log q) \exp\left[\frac{K}{4} - \frac{\mu}{2}\right] \\ \cdot \left\{ \int_0^{2+1/K} \frac{\log(|t| + 2)}{|t| + \frac{1}{2}} dt + \int_{2+1/K}^{\infty} \exp[-Kt^2] dt \right\} \\ \ll \mu \exp[-\mu/3] \left\{ \log^2\left(2 + \frac{1}{K}\right) + \frac{1}{\sqrt{K}} \exp\left[-K\left(2 + \frac{1}{K}\right)^2\right] \right\} \ll \exp[-\mu/4]. \end{aligned}$$

Now, by Cauchy's residues theorem, we get

$$(3.12) \quad \frac{1}{2\pi i} \int_{(2)} \frac{F(s)}{s} \exp[Ks^2 + \mu s] ds = \sum_{\rho} \frac{a_{\rho}}{\rho} \exp[K\rho^2 + \mu\rho] + \mu f_{-1} + f_0 + R$$



and, in view of (3.11),

$$(3.13) \quad \frac{1}{2\pi i} \int_{(2)} \frac{F(s)}{s} \exp(Ks^2 + \mu s) ds = \sum_{\rho} \frac{\alpha_{\rho}}{\rho} \exp[K\rho^2 + \mu\rho] + \mu f_{-1} + f_0 + O(\exp(-\mu/4)).$$

Moreover, using the trivial estimate  $\Delta_4(x) \ll x$ , we obtain

$$(3.14) \quad \frac{1}{2\sqrt{\pi K}} \left| \int_{\exp(\mu+3\sqrt{\mu K})}^{\infty} \frac{\Delta_4(x)}{x} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx \right| \\ \leq \left| \int_{\frac{3\sqrt{\mu}}{2}}^{\infty} \exp\left(-(y - \sqrt{K})^2 + \mu + K\right) dy \right| \ll \exp\left(-\left(\frac{3\sqrt{\mu}}{2} - \sqrt{K}\right)^2 + \mu + K\right) \\ \leq \exp(-\mu/4)$$

where we have introduced in the integral the variable  $y = \log x - \mu/2\sqrt{K}$  and we have used  $3\sqrt{\mu}/2 > \sqrt{K}$ ,  $\mu > 9K$ .

Since a similar estimate holds, completely trivially, also for  $\int_1^{\exp(\mu-3\sqrt{\mu K})}$ , we get

$$(3.15) \quad \frac{1}{2\sqrt{\pi K}} \int_1^{\infty} \frac{\Delta_4(x)}{x} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx \\ = \frac{1}{2\sqrt{\pi K}} \int_{\exp(\mu-3\sqrt{\mu K})}^{\exp(\mu+3\sqrt{\mu K})} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) \frac{\Delta_4(x)}{x} dx + O(\exp(-\mu/4)).$$

Now, our Lemma follows collecting together (3.9), (3.13) and (3.15). ■

Since the main part of (3.6) can be written as a powersum, our problem is reduced to give a good lower bound for it. This is accomplished by means of a « one-sided » powersum theorem of Knapowski and Turàn (see Theorem 4.1 in part III of [3]). In order to obtain sharper estimates, we shall need this result in the following slightly modified form:

LEMMA 2. Let  $b_j, z_j \in \mathbf{C}$  for  $j = 1, 2, \dots, n$ , with

$$(3.16) \quad 0 < \kappa \leq |\arg z_j| < \pi \quad \forall_j$$

$$(3.17) \quad |z_1| \geq |z_2| \geq \dots \geq |z_n|.$$

Then for any  $h$  with  $1 < h < n$  and for any  $m \geq 0$ , there exists an integer  $\nu$  with

$$(3.18) \quad \nu \in \left[ m, m + n \left( 3 + \frac{\pi}{\kappa} \right) \right]$$

such that

$$(3.19) \quad \operatorname{Re} \sum_{j=1}^n b_j z_j^\nu > \frac{E}{2n+1} \left( \frac{24e(m+n(5+\pi/\kappa))}{n} \right)^{-2n} |z|^\nu \left| \frac{z_h}{z_1} \right|^{n(5+\pi/\kappa)}$$

where

$$(3.20) \quad E = \min_{i \geq h} \left| \operatorname{Re} \sum_{j=1}^i b_j \right|.$$

PROOF. Following the lines the theorem of [3] quoted above, we obtain the following inequality, in the case  $|z_1| = 1$ ,

$$(3.21) \quad \operatorname{Re} \sum_{j=1}^n b_j z_j^\nu > \frac{E}{2n+1} \left( \frac{|z_h| - \delta}{48} \right)^{2n} \delta^{m+n(3+\pi/\kappa)}$$

for a suitable  $\nu$  verifying (3.18) and for every  $\delta$  with

$$(3.22) \quad 0 < \delta < |z_h|$$

Then, (3.19) follows by choosing

$$(3.23) \quad \delta = |z_h| \left( 1 - \frac{2n}{m+n(5+\pi/\kappa)} \right)$$

unlike to Knapowski-Turàn's choice  $\delta = |z_h| - 2n/(m+n(3+\pi/\kappa))$ . ■

According to (3.19), in our applications we shall need a non trivial lower estimate for  $E$ . This requires a modification of the coefficients in the power-sum, furnished by the following Lemma:

LEMMA 3. *There exists a prime  $P \equiv l_1 \pmod{q}$  with*

$$(3.24) \quad \frac{D}{2} < P \log^2 P < D$$

such that, for  $P_0 = P + \frac{1}{2}$  or  $P_0 = P - \frac{1}{2}$  we have

$$(3.25) \quad \left| \sum_{\varrho} \frac{a_\varrho}{\varrho} \exp(K_0 \varrho^2 + i\mu_0 \varrho) \right| > \frac{\log P}{3}$$

where

$$(3.26) \quad K_0 = \frac{P_0^2 \log^2 P_0}{1}, \quad \mu_0 = \log P_0.$$

PROOF. Since the finite Riemann-Piltz conjecture is assumed to be true, the prime number formula of arithmetic progressions, truncated at  $D$ , assures the existence of a prime  $P \equiv l_1 \pmod{q}$  verifying (3.24). Then, the function  $\Delta_4(x)$  has a jump  $\log P$  at the point  $P$  and we have

$$(3.27) \quad \Delta_4(P_2) - \Delta_4(P_1) = \log P$$

for  $P_1 = P - \frac{1}{2}$ ,  $P_2 = P + \frac{1}{2}$ .

Now, we use Lemma 1 with  $\mu_i = \log P_i$ ,  $K_i = 1/P_i^2 \log^2 P_i$  for  $i = 1, 2$ . This choice implies

$$(3.28) \quad [\exp(\mu_i - 3\sqrt{\mu_i K_i}), \exp(\mu_i + 3\sqrt{\mu_i K_i})] \subset \left(P_i - \frac{1}{2}, P_i + \frac{1}{2}\right)$$

since  $P$  is large enough by (3.24) and (2.1).

Thus,  $\Delta_4(x) = \Delta_4(P_i)$  in the above interval and, using Lemma 1, and setting  $x = \exp(\mu + 2\sqrt{K}y)$  we obtain:

$$(3.29) \quad \begin{aligned} & \sum_{\varrho} \frac{a_{\varrho}}{\varrho} \exp(K_i \varrho^2 + \mu_i \varrho) + f_{-1} \mu_i + f_0 \\ &= \frac{\Delta_4(P_i)}{\sqrt{\pi}} \int_{\exp(\mu_i - 3\sqrt{\mu_i K_i})}^{\exp(\mu_i + 3\sqrt{\mu_i K_i})} \exp\left(-\frac{(\mu_i - \log x)^2}{4K_i}\right) \frac{dx}{2\sqrt{K_i}x} + O(\exp(-\mu/4)) \\ &= \frac{\Delta_4(P_i)}{\sqrt{\pi}} \int_{-3\sqrt{\mu_i/2}}^{3\sqrt{\mu_i/2}} \exp(-y^2) dy + O(\exp(-\mu/4)) = \Delta_4(P_i) + O(P^{-1/4}). \end{aligned}$$

Owing to  $|f_{-1}| \leq 2$  and (3.27) we get from this

$$(3.30) \quad O\left(\frac{1}{P}\right) + \sum_{\varrho} \frac{a_{\varrho}}{\varrho} (\exp(K_2 \varrho^2 + \mu_2 \varrho) - \exp(K_1 \varrho^2 + \mu_1 \varrho)) = \log P + O(P^{-1/4})$$

which implies Lemma 3. ■

In view of the application of Lemma 2 to the power-sum appearing in (2.6), it is also necessary to assure the argument condition (3.16), and this is made by means of

LEMMA 4. *Let  $a_j$  be real numbers for  $j = 1, \dots, n$  with  $a_j \neq 0$  and*

$$(3.31) \quad \frac{1}{n} \sum_{j=1}^n |a_j|^{-1} \leq \eta.$$

Then, for every  $H$  there exists an  $y_0$  with

$$(3.32) \quad y_0 \in [H, H + \eta]$$

such that, for any integer  $K$  and for every  $j = 1, \dots, n$

$$(3.33) \quad |y_0 a_j - 2K\pi| \geq \frac{1}{4n}.$$

PROOF. For fixed  $j$ , (3.33) can be false for  $K$  in an interval of length at most  $(\eta|a_j| + 1/2n)/2\pi$ ; for fixed  $K$ , leaving  $j$  fixed, this can happen for  $y$  in an interval of length at most  $1/2n|a_j|$ .

Thus, the total Lebesgue measure of  $y$  for which (3.33) is false for fixed  $j$  is majorised by

$$(3.34) \quad \frac{1}{2n|a_j|} \left( \frac{\eta|a_j|}{2} + \frac{3}{2} \right) = \frac{\eta}{4n} + \frac{3}{4} \frac{1}{n|a_j|}.$$

Summing over  $j = 1, \dots, n$ , we obtain our Lemma. ■

Now, we are in good position to apply Lemma 2.

We introduce the following position:

$$(3.35) \quad b_\varrho = \frac{a_\varrho}{\varrho} \exp(K_0 \varrho^2 + \mu_0 \varrho)$$

for  $a_\varrho$  given by (3.5) and  $K_0, \mu_0$  furnished by (3.26).

Furthermore, let

$$(3.36) \quad \lambda > 2D, \quad L > \frac{c q^2}{A(q)} \lambda \log^3 \lambda$$

Let  $B$  be a real number to be chosen later with

$$(3.37) \quad B \in \left[ \frac{c \log \lambda}{A(q) \lambda}, \frac{2c \log \lambda}{A(q) \lambda} \right]$$

and let  $\nu$  be an integer to be chosen later with

$$(3.38) \quad \nu \in \left[ \frac{L - \mu_0}{B} - c q^2 \lambda^2 \log^2 L, \frac{L - \mu_0}{B} \right]$$

$$(3.39) \quad K = K_0 + \frac{B\nu}{\lambda^2} \quad \mu = \mu_0 + B\nu$$

$$(3.40) \quad z_\varrho = \exp \left[ \frac{B\varrho^2}{\lambda^2} + B\varrho \right].$$

LEMMA 5. *With the positions (3.35) to (3.40), we have*

$$(3.41) \quad \operatorname{Re} \sum_{\varrho} \frac{a_{\varrho}}{\varrho} \exp(K\varrho^2 + \mu\varrho) > \exp \left\{ \frac{L}{2} - \frac{LD^2}{\lambda^2} - \frac{cq^2}{A(q)} \lambda \log^3 L \right\}$$

for suitable values of  $\nu$  and  $B$  satisfying (3.38).

PROOF. By our definitions, we have

$$(3.42) \quad \sum_{\varrho} \frac{a_{\varrho}}{\varrho} \exp(K\varrho^2 + \mu\varrho) = \sum_{\varrho} b_{\varrho} z_{\varrho}^{\nu}.$$

Moreover

$$(3.42) \quad \left| \sum_{|\varrho| \geq 2\lambda} \frac{a_{\varrho}}{\varrho} \exp(K\varrho^2 + \mu\varrho) \right| \ll \sum_{m=\lceil 2\lambda \rceil}^{\infty} \exp(\mu + K(1 - m^2)(\log qm)) \ll 1$$

and so we have only to consider

$$(3.43) \quad \sum_{|\varrho| < 2\lambda} b_{\varrho} z_{\varrho}^{\nu}.$$

Here, the number  $n$  of terms is clearly

$$(3.44) \quad c_1 \varphi(q) \lambda \log \lambda \leq n \leq c_2 q \lambda \log \lambda.$$

Now, we apply Lemma 4, setting  $\varrho = \beta + i\gamma$ ,

$$(3.45) \quad a_j = \left( \frac{2\beta}{\lambda^2} + 1 \right) \gamma, \quad H = \frac{c \log \lambda}{A(q) \lambda}.$$

Using Jensen's inequality, we have by  $A(q) < 1$  and  $\lambda > q$ ,

$$(3.46) \quad \sum_{j=1}^n \frac{1}{|a_j|} \ll \sum_{|\varrho| \leq 2\lambda} \frac{1}{|\gamma|} \leq \sum_{\substack{A(q) \leq |\gamma| \leq 1 \\ |\gamma| > 1}} \frac{1}{|\gamma|} + \sum_{\substack{|\varrho| \leq 2\lambda \\ |\gamma| > 1}} \frac{1}{|\gamma|} \ll \frac{\varphi(q) \log q}{A(q)} + \varphi(q) \log^2 \lambda;$$

so, in view of (3.44), condition (3.31) holds with

$$(3.47) \quad \eta = \frac{c \log \lambda}{A(q) \lambda}.$$

Hence, Lemma 4 says that there exists a  $B$  in the interval (3.37) such that, for every  $j = 1, \dots, n$

$$(3.48) \quad |Ba_j - 2K\pi| \geq \frac{1}{4n} = \frac{c}{q\lambda \log \lambda}.$$

Since in our case  $z_{e_j} = |z_{e_j}| \exp(ia_j B)$ , this means that

$$(3.49) \quad |\text{Arg } z_e| > \frac{c}{\eta} = \frac{c}{q\lambda \log \lambda} = \varkappa.$$

Now, we order the numbers  $z_e$  according to (3.17) and, in view of (FR-P), we choose  $h$  of Lemma 2 as the largest index corresponding to a zero  $\rho$  with  $|\rho_0| \leq D$ .

Since

$$(3.50) \quad \left| \sum_{|\rho| > D} \frac{a_\rho}{\rho} \exp(K_0 \rho^2 + \mu_0 \rho) \right| \ll \sum_{m=P \log^2 P}^{\infty} \exp \left[ \frac{1-m^2}{P^2 \log^2 P} \right] P(\log m q) \ll 1$$

with constants implied by the  $\ll$  sign, independent of  $c_0$  appearing in (2.1), we have by Lemma 3

$$(3.51) \quad \left| \sum_{\rho \in S} b_\rho \right| > \frac{\log P}{4}$$

for every set  $S$  containing all zeros  $\rho$  with  $|\rho| \leq D$ .

Finally, we have

$$(3.52) \quad |z_h| \geq \exp \left( \frac{B}{2} - \frac{B}{\lambda^2} D^2 \right)$$

because  $\rho_0$  is on the critical line, and (by (3.40) and  $\lambda > 2D$ )

$$(3.52) \quad \left| \frac{z_h}{z_1} \right| \geq \exp \left( \frac{B}{2} - \frac{B}{\lambda^2} D^2 - B - \frac{B}{\lambda^2} \right) > \exp(-B).$$

Recalling (3.44), (3.49), (3.51), (3.52), (3.53) and choosing  $m$  as  $m = (L - \mu_0)/B$  in view of (3.38), we obtain by Lemma 2 for suitable  $B$  and  $\nu$ :

$$(3.54) \quad \text{Re} \sum_{|\rho| < 2\lambda} b_\rho z_\rho^\nu > \exp \left\{ \frac{B\nu}{2} - B\nu \frac{D^2}{\lambda^2} - c \frac{q^2}{A(q)} \lambda \log^3 \lambda - cq\lambda \log \lambda \log L \right\} \\ \gg \exp \left\{ \frac{L}{2} - \frac{D^2}{\lambda^2} L - \frac{cq^2}{A(q)} \lambda \log^3 L \right\},$$

from which our Lemma immediately follows. ■

**4. We formulate the results of Section 3 as:**

**THEOREM 3.** *Assume (H), (FR-P) and*

$$(4.1) \quad \lambda > 2D, \quad L > \max(cq^2 A^{-1}(q) \lambda \log^2 \lambda, \lambda^2).$$

Then there exist

$$(4.2) \quad \mu \in \left[ L - \frac{cq^2}{A(q)} \lambda \log^3 \lambda, L \right]$$

$$(4.3) \quad K \in \left[ \frac{L - \mu_0}{\lambda^2} - \frac{cq^2}{A(q)} \frac{\log^3 \lambda}{\lambda}, K_0 + \frac{L}{\lambda^2} \right]$$

such that

$$(4.4) \quad \frac{1}{2\sqrt{\pi K}} \int_{\exp(\mu - 3\sqrt{\mu K})}^{\exp(\mu + 3\sqrt{\mu K})} \frac{\Delta_4(x)}{x} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx > \exp\left\{\frac{L}{2} - \frac{LD^2}{\lambda^2} - \frac{cq^2}{A(q)} \lambda \log^3 L\right\}.$$

PROOF. The theorem follows immediately from Lemmas 1 and 5. ■

PROOF OF THEOREM 2. We set, with the notations of Theorem 3

$$(4.5) \quad Y = \exp(L_1), \quad L = L_1 \left(1 - \frac{7}{2\lambda}\right).$$

Then we obtain, by easy calculations from (4.1)-(4.3)  $\sqrt{\mu K} \leq 1$ ,  $1/L \leq \lambda$  and so

$$(4.6) \quad I \subseteq \left[ Y \exp\left(-\frac{7L_1}{\lambda} - \frac{cq^2}{A(q)} \lambda \log^3 \lambda\right), Y \right].$$

We set also

$$(4.7) \quad A = \max_{x \in I} \frac{\Delta_4(x)}{\sqrt{x}}, \quad I = [\exp(\mu - 3\sqrt{\mu K}), \exp(\mu + 3\sqrt{\mu K})].$$

By Theorem 3, we have for suitable  $\mu, K$  verifying (4.2), (4.3),

$$(4.8) \quad \frac{A}{2\sqrt{\pi K}} \int_{x \in I} \frac{1}{\sqrt{x}} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx > \int_{x \in I} \frac{\Delta_4(x)}{x} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx \cdot \frac{1}{2\sqrt{\pi K}} > \exp\left\{\frac{L}{2} - \frac{LD^2}{\lambda^2} - \frac{cq^2}{A(q)} \lambda \log^3 L\right\}.$$

Since, as it is easily verified,

$$(4.9) \quad \frac{1}{2\sqrt{\pi K}} \int_0^\infty \frac{1}{\sqrt{x}} \exp\left(-\frac{(\mu - \log x)^2}{4K}\right) dx = \exp\left(\frac{\mu}{2} + \frac{k}{4}\right)$$

and  $\mu \leq L$ , inequality (4.8) yields

$$(4.10) \quad A > \exp \left\{ -\frac{2L_1 D^2}{\lambda^2} - \frac{cq^2}{A(q)} \lambda \log^3 L_1 \right\}.$$

In order to optimise the lower bound (4.10), we choose

$$(4.11) \quad \lambda = cA^{\frac{1}{3}}(q) q^{\frac{1}{3}} \log^4 q \frac{L_1^{\frac{1}{3}}}{\log L_1}$$

which satisfies (4.1), in view of (2.6) and (4.5).

Thus, by (2.1), (4.5), (4.6), (4.10) we obtain

$$(4.12) \quad A > \exp \left( -\frac{cq^{8/3} \log q}{A^{\frac{1}{3}}(q)} (\log Y)^{\frac{1}{3}} (\log_2 Y)^2 \right)$$

and

$$(4.13) \quad I \subset \left[ Y \exp \left( -\frac{c(\log Y)^{\frac{1}{3}} \log_2 Y}{(DA(q))^{\frac{1}{3}}} \right), Y \right]$$

which proves Theorem 2. ■

**PROOF OF THEOREM 1.** We set now also (4.5) and similarly we obtain (4.6)-(4.10) from (4.1)-(4.3).

In order to optimise localisation in (4.6) we choose

$$(4.14) \quad \lambda = \frac{cA^{\frac{1}{3}}(q)}{q} \frac{L^{\frac{1}{3}}}{(\log L_1)^{\frac{1}{3}}} > D^2,$$

in view of (2.1)-(2.2), and this proves also (4.1).

In such a way we obtain by (4.10)

$$(4.15) \quad I \subset \left[ Y \exp \left( -\frac{cL_1}{\lambda} \right), Y \right]$$

and, for suitable  $x$  in  $I$ , by (4.10), (4.14) and (4.15)

$$(4.16) \quad \Delta_4(x) = A \sqrt{x} > \exp \left( -\frac{cL_1}{\lambda} \right) \sqrt{Y} \exp \left( -\frac{cL_1}{\lambda} \right),$$

which proves Theorem 1, due to the choice of  $\lambda$  in (4.14). ■



## REFERENCES

- [1] R. J. ANDERSON - H. M. STARK, *Oscillation Theorems*, Analytic Number Theory, Proceedings, Philadelphia, 1980, Ed. F. Knopp, Springer, Lecture Notes in Mathematics N. 899, pp. 79-106.
- [2] E. GROSSWALD, *On some generalizations of theorems by Landau and Pòlya*, Israel J. Math., **3** (1965), pp. 211-220.
- [3] S. KNAPOWSKI - P. TURÁN, *Comparative prime number theory I-VIII*, Acta Math. Acad. Sci. Hungar., part I: **13** (1962), pp. 299-314; part II: **13** (1962), pp. 315-342; part III: **13** (1962), pp. 343-346; part IV: **14** (1963), pp. 31-42; part V: **14** (1963), pp. 43-63; part VI: **14** (1963), pp. 65-78; part VII: **14** (1963), pp. 241-250; part VIII: **14** (1963), pp. 251-268.
- [4] E. LANDAU, *Handbuch der Lehre von der Verteilung der Primzahlen*, Teubner, Leipzig und Berlin, 1909.
- [5] K. PRACHAR, *Primzahlverteilung*, Springer, Berlin-Göttingen-Heidelberg, 1957.
- [6] R. SPIRA, *Calculation of Dirichlet L-functions*, Math. Comput., **23**, N. 107 (1969), pp. 484-497.

Mathematical Institute of the  
Hungarian Academy of Sciences  
Budapest  
Realtanoda u. 13-15  
H-1053 Hungary

Istituto di Matematica  
Facoltà di Scienze dell'Università  
84100 Salerno