

Time-Delay Operators in Semiclassical Limit, Finite Range Potentials

XUE PING WANG

1. Introduction

In this paper we consider the semi-classical approximation of time-delay operators in quantum scattering theory. Let us recall briefly some related aspects of scattering theory. Let V be a C^∞ real function on $\mathbb{R}^n$ so that for some $\varepsilon > 0$ and all $\alpha \in \mathbb{N}^n$, one has:

$$(1.1) \quad |\partial_x^\alpha V(x)| \leq C_\alpha (1 + |x|)^{-|\alpha| - 1 - \varepsilon}.$$

For $h > 0$ a small parameter proportional to Planck constant, set: $H_0^h = -\frac{h}{2}\Delta$, $H^h = H_0^h + V_h$ where $V_h(x) = V(h^{1/2}x)$. Define the unitary groups $U_0^h(t)$ and $U^h(t)$ as:

$$U_0^h(t) = \exp(-ih^{-1}tH_0^h), \quad U^h(t) = \exp(-ih^{-1}tH^h), \quad t \in \mathbb{R}.$$

Then the wave operators $\Omega_\pm(h)$:

$$\Omega_\pm(h) = s - \lim_{t \rightarrow \pm\infty} U^h(t)^* U_0^h(t) \quad \text{in } L^2(\mathbb{R}^n)$$

exist and are complete, that is to say, the ranges of $\Omega_+(h)$ and $\Omega_-(h)$ are both identical with the continuous spectral subspace of H^h . The scattering operator $S(h)$ is defined by: $S(h) = \Omega_+(h)^* \Omega_-(h)$. $S(h)$ is a unitary operator in $L^2(\mathbb{R}^n)$. We denote by $F_h : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{$

