

ANNALI DELLA SCUOLA NORMALE SUPERIORE DI PISA *Classe di Scienze*

MAREK JARNICKI

PETER PFLUG

REIN ZEINSTRÄ

Geodesics for convex complex ellipsoids

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 20, n° 4 (1993), p. 535-543

http://www.numdam.org/item?id=ASNSP_1993_4_20_4_535_0

© Scuola Normale Superiore, Pisa, 1993, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Geodesics for Convex Complex Ellipsoids

MAREK JARNICKI - PETER PFLUG - REIN ZEINSTRÄ

Introduction

Given $p = (p_1, \dots, p_n)$ with $p_1, \dots, p_n \geq \frac{1}{2}$, define

$$\mathcal{E}(p) := \left\{ (z_1, \dots, z_n) \in \mathbb{C}^n : \sum_{j=1}^n |z_j|^{2p_j} < 1 \right\}.$$

The aim of this note is to describe all *complex geodesics* (cf. [Ves], [Din]) $\varphi : E \rightarrow \mathcal{E}(p)$, where $E := \{\lambda \in \mathbb{C} : |\lambda| < 1\}$. So far, only special cases of p and φ have been studied, e.g. [Pol], [Gen], [BFKKMP] — cf. Remark 3.

Recall that a holomorphic mapping $\varphi : E \rightarrow \mathcal{E}(p)$ is a complex geodesic if

$$k_{\mathcal{E}(p)}(\varphi(\lambda'), \varphi(\lambda'')) = k_E(\lambda', \lambda''), \quad \lambda', \lambda'' \in E,$$

where k stands for the Kobayashi distance. Let us mention that $\mathcal{E}(p)$ is convex and therefore, by the Lempert theorem (cf. [Lem]), the Kobayashi distance $k_{\mathcal{E}(p)}$ coincides with the Carathéodory distance and the Kobayashi-Royden metric $\kappa_{\mathcal{E}(p)}$ coincides with the Carathéodory-Reiffen metric.

Observe that if $\varphi = (\varphi_1, \dots, \varphi_n) : E \rightarrow \mathcal{E}(p)$ is a complex geodesic with $\varphi_n \equiv 0$ then the mapping $(\varphi_1, \dots, \varphi_{n-1}) : E \rightarrow \mathcal{E}((p_1, \dots, p_{n-1}))$ is a “lower dimensional” complex geodesic. This means that, without loss of generality, we can assume that

$$(1) \quad \varphi_j \not\equiv 0, \quad j = 1, \dots, n.$$

Moreover, using a suitable permutation of coordinates, we can always assume that for some $0 \leq s \leq n$:

$$(2) \quad \varphi_1, \dots, \varphi_s \text{ have zeros in } E \text{ and } \varphi_{s+1}, \dots, \varphi_n \text{ have no zeros in } E.$$

Pervenuto alla Redazione il 9 Gennaio 1992 e in forma definitiva il 25 Marzo 1993.

The main result of the paper is the following:

THEOREM 1. *A mapping $\varphi = (\varphi_1, \dots, \varphi_n) : E \rightarrow \mathbb{C}^n$ enjoying (1) and (2) is a complex geodesic in $\mathcal{E}(p)$ if and only if*

$$(3) \quad \varphi_j(\lambda) = \begin{cases} a_j \frac{\lambda - \alpha_j}{1 - \bar{\alpha}_j \lambda} \left(\frac{1 - \bar{\alpha}_j \lambda}{1 - \bar{\alpha}_0 \lambda} \right)^{1/p_j}, & j = 1, \dots, s \\ a_j \left(\frac{1 - \bar{\alpha}_j \lambda}{1 - \bar{\alpha}_0 \lambda} \right)^{1/p_j}, & j = s+1, \dots, n \end{cases},$$

where

$$(4) \quad a_1, \dots, a_n \in \mathbb{C} \setminus \{0\},$$

$$(5) \quad \alpha_0, \dots, \alpha_s \in E, \quad \alpha_{s+1}, \dots, \alpha_n \in \bar{E},$$

$$(6) \quad \alpha_0 = \sum_{j=1}^n |a_j|^{2p_j} \alpha_j,$$

$$(7) \quad 1 + |\alpha_0|^2 = \sum_{j=1}^n |a_j|^{2p_j} (1 + |\alpha_j|^2),$$

$$(8) \quad \text{the case } s = 0, \alpha_0 = \alpha_1 = \dots = \alpha_n \text{ is excluded.}$$

Moreover, the complex geodesics in $\mathcal{E}(p)$ are uniquely determined mod $\text{Aut}(E)$, i.e. if φ, ψ are complex geodesics in $\mathcal{E}(p)$ with $\varphi(0) = \psi(0)$, $\varphi'(0) = \psi'(0)$ (or $\varphi(0) = \psi(0)$, $\varphi(\sigma) = \psi(\sigma)$ for some $0 < \sigma < 1$) then $\varphi \equiv \psi$.

REMARK. (a) In particular, Theorem 1 shows that the components of a geodesic φ satisfying (1) have at most simple zeros in E .

(b) The case $p_1 = \dots = p_n = 1$, i.e. the case of the unit ball $\mathbb{B}_n$, is well-known — cf. Remark 7. Observe that the general case is a simple “transformation” of the case of $\mathbb{B}_n$.

COROLLARY 2. *Let $\varphi : E \rightarrow \mathcal{E}(p)$ be a complex geodesic.*

- (a) *If $p_1, \dots, p_n \in \left\{ \frac{1}{2}, 1 \right\}$ then φ extends holomorphically to a neighborhood of $\bar{E}$.*
- (b) *If $t := \max\{p_1, \dots, p_n\} > 1$ then φ extends to a mapping of the class $C^{0,1/t}(\bar{E}) := \left\{ \frac{1}{t}\text{-Hölder continuous mappings on } \bar{E} \right\}$.*
- (c) *If $u := \max\{p_j : p_j \neq 1\} < 1$ then φ extends to a mapping of the class $C^{1,(1/u)-1}(\bar{E})$.*

REMARK 3. (a) The case $p_1 = \dots = p_n > 1$ has been studied in [Pol].

(b) The case $p_1 = \dots = p_n = \frac{1}{2}$ under the additional assumption that φ extends continuously on $\bar{E}$ has been discussed in [Gen].

(c) The case $n = 2, p_1 = 1$ can be found in [BFFKKMP].

(d) Notice that in many cases Theorem 1 gives a tool to calculate $\kappa_{\mathcal{E}(p)}$ effectively. For example, if $n = 2$ and $p_1 = 1$ then, using Theorem 1, one can easily verify the formulas for $\kappa_{\mathcal{E}(p)}$ obtained in [BFKKMP].

PROOF OF THEOREM 1. The proof will be based on the following criterion due to H. Royden and P.-M. Wong:

THEOREM 4. *Let $G \subset \mathbb{C}^n$ be a bounded convex domain with $0 \in G$. Then a holomorphic mapping $\varphi : E \rightarrow G$ is a complex geodesic if and only if:*

$$\varphi^*(\lambda) \in \partial G \quad \text{for almost all } \lambda \in \partial E$$

and there exists $h \in H^1(E, \mathbb{C}^n)$, $h \neq 0$, such that

$$\operatorname{Re} \left(\frac{1}{\lambda} \varphi^*(\lambda) \bullet h^*(\lambda) \right) = \hat{q}_G \left(\frac{1}{\lambda} h^*(\lambda) \right) \quad \text{for almost all } \lambda \in \partial E,$$

where:

H^1 denotes the Hardy space,

φ^* , h^* stand for non-tangential boundary values,

$$w \bullet z := \sum_{j=1}^n w_j z_j \quad \text{and}$$

$\hat{q}_G$ is the dual subnorm for the Minkowski function q_G of G , i.e.

$$\hat{q}_G(w) := \max \{ \operatorname{Re}(w \bullet z) : z \in \partial G \}, \quad w \in \mathbb{C}^n.$$

REMARK. The proof presented in [Roy-Won] contains an error in the proof of Proposition 1. Complete proofs may be found in [Aba, Section 2.6] and in [Jar-Pfl, Section 8.2].

REMARK 5. (a) Suppose that $z_0 \in \partial G$ is such that there is a uniquely determined unit outer normal vector $\nu(z_0)$ to ∂G at z_0 . Then for $w_0 \in \mathbb{C}^n \setminus \{0\}$ one has:

$$[\operatorname{Re}(z_0 \bullet w_0) = \hat{q}_G(w_0)] \Leftrightarrow [\exists t > 0 : w_0 = t \overline{\nu(z_0)}].$$

(b) Observe that any point $z_0 \in \partial \mathcal{E}(p)$ has a local peak function. In particular, if $\varphi : E \rightarrow \overline{\mathcal{E}(p)}$ is a non-constant holomorphic mapping then $\varphi : E \rightarrow \mathcal{E}(p)$.

Theorem 4, Remark 5 and the identity theorem for H^1 -functions imply the following useful

COROLLARY 6. *Let $\varphi = (\varphi_1, \dots, \varphi_n) : E \rightarrow \mathbb{C}^n$ be a non-constant bounded holomorphic mapping enjoying (1). Then φ is a complex geodesic in $\mathcal{E}(p)$ if and only if:*

$$(9) \quad \sum_{j=1}^n |\varphi_j^*|^2 p_j = 1 \quad \text{a.e. on } \partial E$$

and there exist $h \in H^1(E, \mathbb{C}^n)$ and $\rho : \partial E \rightarrow \mathbb{R}_{\geq 0}$ such that

$$(10) \quad \frac{1}{\lambda} h_j^* = \rho p_j |\varphi_j^*|^{2(p_j-1)} \overline{\varphi_j^*} \quad \text{a.e. on } \partial E, \quad j = 1, \dots, n.$$

REMARK 7. Let $p_1 = \dots = p_n = 1$ and φ be given by (3) with conditions (4)–(8). Then φ is a complex geodesic in $\mathbb{B}_n$. In fact:

- condition (8) assures that φ is non-constant,
- conditions (6) and (7) imply (9),
- if we put

$$(11) \quad h_j(\lambda) := \begin{cases} \bar{a}_j(1 - \bar{\alpha}_j\lambda)(1 - \bar{\alpha}_0\lambda), & j = 1, \dots, s \\ \bar{a}_j(\lambda - \alpha_j)(1 - \bar{\alpha}_0\lambda), & j = s+1, \dots, n \end{cases},$$

then also condition (10) is fulfilled with $\rho(\lambda) := |1 - \bar{\alpha}_0\lambda|^2$.

Using Corollary 6, one can easily prove the following “transport lemma” for complex geodesics.

LEMMA 8. Let $\varphi = (\varphi_1, \dots, \varphi_n) : E \rightarrow \mathcal{E}(p)$ be a complex geodesic enjoying (1) and let $h = (h_1, \dots, h_n)$ be as in (10). Write

$$\varphi_j = B_j \psi_j,$$

where B_j is the Blaschke product for φ_j and ψ_j is nowhere vanishing (if φ_j has no zeros in E then we put $B_j := 1$), $j = 1, \dots, n$. Let $\tilde{p} = (\tilde{p}_1, \dots, \tilde{p}_n)$, $\tilde{p}_1, \dots, \tilde{p}_n \geq \frac{1}{2}$ and define:

$$\tilde{\varphi}_j := B_j \psi_j^{p_j/\tilde{p}_j},$$

$$\tilde{h}_j := \frac{\tilde{p}_j}{p_j} h_j \frac{\varphi_j}{\tilde{\varphi}_j}.$$

Then the mappings $\tilde{\varphi}$ and $\tilde{h}$ satisfy (9) and (10) with respect to $\mathcal{E}(\tilde{p})$ (with the same function ρ).

In particular, if $\tilde{h} \in H^1(E, \mathbb{C}^n)$ (e.g. if $p_j \leq \tilde{p}_j$, $j = 1, \dots, n$) then $\tilde{\varphi} : E \rightarrow \mathcal{E}(\tilde{p})$ is a complex geodesic.

Consequently, in view of Remark 7, we get:

COROLLARY 9. If φ is given by (3) with (4)–(8) then $\varphi : E \rightarrow \mathcal{E}(p)$ is a complex geodesic.

PROOF. In view of (11), the new mapping $\tilde{h}$ belongs to H^1 . □

COROLLARY 10. If all complex geodesics $\varphi : E \rightarrow \mathcal{E}(p)$ with (1) and (2) are of the form (3) then the same is true for all complex geodesics $\tilde{\varphi} : E \rightarrow \mathcal{E}(\tilde{p})$ with $\tilde{p}_j \leq p_j$, $j = 1, \dots, n$.

Thus, in order to finish the proof of the first part of Theorem 1, it is sufficient to verify the following:

LEMMA 11. *If $p_1 = \dots = p_n =: p_0$ then any complex geodesic $\varphi : E \rightarrow \mathcal{E}(p)$ with (1) and (2) is of the form (3) with (4)–(8).*

PROOF. Let $h = (h_1, \dots, h_n) \in H^1(E, \mathbb{C}^n)$ and ρ be as in (10). Then

$$\frac{1}{\lambda} \varphi_j^* h_j^* \in \mathbb{R}_{>0} \quad \text{a.e. on } \partial E, \quad j = 1, \dots, n.$$

Hence, by [Gen, Lemma 2 and proof of Theorem 5] (see also [Pol, Statement 6]), there exist

$$r_0, \dots, r_n > 0, \quad \alpha_0, \dots, \alpha_n \in \bar{E}$$

such that

$$(12) \quad \varphi_j(\lambda) h_j(\lambda) = r_j(\lambda - \alpha_j)(1 - \bar{\alpha}_j \lambda), \quad \lambda \in E, \quad j = 1, \dots, n,$$

$$(13) \quad \varphi(\lambda) \bullet h(\lambda) = r_0(\lambda - \alpha_0)(1 - \bar{\alpha}_0 \lambda), \quad \lambda \in E.$$

Substituting h by $\frac{1}{r_0} h$ we can always assume that $r_0 = 1$. In view of (2),

$$\alpha_1, \dots, \alpha_n \in E.$$

Moreover, (12) and (13) imply that

$$\alpha_0 = \sum_{j=1}^n r_j \alpha_j \quad \text{and}$$

$$1 + |\alpha_0|^2 = \sum_{j=1}^n r_j (1 + |\alpha_j|^2).$$

Observe that, by (9), (10) and (13), we get

$$(14) \quad p_0 \rho(\lambda) = |1 - \bar{\alpha}_0 \lambda|^2 \quad \text{for almost all } \lambda \in \partial E.$$

In particular, (9), (10) and (14) show that

$$(15) \quad \text{the functions } h_1, \dots, h_n \text{ are bounded}$$

and, in view of (12), that

$$|\varphi_j^*(\lambda)| = r_j^{1/2p_0} \left| \frac{1 - \bar{\alpha}_j \lambda}{1 - \bar{\alpha}_0 \lambda} \right|^{\frac{1}{p_0}} \quad \text{for almost all } \lambda \in \partial E, \quad j = 1, \dots, n.$$

Consequently, since $(1 - \bar{\alpha}_j \lambda)^{1/p_0}$ is an outer function (cf. [Rud, Ch. 17]), we get:

$$(16) \quad \varphi_j(\lambda)(1 - \bar{\alpha}_0 \lambda)^{1/p_0} = r_j^{1/2p_0} e^{i\theta_j} B_j(\lambda) S_j(\lambda) (1 - \bar{\alpha}_j \lambda)^{1/p_0}, \quad \lambda \in E,$$

where

$$\begin{aligned} \theta_j &\in \mathbb{R}, \\ B_j(\lambda) &:= \begin{cases} \frac{\lambda - \alpha_j}{1 - \bar{\alpha}_j \lambda}, & j = 1, \dots, s \\ 1, & j = s+1, \dots, n \end{cases}, \\ S_j(\lambda) &:= \exp \left(- \int_{-\pi}^{\pi} \frac{e^{i\theta} + \lambda}{e^{i\theta} - \lambda} d\sigma_j(\theta) \right) \end{aligned}$$

and where σ_j is a singular non-negative Borel measure, $j = 1, \dots, n$.

Now, in order to complete the proof of the lemma, we only need to show that $\sigma_1 = \dots = \sigma_n = 0$. First, observe that, in view of (12), (15) and (16), we have:

$$(17) \quad |\lambda - \alpha_j|^{p_0} |1 - \bar{\alpha}_j \lambda|^{2p_0-1} |1 - \bar{\alpha}_0 \lambda| |S_j(\lambda)|^{-p_0} \leq M_j, \quad \lambda \in E,$$

for some $M_j > 0$, $j = 1, \dots, n$. On the other hand, by [Gar, Ch. II],

$$S_j^*(\lambda) = 0 \quad \text{for } \sigma_j\text{-almost all } \lambda \in \partial E, \quad j = 1, \dots, n.$$

Moreover, for any $\beta \in \mathbb{R}$, $b > 0$, the function

$$E \ni \lambda \rightarrow |\lambda - 1|^\beta \exp \left(b \frac{1 - |\lambda|^2}{|\lambda - 1|^2} \right)$$

is unbounded. Consequently, condition (17) is fulfilled only when $\sigma_j = 0$ for all $j = 1, \dots, n$. $\square$

PROOF OF THE UNIQUENESS IN THEOREM 1. Let $\varphi, \psi : E \rightarrow \mathcal{E}(p)$ be two complex geodesics with

$$(18) \quad \varphi(0) = \psi(0) \quad \text{and} \quad \varphi'(0) = \psi'(0).$$

$$(18') \quad \varphi(0) = \psi(0) \quad \text{and} \quad \varphi(\sigma) = \psi(\sigma) \quad \text{with} \quad 0 < \sigma < 1,$$

respectively.

So far, Theorem 1 (see also Remark (a) after Theorem 1) shows that: $\varphi_j \equiv 0$ if and only if $\psi_j \equiv 0$. Thus, without loss of generality, we assume that φ and ψ satisfy (1). Moreover, we assume that φ fulfils condition (2). Put $I_0 := \{j : \psi_j \text{ has a zero in } E\}$.

Now, observe that $\chi := (\varphi + \psi)/2$ is again a complex geodesic in $\mathcal{E}(p)$ ($\mathcal{E}(p)$ is convex!). In particular:

$$\chi(\lambda), \varphi(\lambda), \psi(\lambda) \in \partial\mathcal{E}(p) \quad \text{for all } \lambda \in \partial E.$$

Therefore we obtain for $\lambda \in \partial E$:

$$(19) \quad \begin{aligned} \arg \varphi_j(\lambda) &= \arg \psi_j(\lambda) = \arg \chi_j(\lambda) \quad \text{if } \varphi_j(\lambda)\psi_j(\lambda) \neq 0 \quad \text{and} \\ |\varphi_j(\lambda)| &= |\psi_j(\lambda)| = |\chi_j(\lambda)| \quad \text{if } p_j > 1/2. \end{aligned}$$

Hence, if $p_j > 1/2$ then $\varphi_j = \psi_j$ on ∂E and so $\varphi_j \equiv \psi_j$ on E .

It remains to discuss j with $p_j = 1/2$. We fix such an j . Because of (19) we have

$$(20) \quad \varphi_j \bar{\psi}_j = \psi_j \bar{\varphi}_j \quad \text{on } \partial E.$$

First let us assume that $1 \leq j \leq s$, i.e.

$$\varphi_j(\lambda) = a_j \frac{(\lambda - \alpha_j)(1 - \bar{\alpha}_j \lambda)}{(1 - \bar{\alpha}_0 \lambda)^2}.$$

Case $j \in I_0$, i.e.

$$\psi_j(\lambda) = b_j \frac{(\lambda - \beta_j)(1 - \bar{\beta}_j \lambda)}{(1 - \bar{\beta}_0 \lambda)^2}.$$

An application of (20) leads to

$$a_j \bar{b}_j \frac{1}{(1 - \bar{\alpha}_0 \lambda)^2 (\lambda - \beta_0)^2} = \bar{a}_j b_j \frac{1}{(1 - \bar{\beta}_0 \lambda)^2 (\lambda - \alpha_0)^2}, \quad \lambda \in \partial E.$$

So we obtain $\alpha_0 = \beta_0$. Moreover, because of (18) and (18'), we get

$$\begin{aligned} \alpha_j a_j &= \beta_j b_j \quad \text{and} \quad a_j(1 + |\alpha_j|^2) = b_j(1 + |\beta_j|^2), \\ \alpha_j a_j &= \beta_j b_j \quad \text{and} \quad a_j(1 - \bar{\alpha}_j \sigma + |\alpha_j|^2) = b_j(1 - \bar{\beta}_j \sigma + |\beta_j|^2), \quad \text{respectively.} \end{aligned}$$

Hence

$$\Phi(0, \alpha_j) = \Phi(0, \beta_j), \quad \text{resp.} \quad \Phi(\sigma, \alpha_j) = \Phi(\sigma, \beta_j),$$

where

$$\Phi(\tau, \lambda) := \frac{\lambda}{1 - \tau \bar{\lambda} + |\lambda|^2}, \quad \lambda \in \bar{E}, \quad 0 \leq \tau < 1.$$

Since $\Phi(\tau, \cdot)$ is injective on $\bar{E}$, we obtain $\alpha_j = \beta_j$ and so $a_j = b_j$.

Case $j \notin I_0$, i.e.

$$\psi_j(\lambda) = b_j \left(\frac{1 - \bar{\beta}_j \lambda}{1 - \bar{\beta}_0 \lambda} \right)^2.$$

Then it follows from (20) that

$$\bar{b}_j a_j \frac{1}{(1 - \bar{\alpha}_0 \lambda)^2} \left(\frac{\lambda - \beta_j}{\lambda - \beta_0} \right)^2 \equiv \bar{a}_j b_j \left(\frac{1 - \bar{\beta}_j \lambda}{1 - \bar{\beta}_0 \lambda} \right)^2 \frac{1}{(\lambda - \alpha_0)^2}$$

as meromorphic functions on $\mathbb{C}$.

So we can conclude that $\alpha_0 = \beta_0$ and so

$$\bar{b}_j a_j (\lambda - \beta_j)^2 \equiv \bar{a}_j b_j (1 - \bar{\beta}_j \lambda)^2.$$

Hence we have $|\beta_j| = 1$. Again using (18) and (18') gives:

$$\alpha_j a_j = b_j \quad \text{and} \quad a_j (1 + |\alpha_j|^2) = -2b_j \bar{\beta}_j$$

$$\alpha_j a_j = b_j \quad \text{and} \quad a_j (1 - \bar{\alpha}_j \sigma + |\alpha_j|^2) = -b_j (2\bar{\beta}_j - \bar{\beta}_j^2 \sigma), \quad \text{respectively.}$$

Then

$$\Phi(0, \alpha_j) = \Phi(0, \beta_j), \quad \text{respectively.} \quad \Phi(\sigma, \alpha_j) = \Phi(\sigma, \beta_j)$$

contradiction.

The remaining case $s < j$ can be solved using similar reasonings as above.

□

Acknowledgment

The first two authors would like to express their gratitude to the Jagiellonian University (Kraków) and the University of Osnabrück (Vechta) for their hospitality during this research work.

Added in proof

See also S. Dineen & R.M. Timoney, *Complex geodesics on convex domains* in "Progress in Functional Analysis", K.D. Bierstedt, J. Bonet, J. Horváth & M. Maestre (eds.), Elsevier Science Publishers B.V., 1992.

REFERENCES

- [Aba] M. ABATE, *Iteration Theory of Holomorphic Maps on Taut Manifolds*, Mediterranean Press, 1989.
- [BFKKMP] B.E. BLANK - D. FAN - D. KLEIN - S.G. KRANTZ - D. MA - M.-Y. PANG, *The Kobayashi metric of a complex ellipsoid in $\mathbb{C}^2$* , preprint (1991).
- [Din] S. DINEEN, *The Schwarz Lemma*, Clarendon Press, Oxford, 1989.
- [Gar] J. GARNETT, *Bounded Analytic Functions*, Academic Press, New York, 1981.
- [Gen] G. GENTILI, *Regular complex geodesics in the domain $D_n = \{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_1| + \dots + |z_n| < 1\}$* , Springer Lecture Notes in Math. **1275** (1987), 235-252.
- [Jar-Pfl] M. JARNICKI - P. PFLUG, *Invariant Distances and Metrics in Complex Analysis*, W. de Gruyter & Co. (to appear).
- [Lem] L. LEMPert, *Holomorphic retracts and intrinsic metrics in convex domains*, Anal. Math. **8** (1982), 257-261.
- [Pol] E.A. POLETSKIĬ, *The Euler-Lagrange equations for extremal holomorphic mappings of the unit disc*, Michigan Math. J. **30** (1983), 317-333.
- [Roy-Won] H. ROYDEN, P.-M. WONG, *Carathéodory and Kobayashi metric on convex domains*, preprint (1983).
- [Rud] W. RUDIN, *Real and Complex Analysis*, McGraw-Hill, 1974.
- [Ves] E. Vesentini, *Complex geodesics*, Compositio Math. **44** (1981), 375-394.

Uniwersytet Jagielloński
 Instytut Matematyki
 30-059 Kraków
 Reymonta 4
 Poland

Universität Osnabrück
 Standort Vechta,
 Fachbereich Naturwissenschaften
 Mathematik
 Postfach 15 53, D-49364 Vechta
 Germany