ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

LEV AIZENBERG
LAWRENCE ZALCMAN

Instability phenomena for the moment problem

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4e série, tome 22, nº 1 (1995), p. 95-107
<http://www.numdam.org/item?id=ASNSP_1995_4_22_1_95_0>

© Scuola Normale Superiore, Pisa, 1995, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (http://www.sns.it/it/edizioni/riviste/annaliscienze/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Instability Phenomena for the Moment Problem

LEV AIZENBERG - LAWRENCE ZALCMAN

Suppose K is a compact set in $\mathbb{C}^n$ or $\mathbb{R}^n$, and let μ be a finite complex Borel measure on K. In this paper we show, under appropriate conditions on K, that if the analytic or harmonic moments of μ decrease sufficiently rapidly (or grow sufficiently slowly) in a certain precise sense dependent on K, then these moments vanish identically. In the most favorable cases, it is then possible to conclude that $\mu = 0$. This phenomenon does not seem to have been noticed previously, even in the classical case of the power moment problem for a finite interval in $\mathbb{R}$.

In the sequel all measures are Borel.

1. - Holomorphic moments, $n = 1$

We begin with a discussion of the situation for $n = 1$.

THEOREM 1. *Let K be a compact set in the plane which does not contain the origin, and let μ be a finite complex measure on K with moments*

$$a_j = \int_K \frac{d\mu(\varsigma)}{\varsigma^{j+1}} \quad j = 0, 1, 2, \cdots. \tag{1}$$

If

$$\overline{\lim_{j\to\infty}} \sqrt[j]{|a_j|} < \frac{1}{\max_K |z|} \tag{2}$$

and K does not separate 0 from ∞ (i.e., 0 belongs to the unbounded component of $\mathbb{C}\backslash K$), then $a_j = 0$ for $j = 0, 1, 2, \cdots$.

If K does separate 0 from ∞, then for each sequence $\{a_j\}$ satisfying (2) *there is a measure μ on K having $\{a_j\}$ as its moment sequence, i.e., such that* (1) *holds.*

Pervenuto alla Redazione il 23 Giugno 1994.

PROOF. Suppose that K does not separate 0 and ∞. Let

$$F(z) = \int_K \frac{d\mu(\zeta)}{\zeta - z}.$$

Clearly, F is analytic off K. Expanding F in a power series about 0, we find

$$F(z) = \sum_{j=0}^{\infty} a_j z^j,$$

where the a_j are given by (1). If (2) holds, this series converges uniformly on an open disc containing K. Thus, the restriction of F to the unbounded component of $\mathbb{C}\backslash K$ extends to be analytic on the entire complex plane. Since $F(\infty) = 0$, F vanishes identically, so $a_j = 0$ for $j = 0, 1, 2, \cdots$.

Now suppose that K separates 0 from ∞. Denote by U the unbounded component of $\hat{\mathbb{C}}\backslash K$ and consider the space A of all continuous functions on $K \cup U$ which are analytic on U. By the maximum modulus principle, $\|f\|_A = \max_K |f(z)|$ defines a norm on A under which it is identified with a closed subspace of $C(K)$. Let $\{a_n\}$ be a sequence which satisfies (2), so that

$$R = \frac{1}{\overline{\lim_{j\to\infty}} \sqrt[j]{|a_j|}} > \max_K |z|$$

Then the function $\varphi(z) = \sum_{n=0}^{\infty} a_n z^n$ is analytic on $\{z : |z| < R\}$, and we may choose ρ so that $\max_K |z| < \rho < R$. Now set

$$L(f) = \frac{1}{2\pi i} \int_\Gamma f(z)\varphi(z)dz \tag{3}$$

where Γ is the positively oriented circle of radius ρ about the origin. Clearly,

$$|L(f)| \leq \rho \max_\Gamma |\varphi(z)| \max_K |f(z)| \leq C(\rho, \varphi)\|f\|_A.$$

Thus L defines a continuous linear functional on A, which (by the Hahn-Banach Theorem) extends to all of $C(K)$. Thus there exists a finite complex measure μ on K such that

$$L(f) = \int_K f d\mu \quad f \in A. \tag{4}$$

On the other hand, we have

$$a_j = \frac{1}{2\pi i} \int_\Gamma \frac{\varphi(z)}{z^{j+1}} dz.$$

Since the functions $f_j(z) = z^{-(j+1)}$, $j = 0, 1, 2, \cdots$, all belong to A, (3) yields $L(f_j) = a_j$. It then follows from (4) that the measure μ has moment sequence $\{a_j\}$, i.e., that (1) holds. □

Note, in particular, that one may choose $\{a_j\}$ such that (2) holds but $a_j \neq 0$ for all j.

An analogous reasoning, based on expanding the function $F(z)$ in a Laurent series about the point $z = \infty$, yields:

THEOREM 1′. *Let K be a compact set in the plane which does not contain the origin, and let μ be a finite complex measure on K with moments*

$$b_j = \int_K \varsigma^j d\mu(\varsigma) \quad j = 0, 1, 2, \cdots. \tag{1'}$$

If

$$\overline{\lim_{j\to\infty}} \sqrt[j]{|b_j|} < \min_K |z| \tag{2'}$$

and K does not separate 0 from ∞, then $b_j = 0$ for all $j = 0, 1, 2, \cdots$.

If K does separate 0 from ∞, then for each sequence $\{b_j\}$ satisfying (2′) there is a measure μ on K having $\{b_j\}$ as its moment sequence, i.e., such that (1′) holds.

COROLLARY 1. *Suppose K has empty interior, $\mathbb{C}\backslash K$ is connected, and $0 \notin K$. Then (2) and (2′) each imply that $\mu = 0$.*

PROOF. Suppose that (2′) holds. Then, by Theorem 1′, all analytic moments (1′) vanish. Taking linear combinations shows that μ is orthogonal to all analytic polynomials. By Mergelyan's Theorem, any function in $C(K)$ can be uniformly approximated on K by such polynomials. Thus μ annihilates all elements of $C(K)$ and hence vanishes identically. If (2) holds, the proof of Theorem 1 shows that $F(z)$ vanishes identically. Thus the coefficients of its Laurent expansion about ∞ (given by (1′)) are identically zero, so again $\mu = 0$. □

When K is an interval that does not contain the origin, it is possible to strengthen Corollary 1 under the assumption that μ does not place any mass at the (relevant) endpoint of $[a, b]$. Specifically, we have

COROLLARY 2. *Let μ be a finite complex measure on $[a, b] \subset \mathbb{R}$, where $0 < a < b$. If either $\mu(\{b\}) = 0$ and*

$$\overline{\lim_{j\to\infty}} \left(\left| \int_a^b \frac{d\mu(t)}{t^{j+1}} \right| \right)^{1/j} \leq \frac{1}{b}$$

or $\mu(\{a\}) = 0$ *and*

$$\overline{\lim_{j\to\infty}} \left(\left| \int_a^b t^j d\mu(t) \right| \right)^{1/j} \leq a,$$

then $\mu = 0$.

PROOF. Suppose

$$\overline{\lim_{j\to\infty}} \left(\left| \int_a^b \frac{d\mu(t)}{t^{j+1}} \right| \right)^{1/j} \leq \frac{1}{b}.$$

Let

$$F(z) = \frac{1}{2\pi i} \int_a^b \frac{d\mu(t)}{t-z},$$

so that $F(z)$ is analytic on $\mathbb{C}\backslash\{b\}$. For $y > 0$ we have

$$F(x+iy) - F(x-iy) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y d\mu(t)}{(x-t)^2 + y^2} = (P_y * \mu)(x),$$

where P_y is the Poisson kernel for the upper half plane. Since $(P_y * \mu)(x)dx$ converges weak$*$ to μ as $y \to 0$, we have

$$\lim_{y\to 0} \int_{-\infty}^{\infty} f(x)[F(x+iy) - F(x-iy)]dx = \int_a^b f(x)d\mu(x)$$

for any function $f \in C_0(\mathbb{R})$.

Now fix $c \in (a,b)$ and let $\varphi \in C_0(\mathbb{R})$ be supported in $[0,c]$. We have

$$\int_a^c \varphi(x)d\mu(x) = \lim_{y\to 0} \int_0^c \varphi(x)[F(x+iy) - F(x-iy)]dx = 0$$

since F is analytic on $[0,c]$. Taking the sup over all such φ satisfying $|\varphi(x)| \leq 1$, we obtain $|\mu|([a,c)) = 0$. It follows that $|\mu|([a,b)) = 0$; and, since $\mu(\{b\}) = 0$, $\mu = 0$. □

This result applies in particular to absolutely continuous measures, i.e., functions in $L^1([a,b])$.

REMARK. The first part of Theorems 1 and 1′ hold not only for measures but for distributions of compact support and for analytic functionals as well; the proofs remain the same. As a consequence of this, we have:

COROLLARY 3. *Let f be an entire function such that*

$$|f(z)| \leq \gamma(1+|z|)^N e^{r|\operatorname{Im} z|}, \quad z \in \mathbb{C}. \tag{5}$$

Suppose that for some $\rho > 0$ one has

$$\overline{\lim_{j\to\infty}} \sqrt[j]{\left|\sum_{k=0}^{j} \frac{j!}{k!(j-k)!} \frac{f^{(k)}(0)}{[-i(r+\rho)]^k}\right|} < \frac{\rho}{r+\rho}. \tag{6}$$

Then $f \equiv 0$.

If, in addition, $f \in L^2(\mathbb{R})$ (i.e., f is in the Wiener class [A1, p. 166]*), we may relax the inequality in* (6) *to allow equality.*

PROOF. If (5) holds, then $f(z) = u(e^{-ixz})$, where u is a distribution supported on $[-r, r]$, cf. [Ru, p. 183]. Now

$$f(z)e^{-i(r+\rho)z} = u(e^{-i(x+r+\rho)z}) = u_1(e^{-ixz}),$$

where u_1 is a distribution whose support lies in $[\rho, \rho+2r]$. It follows that

$$\begin{aligned} u_1(x^j) &= \frac{1}{(-i)^j} \frac{d^j}{dz^j} [f(z)e^{-i(r+\rho)z}]\Big|_{z=0} \\ &= i^j \sum_{k=0}^{j} \frac{j!}{k!(j-k)!} f^{(k)}(0)[-i(r+\rho)]^{j-k}. \end{aligned}$$

By Corollary 1 (and the previous Remark), it follows that $u_1 = 0$ and hence $f \equiv 0$. When $f \in L^2(\mathbb{R})$, the distribution u_1 is a function in $L^2([\rho, \rho+2r])$, so the result follows from Corollary 2. □

2. - Holomorphic moments, $n > 1$

We shall follow the conventional notations for multi-indices. Thus, for $\alpha = (\alpha_1, \alpha_2, \ldots, \alpha_n)$ an n-tuple of non-negative integers and $z = (z_1, z_2, \ldots, z_n) \in \mathbb{C}^n$ we shall write $|\alpha| = \alpha_1 + \alpha_2 + \cdots + \alpha_n$ and $z^\alpha = z_1^{\alpha_1} z_2^{\alpha_2} \cdots z_n^{\alpha_n}$. For K a compact set in $\mathbb{C}^n$, set $d_\alpha(K) = \max_K |z^\alpha|$ and $\tilde{K} = \bigcup_{j=1}^{n} \{w : \Gamma_{z_j} \cap K \neq \emptyset\}$, where $\Gamma_{z_j} = \{w \in \mathbb{C}^n : w_j = z_j\}$. Put $\hat{\mathbb{C}}^n = \hat{\mathbb{C}} \times \hat{\mathbb{C}} \times \cdots \times \hat{\mathbb{C}}$ (n times).

THEOREM 2. *Let K be a compact set in $\mathbb{C}^n$ such that $\varsigma_j \neq 0$, $j = 1, \ldots, n$, for $\varsigma \in K$ and the points 0 and $(\infty, \ldots, \infty)$ belong to the same connected component of $\hat{\mathbb{C}}^n \backslash \tilde{K}$. Let μ be a finite complex measure on K with moments*

$$a_\alpha = \int_K \frac{d\mu}{\varsigma^{\alpha+I}}, \tag{7}$$

where $I = (1, \ldots, 1)$.

If

$$\overline{\lim_{|\alpha|\to\infty}} \sqrt[|\alpha|]{|\alpha_\alpha| d_\alpha(K)} < 1, \tag{8}$$

then $a_\alpha = 0$ *for all* $\alpha \in (\mathbb{Z}^+)^n$.

PROOF. Let

$$F(z) = \int_K \frac{d\mu(\zeta)}{(\zeta_1 - z_1)\dots(\zeta_n - z_n)}.$$

Clearly, F is analytic on $\hat{\mathbb{C}}^n \backslash \tilde{K}$. Expanding F in a power series about 0, we find

$$F(z) = \sum_{|\alpha|\geq 0} a_\alpha z^\alpha,$$

where the a_α are given by (7). Since (8) holds, the series converges uniformly on a complete Reinhardt domain $\mathcal{D}$ containing K ([AM] cf. [F, pp. 49-51]). Denote by U the component of $\hat{\mathbb{C}}^n \backslash \tilde{K}$ containing 0 and $(\infty, \dots, \infty)$. Clearly, F is analytic on $\mathcal{D} \cup U$. In particular, it is analytic on the union of n polydiscs about the origin, the j-th of which has j-radius equal ∞. Since the envelope of holomorphy of this union is clearly all of $\mathbb{C}^n$, the series for F converges everywhere and thus defines an entire function. But

$$\lim_{|z|\to\infty} F(z) = 0.$$

Thus F vanishes identically, so $a_\alpha = 0$ for all α such that $\alpha_j \geq 0$, $j = 1, 2, \dots, n$. □

COROLLARY 4. *Let K and μ be as in Theorem 2. If every function in $C(K)$ is uniformly approximable on K by polynomials in $1/z_1, \dots, 1/z_n$, then* (8) *implies* $\mu = 0$.

PROOF. According to Theorem 2, all $a_\alpha = 0$. Consequently, for every polynomial P one has

$$\int_K P\left(\frac{1}{z_1}, \dots, \frac{1}{z_n}\right) d\mu = 0.$$

Thus, for every $\varphi \in C(K)$,

$$\int_k \varphi d\mu = 0,$$

so $\mu = 0$. □

When K is a subset of the real subspace $\mathbb{R}^n$ of $\mathbb{C}^n$ which does not intersect the coordinate planes, the approximation condition of Corollary 4 holds (by the Stone Weierstrass Theorem). Thus we have:

COROLLARY 5. *Let K be a compact set in $\mathbb{R}^n \subset \mathbb{C}^n$ which does not intersect the coordinate planes. If* (8) *holds, then $\mu = 0$.*

PROOF. In this case, the projection of K onto each complex coordinate plane does not separate 0 and ∞ there. Thus, the points 0 and $(\infty, \dots, \infty)$ belong to the same connected component of $\hat{\mathbb{C}}^n \backslash \tilde{K}$, so the hypotheses of Theorem 2 and Corollary 4 hold, and hence $\mu = 0$. □

In analogy with Corollary 3, we have also:

COROLLARY 6. *Let f be an entire function in $\mathbb{C}^n$ such that*

$$|f(z)| \leq \gamma(1 + |z|)^N e^{r|\operatorname{Im} z|}, \quad z \in \mathbb{C}^n.$$

Suppose that for some $\rho > 0$ we have

$$\overline{\lim_{|\alpha| \to \infty}} \left(d_\alpha(B) \left| \sum_{\substack{\beta_k = 0 \\ k=1,\dots,n}}^{\alpha_k} \frac{\alpha!}{\beta!(\alpha - \beta)!} \frac{\partial^{|\beta|} f(0)}{\partial z_n^{\beta_n} \cdots \partial z_1^{\beta_1}} [-i(r + \rho)]^{|\alpha - \beta|} \right| \right)^{1/|\alpha|} < 1, \tag{9}$$

where

$$B = \left\{ x \in \mathbb{R}^n \subset \mathbb{C}^n : \left| \frac{1}{x_1} - r - \rho \right|^2 + \cdots + \left| \frac{1}{x_n} - r - \rho \right|^2 \leq r^2 \right\}$$

and $\alpha! = \alpha_1! \dots \alpha_n!$. Then $f \equiv 0$.

If, in addition, $f \in L^2(\mathbb{R}^n)$, we may relax the inequality in (9) *to allow equality.*

Denote by J the collection of 2^n vectors of the form $p = (p_1, p_2, \dots, p_n)$ where $p_j = \pm 1$ for each j. A vector $p \in J$ operates on a point $z = (z_1, \dots, z_n)$ with nonzero coordinates via $p(z) = (z_1^{p_1}, z_2^{p_2}, \dots, z_n^{p_n})$. Let pK be the image of K under this mapping and set

$$a_\alpha^p = \int_{pK} \frac{d(\mu \circ p^{-1})}{\varsigma^{\alpha + I}}.$$

The following extension of Theorem 2 obtains.

THEOREM 3. *Let K and μ be as in Theorem* 2 *with $\mathbb{C}^n \backslash \tilde{K}$ connected. If*

$$\min_{p \in J} \overline{\lim_{|\alpha| \to \infty}} \sqrt[|\alpha|]{|a_\alpha^p| d_\alpha(pK)} < 1 \tag{10}$$

holds in place of (8), *then all the moments $a_\alpha^p = 0$ (so that $a_\beta = 0$ for all $\beta \in \mathbb{Z}^n$).*

COROLLARY 7. *If, in addition, every function in $C(K)$ is uniformly approximable on K by polynomials in $z_1, \dots, z_n$ and $1/z_1, \dots, 1/z_n$, then* (10) *implies $\mu = 0$.*

Denoting by $R(K)$ the uniform closure on K of the rational functions which are holomorphic on K, we see that when K satisfies the conditions of Corollary 7, one has $C(K) = R(K)$. It does not seem easy to find general conditions which insure this. For instance, one may have $R(K) \neq C(K)$ even when K is polynomially convex and contains no ordinary analytic structure.

EXAMPLE. Let X be a Swiss Cheese [Z1, pp. 69-70], i.e., a compact set in $\mathbb{C}$ without interior such that $R(X) \neq C(X)$. It is well-known that there exists $\varphi \in R(X)$ such that z and φ generate $R(X)$, i.e., polynomials in z and φ are uniformly dense in $R(X)$. (This is the Bishop-Hoffman Theorem; for the proof, cf. [Ro, Theorem 3.6].) We may clearly choose X and φ so that 0 belongs to the unbounded component of X and $\varphi(z) \neq 0$ on X. Now set $\Phi(z) = (z, \varphi(z))$ and put $K = \Phi(X)$. Evidently, K contains no analytic discs. Let $P(K)$ be the uniform algebra of functions uniformly approximable on K by polynomials. Now $\Phi^* F = F \circ \Phi$ defines an algebra isomorphism of $C(K)$ onto $C(X)$ which maps $P(K)$ onto $R(X)$. Since $R(X) \neq C(X)$ we have $P(K) \neq C(K)$. On the other hand, since the spectrum of the Banach algebra $R(X)$ is X, the spectrum of $P(K)$ is K, i.e., K is polynomially convex. Hence, by the Oka-Weil Theorem, $R(K) = P(K)$, so that $R(K) \neq C(K)$.

3. - Harmonic moments

In discussing harmonic moments, it will be convenient to consider the cases $n = 2$ and $n \geq 3$ separately. We begin with $n \geq 3$. Denote by B_1 the open unit ball in $\mathbb{R}^n$ and by ∂B_1 its boundary, the unit sphere. Let $\{P_{j,s}\}$ be an orthonormal basis of homogeneous harmonic polynomials in $L^2(\partial B_1)$, where j is the degree of $P_{j,s}$ and $s = 1, \dots, \sigma(j, n)$.

THEOREM 4. *Let K be a compact set in $\mathbb{R}^n$ $(n \geq 3)$ which does not contain the origin, and let μ be a finite complex measure on K with moments*

$$a_{j,s} = \int_K \frac{\overline{P_{j,s}(x)} d\mu(x)}{|x|^{n+2j-2}}. \tag{11}$$

If

$$\overline{\lim_{j \to \infty}} \max_s \sqrt[j]{|a_{j,s}|} < \frac{1}{\max_K |x|}, \tag{12}$$

and K does not separate 0 from ∞, then $a_{j,s} = 0$ for all j, s.

If K does separate 0 from ∞, then for each collection of numbers $\{a_{j,s}\}$ satisfying (12) *there is a measure on K such that* (11) *holds.*

PROOF. Consider the Newtonian potential

$$F(y) = \int_K \frac{d\mu(x)}{|x-y|^{n-2}}.$$

Clearly, F is harmonic on $\mathbb{R}^n \backslash K$. For the fundamental solution of the Laplace equation in $\mathbb{R}^n$ ($n \geq 3$), one has the expansion ([D], cf. [A1, Lemmas 36.3 and 38.5])

$$\frac{1}{|x-y|^{n-2}} = \frac{1}{|x|^{n-2}} + \Omega_n(n-2) \sum_{j=1}^{\infty} \sum_{s=1}^{\sigma(j,n)} \frac{P_{j,s}(y)\overline{P_{j,s}(x)}}{(n+2j-2)|x|^{n+2j-2}}, \tag{13}$$

where Ω_n is the area of the unit sphere ∂B_1, and the series on the right-hand side of (13) converges uniformly together with all derivatives on compact subsets of the cone $\{(x,y) \in \mathbb{R}^{2n} : |y| < |x|\}$. Expanding F in a series of homogeneous polynomials in a neighborhood of 0, we find

$$F(y) = \int_K \frac{d\mu(x)}{|x|^{n-2}} + \Omega_n(n-2) \sum_{j=1}^{\infty} \sum_{s=1}^{\sigma(j,n)} \frac{a_{j,s}}{n+2j-2} P_{j,s}(y),$$

where the $a_{j,s}$ are given by (11). If (12) holds, then the series converges uniformly on some ball B_R with radius $R > \max_K |x|$. Then the restriction of F to the unbounded component of $\mathbb{R}^n \backslash K$ extends to be harmonic on all of $\mathbb{R}^n$. Since $F(\infty) = 0$, it follows that $F \equiv 0$, so $a_{j,s} = 0$ for all j, s.

Now suppose that K separates 0 from ∞. Denote by U the unbounded component of $(\mathbb{R}^n \cup \{\infty\}) \backslash K$ and consider the space H of all continuous functions on $K \cup U$ which are harmonic on U. By the maximum principle, $\|f\|_H = \max_K |f(x)|$ defines a norm on H under which H is (identified with) a closed subspace of $C(K)$. Now suppose that the $a_{j,s}$ satisfy (12), so that

$$R = \frac{1}{\overline{\lim_{j\to\infty}} \max_s \sqrt[j]{|a_{j,s}|}} > \max_K |x|.$$

Choose ρ such that $\max_K |x| < \rho < R$ and set

$$\varphi(x) = \sum_{j,s} a_{j,s}\rho^{-1}P_{j,s}(x). \tag{14}$$

This series converges uniformly on each ball $\overline{B}_r$, $r < R$, and defines a harmonic

function on B_R. Put

$$L(f) = \int_{\partial B_\rho} f(x)\varphi(x)d\sigma. \tag{15}$$

Clearly,

$$L(f) \leq \Omega_n \rho^{n-1} \max_{\partial B_\rho} |\varphi(x)| \max_K |f(x)| \leq C(\rho, \varphi)\|f\|.$$

Thus L defines a continuous linear functional on H, which (by the Hahn-Banach Theorem) extends to all of $C(K)$. Thus there exists a finite complex measure μ on K such that

$$L(f) = \int_K f d\mu \quad f \in H. \tag{16}$$

Now the functions

$$f_{j,s}(x) = \frac{\overline{P_{j,s}(x)}}{|x|^{n+2j-2}} \quad s = 1, 2, \cdots, \sigma(j, n) \quad j = 0, 1, 2, \cdots,$$

obtained by applying the Kelvin transformation with respect to ∂B_1 to $P_{j,s}(x)$, all belong to H. Thus by (14) and (15) we have

$$\begin{aligned}
L(f_{j,s}) &= \int_{\partial B_\rho} \frac{\overline{P_{j,s}(x)}}{|x|^{n+2j-2}} \varphi(x)d\sigma \\
&= \int_{\partial B_\rho} \sum_{k,t} \frac{a_{k,t}}{\rho} \frac{1}{|x|^{n+2j-2}} \overline{P_{j,s}(x)} P_{k,t}(x)d\sigma \\
&= \int_{\partial B_1} \sum_{k,t} \frac{a_{k,t}}{\rho} \frac{1}{\rho^{n+2j-2}} \rho^j \overline{P_{j,s}(y)} \rho^k P_{k,t}(y) \rho^{n-1} d\sigma \\
&= \sum_{k,t} a_{k,t} \rho^{k-j} \int_{\partial B_1} \overline{P_{j,s}(y)} P_{k,t}(y) d\sigma \\
&= a_{j,s}
\end{aligned}$$

by the orthonormality of $\{P_{j,s}\}$ on ∂B_1. It now follows from (16) that μ has the $a_{j,s}$ as its harmonic moments, i.e., that (11) holds. □

In analogy with Theorem 1′ we have the following analogue of Theorem 4.

THEOREM 4′. *Let K be a compact set in $\mathbb{R}^n$ ($n \geq 3$) which does not contain the origin and let μ be a finite complex measure on K with moments*

$$a_{j,s} = \int_K P_{j,s}(x)d\mu(x). \tag{11′}$$

If

$$\overline{\lim_{j\to\infty}} \max_s \sqrt[j]{|a_{j,s}|} < \min_K |x|, \tag{12′}$$

and K does not separate 0 from ∞, then all the moments $a_{j,s} = 0$.

If K does separate 0 from ∞, then for each collection $\{a_{j,s}\}$ of numbers satisfying (12′) there is a measure on K such that (11′) holds.

The two-dimensional analogues of Theorems 4 and 4′ are also valid. Here one has $\sigma(j,2) = 2$ for all $j \geq 1$. One may take $P_{j,1}(x_1,x_2) = \mathrm{Re}(x_1 + ix_2)^j$ and $P_{j,2}(x_1,x_2) = \mathrm{Im}(x_1 + ix_2)^j$, up to a normalizing constant. Writing $x_1 + ix_2 = z$, one considers, in place of the Newtonian potential, the logarithmic potential

$$F(z) = \int_K \log|\varsigma - z|d\mu(\varsigma),$$

which is again harmonic on $\mathbb{R}^2\backslash K$. If (12) or (12′) holds, the restriction of F to the unbounded component of $\mathbb{R}^2\backslash K$ extends to be harmonic on all of $\mathbb{R}^2$. Evidently, $|F(z)| \leq C \log|z|$ for large $|z|$. It follows from a version of Liouville's Theorem [Bu, Corollary 6.33] that F is constant. To see that $F = 0$, observe that

$$\begin{aligned} F(\infty) = \lim_{z\to\infty} F(z) &= \lim_{z\to\infty} \int_K \log|\varsigma - z|d\mu(\varsigma) \\ &= \lim_{z\to\infty} \int_K \log|z|d\mu(\varsigma) + \lim_{z\to\infty} \int_K \log\left|1 - \frac{\varsigma}{z}\right| d\mu(\varsigma) \\ &= \mu(K) \lim_{z\to\infty} \log|z|, \end{aligned}$$

since $\log\left|1 - \frac{\varsigma}{z}\right|$ tends uniformly to 0 on K as $z \to \infty$. But the left hand side is finite, so we must have $\mu(K) = 0$. Thus $F(\infty) = 0$ and F vanishes identically.

Denote by $h(K)$ the hull of K, i.e., the union of K with all the bounded components of its complement. Recall ([L, p. 307]) that a set E is *thin* at $x_0 \in \mathbb{R}^n$ if either E does not have x_0 as a limit point or there exists a function v superharmonic on $\mathbb{R}^n$ such that

$$v(x_0) < \varliminf_{x\to x_0} v(x) \quad x \in E\backslash\{x_0\}. \tag{17}$$

COROLLARY 8. *Suppose that* $\operatorname{int} K = \emptyset$ *and* $\mathbb{R}^n \backslash h(K)$ *is not thin at any point of* K. *Then if* $0 \notin K$ *and* K *does not separate* 0 *from* ∞, (12) *and* (12′) *each imply that* $\mu \equiv 0$.

This follows from Theorem 4 and 4′ together with the fact ([Br], cf. [D]) that in this case every function in $C(K)$ is the uniform limit on K of harmonic polynomials.

As an immediate consequence we have:

COROLLARY 9. *Suppose that* K *has zero Lebesgue measure in* $\mathbb{R}^n$, $\mathbb{R}^n \backslash K$ *is connected, and* $0 \notin K$. *Then* (12) *and* (12′) *each imply that* $\mu \equiv 0$.

Indeed, since $\mathbb{R}^n \backslash K$ is connected, $h(K) = K$. But $\mathbb{R}^N \backslash K$ is nowhere thin if K has Lebesgue measure 0, since otherwise (17) would be inconsistent with the super-mean-value property of superharmonic functions for balls around x_0.

4. - Final comments

This work is in large measure a continuation of [A2], where problems of Morera type were considered for non-closed curves and pieces of hypersurfaces. For further information on such problems see [Z2], [Z3], and [BCPZ]. Additional motivation for the questions considered here is in [AR] and the papers listed there.

We are grateful to Paul Gauthier for enlightening discussions on harmonic approximation and to T.W. Gamelin for providing the reference [Ro].

REFERENCES

[A1] L. AIZENBERG, *Carleman's Formulas in Complex Analysis*. Kluwer Acad. Publ., Dordrecht, 1993.

[A2] L. AIZENBERG, *Variations on the theorem of Morera and the problem of Pompeiu*. Dokl. Akad. Nauk (1994) (to appear).

[AM] L.A. AIZENBERG - B.C. MITYAGIN, *The spaces of functions analytic in multicircular domains*. Sibirsk. Mat. Zh. **1** (1960), 153-170, (Russian).

[AR] L.A. AIZENBERG - C. REA, *The moment-condition for the free boundary problem for CR functions*. Ann. Scuola Norm. Sup. Pisa Cl. Sci (4) **20** (1993), 313-322.

[BCPZ] C. BERENSTEIN - D.C. CHANG - D. PASCUAS - L. ZALCMAN, *Variations on the theorem of Morera*. Contemp. Math. **137** (1992), 63-78.

[Br] M. BRELOT, *Sur l'approximation et la convergence dans le théorie des fonctions harmoniques ou holomorphes*. Bull. Soc. Math. France **73** (1945), 55-70.

[Bu] R.B. BURCKEL, *An Introduction to Classical Complex Analysis*. Birkhäuser Verlag, Boston, 1979.

[D] J. DENY, *Systèmes totaux de fonctions harmoniques*. Ann. Inst. Fourier **1** (1949), 103-112.

[F] B.A. FUKS, *Introduction to the Theory of Analytic Functions of Several Complex Variables*. Amer. Math. Soc. Providence RI, 1963.

[L] N.S. LANDKOF, *Foundations of Potential Theory*. Springer-Verlag, Berlin, 1972.

[Ro] H. ROSSI, *Holomorphically convex sets in several complex variables*. Ann. Math. (2) **74** (1961), 470-493.

[Ru] W. RUDIN, *Functional Analysis*. McGraw-Hill, New York, 1973.

[Z1] L. ZALCMAN, *Analytic Capacity and Rational Approximation*. Springer-Verlag, Berlin, 1968.

[Z2] L. ZALCMAN, *Offbeat integral geometry*. Amer. Math. Monthly **87** (1980), 161-175.

[Z3] L. ZALCMAN, *A bibliographic survey of the Pompeiu problem*. In: "Approximation by Solutions of Partial Differential Equations" (B. Fuglede et al, eds.), Kluwer Acad. Publ., Dordrecht, 1992, pp. 185-194.

Department of Mathematics & Computer Science
The Gelbart Research Institute for Mathematical Sciences
Bar-Ilan University
Ramat-Gan, Israel