

ANNALI DELLA SCUOLA NORMALE SUPERIORE DI PISA *Classe di Scienze*

SERGIO VENTURINI

Continuous dynamical systems on Taut complex manifolds

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 24, n° 2 (1997), p. 291-298

http://www.numdam.org/item?id=ASNSP_1997_4_24_2_291_0

© Scuola Normale Superiore, Pisa, 1997, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Continuous Dynamical Systems on Taut Complex Manifolds

SERGIO VENTURINI

1. – Introduction and statements of main results

Let X be a smooth vector field on a smooth manifold M . In order to fix some notation, let us recall that the flow associated to X is a pair (V, φ) , where $V \subset \mathbb{R} \times M$ is open and $\varphi : V \rightarrow M$, $(t, m) \mapsto \varphi_t(m)$, is a smooth map such that

- (i) for any $m \in M$, $(0, m) \in V$, and $\varphi_0(m) = m$;
- (ii) $\varphi_t(\varphi_s(m)) = \varphi_{t+s}(m)$ whenever $(s, m), (t + s, m), (t, \varphi_s(m)) \in V$;
- (iii) for every $m \in M$

$$\left. \frac{d}{dt} \varphi_t(m) \right|_{t=0} = X(m);$$

- (iv) the pair (V, φ) is maximal, i.e. if (V', φ') is an other pair satisfying (i), (ii) and (iii) then $V' \subset V$ and $\varphi' = \varphi|_{V'}$.

It is a standard result that any smooth vector field has a unique associated flow (and every flow comes from a vector field).

The vector field X with associated flow $\varphi_t(\cdot)$ is *complete* (respectively *right complete*) if for every $m \in M$ $\varphi_t(m)$ is defined for every $t \in \mathbb{R}$ (respectively $t > 0$).

In the paper [W] Wu introduced the notion of *taut* complex manifold. Let us recall briefly such a definition. Let M be a complex manifold and let

$$\Delta = \{z \in \mathbb{C} \mid |z| < 1\}$$

be the open unit disk. A sequence $f_n : \Delta \rightarrow M$ is said to be *compactly divergent* if for any choice of a compact set $H \subset \Delta$ and $K \subset M$, $f_n(H) \cap K = \emptyset$ for n large enough. The complex manifold M is *taut* if given any sequence $f_n : \Delta \rightarrow M$ of holomorphic maps admits a subsequence converging to a holomorphic map $f : \Delta \rightarrow M$ (uniformly on the compact subset of Δ) or a subsequence compactly divergent.

The purpose of this paper is to describe the behaviour as $t \rightarrow +\infty$ of the flow $\varphi_t(\cdot)$ associated to a right complete holomorphic vector field X on a taut complex manifold M and to investigate the link between the structure of the zeroes of the vector field X (i.e. the fixed point set for the flow $\varphi_t(\cdot)$) and the topology of the manifold M .

If the manifold M is compact then the theory is quite trivial, since such a manifold does not admit any non trivial holomorphic vector field. So, throughout the paper M will be a non compact connected taut complex manifold, X a holomorphic vector field on M with associated flow $\varphi_t(\cdot)$.

In order to explain our main results we need to introduce some notations.

We denote by S^1 the circle group (the topological boundary of Δ), by $T^r = S^1 \times \cdots \times S^1$ the standard r -dimensional torus Lie group and by S^n the unit sphere in $\mathbb{R}^{n+1}$.

We denote by $\text{Hol}(M, N)$ the space of all holomorphic maps between two complex manifold M and N endowed with the compact open topology, and by $\text{Aut}(M)$ the group of all holomorphic automorphisms of M .

Let X be right complete vector field on a complex manifold M with associated flow $\varphi_t(\cdot)$. We set

$$Z(X) = \{m \in M \mid X(m) = 0\}.$$

The flow $\varphi_t(\cdot)$ on M is said

- (i) *compact* if the family $\{\varphi_t(\cdot)\}_{t \geq 0}$ is relatively compact in $\text{Hol}(M, M)$,
- (ii) *compactly divergent* if for each pair of compact subsets $H, K \subset M$ for some t_0 one has $\varphi_t(H) \cap K = \emptyset$ if $t \geq t_0$.

For every $m \in M$, $s \geq 0$, set

$$\Gamma_m(s) = \{\varphi_t(m) \mid t \geq s\},$$

$$\Gamma_m = \bigcap_{s \geq 0} \overline{\Gamma_m(s)};$$

we also set

$$E(X) = \{m \in M \mid m \in \Gamma_m\},$$

$$T(X) = M \setminus E(X).$$

(“ E ” and “ T ” stand respectively for “ergodic” and “transient”).

Let us recall that the Kuratowsky limits of a family of subset $A_t \subset M$, $t \geq 0$ are defined as

$$K - \liminf_{t \rightarrow +\infty} A_t =$$

$$\{m \in M \mid \forall U \text{ neighbourhood of } m \exists t_0 \geq 0 \text{ s.t. } \forall t \geq t_0 \ A_t \cap U \neq \emptyset\},$$

$$K - \limsup_{t \rightarrow +\infty} A_t =$$

$$\{m \in M \mid \forall U \text{ neighbourhood of } m \forall t \geq 0 \exists t_0 \geq t \text{ s.t. } A_{t_0} \cap U \neq \emptyset\},$$

If $u : [0, +\infty[\rightarrow M$ is a map then we will write $K - \liminf_{t \rightarrow +\infty} u(t)$ (respectively $K - \limsup_{t \rightarrow +\infty} u(t)$) instead of $K - \liminf_{t \rightarrow +\infty} \{u(t)\}$ (respectively $K - \limsup_{t \rightarrow +\infty} \{u(t)\}$).

Finally, as usual, $\pi_p(M)$, $H_p(M, G)$ and $H^p(M, G)$ will stand respectively for the p -th homotopy group (with respect to some base point in M), the p -th homology and cohomology groups of M with coefficients in the abelian group G . The manifold M is of *finite topological type* if for every $p \geq 0$

$$\dim_{\mathbb{Q}} H^p(X, \mathbb{Q}) < +\infty$$

and for such a manifold the *Euler characteristic* is

$$\chi(M) = \sum_{i=0}^{\dim M} (-1)^i \dim_{\mathbb{Q}} H^i(X, \mathbb{Q}).$$

Our main results on flows on taut manifolds are described by the followings theorems:

THEOREM 1.1. *Let X be a right complete vector field on a (non compact) connected taut complex manifold M with associated flow $\varphi_t(\cdot)$. Then $\varphi_t(\cdot)$ is either compact or compactly divergent. If it is compact then the following assertions hold:*

- (i) $E(X)$ is a not empty closed integral submanifold of M ;
- (ii) the restriction of X to $E(X)$ is a complete vector field on $E(X)$;
- (iii) if L is an integral submanifold of X and the restriction of X to L is a complete vector field on L then $L \subset E(X)$;
- (iv) the inclusion map $i : E(X) \rightarrow M$ is a homotopical equivalence between $E(X)$ and M ;
- (v) there exist a smooth toral action on $E(X)$

$$T^r \times E(X) \rightarrow E(X)$$

such that for all $m \in E(X)$ and any $s \geq 0$

$$\Gamma_m = \overline{\Gamma_m(s)} = T^r m.$$

In particular, $X(m) = 0$ (i.e. $\varphi_t(m) = m$) if, and only if, m is a fixed point for such a toral action;

- (vi) there exist a holomorphic retraction $\rho : M \rightarrow E(X)$ such that for all $m \in T(X)$

$$K - \limsup_{t \rightarrow +\infty} \varphi_t(m) = \Gamma_{\rho(m)}.$$

In particular $\varphi_t(m)$ converges to $\bar{m} \in M$ as $t \rightarrow +\infty$ if, and only if, $\bar{m} = \rho(m)$ and $X(\bar{m}) = 0$.

THEOREM 1.2. *Let X , M and $\varphi_t(\cdot)$ be as in Theorem 1.1 and assume further that M is of finite topological type and the flow $\varphi_t(\cdot)$ is compact. Then $Z(X)$ is a closed complex submanifold of finite topological type and*

$$\chi(Z(X)) = \chi(M).$$

In particular, if $\chi(M) \neq 0$ then $Z(X) \neq \emptyset$.

We recall that a topological space S is *acyclic* (over $\mathbb{Q}$) if $\dim_{\mathbb{Q}} H^i(X, \mathbb{Q}) = 0$ for every $i > 0$.

THEOREM 1.3. *Let X , M and $\varphi_t(\cdot)$ be as in Theorem 1.2 and assume further M is a connected acyclic manifold. Then $Z(X)$ also is a connected acyclic connected (closed) submanifold of M .*

THEOREM 1.4. *Let X , M and $\varphi_t(\cdot)$ be as in Theorem 1.2. If the even rational homotopy groups of M vanish then $Z(X)$ is either empty or a connected complex submanifold of M .*

I thank Prof. Angelo Vistoli for some helpful conversation on the subject.

2. – Proofs

Let X , M and $\varphi_t(\cdot)$ be as in Theorem 1.1. Since the law of composition is continuous in $\text{Hol}(M, M)$ then the family

$$\mathcal{F} = \{\varphi_t(\cdot) \mid t \in [0, 1]\}.$$

is compact (with respect to the compact open topology on $\text{Hol}(M, M)$). Set also $f = \varphi_1 : M \rightarrow M$; then the sequence $f^n = f \circ \dots \circ f = \varphi_n : M \rightarrow M$ of the iterates of f is either relatively compact in $\text{Hol}(M, M)$ or compactly divergent ([A1]).

Assume the sequence f^n relatively compact. Since every $\varphi_t(\cdot)$ is of the form $f^n \circ \varphi_s$ with $\varphi_s \in \mathcal{F}$, then $\{\varphi_t\}$ is a relatively compact family in $\text{Hol}(M, M)$. Assume now that the sequence f^n is compactly divergent. Let H and K be any two compact subset of M . Then, being $\mathcal{F}$ a compact family,

$$H' = \bigcup_{s \in [0, 1]} \varphi_s(H)$$

is a compact subset of M . Since, by assumption, the sequence f^n is compactly divergent then there exists n_0 such that $f^n(H') \cap K = \emptyset$ for all $n \geq n_0$. Then, for every $t \geq n_0$, writing $t = n + s$ with $n \in \mathbb{N}$ and $s \in [0, 1]$,

$$\varphi_t(H) \cap K = f^n(\varphi_s(H)) \cap K \subset f^n(H') \cap K = \emptyset,$$

which shows that the flow $\varphi_t(\cdot)$ is compactly divergent. The first statement of Theorem 1.1 is thus proved.

Assume now that the flow $\varphi_t(\cdot)$ is compact. Let S be the closure of the family $\{\varphi_t(\cdot)\}_{t \geq 0}$ in $\text{Hol}(M, M)$. Then S is a compact topological abelian semigroup with respect to the law of composition of (holomorphic) maps. It is a standard fact of the theory of compact (semi-)topological semi-groups that S contain an idempotent element ρ (i.e. ρ satisfies $\rho^2 = \rho$) such that the semigroup $G = \rho S$ is a compact topological abelian group with identity ρ ([Ru]).

The map $\rho : M \rightarrow M$ is therefore a retraction of M onto its image $N = \rho(M)$. It follows from a result of Rossi ([R]) that N is a closed complex submanifold of M .

Let $m \in N$. Then, since S is abelian,

$$\varphi_t(m) = \varphi_t(\rho(m)) = \rho(\varphi_t(m)) \Rightarrow \varphi_t(m) \in N,$$

that is, N is invariant under each φ_t , or, equivalently, N is an integral submanifold of X .

Moreover, since G is a group, it easily follows that the restriction of each φ_t to N is an automorphism of N , and setting

$$\psi_t(m) = \begin{cases} \varphi_t(m) & \text{if } t \geq 0 \\ \varphi_{-t}^{-1}(m) & \text{if } t < 0, \end{cases}$$

then $\psi_t(\cdot)$ is a smooth one parameter group of automorphisms of N such that

$$\left. \frac{d}{dt} \psi_t(m) \right|_{t=0} = X(m).$$

It follows that the restriction of X to N is a complete vector field on N .

Before going further we need to recall some basic fact on the "Kobayashi distance". Such a pseudodistance

$$k_M : M \times M \rightarrow [0, +\infty[$$

is defined for every (connected) complex manifold M , and can be characterized as the biggest one among all the pseudodistances $\delta : M \times M \rightarrow [0, +\infty[$ such that for all $f \in \text{Hol}(\Delta, M)$ and all $z, w \in \Delta$

$$\delta(f(z), f(w)) \leq \omega(z, w),$$

where $\omega(z, w)$ is the Poincaré distance on Δ , i.e. the integrated form of the Poincaré metric

$$\frac{dz \overline{dz}}{(1 - z\overline{z})^2}.$$

If it happens that k_M is a distance, that is $k_M(m, m') > 0$ if $m \neq m'$ then the complex manifold M is said *hyperbolic*.

We summarize here the results on hyperbolic manifold that we need in the sequel. For references see e.g. [K], [A2].

(i) if $f \in \text{Hol}(M, N)$ then, for each pair of points m and $m' \in M$,

$$k_N(f(m), f(m')) \leq k_M(m, m');$$

- (ii) if N is a complex submanifold of M and M is hyperbolic then N is hyperbolic;
- (iii) the group $\text{Aut}(M)$ of an hyperbolic complex manifold M is a Lie group acting smoothly on M ;
- (iv) every taut manifold is hyperbolic.

Coming back to the proof of Theorem 1.1, set

$$K = \{u|_N \mid u \in S\}$$

Then K , being a compact connected abelian subgroup of $\text{Aut}(N)$, which by the assertions above is a Lie group, it is a Lie group. But a compact connected abelian Lie group is isomorphic (as Lie group) to a standard real torus T^r for some $r \geq 0$. Thus we have a toral action

$$T^r \times E(X) \rightarrow E(X).$$

Since every $u \in K$ is limit of a sequence of maps in the flow φ_t , it easily follows that for every $m \in N$ and $s \geq 0$

$$\overline{\Gamma_m(s)} = \Gamma_m = T^r m,$$

which clearly implies that $N \subset E(X)$.

Now let $m \in M \setminus N$. Choose a sequence $f_n = \varphi_{t_n}$ converging in $\text{Hol}(M, M)$ to ρ . Given $\varepsilon > 0$, pick n large enough in such a way that $k_M(f_n(m), \rho(m)) < \varepsilon$. Then, for every $t \geq t_n$,

$$\begin{aligned} k_M(\varphi_t(m), \varphi_{t-t_n}(\rho(m))) \\ = k_M(\varphi_{t-t_n}(f_n(m)), \varphi_{t-t_n}(\rho(m))) \leq k_M(f_n(m), \rho(m)) < \varepsilon, \end{aligned}$$

whence

$$k_M(\varphi_t(m), T^r \rho(m)) \leq k_M(\varphi_t(m), \varphi_{t-t_n}(\rho(m))) < \varepsilon.$$

(Where, by definition, for a subset $A \subset M$ we set $k_M(m, A) = \inf_{m' \in A} \{k_M(m, m') \mid m' \in A\}$).

The estimates above, together with the fact that $\{\varphi_t(\rho(m))\}_{t \geq 0}$ is dense in $T^r \rho(m)$, imply that for every $s \geq 0$

$$\overline{\Gamma_{\rho(m)}(s)} = \Gamma_m = T^r \rho(m) \subset N,$$

and also $m \in T(X)$, being $m \in M \setminus N$.

Since $m \in M \setminus N$ is arbitrary, it follows then that $M \setminus N \subset T(X)$, which combined with the previously proved inclusion $N \subset E(X)$ yields the equalities $N = E(X)$ and $M \setminus N = T(X)$.

All that proves statements (i), (ii), (v) and (vi) of Theorem 1.1.

Let us prove statement (iii).

Let $L \subset M$ be an integral submanifold of the vector field X , and assume that the restriction of X to L is complete. Let $\psi_t(\cdot)$ be the flow on L associated to X . It is an one parameter group of automorphisms of L , which is a hyperbolic complex manifold, relatively compact in $\text{Aut}(L)$. Its closure G in $\text{Aut}(L)$ is therefore a compact connected abelian Lie group, whence it is isomorphic to a r -dimensional torus T^r . As before, given $m \in L$, we obtain $\Gamma_m = T^r m$, whence $m \in \Gamma_m$, that is $m \in E(X)$. Since $m \in L$ is arbitrary the inclusion $L \subset E(X)$ follows.

It remains to prove statement (iv) of Theorem 1.1.

Because the spaces M and $E(X) = \rho(M)$ are connected (complex) manifold, they have some CW -complex structure, and hence it suffices to prove that the induced group homomorphisms

$$i_* : \pi_k(E(X)) \rightarrow \pi_k(M)$$

are isomorphisms in each dimension k .

The identity $\rho \circ i = id_N$ yields $\rho_* \circ i_* = id_{N*}$, and hence i_* is a monomorphism. We will prove that i_* is an epimorphism showing that ρ_* is a monomorphism.

Let $[u] \in \pi_k(E(X))$, represented by the continuous map $u : S^k \rightarrow E(X)$, be in the kernel of ρ_* . Let $f^n = \varphi_{t_n}$ (here the “ n ” in “ f^n ” is an index written as superscript for “typographical” reason) be a sequence converging to ρ in $\text{Hol}(M, M)$. Then, since M is a manifold, $f_*^n([u]) = [u \circ f^n] = [u \circ \rho] = \rho_*([u]) = 0$ for n large enough. Obviously $f^n = \varphi_{t_n}$ is homotopic to φ_0 , which is the identity map on M , and hence $[u] = [u \circ \varphi_0] = [u \circ f^n] = \rho_*([u]) = 0$.

The proof of Theorem 1.1 is complete.

Let now X , M , $\varphi_t(\cdot)$ and F . By Theorem 1.1, after replacing M with a closed submanifold homotopically equivalent to M if necessary, we may assume that X is a complete vector field, that is $E(X) = M$. Again by Theorem 1.1 there exists a toral action

$$T^r \times M \rightarrow M$$

such that $\Gamma_m = T^r m$ for each $m \in M$. Obviously the set F is then the fixed point set of such an action. Theorem 1.2 then follows immediatly from the following

THEOREM 2.1. *Let $T^r \times V \rightarrow V$ be a smooth action on the (not necessarily connected) smooth manifold V , and let F be the fixed point set for such an action. If V is a manifold of finite topological type then also F is and*

$$\chi(F) = \chi(V).$$

PROOF. Let us write $T^r = S^1 \times T^{r-1}$, and let F_1 be the fixed point set for S^1 . Since a smooth action of a compact Lie group is linearizable in a neighbourhood of each fixed point, it follows that F_1 is a smooth (obviously closed) submanifold of V . Moreover T^{r-1} acts on F_1 . After r steps we so obtain a chain

$$F = F_r \subset \cdots \subset F_1 \subset V$$

of closed submanifold with F_{i+1} the fixed point set of a circle action on F_i .

Then, clearly it suffices to prove the theorem for $r = 1$, that is for a circle action, which is done, e.g., in [Br] Theorem 10.9 (and remark below). $\square$

Theorems 1.3 and 1.4 follow immediatly from the corresponding statements on toral actions (see, e.g., respectively [B, Theorem 5.3 (a)] and [H, Theorem (IV.5)]).

REFERENCES

- [A1] M. ABATE, *Iteration Theory of Holomorphic Maps on Taut Manifolds*, Meditrrerlean Press, Cosenza, 1989.
- [A2] M. ABATE, *Iteration theory, compactly divergent sequences and commuting holomorphic maps*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **18** (1991), 167-191.
- [Bo] A. BOREL, *Seminar on Transformations Groups*, Ann. of Math. Studies (46), Princeton, 1960.
- [Br] G.E. BREDON, *Introduction to compact transformation groups*, Academic Press, New York, 1972.
- [H] W. Y. HSIANG, *Cohomological Theory of Topological Transformations Groups*, Springer-Verlag, 1975.
- [K] S.KOBAYASHI, *Intrinsic distances, measures and geometric function theory*, Bull. Amer. Math. Soc. **82** (1976), 357-416.
- [R] H. ROSSI, *Vector fields on analytic spaces*, Ann. Math. **78** (1963), 455-467.
- [Ru] W. RUPPERT, *Compact Semitopological Semigroups: An Intrinsic Theory*, Lect. Notes in Math. 1079, Springer, 1984.
- [W] H. WU, *Normal families of holomorphic mappings*, Acta Math. **119** (1967), 193-233.

Dipartimento di Matematica
Università di Bologna
Piazza di Porta S. Donato 5
40127 Bologna, Italia