

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

STEFAN MÜLLER

MICHAEL STRUWE

VLADIMIR ŠVERÁK

Harmonic maps on planar lattices

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 25, n° 3-4 (1997), p. 713-730

http://www.numdam.org/item?id=ASNSP_1997_4_25_3-4_713_0

© Scuola Normale Superiore, Pisa, 1997, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Harmonic Maps on Planar Lattices

STEFAN MÜLLER – MICHAEL STRUWE – VLADIMIR ŠVERÁK

Abstract. We show that a sequence of harmonic maps from a square, two-dimensional lattice of mesh-size h to a compact Riemannian manifold N with uniformly bounded energy as $h \rightarrow 0$ weakly accumulates at a harmonic map $u: T^2 \rightarrow N$ on the flat two-dimensional torus.

1. – Introduction

Let N be a smooth, compact Riemannian manifold without boundary of dimension k . By Nash's embedding theorem we may assume that $N \subset \mathbb{R}^n$ isometrically for some n . We are interested in understanding the relation between (smooth) harmonic maps $u: T^2 = \mathbb{R}^2/\mathbb{Z}^2 \rightarrow N \subset \mathbb{R}^n$ on the torus, characterized as critical points of the energy

$$(1) \quad E(v) = \int_{T^2} e(v) dx$$

with density

$$(2) \quad e(v) = \frac{1}{2} |\nabla v|^2$$

subject to the "target constraint" $v(T^2) \subset N$, and their counterparts on a discrete domain.

Our main result in this paper states that harmonic maps with uniformly bounded energy on a lattice, as the mesh-size $h \rightarrow 0$, $h^{-1} \in \mathbb{N}$, weakly accumulate at a harmonic map on the torus.

The above result may be of relevance for numerical purposes and may have implications for questions regarding existence of harmonic maps under constraints.

In fact, in a sequel to this paper [12] we will use a spatially discrete ansatz to give an alternative proof for the existence of global weak solutions to the Cauchy problem for wave maps on $(1+2)$ -dimensional Minkowski space, established in Müller-Struwe [11] by a different method.

A core ingredient in the analysis - both in the stationary and in the time-dependent case - are the weak compactness results of Freire-Müller-Struwe [7], [8] which make contact with work of Bethuel [1], [2], Evans [5], Hélein [9] and stress the importance of Hardy space estimates for Jacobians, due to Coifman-Lions-Meyer-Semmes [3], and the \mathcal{H}^1 -BMO-duality, due to Fefferman-Stein [6].

2. – Notation

Denote as $T = \mathbb{R}^2/\mathbb{Z}^2$ the flat 2-dimensional torus, and let $x = (x^1, x^2)$ denote a generic point in T ; also let $e_\alpha, \alpha = 1, 2$, denote the standard basis for \mathbb{R}^2 .

For $h > 0$ with $h^{-1} \in \mathbb{N}$ consider the lattice $T_h = (h\mathbb{Z})^2/\mathbb{Z}^2$ with generic point $x_h = (x_h^1, x_h^2)$. For $l \in \mathbb{N}$ and $x_h \in T_h$ also let

$$Q_{lh}(x_h) = \{x \in T; x_h^\alpha \leq x^\alpha < x_h^\alpha + lh, \alpha = 1, 2\}$$

denote the square of edge-length lh with lower left corner x_h and for $x \in T$ denote as $[x]_h$ the unique point $[x]_h \in T_h$ such that $x \in Q_h([x]_h)$.

Given a discretely defined map $u^h: T_h \rightarrow \mathbb{R}^n$, we may extend u^h to T by letting

$$(3) \quad u^h(x) = u^h([x]_h) \quad \text{for } x \in T.$$

The forward and backward difference quotients in direction e_α , defined as

$$\partial_\alpha^{\pm h} u^h(x) = \frac{u^h(x \pm h e_\alpha) - u^h(x)}{\pm h}$$

for $x \in T_h, \alpha = 1, 2$, then also trivially satisfy the relation

$$(4) \quad \partial_\alpha^{\pm h} u^h(x) = \partial_\alpha^{\pm h} u^h([x]_h)$$

for $x \in T$.

Finally, we also introduce forward and backward means

$$(5) \quad m_\alpha^{\pm h} u^h(x) = \frac{u^h(x \pm h e_\alpha) + u^h(x)}{2} \equiv m_\alpha^{\pm h} u^h([x]_h),$$

and translates

$$(6) \quad \tau_\alpha^{\pm h} u^h(x) = u^h(x \pm h e_\alpha) \equiv \tau_\alpha^{\pm h} u^h([x]_h)$$

for $x \in T, \alpha = 1, 2$.

Observe that, in particular, there holds

$$(7) \quad \partial_\alpha^{-h} \left(\tau_\alpha^h u^h \right) = \tau_\alpha^h \left(\partial_\alpha^{-h} u^h \right) = \partial_\alpha^h u^h;$$

moreover, we have

$$(8) \quad \partial_\alpha^h \partial_\alpha^{-h} = \partial_\alpha^{-h} \partial_\alpha^h$$

for any $\alpha = 1, 2$.

3. – Difference calculus

For completeness, we quickly review the basic formulas from difference calculus that we will need.

3.1. – Product rule

Let $u^h, v^h: T_h \rightarrow \mathbb{R}$. Then

$$\begin{aligned}
 (9) \quad \partial_\alpha^h (u^h v^h) &= \partial_\alpha^h u^h \tau_\alpha^h v^h + u^h \partial_\alpha^h v^h \\
 &= \partial_\alpha^h u^h v^h + \tau_\alpha^h u^h \partial_\alpha^h v^h \\
 &= \partial_\alpha^h u^h m_\alpha^h v^h + m_\alpha^h u^h \partial_\alpha^h v^h,
 \end{aligned}$$

and similarly for $\partial_\alpha^{-h}, \alpha = 1, 2$.

In particular, by (7) we have

$$(10) \quad \partial_\alpha^h (\partial_\alpha^{-h} u^h v^h) = \partial_\alpha^h \partial_\alpha^{-h} u^h v^h + \partial_\alpha^h u^h \partial_\alpha^h v^h = \partial_\alpha^{-h} (\partial_\alpha^h u^h \tau_\alpha^h v^h).$$

In view of (10), later we will be able to avoid unnecessary shifting of arguments by working with backward differences.

3.2. – Discrete integration

For $u^h: T_h \rightarrow \mathbb{R}$ we define the integral of u^h as

$$\int_{T_h} u^h = h^2 \sum_{x \in T_h} u^h(x) = \int_T u^h dx,$$

where in the latter integral u^h denotes the piecewise constant extension of u^h as in (3). Obviously, we have

$$\int_{T_h} \partial_\alpha^h u^h = 0, \quad \int_{T_h} \tau_\alpha^h u^h = \int_{T_h} u^h$$

for any $u^h: T_h \rightarrow \mathbb{R}, \alpha = 1, 2$. Hence from (9) and (7) we obtain the following formula for integrating by parts

$$\begin{aligned}
 (11) \quad 0 &= \int_{T_h} \partial_\alpha^h (u^h v^h) = \int_{T_h} \partial_\alpha^h u^h \tau_\alpha^h v^h + \int_{T_h} u^h \partial_\alpha^h v^h \\
 &= \int_{T_h} \partial_\alpha^{-h} u^h v^h + \int_{T_h} u^h \partial_\alpha^h v^h;
 \end{aligned}$$

in particular,

$$\int_{T_h} \partial_\alpha^h u^h \partial_\alpha^h v^h = - \int_{T_h} \partial_\alpha^{-h} \partial_\alpha^h u^h v^h.$$

3.3. – Dirichlet integral

For u^h as above, denote its energy density at a point $x \in T_h$ as

$$(12) \quad e_h(u^h)(x) = \frac{1}{4} \sum_{\alpha=1,2} \left\{ |\partial_\alpha^h u^h(x)|^2 + |\partial_\alpha^{-h} u^h(x)|^2 \right\}$$

and define

$$(13) \quad E_h(u^h) = \int_{T_h} e_h(u^h).$$

We compute the first variation of E_h at u^h in direction v^h as

$$(14) \quad \begin{aligned} \langle dE_h(u^h), v^h \rangle &= \frac{d}{d\varepsilon} E_h(u^h + \varepsilon v^h)|_{\varepsilon=0} \\ &= \frac{1}{2} \sum_{\alpha} \int_{T_h} \left\{ \partial_\alpha^h u^h \partial_\alpha^h v^h + \partial_\alpha^{-h} u^h \partial_\alpha^{-h} v^h \right\} \\ &= \sum_{\alpha} \int_{T_h} \partial_\alpha^h u^h \partial_\alpha^h v^h = - \int_{T_h} \Delta^h u^h v^h, \end{aligned}$$

where $\Delta^h = \sum_{\alpha} \partial_\alpha^{-h} \partial_\alpha^h$ is the discrete (5-point) Laplacian.

3.4. – Exterior calculus

Differential forms on T_h can be most easily expressed in terms of the standard basis $dx^\alpha, \alpha = 1, 2, dx^1 \wedge dx^2$, respectively. A 1-form φ^h on T_h then can be identified with a pair of functions $\varphi_\alpha^h, \alpha = 1, 2$, such that $\varphi^h = \sum_{\alpha} \varphi_\alpha^h dx^\alpha$; similarly a 2-form b^h on T_h can be identified with a function β^h such that $b^h = \beta^h dx^1 \wedge dx^2$.

Two 1-forms $\varphi^h = \sum_{\alpha} \varphi_\alpha^h dx^\alpha$ and $\psi^h = \sum_{\alpha} \psi_\alpha^h dx^\alpha$ are contracted by letting

$$\varphi^h \cdot \psi^h = \sum_{\alpha} \varphi_\alpha^h \psi_\alpha^h;$$

also let

$$\varphi^h \cdot \varphi^h = |\varphi^h|^2.$$

The Hodge $*$ -operator acts on the basis elements as

$$*1 = dx^1 \wedge dx^2, *dx^1 = dx^2, *dx^2 = -dx^1, *dx^1 \wedge dx^2 = 1,$$

and $*$ is linear with respect to multiplication by functions. In particular, there holds

$$\varphi^h \wedge (*\varphi^h) = |\varphi^h|^2 dx^1 \wedge dx^2$$

and

$$**\varphi^h = (-1)^p \varphi^h$$

for any p -form φ^h on T_h .

For a function $u^h: T_h \rightarrow \mathbb{R}$ or a 1-form $\varphi^h = \sum_{\alpha} \varphi_{\alpha}^h dx^{\alpha}$ the discrete differential is defined as

$$d^{\pm h} u^h = \sum_{\alpha} \partial_{\alpha}^{\pm h} u^h dx^{\alpha},$$

$$d^{\pm h} \varphi^h = \sum_{\alpha, \beta} \partial_{\alpha}^{\pm h} \varphi_{\beta}^h dx^{\alpha} \wedge dx^{\beta} = \left(\partial_1^{\pm h} \varphi_2^h - \partial_2^{\pm h} \varphi_1^h \right) dx^1 \wedge dx^2.$$

The co-differential operator is given by

$$\delta^{\pm h} = * \circ d^{\mp h} \circ *.$$

Note that for any $u^h: T_h \rightarrow \mathbb{R}$ and any 1-form φ^h on T_h we have

$$\int_{T_h} d^{\pm h} u^h \cdot \varphi^h = - \int_{T_h} u^h \delta^{\pm h} \varphi^h;$$

that is, $-\delta^{\pm h}$ is the adjoint of $d^{\pm h}$ with respect to the inner product on forms defined by contraction and the L^2 -inner product on T_h .

Moreover, a direct computation shows that

$$d^h \circ d^h = 0, \delta^h \circ \delta^h = 0.$$

The Laplacian on p -forms (with the analysts' sign convention) is defined as

$$\Delta^h = d^h \circ \delta^h + \delta^h \circ d^h.$$

By (8) there holds $\Delta^h = \Delta^{-h}$. Moreover, it is easy to check that Δ^h acts as a diagonal operator with respect to the standard basis on forms; in particular, for a 1-form $\varphi^h = \sum_{\alpha} \varphi_{\alpha}^h dx^{\alpha}$ we have

$$\Delta^h \varphi^h = \sum_{\alpha} (\Delta^h \varphi_{\alpha}^h) dx^{\alpha}.$$

3.5. – Hodge decomposition

As in the continuous case, the following result holds.

PROPOSITION 3.1. Any 1-form $\varphi^h = \sum_{\alpha} \varphi_{\alpha} dx^{\alpha}$ may be decomposed uniquely as

$$\varphi^h = d^h a^h + \delta^h b^h + c^h,$$

where

$$(15) \quad \int_{T_h} a^h = 0, \int_{T_h} b^h = 0, d^h c^h = 0, \delta^h c^h = 0.$$

PROOF. Letting a^h, b^h be the unique solutions to the equations

$$(16) \quad \Delta^h a^h = \delta^h \varphi^h, \Delta^h b^h = d^h \varphi^h,$$

normalized by (15), for the remainder $c^h = \varphi^h - d^h a^h - \delta^h b^h$ we obtain

$$d^h c^h = d^h \varphi^h - \Delta^h b^h = 0, \delta^h c^h = \delta^h \varphi^h - \Delta^h a^h = 0,$$

as claimed. \square

REMARK 3.2. Solving (16) for a^h can be achieved, for instance, by minimizing the functional

$$F(a^h) = \int_{T_h} \{e_h(a^h) + a^h \delta^h \varphi^h\},$$

confer (14), and similarly for b^h .

4. – Interpolation and discretization

In addition to the trivial extension of a map $u^h: T_h \rightarrow \mathbb{R}$ to the torus, defined by 3, we introduce the bilinear extension of u^h , defined by letting

$$\bar{u}^h(x) = u^h(x) + \sum_{\alpha} \xi^{\alpha} \partial_{\alpha}^h u^h(x) + \xi^1 \xi^2 \partial_1^h \partial_2^h u^h(x),$$

for $x = [x]_h + \xi \in T$.

The following result is immediate from the definition.

LEMMA 4.1. We have $\bar{u}^h \in H^{1,2} \cap L^{\infty}(T)$, and with a uniform constant C there holds

- i) $\|\bar{u}^h - u^h\|_{L^{\infty}(Q_h(x_h))}^2 \leq C \int_{Q_{2h}(x_h)} e_h(u^h), \quad \text{for all } x_h \in T_h;$
- ii) $\|\bar{u}^h - u^h\|_{L^2(T)}^2 \leq Ch^2 E_h(u^h);$
- iii) $C^{-1} E_h(u^h) \leq E(\bar{u}^h) \leq CE(u^h).$

In view of Lemma 4.1 iii) we will say that $u^h \rightarrow u$ weakly in $H^{1,2}(T)$ as $h \rightarrow 0$ if $\bar{u}^h \rightarrow u$ weakly in $H^{1,2}(T)$, and similarly for vector-valued maps.

Observe, however, that for $u^h: T_h \rightarrow N \subset \mathbb{R}^n$ the range of \bar{u}^h in general will not lie in N . On the other hand, we have

LEMMA 4.2. *Suppose $u^h \rightarrow u$ weakly in $H^{1,2}(T; \mathbb{R}^n)$ as $h \rightarrow 0$, where $u^h: T_h \rightarrow N$. Then $u(x) \in N$ for almost every $x \in T$.*

PROOF. For $\delta > 0, h > 0$ let

$$\Sigma_h^\delta = \left\{ x \in T_h; \|u^h - \bar{u}^h\|_{L^\infty(Q_h(x))} \geq \delta \right\}.$$

By Lemma 4.1 i) then there holds

$$\delta |\Sigma_h^\delta| \leq C E_h(u^h) \leq C < \infty,$$

and hence

$$\mathcal{L}^2 \left(\left\{ x \in T; \text{dist}(\bar{u}^h(x), N) \geq \delta \right\} \right) \leq C \delta^{-1} h^2.$$

Here, \mathcal{L}^2 denotes 2-dimensional Lebesgue measure. Since $\bar{u}^h \rightarrow u$ weakly in $H^{1,2}(T; \mathbb{R})$ and hence strongly in $L^2(T; \mathbb{R}^n)$, after passing to a sub-sequence, if necessary, we may assume that $\bar{u}^h \rightarrow u$ almost everywhere as $h \rightarrow \infty$. Thus, for any $\delta > 0$ we infer

$$\mathcal{L}^2 \left(\left\{ x \in T; \text{dist}(u(x), N) \geq \delta \right\} \right) = 0,$$

as claimed. □

In contrast to interpolating functions $u^h: T_h \rightarrow \mathbb{R}$, discretizing functions $\varphi \in H^{1,2}(T)$ is somewhat subtle, as such maps, for instance, are only defined pointwise almost everywhere. Moreover, interpolating the discretized map φ should recover the regularity properties of φ as much as possible.

Of the many possible choices we define as a discretized function φ the map

$$\varphi^h(x) = h^{-2} \int_{Q_h(x)} \varphi \, dx, \quad x \in T_h.$$

Note that if φ is the trivial extension of a map $\psi^h: T_h \rightarrow \mathbb{R}$, defined by (3), then $\varphi^h = \psi^h$; however, in general, and even if φ is piecewise bilinear, $\varphi \neq \bar{\varphi}^h$, the bilinearly interpolated discretized map.

LEMMA 4.3. *Let $\varphi \in H^{1,2}(T)$. Then with a uniform constant C there holds*

i) $\|\varphi^h - \varphi\|_{L^2(T)}^2 \leq C h^2 E(\varphi);$

ii) $\|\partial_\alpha^h \varphi^h - \partial_\alpha \varphi\|_{L^2(T)} \rightarrow 0$ as $h \rightarrow 0$.

PROOF. i) Estimating, for $x \in T$,

$$|\varphi^h(x) - \varphi(x)|^2 \leq h^{-2} \int_{Q_h([x]_h)} |\varphi(y) - \varphi(x)|^2 dy,$$

we obtain

$$\begin{aligned} \|\varphi^h - \varphi\|_{L^2(T)}^2 &\leq h^{-2} \int_T \int_{Q_h([x]_h)} |\varphi(y) - \varphi(x)|^2 dy dx \\ &\leq \int_T \int_{Q_h([x]_h)} \int_0^1 |\nabla \varphi(x + \vartheta(y-x))|^2 d\vartheta dy dx \leq Ch^2 E(\varphi). \end{aligned}$$

ii) By the Lebesgue differentiability theorem

$$\begin{aligned} \partial_\alpha^h \varphi^h(x) &= h^{-2} \int_{Q_h([x]_h)} \frac{\varphi(y + h\underline{e}_\alpha) - \varphi(y)}{h} dy \\ &= h^{-3} \int_0^h \int_{Q_h([x]_h)} \partial_\alpha \varphi(y + \vartheta \underline{e}_\alpha) d\vartheta dy \\ &\rightarrow \partial_\alpha \varphi(x) \quad \text{as } h \rightarrow 0 \end{aligned}$$

for almost every $x \in T$. Moreover, for any $\Omega \subset T$ we can estimate

$$\begin{aligned} \int_\Omega |\partial_\alpha^h \varphi^h(x)|^2 dx &\leq \int_\Omega \left\{ h^{-3} \int_0^h \int_{Q_h([x]_h)} |\partial_\alpha \varphi(y + \vartheta \underline{e}_\alpha)|^2 d\vartheta dy \right\} dx \\ &\leq h^{-3} \int_0^h \int_{Q_{2h}((-h, -h))} \left\{ \int_\Omega |\partial_\alpha \varphi(x + y + \vartheta \underline{e}_\alpha)|^2 dx \right\} dy d\vartheta \\ &\leq 4 \sup_{y \in T} \left\{ \int_{\Omega+y} |\partial_\alpha \varphi|^2 dx \right\} < \delta \end{aligned}$$

if $\mathcal{L}^2(\Omega) < \mu_0(\delta)$, by absolute continuity of the Lebesgue integral. Thus, the family of indefinite integrals $(\int |\partial_\alpha^h \varphi^h|^2)_{h>0}$ is uniformly absolutely continuous, and the assertion follows from Vitali's convergence theorem. \square

5. – Harmonic maps

A map $u^h: T_h \rightarrow N \subset \mathbb{R}^n$ by definition is *harmonic* if u^h is a critical point for E_h among maps $v^h: T \rightarrow N$; that is, if the first variation

$$(17) \quad \langle dE_h(u^h), \varphi^h \rangle = 0$$

for all $\varphi^h \in (u^h)^{-1}TN$, where

$$(u^h)^{-1}TN = \{\varphi^h: T_h \rightarrow \mathbb{R}^n; \varphi^h(x) \in T_{u^h(x)}N \text{ for all } x \in T_h\}$$

is the pullback tangent bundle, with T_pN denoting the tangent space to N at a point $p \in N$. By (14), equation (17) is equivalent to the relation

$$(18) \quad -\Delta^h u^h \perp T_{u^h}N,$$

where “ \perp ” means orthogonal with respect to the scalar product $\langle \cdot, \cdot \rangle$ in the ambient \mathbb{R}^n .

Introducing a local frame v_{k+1}, \dots, v_n for the normal bundle TN^\perp near a point $p = u^h(x) \in N$, we can also locally express (18) in the form

$$(19) \quad -\Delta^h u^h = \sum_l \lambda^l v_l \circ u^h$$

The coefficient functions λ^l can be determined as

$$(20) \quad \begin{aligned} \lambda^l &= -\langle \Delta^h u^h, v_l \circ u^h \rangle \\ &= -\sum_\alpha \partial_\alpha^{\mp h} \langle \partial_\alpha^{\pm h} u^h, v_l \circ u^h \rangle + \sum_\alpha \langle \partial_\alpha^{\pm h} u^h, \partial_\alpha^{\pm h} (v_l \circ u^h) \rangle \end{aligned}$$

Also recall that a smooth map $u: T \rightarrow N \subset \mathbb{R}^n$ is harmonic if u is critical for E among maps $v: T \rightarrow N$, or, equivalently, if

$$(21) \quad -\Delta u = A(u)(\nabla u, \nabla u) \perp T_u N,$$

where $A(p): T_p N \times T_p N \rightarrow T_p N^\perp$ is the second fundamental form of $N \subset \mathbb{R}^n$. Locally, with respect to a smooth local frame v_{k+1}, \dots, v_n for TN^\perp we have

$$A(p)(\xi, \eta) = \sum_l A^l(p)(\xi, \eta) v_l(p),$$

where

$$A^l(p)(\xi, \eta) = \langle \xi, dv_l(p) \cdot \eta \rangle$$

denotes the second fundamental form of $v_l, k < l \leq n$.

Our main result then is the following.

THEOREM 5.1. *For a sequence of numbers $h \rightarrow 0, h^{-1} \in \mathbb{N}$, suppose $u^h: T_h \rightarrow N \subset \mathbb{R}^n$ is harmonic and $u^h \rightarrow u$ weakly in $H^{1,2}(T; N)$ as $h \rightarrow 0$. Then u is harmonic.*

6. – Equivalent Hodge system

As observed by Christodoulou-Tahvildar-Zadeh [4] and Hélein [9], we may assume that TN is parallelizable. Let e_1, \dots, e_k be a smooth orthonormal frame field on N , such that $(e_1(p), \dots, e_k(p))$ is an orthonormal basis for T_pN at any $p \in N$.

Given $u^h: T_h \rightarrow N$, and a family of rotations $R^h: T_h \rightarrow SO(k)$, then

$$e_i^h = \sum_j R_{ij}^h(e_j \circ u^h)$$

is a frame for $(u^h)^{-1}TN$.

Let

$$\begin{aligned} \vartheta_{i,\alpha}^h &= \langle \partial_\alpha^h u^h, \tau_\alpha^h e_i^h \rangle dx^\alpha, \\ \vartheta_{i,\alpha}^{-h} &= \tau_\alpha^{-h} \vartheta_{i,\alpha}^h = \langle \partial_\alpha^{-h} u^h, e_i^h \rangle dx^\alpha, \\ \omega_{ij,\alpha}^{\pm h} &= \langle \partial_\alpha^{\pm h} e_k^h, m_\alpha^{\pm h} e_j^h \rangle dx^\alpha. \end{aligned}$$

Then

$$\begin{aligned} \delta^h \vartheta_i^h &= \sum_\alpha \partial_\alpha^{-h} \vartheta_{i,\alpha}^h = \sum_\alpha \partial_\alpha^h \vartheta_{i,\alpha}^{-h} \\ &= \sum_\alpha \langle \partial_\alpha^h \partial_\alpha^{-h} u^h, e_i^h \rangle + \sum_\alpha \langle \partial_\alpha^h u^h, \partial_\alpha^h e_i^h \rangle. \end{aligned}$$

It follows that $u^h: T_h \rightarrow N$ is discretely harmonic, if and only if

$$\delta^h \vartheta_i^h = \sum_\alpha \langle \partial_\alpha^h u^h, \partial_\alpha^h e_i^h \rangle = \sum_\alpha \tau_\alpha^h \langle \partial_\alpha^{-h} u^h, \partial_\alpha^{-h} e_i^h \rangle.$$

Letting v_{k+1}, \dots, v_n be a smooth local frame for the normal bundle, we can expand

$$\begin{aligned} \langle \partial_\alpha^{-h} u^h, \partial_\alpha^{-h} e_i^h \rangle &= \sum_j \langle \partial_\alpha^{-h} u^h, e_j^h \rangle \langle e_j^h, \partial_\alpha^{-h} e_i^h \rangle \\ &\quad + \sum_l \langle \partial_\alpha^{-h} u^h, v_l \circ u^h \rangle \langle v_l \circ u^h, \partial_\alpha^{-h} e_i^h \rangle \\ &= \sum_j \vartheta_{j,\alpha}^{-h} \omega_{ij,\alpha}^{-h} + \sum_j \vartheta_{j,\alpha}^{-h} \left\langle \frac{e_j^h - \tau_\alpha^{-h} e_j^h}{2}, \partial_\alpha^{-h} e_i^h \right\rangle \\ &\quad + \sum_l \langle \partial_\alpha^{-h} u^h, v_l \circ u^h \rangle \langle v_l \circ u^h, \partial_\alpha^{-h} e_i^h \rangle \\ &=: \sum_j \vartheta_{j,\alpha}^{-h} \omega_{ij,\alpha}^{-h} + \eta_{li,\alpha}^{-h}. \end{aligned}$$

The presence of the error term $\eta_{li,\alpha}^{-h}$ marks one major difference between the discrete and continuous cases.

LEMMA 6.1. *There is a constant $C = C(N)$ such that*

$$|\eta_{1i,\alpha}^{-h}| \leq C|u^h - \tau_\alpha^{-h}u^h| \left(|\partial_\alpha^{-h}u^h|^2 + \sum_j |\partial_\alpha^{-h}e_j^h|^2 \right).$$

PROOF. Since $u^h: T_h \rightarrow N$ and since for $p, q \in N$, with $C = C(N)$ we have

$$|\langle p - q, v \rangle| \leq C|p - q|^2, \quad \text{for all } v \in T_pN^\perp,$$

it follows that

$$\begin{aligned} |\langle \partial_\alpha^{-h}u^h, v_l \circ u^h \rangle| &\leq Ch^{-1}|u^h - \tau_\alpha^{-h}u^h|^2 \\ &= C|u^h - \tau_\alpha^{-h}u^h| |\partial_\alpha^{-h}u^h|. \end{aligned}$$

Hence the second term defining η_1^{-h} can be estimated as claimed. For the first it suffices to note that

$$|\partial_\alpha^{-h}u||e_j^h - \tau_\alpha^{-h}e_j^h| = |u^h - \tau_\alpha^{-h}u^h| |\partial_\alpha^{-h}e_j^h|. \quad \square$$

In the following we denote

$$\eta_{1i,\alpha}^h = \tau_\alpha^h \eta_{1i,\alpha}^{-h}, \quad \eta_1^h = (\eta_{1i,\alpha}^h)_{1 \leq i \leq k, 1 \leq \alpha \leq 2}.$$

With this notation then u^h is discretely harmonic if and only if there holds

$$(22) \quad \delta^h \vartheta_i^h = \sum_j \vartheta_j^h \cdot \omega_{ij}^h + \eta_{1i}^h.$$

Observe that $\tau_\alpha^h (\vartheta_{j\alpha}^{-h} \omega_{ij,\alpha}^{-h}) = \vartheta_{j,\alpha}^h \cdot \omega_{ij,\alpha}^h$.

Similarly, if $u \in H^{1,2}(T; N)$ is weakly harmonic and if $e_i = R_{ij}\bar{e}_j \circ u$ is a frame for $u^{-1}TN$ with connection 1-forms

$$\omega_{ij} = \langle e_j, de_i \rangle, \quad 1 \leq i, j \leq k,$$

then, letting

$$\vartheta_i = \langle du, e_i \rangle, \quad 1 \leq i \leq k,$$

the equation holds

$$(23) \quad \delta \vartheta_i = \sum_j \vartheta_j \cdot \omega_{ij},$$

and conversely.

7. – Coulomb gauge

Equations (22), (23) involve an arbitrary choice of frame for the pull-back bundle. We may use this gauge freedom of rotating the frames to fix a frame with particularly nice analytic properties. Following Hélein [9], we impose Coulomb gauge, as follows.

For each h choose $R^h: T_h \rightarrow SO(k)$ such that

$$(24) \quad E_h \left(R^h(\underline{e} \circ u^h) \right) := \sum_i E_h \left(\sum_j R_{ij}^h(\underline{e}_j \circ u^h) \right) = \inf_{S^h: T_h \rightarrow SO(k)} E_h(S^h(\underline{e} \circ u^h)).$$

Observe that trivially

$$E_h(R^h(\underline{e} \circ u^h)) \leq E_h(\underline{e} \circ u^h) \leq CE_h(u^h).$$

LEMMA 7.1. *Let $R^h: T_h \rightarrow SO(k)$ satisfy (24). Then there holds the equation $\delta^h \omega_{ij}^h = \sum_\alpha \partial_\alpha^{-h} \omega_{ij,\alpha}^h = 0$ for the connection 1-form ω_{ij}^h associated with the frame $e_i^h = \sum_j R_{ij}^h(\underline{e}_j \circ u^h)$, $1 \leq i \leq k$.*

PROOF. For $r^h = r_{ij}^h: T_h \rightarrow \text{Id}SO(k) = \mathfrak{so}(k)$, letting $e_i^h = \sum_j R_{ij}^h(\underline{e}_j \circ u^h)$ and using (9), (11), and (10) we compute

$$\begin{aligned} 0 &= \frac{d}{d\varepsilon} E_h((id + \varepsilon r^h) \cdot R^h(\underline{e} \circ u^h)) \Big|_{\varepsilon=0} \\ &= \frac{1}{2} \sum_{\alpha,i,j} \int_{T_h} \left\{ \langle \partial_\alpha^h e_i^h, \partial_\alpha^h(r_{ij}^h e_j^h) \rangle + \langle \partial_\alpha^{-h} e_i^h, \partial_\alpha^{-h}(r_{ij}^h e_j^h) \rangle \right\} \\ &= \frac{1}{2} \sum_{\alpha,i,j} \int_{T_h} \left\{ \langle \partial_\alpha^h e_i^h, \partial_\alpha^h e_j^h \rangle + \langle \partial_\alpha^{-h} e_i^h, \partial_\alpha^{-h} e_j^h \rangle \right\} r_{ij}^h \\ &\quad + \frac{1}{2} \sum_{\alpha,i,j} \int_{T_h} \left\{ \langle \partial_\alpha^h e_i^h, \tau_\alpha^h e_j^h \rangle \partial_\alpha^h r_{ij}^h + \langle \partial_\alpha^{-h} e_i^h, \tau_\alpha^{-h} e_j^h \rangle \partial_\alpha^{-h} r_{ij}^h \right\} \\ &= -\frac{1}{2} \sum_{\alpha,i,j} \int_{T_h} \left\{ \partial_\alpha^h \langle \partial_\alpha^{-h} e_i^h, e_j^h \rangle + \partial_\alpha^{-h} \langle \partial_\alpha^h e_i^h, e_j^h \rangle \right\} r_{ij}^h, \end{aligned}$$

where we also used anti-symmetry $r_{ij}^h = -r_{ji}^h$ to obtain the last equation.

Recalling that

$$\partial_\alpha^h \langle \partial_\alpha^{-h} e_i^h, e_j^h \rangle = \partial_\alpha^{-h} \langle \partial_\alpha^h e_i^h, \tau_\alpha^h e_j^h \rangle,$$

we infer that

$$\sum_{\alpha,i,j} \int_{T_h} \partial_\alpha^{-h} \langle \partial_\alpha^h e_i^h, m_\alpha^h e_j^h \rangle r_{ij}^h = 0$$

for all $r^h \in \mathfrak{so}(k)$. Since by (9) we have

$$\omega_{ij,\alpha}^h = \langle \partial_\alpha^h e_i^h, m_\alpha^h e_j^h \rangle = -\omega_{ji,\alpha}^h,$$

we conclude that

$$\delta^h \omega_{ij}^h = \sum_\alpha \partial_\alpha^{-h} \omega_{ij,\alpha}^h = 0,$$

as claimed. \square

In view of the gauge condition we may assume that

- (25) $e_i^h \rightarrow e_i$ weakly in $H^{1,2}$,
- (26) $\vartheta_i^h \rightarrow \vartheta_i$ weakly in L^2 ,
- (27) $\omega_{ij}^h \rightarrow \omega_{ij}$ weakly in L^2 .

Let

$$\omega_{ij}^h = d^h a_{ij}^h + \delta^h b_{ij}^h + c_{ij}^h$$

be the Hodge decomposition of ω_{ij}^h . In view of the gauge condition $\delta^h \omega_{ij}^h = 0$, it follows that

$$a_{ij}^h = 0.$$

Moreover, we have

$$E_h(b_{ij}^h) \leq C \int_{T_h} |\omega_{ij}^h|^2 \leq C, |c_{ij}^h| \leq C \int_{T_h} |\omega_{ij}^h|^2 \leq C.$$

Hence we may assume

$$b_{ij}^h \xrightarrow{w} b_{ij} \text{ in } H^{1,2}, c_{ij}^h \rightarrow c_{ij} (h \rightarrow 0),$$

where

$$\omega_{ij} = \delta b_{ij} + c_{ij}.$$

Let $\beta_{ij}^h = *b_{ij}^h, \beta_{ij} = *b_{ij}$. Then

$$\omega_{ij}^h = *d^{-h} \beta_{ij}^h + c_{ij}^h, \omega_{ij} = *d\beta_{ij} + c_{ij}.$$

Note that

$$d^h u^h \cdot *d^{-h} \beta_{ij}^h = *(d^h u^h \wedge d^{-h} \beta_{ij}^h), \text{ etc.}$$

Thus, we have

$$\vartheta_j^h \cdot \omega_{ij}^h = *\langle d^h u^h \wedge d^{-h} \beta_{ij}^h, e_j^h \rangle + \sum_{\alpha} \langle \partial_{\alpha}^h u^h, \tau_{\alpha}^h e_j^h - e_j^h \rangle \omega_{ij,\alpha}^h + \vartheta_j^h \cdot c_{ij}^h.$$

8. – Convergence

Passing to the limit $h \rightarrow 0$ in the distribution sense is no problem for the terms $\delta^h \vartheta_j^h, \vartheta_j^h \cdot c_{ij}^h$. The term

$$\eta_{2i,\alpha}^h = \langle \partial_\alpha^h u^h, \tau_\alpha^h e_j^h - e_j^h \rangle \omega_{ij,\alpha}^h$$

is easily dominated

$$|\eta_{2i,\alpha}^h| \leq C |\tau_\alpha^h u^h - u^h| \cdot \sum_j |\partial_\alpha^h e_j^h|^2$$

in the same way as $\eta_{1i,\alpha}^h$; see Lemma 6.1. These and similar error terms will be dealt with later.

Now we concentrate on the term

$$J^h(u^h, \beta_{ij}^h, e_j^h) = \langle d^h u^h \wedge d^{-h} \beta_{ij}^h, e_j^h \rangle.$$

This term has the structure of a Jacobian determinant. In the continuous limit $h = 0$ concentration-compactness arguments yield weak convergence results for such terms. Our aim in the following will be to reduce the discrete case to the continuous one.

Choose a smooth testing function $\varphi \in C^\infty(T), 0 \leq \varphi \leq 1$. Discretize φ to obtain a map $\varphi^h: T_h \rightarrow \mathbb{R}$. We intend to prove

$$\int_{T_h} J^h(u^h, \beta_{ij}^h, e_j^h) \varphi^h \rightarrow \int_T \langle du \wedge d\beta_{ij}, e_j \varphi \rangle$$

as $h \rightarrow 0$.

Let $v^h = \beta_{ij}^h, w^h = e_j^h \cdot \varphi^h$ for brevity. Note that $u^h \rightarrow u, v^h \rightarrow v = \beta_{ij}, w^h \rightarrow w = e_j \varphi$ weakly in $H^{1,2}$ as $h \rightarrow 0$.

LEMMA 8.1. *For any $u^h, v^h, w^h: T_h \rightarrow \mathbb{R}^n$ there holds*

$$\begin{aligned} \int_{T_h} J^h(u^h, v^h, w^h) &= - \int_{T_h} J^h(u^h, w^h, v^h) + \int_{T_h} \eta_3^h \\ &= - \int_{T_h} J^h(w^h, v^h, u^h) + \int_{T_h} \eta_4^h \end{aligned}$$

where

$$|\eta_{3,4}^h| \leq C \sum_\alpha |\tau_\alpha^h u^h - u^h| \left(|\partial_\alpha^{\pm h} v^h|^2 + |\partial_\alpha^{\pm h} w^h|^2 \right)$$

with an absolute constant C .

PROOF. It suffices to prove the first estimate; the second is obtained replacing forward by backward difference quotients. Moreover, considering each triple of components separately, we may assume that $n = 1$ to simplify the notation.

We have

$$\begin{aligned} \tau_2^h (\partial_1^h u^h \partial_2^{-h} v^h w^h) &= \partial_2^h (\partial_1^h u^h v^h w^h) - \partial_2^h \partial_1^h u^h v^h w^h - \partial_1^h \tau_2^h u^h v^h \partial_2^h w^h, \\ \tau_1^h (\partial_2^h u^h \partial_1^{-h} v^h w^h) &= \partial_1^h (\partial_2^h u^h v^h w^h) - \partial_1^h \partial_2^h u^h v^h w^h - \partial_2^h \tau_1^h u^h v^h \partial_1^h w^h. \end{aligned}$$

Taking the difference of these two equations, since $\partial_1^h \partial_2^h u^h = \partial_2^h \partial_1^h u^h$ the middle terms on the right cancel. Integrating the resulting identity and shifting, we thus obtain

$$\begin{aligned} \int_{T_h} J(u^h, v^h, w^h) &= - \int_{T_h} \left\{ \partial_1^h u^h \tau_2^{-h} v^h \partial_2^{-h} w^h - \partial_2^h u^h \tau_1^{-h} v^h \partial_1^{-h} w^h \right\} \\ &= - \int_{T_h} J(u^h, w^h, v^h) + \int_{T_h} \eta_3^h, \end{aligned}$$

where

$$\begin{aligned} \eta_3^h &= \partial_1^h u^h (v^h - \tau_2^{-h} v^h) \partial_2^{-h} w^h - \partial_2^h u^h (v^h - \tau_1^{-h} v^h) \partial_1^{-h} w^h \\ &= (\tau_1^h u^h - u^h) \partial_2^{-h} v^h \partial_2^{-h} w^h + (\tau_2^h u^h - u^h) \partial_1^{-h} v^h \partial_1^{-h} w^h \end{aligned}$$

can be estimated as claimed. □

Let $\bar{u}^h, \bar{v}^h, \bar{w}^h$ be the bilinearly interpolated functions u^h , etc. Observe the following:

$$\begin{aligned} \partial_1 \bar{u}^h(x_h + h\xi) &= (1 - \xi_2) \partial_1^h u^h(x_h) + \xi_2 \partial_1^h u^h(x_h + h e_2) \\ \partial_2 \bar{u}^h(x_h + h\xi) &= (1 - \xi_1) \partial_2^h u^h(x_h) + \xi_1 \partial_2^h u^h(x_h + h e_1) \end{aligned}$$

for $x_h \in T_h, \xi \in Q_1(0)$, and similarly for \bar{v}^h and \bar{w}^h . In particular,

$$\begin{aligned} \partial_1 \bar{u}^h(x_h + \xi) &= \partial_1 \bar{u}^h(x_h + \xi_2 e_2), \\ \partial_2 \bar{u}^h(x_h + \xi) &= \partial_2 \bar{u}^h(x_h + \xi_1 e_1) \end{aligned}$$

for all $x_h \in T_h, \xi \in Q_h(0)$.

Thus, for instance,

$$\begin{aligned} \int_{Q_h(x_h)} \langle \partial_1 \bar{u}^h \partial_2 \bar{v}^h, w^h \rangle d\xi_1 d\xi_2 &= \int_{Q_h(x_h)} \langle m_2^h \partial_1^h u^h m_1^h \partial_2^h v^h, w^h \rangle \\ &= \int_{Q_h(x_h)} \langle \partial_1^h (m_2^h u^h) \partial_2^h (m_1^h v^h), w^h \rangle. \end{aligned}$$

In view of this identity,

$$\begin{aligned} &\int_T (\partial_1 \bar{u}^h \partial_2 \bar{v}^h - \partial_2 \bar{u}^h \partial_1 \bar{v}^h, w^h) dx_1 dx_2 \\ &= \int_{T_h} \langle \partial_1^h (m_2^h u^h) \partial_2^h (m_1^h v^h) - \partial_2^h (m_1^h u^h) \partial_1^h (m_2^h v^h), w^h \rangle \end{aligned}$$

Shifting coordinates,

$$\begin{aligned} \int_{T_h} \langle \partial_1^h(m_2^h u^h) \partial_2^h(m_1^h v^h), w^h \rangle &= \int_{T_h} \langle \partial_1^h(m_2^{-h} u^h) \partial_2^{-h}(m_1^h v^h), w^h \rangle + \int_{T_h} \eta_5^h \\ &= \int_{T_h} \langle \partial_1^h(m_2^{-h} u^h), m_1^h(\partial_2^{-h} v^h w^h) \rangle + \int_{T_h} \eta_6^h \\ &= \int_{T_h} \langle \partial_1^h(m_1^{-h} m_2^{-h} u^h) \partial_2^{-h} v^h, w^h \rangle + \int_{T_h} \eta_7^h, \end{aligned}$$

with η_j^h bounded like $\eta_{3,4}^h$, and similarly with ∂_1^h and ∂_2^h exchanged.

Thus

$$\int_T d\bar{u}^h \wedge d\bar{v}^h w^h = \int_{T_h} J^h(m_1^{-h} m_2^{-h} u^h, v^h, w^h) + \int_{T_h} \eta_8^h,$$

and by Lemma 8.1 the latter

$$\begin{aligned} &= - \int_{T_h} J^h(w^h, v^h, m_1^{-h} m_2^{-h} u^h) + \int_{T_h} \eta_9^h \\ &= - \int_{T_h} J^h(w^h, v^h, u^h) + \int_{T_h} \eta_{10}^h \\ &= \int_{T_h} J^h(u^h, v^h, w^h) + \int_{T_h} \eta_{11}^h. \end{aligned}$$

It follows that

$$\int_{T_h} J^h(u^h, v^h, w^h) = \int_T \langle d\bar{u}^h \wedge d\bar{\beta}_{ij}^h, \bar{e}_j^h \rangle \varphi + \int_{T_h} \eta_{12}^h,$$

where the η_j^h are all bounded as $\eta_{3,4}^h$ above.

We can now use the concentration-compactness argument from [8], proof of Theorem 1.1, based on [10], Lemma 4.3, to pass to the limit in (22). In fact, [10] implies that, as $h \rightarrow 0$,

$$\langle d\bar{u}^h \wedge d\beta_{ij}^h, e_j^h \rangle \rightarrow \langle du \wedge d\beta_{ij}, e_j \rangle + \sum_{k \in K} \nu_k \delta_{\{\bar{\nu}_k\}}$$

in the sense of distributions, where K is at most countable and $\sum_{k \in K} |\nu_k| < \infty$.

For the error terms η_j^h we have a similar result. We combine all these terms in a single term η^h , satisfying the estimate

$$(28) \quad |\eta^h| \leq C \sum_{\alpha} \left(h \|\nabla \varphi\|_{L^\infty} + |\varphi| \cdot |\tau_{\alpha}^h u^h - u^h| \right) \left(|\partial_{\alpha}^h u^h|^2 + \sum_{i,j} |\partial_{\alpha}^h e_j^h|^2 + |\partial_{\alpha}^h \beta_{ij}^h|^2 \right)$$

with a constant $C = C(N)$ independent of φ .

LEMMA 8.2. *There exists an at most countable set $\{\bar{x}_j\}_{j \in J}$ in T and numbers $\eta_j > 0$ such that, as $h \rightarrow 0$ suitably, $\eta^h \rightarrow \sum_j \eta_j \delta_{\{\bar{x}_j\}}$ in measure, where $\eta_j = \mu_j \varphi(\bar{x}_j)$ and $\sum_j |\mu_j|^{2/3} < \infty$.*

PROOF. Passing to a subsequence, if necessary, we may assume that $|\eta^h| \rightarrow \eta$ in measure. Let $\delta > 0$. As observed in Lemma 4.2 above, denoting

$$\Sigma_h^\delta = \left\{ x_h \in T_h; \sum_\alpha |(\tau_\alpha^h u^h - u^h)(x_h)|^2 > \delta \right\},$$

there holds

$$\delta |\Sigma_h^\delta| \leq C E_h(u^h) \leq C < \infty.$$

Possibly passing to a further subsequence, we may assume $\Sigma_h^\delta \rightarrow \Sigma^\delta$, where $|\Sigma^\delta| < \infty$.

Moreover, for $h > 0$ we have

$$\int_T |\eta^h| = \int_{T \setminus \cup_{x_h \in \Sigma_h^\delta} Q_h(x_h)} |\eta^h| + \sum_{x_h \in \Sigma_h^\delta} \int_{Q_h(x_h)} |\eta^h| = I_h^\delta + II_h^\delta,$$

and for fixed $\varphi \in C^\infty(T)$ we can estimate

$$\begin{aligned} I_h^\delta &\leq C \left(h \|\nabla \varphi\|_{L^\infty} + \sqrt{\delta} \|\varphi\|_{L^\infty} \right) \int_{T_h} \sum_{\alpha, i, j} (|\partial_\alpha^h u^h|^2 + |\partial_\alpha^h e_j^h|^2 + |\partial_\alpha^h \beta_{ij}^h|^2) \\ &\leq C \left(h + \sqrt{\delta} \right) \end{aligned}$$

while for each $x_h \in \Sigma_h^\delta$ in view of Lemma 4.1. i) we find

$$\begin{aligned} \int_{Q_h(x_h)} |\eta^h| &\leq C \left\{ h \|\nabla \varphi\|_{L^\infty} + |\varphi(x_h)| \left(\int_{Q_{2h}(x_h)} e_h(u^h) \right)^{1/2} \right\} \\ &\quad \int_{Q_h(x_h)} \sum_{\alpha, i, j} (|\partial_\alpha^h u^h|^2 + |\partial_\alpha^h e_j^h|^2 + |\partial_\alpha^h \beta_{ij}^h|^2). \end{aligned}$$

Thus, for $\delta > 0$ we may decompose $\eta = \eta_0^\delta + \sum_{\bar{x}_j \in \Sigma^\delta} \eta_j^\delta \delta_{\{\bar{x}_j\}}$, where

$$\int_T \eta_0^\delta \leq \lim_{r \rightarrow 0} \lim_{h \rightarrow 0} \int_{T \setminus \cup_{\bar{x}_j \in \Sigma^\delta} B_r(\bar{x}_j)} \eta^h \leq C \sqrt{\delta}$$

and where for each $\bar{x}_j \in \Sigma^\delta$ we have

$$\eta_j = \mu_j \varphi(\bar{x}_j)$$

with

$$\sum_j \mu_j^{2/3} \leq C$$

independent of $\delta > 0$. (In the latter estimate we also used concavity of the function $t \mapsto t^{2/3}$ to obtain the desired bound in case different sequences (x_h) in Σ_h^δ should converge to the same limit $\bar{x}_j \in \Sigma^\delta$.) Passing to the limit $\delta \rightarrow 0$, the assertion follows. \square

Passing to the limit $h \rightarrow 0$ in (22), thus we obtain the equation

$$\delta\vartheta_i - \sum_j \omega_{ij} \cdot \vartheta_j = \sum_{j \in J} \mu_j \delta_{\{\bar{x}_j\}} + \sum_{k \in K} \nu_k \delta_{\{\bar{y}_k\}}.$$

But the left hand side belongs to the space $H^{-1} + L^1$ which does not contain any atoms. Therefore all μ_j and ν_k must vanish and we find (23), as desired.

REFERENCES

- [1] F. BETHUEL, *Weak convergence of Palais-Smale sequences for some critical functionals*, Calc. Var. **1** (1993), 267-310.
- [2] F. BETHUEL, *On the singular set of stationary harmonic maps*, Manuscr. Math. **78** (1993), 417-443.
- [3] R. R. COIFMAN – P.-L. LIONS – Y. MEYER – S. SEMMES, *Compensated compactness and Hardy spaces*, J. Math. Pures Appl. **72** (1993), 247-286.
- [4] D. CHRISTODOULOU – S. TAHVILDAR-ZADEH, *On the regularity of spherically symmetric wave maps*, Comm. Pure Appl. Math. **46** (1993), 1041-1091.
- [5] L. C. EVANS, *Partial regularity for stationary harmonic maps into spheres*, Arch. Rat. Mech. Anal. **116** (1991), 101-113.
- [6] C. FEFFERMAN – E. M. STEIN, *H^p spaces of several variables*, Acta Math. **129** (1972), 137-193.
- [7] A. FREIRE – S. MÜLLER – M. STRUWE, *Weak Convergence of Wave Maps from (1+2)-Dimensional Minkowski Space to Riemannian Manifolds*, Invent. Math. (to appear).
- [8] A. FREIRE – S. MÜLLER – M. STRUWE, *Weak Compactness of Wave Maps and Harmonic Maps*, preprint (1996).
- [9] F. HÉLEIN, *Regularité des applications faiblement harmoniques entre une surface et une variété Riemannienne*, C. R. Acad. Sci. Paris Ser. I Math. **312** (1991), 591-596.
- [10] P.L. LIONS, *The concentration compactness principle in the calculus of variations, the limit case, part II*, Rev. Mat. Iberoam. **12** (1985), 45-121.
- [11] S. MÜLLER – M. STRUWE, *Global Existence of Wave Maps in 1 + 2 Dimensions for Finite Energy Data*, Top. Methods Nonlinear Analysis, **7** (1996), 245-259.
- [12] S. MÜLLER – M. STRUWE, *Spatially Discrete Wave Maps in 1+2 Dimensions*, in preparation.

Mathematics in the Sciences
Max-Planck Institute
Inselstrasse 22-26
D-04103 Leipzig
sm@mis.mpg.de

Mathematik
ETH-Zentrum
CH-8092, Zürich
struwe@math.ethz.ch

School of Mathematics
University of Minnesota
Minneapolis, MN 55455, USA
sverak@math.umn.edu