

Astérisque

NOBUO AOKI

**Ergodic automorphisms of compact metric groups
are isomorphic to Bernoulli shifts**

Astérisque, tome 40 (1976), p. 5-10

<http://www.numdam.org/item?id=AST_1976__40__5_0>

© Société mathématique de France, 1976, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ERGODIC AUTOMORPHISMS OF COMPACT METRIC GROUPS
ARE ISOMORPHIC TO BERNOULLI SHIFTS

Nobuo Aoki

I wish to discuss its title. Let X be a compact, metric group and μ be its normalized Haar measure. Then (X, μ) is a Lebesgue space. Let σ be an automorphism of X , then σ is an invertible measure preserving transformation from X onto X . Our problem is concerned with measure theoretic properties of σ .

Today we will outline a proof of the following

Theorem.

An ergodic automorphism of a compact, metric, abelian group is a Bernoulli shift.

In order to outline the Theorem, we prepare the following

Lemma 1. Let X be a compact, metric; abelian group and σ be an ergodic automorphism of X . Then there exists a sequence of subgroup $X \supset X_1 \supset \dots \bigcap_{n=1}^{\infty} X_n = \{e\}$, invariant with respect to σ , such that there are a σ -invariant totally disconnected subgroup $X_D^{(n)}$, a σ -invariant infinite dimensional connected subgroup $X_B^{(n)}$ and a σ -invariant finite dimensional connected subgroup $X_A^{(n)}$, so that $\sigma|_{X_D^{(n)}}$ is isomorphic to a Bernoulli shift, $\sigma|_{X_B^{(n)}}$ is a Bernoulli automorphism (see [28]) and $\sigma|_{X_A^{(n)}}$ is ergodic, and $(X/X_n, \sigma)$ is an algebraic factor of $(X_A^{(n)} \times X_B^{(n)} \times X_D^{(n)}, \sigma \times \sigma \times \sigma)$.

To show the ergodic system $(X_A^{(n)}, \sigma)$, $n \geq 1$, have the

Bernoulli properties, let $\bar{T}^r$ be a compact, connected, abelian group with the character group Q^r , being the r -vector of rational numbers and σ be an automorphism of $\bar{T}^r$, then it is sufficient to prove that if $(\bar{T}^r, \sigma)$ is ergodic then $(\bar{T}^r, \sigma)$ is Bernoullian.

We write $Q^r = \{g_1, g_2, \dots\}$. Let Q_n denote a subgroup generated by $\{g_k \circ \sigma^j : j \in \mathbb{Z}, 1 \leq k \leq n\}$ for $n \geq 1$. Since $\text{rank}(Q^r) = r$, there exists an integer $m > 0$ such that $\text{rank}(Q_n) = \text{rank}(Q^r)$ for $n \geq m$.

We denote by $\bar{T}(Q_n)$ the annihilator of Q_n in $\bar{T}^r$ for $n \geq 1$.

Then we have

Lemma 2. $(\bar{T}^r / \bar{T}(Q_n), \sigma)$ is Bernoullian for $n \geq 1$.

The proof is done by ideas of Katznelson together with the following facts.

We may consider σ as operating on R^r . So we have the decomposition

$$R^r = V_{-p} \oplus \dots \oplus V \oplus \dots \oplus V_q$$

such that $V_j, -p \leq j \leq q$, are σ -invariant, all the eigenvalues of $\sigma|_{V_0}$ have modulus 1, and all the eigenvalues of $\sigma|_{V_j}, j \neq 0$, have the same modulus ρ_j where

$$\rho_{-p} < \dots < 1 = \rho_0 < \dots < \rho_q.$$

Lemma 3. If $\{0\} \neq Q^r \cap \tilde{V}$ where $\tilde{V} = V_{-p} \oplus \dots \oplus V_0$, then R^r is decomposed by subspaces $R^{r_1}_{(\sigma R^r_1 = R^r_1)}$ and R^{r-r_1}

with the following properties: there exists an automorphism σ' of R^{r-r_1} and a homomorphism λ from R^{r-r_1} into R^{r_1} such that $\sigma(\chi_1, \chi_2) = (\sigma' \chi_1, \lambda(\chi_1) + \sigma \chi_2)$, $(\chi_1, \chi_2) \in R^{r-r_1} \oplus R^{r_1}$, and $\sigma|_{R^{r_1}}$ gives $R^{r_1} = V'_{-p} \oplus \dots \oplus V'_0$

where $V_j' \subset V_j$ for $j = -p, \dots, 0$, and σ' gives
 $R^{r-r_1} = V_{-p}'' \oplus \dots \oplus V'' \oplus \dots \oplus V_q''$

where $V_j'' \subset V_j$ for $j = -p, \dots, q$, such that

$$(V_{-p}'' \oplus \dots \oplus V'') \cap Q^r = \{0\}.$$

Lemma 4. If σ acts ergodically on Q^r , then so is

$\sigma|_{Q^{r_1}}$ and $Q^r \cap V' = \{0\}$.

If $Q^{r-r_1} \cap (V'' \oplus \dots \oplus V_q'') \neq \{0\}$, then we again decompose R^{r-r_1} as in Lemma 3. However, since the dimension of R^r is finite, such the argument ends in finite time.

For $m \geq 0$, we denote by $V^{(m)}$ the set of all r -vectors such that each component of a vector consists of an element of $\{0, \pm 1, \dots, \pm m^{10}\}$. We define

$$V(1, M) = \left\{ \sum_{m=1}^M \sigma^{-m} \tilde{\alpha}_m : \tilde{\alpha}_m \in V^{(M)}, 1 \leq m \leq M \right\},$$

$$V(K, K+M) = \left\{ \sum_{m=K}^{K+M} \sigma^m \tilde{\alpha}_m : \tilde{\alpha}_m \in V^{(M)}, K \leq m \leq K+M \right\}$$

for $K > 0$ and $M > 0$. Then $V(1, M)$ and $V(K, K+M)$ are subsets of R^r .

Lemma 5. Assume that R^r is decomposed as follows. If

$R^r = V_{-p} \oplus \dots \oplus V$, then $V_0 \cap Q^r = \{0\}$, and if $R^r = V_{-p} \oplus \dots \oplus V_0 \oplus \dots \oplus V_q$, then $(V_{-p} \oplus \dots \oplus V) \cap Q^r = \{0\}$ and $(V_0 \oplus \dots \oplus V_q) \cap Q^r = \{0\}$. Then there exists $K_1 > 0$ such that

$$V(1, M) \cap V(K, K+M) = \{0\}$$

for $K \geq K_1$ and an odd integer $M > 0$.

In the proofs of Lemmas 3, 4 and 5, we use no results of number theory.

We now conclude that (X, σ) is Bernoullian. Using theorem and structures of skew product transformations, I can prove the result for the case of non-abelian. But, proofs of these results will appear elsewhere.

REFERENCES

- [1] L.M. Abramov, The entropy of an automorphism of a solenoidal group, Theory of Prob. and its Appl. 4 (1959), 231 - 236.
- [2] R.L. Adler and B. Weiss, Similarity of automorphisms of the torus, Mem. Amer. Soc. 98 (1970).
- [3] N.Aoki and H. Totoki, Ergodic automorphisms of T are Bernoulli transformations, Publ. RIMS, Kyoto Univ. 10 (1975), 535 - 544.
- [4] D.Z. Arov, The computation of entropy for one class of group endomorphisms, Zap.Meh.Mat.Fak.Kharov, Math. Obshch 30 (1964), 48 - 69.
- [5] K.R. Berg, Entropy of torus automorphisms, Topological Dynamics (edited by W.H. Gottschalk and F. Hahn), Benjamin New York, (1969), 516 - 519.
- [6] D. Lind, Ergodic automorphisms of the infinite torus are Bernoulli, Israel J. Math. 10 (1974), 186 - 195.
- [7] N.A. Friedman and D.S. Ornstein, On isomorphism of weak Bernoulli transformations, Advances in Math. 5 (1971), 365 - 394.
- [8] P.R.Halmos and H.Samelson, On monothetic groups, Proc.Nat. Acad.Sci. U.S.A. 28 (1942), 254 - 258.
- [9] G. Hochschild, The structure of Lie Groups, Holden-Day, San Francisco, 1965.
- [10] I. Kaplansky, Groups with representations of bounded degree, Canad.J.Math. 1 (1949); 105 - 112.
- [11] Y. Katznelson, Ergodic automorphisms of T^n are Bernoulli shifts, Israel J.Math. 10 (1971), 186 - 195.

- [12] A.G. Kurosch, The theory of Groups I, Chelsea, New York, 1960.
- [13] D.S. Ornstein, Bernoulli shifts with the same entropy are isomorphic, Advances in Math. 4 (1970), 337 - 352.
- [14] ———, Two Bernoulli shifts with infinite entropy are isomorphic, ibid. 5 (1971), 339 - 348.
- [15] ———, Factors of Bernoulli shifts are Bernoulli shifts, ibid. 5 (1971), 349 - 364.
- [16] W. Parry, Intrinsic Markov chains, Trans.Amer. Math.Soc. 112 (1964), 55 - 66.
- [17] M.S. Pinsker, Dynamical systems with completely positive and zero entropy, Soviet Mathematics Doklady, 1 (1960), 937 - 938.
- [18] L. Pontjagin, Topological Groups, Princeton Univ.Press. Princeton, 1946.
- [19] V.A. Rohlin, Metric properties of endomorphisms of compact commutative groups, Amer.Math.Soc. Transl. 64 (1967), 244 - 272.
- [20] ———, On the fundamental ideas of measure theory. Amer.Math.Soc. Transl. 10 (1962), 1 - 54.
- [21] ———, Exact endomorphisms of a Lebesgue space, Amer. Math.Soc. Transl. 39 (1969), 1 - 36.
- [22] ———, Selected topics from the metric theory of dynamical systems, Amer.Math.Soc. Transl. 49 (1966), 171 - 240.
- [23] Ya.G. Sinai, Markov partitions and C-diffeomorphisms, Fanc.Anal. and its Appl. 2 (1968), 61 - 82.

- [24] ———, On the concept of entropy of a dynamical system,
Dokl.Akad.Nauk. 124 (1959), 768 - 771.
- [25] H. Totoki, Introduction to Ergodic Theory, Kyoritsu
Shuppan, Tokyo, 1971 (in Japanese).
- [26] ———, Ergodic Theory, Lecture notes series 14,
Aarhus Univ. 1969.
- [27] A. Weil, L'integration dans les Groups Topologique et ses
Applications, Actualites Sci, Indust. 869, Hermann, Paris,
1940, 2nd ed. 1951.
- [28] S.A. Yuzvinskii, Metric properties of endomorphisms of
compact groups, Amer.Math.Transl. 66 (1968), 63 - 98.

Department of Mathematics
Tokyo Metropolitan University
Tokyo, Japan