

# *Astérisque*

JEFF CHEEGER

## **Hodge theory of complex cones**

*Astérisque*, tome 101-102 (1983), p. 118-134

[http://www.numdam.org/item?id=AST\\_1983\\_\\_101-102\\_\\_118\\_0](http://www.numdam.org/item?id=AST_1983__101-102__118_0)

© Société mathématique de France, 1983, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# Hodge Theory of Complex Cones

by

Jeff Cheeger

## 0. Introduction

In this note, we amend and continue the final portion of the discussion of [2]. In the initial paragraph of Section 7 of [2], we stated that for "admissible" riemannian pseudomanifolds (in the sense of that paper) for which the metric is Kähler, the standard consequences of Hodge theory (the "Kähler package") follow from the Strong Hodge Theorem. In particular, it was asserted that the properties of the Kähler metric on which these results depend are purely local on the nonsingular part. However, as we realized shortly after the publication of [2], and pointed out in [5] (see pp. 307 and 317) this assertion is incorrect. Let  $J$  be the almost complex structure and  $\Delta$  the Laplacian. Then the relation  $\Delta J = \Delta J$  holds locally; see [6]. Since  $J$  is an isometry, it follows that  $h \in L^2$  implies  $Jh \in L^2$  and  $\Delta h = 0$  implies  $Jh = 0$ . However, in the present context (of incomplete riemannian manifolds) the global relations  $Jh \in L^2$ ,  $\Delta Jh = 0$  do not in general imply  $dJh = \delta Jh = 0$ , even if  $dh = \delta h = 0$ . This is the additional fact which is actually required for the applications to Kähler geometry; see [6], pp. 109-111. Of course it fails locally, even in  $\mathbb{C}^N$ . More generally however, let  $X$  be a suitable space with singularities  $\Sigma$ . Then there does exist a criterion which is local near  $\Sigma$ , and which implies that if  $h \in L^2$ ,  $dh = \delta h = 0$  on  $X \setminus \Sigma$ , then  $dJh = \delta Jh = 0$ . Consider the Laplacian  $\Delta_G$ , such that a smooth form  $\theta$  is in  $\text{dom } \Delta_G$  if  $\theta, d\theta, \delta\theta, d\delta\theta, \delta d\theta \in L^2$ . For spaces with conical singularities,  $\bar{d}^* = \bar{\delta}$  (apart from the case of ideal boundary conditions) and we assume this from now on; see [2]. Then

$\Delta_G$  is essentially self adjoint and its kernel contains precisely those  $h \in L^2$  with  $dh = \delta h = 0$ . If one assumes that away from the singular set,  $X$  is a complete manifold without boundary, and  $\theta \in L^2$  is smooth, then to check if  $\theta \in \text{dom } \Delta_G$ , it suffices to check the above conditions near  $\Sigma$ .

The purpose of the present paper is to show that with suitable extra hypotheses,  $J$  does preserve  $\text{dom } \Delta_G$ , and hence  $\ker \Delta_G$  as well. Our argument is computational and requires rather strong assumptions. However, it does apply in some cases where  $\Delta_0$ , the Laplacian on forms of compact support, is not essentially self adjoint. This possibility, which is not an invariant of quasi-isometry type, is the essential complication in the general problem (which is still open).

In Section 1, we consider a metric cone  $C(N^m)$ , and a bounded map  $J: \Lambda^i(C(N)) \rightarrow \Lambda^i(C(N))$ , such that

- 1)  $J$  is linear with respect to functions of  $r$ ,
- 2)  $J$  commutes with dilations,
- 3)  $\Delta J = J \Delta$ .

We call such a map  $J$  conical. We show that subject to these assumptions,  $J$  preserves  $\text{dom } \Delta_G$  except for a small number of possibilities. By 1) and 3) it is easy to see that these can occur only in dimensions  $i = \frac{m+1}{2}$ ,  $\frac{m+1}{2} \pm 1$ , ( $m+1$  even), or  $i = \frac{m+1}{2} \pm \frac{1}{2}$ ,  $\frac{m+1}{2} \pm \frac{3}{2}$ , ( $m+1$  odd). Using the strong hypothesis 2), we show that in fact, only  $i = \frac{m+1}{2}$  or  $i = \frac{m+1}{2} \pm \frac{1}{2}$  are possible.

In Section 2, we consider an arbitrary Kähler manifold  $Y^{m+1}$  with the property that an  $L^2$ -harmonic form is in  $\text{dom } \Delta_G$  except possibly when  $i = \frac{m+1}{2}$ ,  $\frac{m+1}{2} \pm 1$ . By an argument which fails for  $i = \frac{m+1}{2} \pm 1$ , we rule out the case  $i = \frac{m+1}{2}$ .

As announced in [5], the proof can then be generalized to case of piecewise flat spaces, but we will give the details of this elsewhere (see Theorem 2.3 for the general result proved here).

We mention that the exceptional cases of Section 1 actually occur as formal possibilities for the double cover of the punctured complex plane (but, as allowed in Section 1,  $J$  is not an almost complex structure). This space also provides a simple example for which  $\Delta_0$  fails to be essentially self adjoint.

We conclude this section by observing that for spaces with (not necessarily isolated) conical singularities, the "Kähler package" can fail to hold if the singularities are not complex analytic. According to [1], an arbitrary orientable manifold  $X^{2n}$ , can be realized as an  $r$ -fold branched cover of the standard  $p.1.$  sphere,  $S^{2n}$ . By forming the connected sum  $S^{2n} \# \mathbb{C}P(n)$  (removing a ball which is disjoint from the branch locus  $L$ ) it follows that  $X' = X \# r\mathbb{C}P(n)$  is an  $r$ -fold branched cover of  $\mathbb{C}P(n)$ . By pulling back the metric on  $\mathbb{C}P(n)$ , one obtains a Kähler metric on  $X'$  which has conical singularities. The Mayer-Vietoris sequence shows  $b_1(X) = b_1(X')$ . In particular, if  $b_1(X)$  is odd,  $H^1(X', \mathbb{C})$  does not have a Hodge structure. Of course, metrics constructed via branched covers are of considerable interest for  $L^2$ -cohomology. But in the present context one should require that the branch locus be a complex subvariety.

1.  $\Delta J = J \Delta$  on cones

In this section we consider the metric cone  $C(N^m)$  on a compact smooth manifold  $N^m$  with empty boundary (see [2], [4] for background and notation). We begin by deriving the formula for the Laplacian of an  $i$ -form,

$$(1.1) \quad \theta = g(r)\phi(x) + f(r)dr \wedge \omega(x),$$

on  $C(N^m)$ . Intrinsic operations on  $N^m$  are denoted by adding a tilde. Let  $\Omega, dr \wedge \Omega$  denote the volume forms on  $N^m, C(N^m)$  respectively.

$$(1.2) \quad d(g\phi) = g d\phi + g' dr \wedge \phi,$$

$$(1.3) \quad *d(g\phi) = r^{m-2i} g' \tilde{*}\phi + (-1)^{i+1} r^{m-2i-2} g dr \wedge \tilde{*}d\phi,$$

$$(1.4) \quad d*d(g\phi) = (r^{m-2i} g'' + (m-2i)r^{m-2i-1} g') dr \wedge \tilde{*}\phi + r^{m-2i} g' d\tilde{*}\phi \\ + (-1)^i r^{m-2i-2} g dr \wedge d\tilde{*}d\phi$$

$$(1.5) \quad *d*d(g\phi) = (-1)^{i(m-i)} (g'' + (m-2i)r^{-1} g') \phi \\ + (-1)^{i+1} r^{-2} g \tilde{*}d\tilde{*}d\phi + (-1)^{m-i+1} r^{-2} g' dr \wedge \tilde{*}d\tilde{*}\phi,$$

$$(1.6) \quad \delta d(g\phi) = (-g'' - (m-2i)r^{-1} g') \phi + r^{-2} g \delta d\phi - r^{-2} g' dr \wedge \delta \phi,$$

Similarly,

$$(1.7) \quad *(g\phi) = (-1)^i r^{m-2i} g dr \wedge \tilde{*}\phi,$$

$$(1.8) \quad d*(g\phi) = (-1)^{i+1} r^{m-2i} g dr \wedge d\tilde{*}\phi,$$

$$(1.9) \quad *d*(g\phi) = (-1)^{i+1} r^{-2} g \tilde{*}d\tilde{*}\phi,$$

$$(1.10) \quad d*d*(g\phi) = (-1)^{i+1} r^{-2} g d\tilde{*}d\tilde{*}\phi + (-1)^{i+1} (-2r^{-3} + r^{-2} g') dr \wedge \tilde{*}d\tilde{*}\phi,$$

$$(1.11) \quad d\delta(g\phi) = (-2r^{-3} g + r^{-2} g') dr \wedge \delta \phi + r^{-2} g d\delta \phi.$$

Also,

$$(1.12) \quad d(fdr_{\wedge}\omega) = -fdr_{\wedge}d\omega,$$

$$(1.13) \quad *d(fdr_{\wedge}\omega) = -r^{m-2i}f\tilde{*}d\omega,$$

$$(1.14) \quad d*d(fdr_{\wedge}\omega) = (-r^{m-2i}f' - (m-2i)r^{m-2i-1}f)dr_{\wedge}\tilde{*}d\omega \\ - r^{m-2i}fd\tilde{*}d\omega,$$

$$(1.15) \quad *d*d(fdr_{\wedge}\omega) = (-1)^{m-i}r^{-2}fdr_{\wedge}\tilde{*}d\tilde{*}d\omega + (-1)^i(m-i)+1(f' + (m-2i)r^{-1}f)d\omega,$$

$$(1.16) \quad \delta d(fdr_{\wedge}\omega) = r^{-2}fdr_{\wedge}\tilde{\delta}d\omega + ((m-2i)r^{-1}f + f')d\omega,$$

and,

$$(1.17) \quad *(fdr_{\wedge}\omega) = r^{m-2i+2}f\tilde{*}\omega,$$

$$(1.18) \quad d*(fdr_{\wedge}\omega) = (r^{m-2i+2}f' + (m-2i+2)r^{m-2i+1}f)dr_{\wedge}\tilde{*}\omega + r^{m-2i+2}fd\tilde{*}\omega,$$

$$(1.19) \quad *d*(fdr_{\wedge}\omega) = (-1)^{m-i+1}r^{-2}fdr_{\wedge}\tilde{*}d\tilde{*}\omega \\ + (-1)^{(m-i+1)(i-1)}(f' + (m-2i+2)r^{-1}f)\omega,$$

$$(1.20) \quad d*d*(fdr_{\wedge}\omega) = (-1)^{(m-i+1)(i-1)}(f'' + (m-2i+2)r^{-1}f' - (m-2i+2)r^{-2}f)dr_{\wedge}\omega \\ + (-1)^{m-i}r^{-2}fdr_{\wedge}d\tilde{*}d\tilde{*}\omega \\ + (-1)^{(m-i+1)(i-1)}(f' + (m-2i+2)r^{-1}f)d\omega,$$

$$(1.21) \quad d\delta(\omega) = (-f'' - (m-2i+2)r^{-1}f' + (m-2i+2)r^{-2}f)dr_{\wedge}\omega \\ + r^{-2}fdr_{\wedge}d\tilde{\delta}\omega + (-f' - (m-2i+2)r^{-1}f)d\omega.$$

Thus, on  $i$ -forms,

$$(1.22) \quad \Delta(g\phi + fdr_{\wedge}\omega) = (-g'' - (m-2i)r^{-1}g')\phi + r^{-2}g\tilde{\Delta}\phi \\ - 2r^{-3}gdr_{\wedge}\tilde{\delta}\phi + (-f'' - (m-2i+2)r^{-1}f' + (m-2i+2)r^{-2}f)dr_{\wedge}\omega \\ + r^{-2}fdr_{\wedge}\tilde{\Delta}\omega - 2r^{-1}fd\omega.$$

Following [3], for  $\mu \geq 0$ , we set

$$(1.23) \quad \alpha(i) = \frac{1+2i-m}{2}$$

$$(1.24) \nu(i, \mu) = \sqrt{\alpha^2(i) + \mu}$$

$$(1.25) a^\pm(i, \mu) = \alpha(i) \pm \nu(i, \mu)$$

Note that  $a^+(i, \mu) \geq 0$ ,  $a^-(i, \mu) \leq 0$ . Let  $\phi(x)$  be a coclosed  $i$ -form and  $\psi$  a closed  $(i+1)$ -form, with

$$(1.26) \tilde{\Delta}\phi = \mu\phi,$$

$$(1.27) \tilde{\Delta}\psi = \mu\psi,$$

(where the  $\mu$ 's in (1.26), (1.27) may be distinct). Then as in [4], for  $0 < r_1 < r_2 < \infty$ , an harmonic form on  $\overline{C_{r_1, r_2}(N^m)}$  can be written as a convergent series on  $C_{r_1, r_2}(N^m)$  of forms of the following four types (the forms in (1.30) were missed in [3]; see [4] for corrections).

$$(1.28) r^{a^\pm(i)} \phi,$$

$$(1.29) r^{a^\pm(i)} d\phi + a^\pm(i) r^{a^\pm(i)-1} dr \wedge \phi,$$

$$(1.30) r^{a^\pm(i)+2} d\phi + a^\mp(i) r^{a^\pm(i)+1} dr \wedge \phi,$$

$$(1.31) r^{a^\pm(i)+1} dr \wedge \psi.$$

In (1.29), (1.30)  $\phi$  can be taken coexact. If  $a^\pm(i, \mu) = 0$ , we must also introduce  $-()$  solutions  $\log r h_{\frac{m-1}{2}}$ ,  $r \log r dr \wedge h_{\frac{m+1}{2}}$ , where  $\tilde{\Delta}h_j = 0$ .

We say that the forms in (1.28)-(1.30) are of types  $\pm(1)$ ,  $\pm(2+3)$ ,  $\pm(2-3)$ ,  $\pm(4)$  respectively. Those of type (1.28), (1.30) are coclosed and closed respectively. Those of type (1.29) are exact and coexact while those of type (1.30) are neither closed nor coclosed for  $\nu(i) \neq 1$ . For  $\nu(i) = 1$ , the  $+(2+3)$  and  $-(2-3)$  solutions coincide.

As noted in [3] (since  $\bar{d}^* = \bar{\delta}$ ) the sums of  $+( )$  solutions are precisely those harmonic forms which are locally in  $\text{dom } \Delta_G$  near the vertex.

Specifically,

A) Type  $-(1) \in L^2(C_{0,1}(N^m))$ , means  $2\alpha(i)-2\nu(i)-2i+m > -1$ , or

$$(1.32) \nu(i) < 1.$$

If we apply  $d$  to type  $-(1)$ , the resulting type  $-(2+3)$  form is never in  $L^2(C_{0,1}(N^m))$ , (unless we have nonempty ideal boundary).

B) Type  $-(2+3)$  is never in  $L^2(C_{0,1}(N^m))$ .

C) Type  $-(2-3)$  is in  $L^2(C_{0,1}(N^m))$  means  $2\alpha(i)-2\nu(i)+2-2i+m > -1$ , or

$$(1.33) \nu(i) < 2.$$

If  $d, \delta$  of type  $-(2-3)$  (types  $-(1)$ ,  $-(4)$  respectively) are in  $L^2$

$$(1.34) \nu(i) < 1.$$

D) Type  $-(4)$  is in  $L^2(C_{0,1}(N^m))$  means

$$(1.35) \nu(i) < 1.$$

Now assume that we are given  $J$  satisfying 1)-3) of Section 0. In particular,  $J$  preserves homogeneity,  $\phi, \omega$  have homogeneity 0 and  $dr$  has homogeneity 1. Thus there exist linear maps

$$J_+ : \Lambda^i(N^m) \rightarrow \Lambda^{i+1}(N^m),$$

$$J_0 : \Lambda^i(N^m) \rightarrow \Lambda^i(N^m),$$

$$J_0 : \Lambda^i(N^m) \rightarrow \Lambda^i(N^m),$$

$$J_- : \Lambda^i(N^m) \rightarrow \Lambda^{i-1}(N^m),$$

such that if we set

$$(1.36) \begin{aligned} J(\phi(r,x)) + dr \wedge \omega(r,x) &= rJ_+(\omega(r,x)) \\ &+ J_0(\phi(r,x) + dr \wedge J_0(\omega(r,x))) \\ &+ r^{-1} dr \wedge J_-(\phi(r,x)), \end{aligned}$$

then we have

$$(1.37) \Delta J = J \Delta.$$



Of course, we are motivated by the case in which  $J$  is a conical almost complex structure with respect to which the metric is Kähler. But we do not assume this yet.

In view of (1.36), (1.37),  $J$  takes harmonic forms to harmonic forms, preserving homogeneity with respect to  $r$ . We consider then the possibility that the expansion in terms of harmonic forms of  $J(\theta)$  contains a term of the form  $\theta_2$  where  $J(\theta_1)$  and  $\theta_2$  are of two particular pure types  $\pm(1)$ ,  $\pm(2+3)$ ,  $\pm(2-3)$ ,  $\pm(4)$ . This leads to the equations

$$(1.38) \quad a^\pm(i, \mu_\alpha^\pm) = a^\pm(i-1, \mu_\beta^\pm) = a^\pm(i-1, \mu_\gamma^\pm) + 2 = a^\pm(i-2, \mu_\delta^\pm) + 2.$$

For each particular pair of choices, say  $(+, \beta)$  and  $(-, \gamma)$ , we can solve uniquely for say  $\mu_\gamma^-$  in terms of  $\mu_\beta^+$ . This illuminates to some extent the action of  $J$  on harmonic forms for the special case,  $C(S_1^{2n-1}) = \mathbb{C}^n$ . Of course, this action can be calculated explicitly by other methods.

We now make explicit the possible solutions of (1.38) corresponding to pairs containing a + sign and a - sign. The other cases, which are straightforward will be omitted. View each of the expressions in (1.38) as a function of  $\mu$  with  $\alpha$  held fixed. In the first and fourth cases we have  $|\alpha| + \mu > 0$  while in the middle two cases  $\mu > 0$ . Observe that in all cases  $\mu = 0$  is a minimum for functions corresponding to a + sign and a maximum for those corresponding to a - sign. Since  $+( ) \rightarrow +\infty$ , and  $-( ) \rightarrow -\infty$  as  $\mu \rightarrow \infty$ , it follows by inspection that the only possibilities for the degree of homogeneity of a  $+( )$  solution to equal that of a  $-( )$  solution are

$$(1.39) \quad v(i, \mu_\alpha^-) + v(i-1, \mu_\beta^+) = 1,$$

$$(1.40) \quad v(i-2, \mu_{\delta}^{-}) + v(i-1, \mu_{\beta}^{+}) = 1,$$

$$(1.41) \quad v(i-1, \mu_{\gamma}^{-}) + v(i, \mu_{\alpha}^{+}) = 1,$$

$$(1.42) \quad v(i-1, \mu_{\gamma}^{-}) + v(i-2, \mu_{\delta}^{+}) = 1,$$

$$(1.43) \quad v(i-1, \mu_{\gamma}^{-}) + v(i-1, \mu_{\beta}^{+}) = 2.$$

The possibilities (1.39)-(1.42) occur if respectively

$$(1.44) \quad i = \frac{m}{2}, \quad i-1 = \frac{m}{2} - 1,$$

$$(1.45) \quad i-1 = \frac{m}{2}, \quad i-2 = \frac{m}{2} - 1,$$

$$(1.46) \quad i = \frac{m}{2}, \quad i-1 = \frac{m}{2} - 1,$$

$$(1.47) \quad i-1 = \frac{m}{2}, \quad i-2 = \frac{m}{2} - 1.$$

If  $m+1$  is odd, (1.43) corresponds to

$$(1.48) \quad i-1 = \frac{m+1}{2} - \frac{1}{2}, \quad \frac{m+1}{2} - \frac{3}{2}.$$

If  $m+1$  is even, (1.43) corresponds to

$$(1.49) \quad i-1 = \frac{m-1}{2}.$$

It will be important to note that in either of these cases, if

$$(1.50) \quad J(r^{a^{+}(i-1, \mu)} d\phi + a^{+}(i-1, \mu) r^{a^{+}(i-1, \mu)-1} dr \wedge \phi) \\ = r^{a^{-}(i-1, \lambda)+2} d\eta + a^{+}(i-1, \lambda) r^{a^{-}(i-1, \lambda)+1} dr \wedge \eta + h,$$

then multiplying through by  $r^{2-2v(i-1, \mu)} = r^{2v(i-1, \lambda)-2}$ , gives also

$$(1.51) \quad J(r^{a^{-}(i-1, \mu)+2} d\phi + a^{+}(i-1, \mu) r^{a^{-}(i-1, \mu)+1} dr \wedge \phi) \\ = r^{a^{+}(i-1, \lambda)} d\eta + a^{+}(i-1, \lambda) r^{a^{+}(i-1, \lambda)-1} dr \wedge \eta + r^{2-2v(i-1, \mu)} h$$

If we now restrict attention to the case  $m+1$  even we have

Proposition 1.1. Let  $J$  on  $C(N^m)$  be conical. Let the  $i$ -forms  $\theta_1^{+}$  and  $\theta_2^{-}$  be  $+( )$  and  $-( )$  solutions as in (1.28)-(1.31). If the expansion of  $J(\theta_1^{+})$  contains  $\theta_2^{-}$ , then

$$(1.52) \theta_1^+ = r^{a^+(i-1,\mu)} d\phi^+ + r^{a^+(i-1,\mu)-1} dr \wedge \phi_+^+,$$

$$(1.53) \theta_2^- = r^{a^-(i-1,\lambda)+2} d\phi^- + r^{a^-(i-1,\lambda)+1} dr \wedge \phi_-^-,$$

$$(1.54) \nu(i-1,\mu) + \nu(i-1,\lambda) = 2,$$

$$(1.55) i = \frac{m+1}{2}.$$

Example 1.1 Consider the cone on a circle of length  $4\pi$ , the double cover of the complex plane minus the origin. Then  $\nu(0,\mu) = 0, \frac{1}{2}, 1, \frac{3}{2}$ , - where the nonzero values have multiplicity 2. Then we have the possibilities  $\frac{1}{2} + \frac{3}{2} = 2, 1+1 = 2$ .

We now consider briefly  $\Delta_F$ , the self adjoint Friedrichs extension of  $\Delta_0$ . Recall that  $\theta \in \text{dom } \Delta_F$  if  $\theta, \Delta\theta \in L^2$  and there exist compactly supported forms  $\{\theta_j\}$ , such that  $\theta_j \rightarrow \theta, d\theta_j \rightarrow d\theta, \delta\theta_j \rightarrow \delta\theta$ . Whereas for  $\Delta_G$  we had  $d\delta\theta, \delta d\theta \in L^2$ , this is not required for  $\Delta_F$ . However, for  $\Delta_G$  there need only exist possibly distinct sequences of compactly supported forms  $\theta_j \rightarrow \theta, \bar{\theta}_j \rightarrow \bar{\theta}$ , with  $d\theta_j \rightarrow d\theta, \delta\bar{\theta}_j \rightarrow \delta\bar{\theta}$ . It follows easily that  $\theta \in \text{ker } \Delta_F$  implies  $d\theta = \delta\theta = 0$  (but not conversely). It is also easy to check (with the help of a cut-off function) that for  $\nu(i-1,\mu) < 1$  the  $+(2+3)$  solution is in  $\text{dom } \Delta_G \setminus \text{dom } \Delta_F$  and the  $-(2-3)$  solution is in  $\text{dom } \Delta_F \setminus \text{dom } \Delta_G$ . For  $\nu(i-1,\mu) > 1$ ,  $\Delta_F$  and  $\Delta_G$  coincide. But if  $\nu(i-1,\mu) < 2$ ,  $\Delta_0$  is still not essentially self adjoint (this is the "limit circle" case for forms of type  $-(2-3)$ ).

$\Delta J = J\Delta$  on Complex Cones

In the previous section we saw that for even dimensional cones and  $J$  as in Proposition 1.1, the possibility that  $J$  takes  $+$ ( ) solutions to  $-$ ( ) solutions could be ruled out unless  $i = \frac{m+1}{2}$ . In this section we want to show that in case  $J$  is the almost complex structure associated to a Kähler metric with conical singularities then this possibility can be ruled out as well.

Let  $Y^{m+1}$  be a (possibly incomplete) Kähler manifold and suppose that with the possible exception of dimensions  $\frac{m+1}{2}, \frac{m+1}{2} \pm 1$ , every  $L^2$ -harmonic form is in  $\text{dom } \Delta_G$  (see A)-D) of Section 2).

Let  $h \in \Lambda^{(m+1)/2}(Y^{m+1}) \cap L^2$ ,  $\Delta h = 0$  and put

$$(2.1) \quad h = Lx + p$$

where  $p$  is primitive. If  $\Lambda^2 y = x$ , then also

$$(2.2) \quad h = L\Lambda^2 y + p = \Lambda L\Lambda y + p.$$

As is well known, it follows that  $x, y, p \in L^2$ ; see [6]. Since  $Jx$  is an  $(\frac{m+1}{2} - 2)$ -form,  $Jx \in \text{dom } \Delta_G$  and

$$(2.3) \quad \begin{aligned} dJLx &= dLJx \\ &= LdJx \in L^2 \end{aligned}$$

Similarly,

$$(2.4) \quad \begin{aligned} \delta J\Lambda L\Lambda y &= \delta\Lambda J L\Lambda y \\ &= \Lambda\delta J L\Lambda y \in L^2 \end{aligned}$$

Thus  $dJ(h-p), \delta J(h-p) \in L^2$  and in the same way  $d(h-p), \delta(h-p) \in L^2$ .

Thus, if we also assume  $dh, \delta h \in L^2$ , then  $dp, \delta p \in L^2$ . Since

$Lp = \Lambda p = 0$ , by [6], p. 109, we have

$$(2.5) \quad L^2 \ni (d\Lambda - \Lambda d)p \\ = -\delta^c p,$$

$$(2.6) \quad L^2 \ni (\delta L - L\delta)p \\ = d^c p.$$

Thus  $dJp, \delta Jp \in L^2$ . This, together with (2.3), (2.4) implies

$$(2.7) \quad dJh, \delta Jh \in L^2.$$

Essentially the same argument shows that if we make the (global) hypothesis that  $L^2$ -harmonic forms on  $Y^{m+1}$  are closed and coclosed, except perhaps in dimensions  $\frac{m+1}{2}, \frac{m+1}{2} \pm 1$ , then in fact the same holds in dimension  $\frac{m+1}{2}$ . But for primitive  $(\frac{m+1}{2} - 1)$ -forms a similar argument shows only  $dJp \in L^2$  and not  $\delta Jp \in L^2$ .

Now in the previous section, we saw that if the metric and  $J$  are conical, then  $J$  carries  $+( )$  solutions to  $+( )$  solutions with the only possible exception corresponding to (1.43), (1.49). Moreover, if this exception occurs, we can assume that  $J$  of a  $+(2+3)$  solution has a component corresponding to a  $-(2-3)$  solution,  $d$  and  $\delta$  of which are not in  $L^2$  on  $C_{0,\epsilon}(N^m)$ . Indeed, by (1.43), (1.50), (1.51) we can choose the  $-(2-3)$  solution to correspond to  $v \leq 1$ , and our claim then follows from A), D) of Section 1. Thus, applying (2.7), we see that  $J$  carries  $+( )$  harmonic forms to  $+( )$  harmonic forms in all cases. This is the crux of what we wanted to prove.

We now show that subject to the same assumptions,  $J$  actually preserves  $\text{dom } \Delta_G$ . We recall the easily proved statement from [3] that an arbitrary eigenform of  $\Delta$  which is in  $L^2(C_{0,\epsilon}(N^m))$ , admits a convergent series representation, which can be grouped so that the first term is a harmonic form  $h$ . Also, the notions of  $+( )$

and  $-()$  solutions are defined for arbitrary eigenforms. Then our previous analysis immediately implies that for arbitrary  $\lambda$ ,  $J$  as above takes  $+$  eigenforms to  $+$  eigenforms. However, the  $+$  solutions are precisely the eigenforms in the domain of the Laplacian  $\Delta_G$ . Since whether a form is in  $\text{dom } \Delta_G$  can be checked by examining its spectral representation, we now have the following result.

Theorem 2.1. Let  $Y$  be a complete Kähler manifold with isolated metrically conical singularities, and suppose that near the singularities the almost complex structure  $J$  is also conical. Then  $J$  preserves  $\text{dom } \Delta_G$ . In particular,  $J$  preserves  $\ker \Delta_G$ , the space of closed and coclosed harmonic  $L^2$ -forms.

Finally, we observe that the conclusions of Theorem 2.1 remain true, if the hypotheses are only satisfied up to sufficiently high order. In the proposition that follows, we will assume for simplicity that  $\bar{d}^* = \bar{\delta}$ . Clearly it would suffice to only assume this near the singularities, and the case of ideal boundary conditions could also be included.

Proposition 2.2 Let  $X^{m+1}$  have isolated metrically conical singularities, with respect to the metric  $g$ . Let  $g'$  be a second metric on  $X^{m+1}$  for which, in polar coordinates, we have

- i)  $\|g - g'\| = O(r^2)$ ,
- ii)  $\|\nabla(g - g')\| = O(r)$ ,
- iii)  $\|\nabla^2(g - g')\| = O(1)$ ,

where  $\nabla$ , denotes covariant differentiation with respect to  $g$ .

Assume that  $\bar{d}^* = \bar{\delta}$  and let  $\Delta_G, \Delta'_G$  denote the corresponding Laplacians. Then  $\text{dom } \Delta_G = \text{dom } \Delta'_G$ .

Proof: We can assume that near some  $p \in \Sigma$ ,  $g$  is of the form  $C_{0,2}(N^m)$ . Set

$$(2.10) \quad U_j = C_{2^{-(j+2)}, 2^{-(j-1)}}(N^m),$$

$$(2.11) \quad V_j = C_{2^{-(j+1)}, 2^{-j}}(N^m).$$

It follows from the usual elliptic estimate and an obvious scaling argument, that for  $k=0,1,2$ ,

$$(2.12) \quad \|\nabla^k \theta\|_{V_j} \leq c(2^j)^k (\|\theta\|_{U_j} + \|\Delta \theta\|_{U_j})$$

Also, it is easy to check that

$$(2.13) \quad \|(d\delta - d\delta')\theta\|_{V_j} \leq c_2 (\|\theta\|_{V_j} + 2^{-j} \|\nabla \theta\|_{V_j} + 2^{-2j} \|\nabla^2 \theta\|_{V_j})$$

with similar estimates, for  $\|(\delta - \delta')\theta\|, \|(\delta d - \delta' d)\theta\|$ .

Thus,

$$\begin{aligned} (2.14) \quad \|d\delta'\theta\|_{C_{0,1}(N^m)} &\leq \|d\delta\theta\|_{C_{0,1}(N^m)} + \|(d\delta - d\delta')\theta\|_{C_{0,1}(N^m)}, \\ &= \|d\delta\theta\|_{C_{0,1}(N^m)} + \sum_{j=1}^{\infty} \|(d\delta - d\delta')\theta\|_{V_j}, \\ &\leq \|d\delta\theta\|_{C_{0,1}(N^m)} + 9c_1 c_2 (\|\theta\|_{C_{0,2}(N^m)} + \|\Delta \theta\|_{C_{0,2}(N^m)}) \end{aligned}$$

where the last inequality follows from (2.12), (2.13). This together with the corresponding arguments for  $(\delta-\delta'), (\delta d-\delta' d)$  shows that  $\text{dom } \Delta_G' \subseteq \text{dom } \Delta_G$ . However, it is straightforward to check that conditions i)-iii) imply the same conditions with the roles of  $g, g'$  interchanged. Then in the same way one shows that  $\text{dom } \Delta_G \subseteq \text{dom } \Delta_G'$

In the same way (using [2]) we have

Theorem 2.3 Let  $Y$  be a Kähler manifold with isolated singularities. Let  $g'$  denote the metric and  $J'$  the almost complex structure. Assume that near the singularities there exist a conical metric  $g$  and almost complex structure  $J$  such the  $g, g'$  are related as in Proposition 2.2 and

- i)  $\|J-J'\| = O(r^2),$
- ii)  $\|\nabla(J-J')\| = O(r),$
- iii)  $\|\nabla^2(J-J')\| = O(1).$

Then  $J'$  preserves  $\text{dom } \Delta_G'$ .

It is not difficult to check that Theorem 2.3 applies to a complex submanifold  $Y$  of a Kähler manifold  $X$  such that the singularities of  $Y$  are isolated and metrically conical up to higher order terms in local normal coordinates (for example complex projective cones with nonsingular base). It then follows from [2] that for such  $Y$ , the "Kähler package" holds for the middle intersection cohomology groups,  $IH^*(Y)$ . Of course, by now there has been considerable progress on the conjecture that this holds



for arbitrary singular projective complex algebraic varieties; see [2] p. 137 and [5]. However, our purpose here was to show that at least in the simplest cases, this conjecture can be explained by considerations based on the induced metric, and that even for these cases the situation is quite delicate; compare [5], section 4, Conjectures B and C. In fact, the existence of a Hodge theory for the induced metric in the conical case was the original motivation for the conjecture that  $IH^*$  satisfies the Kähler package in general.

Bibliography

- [1] J. W. Alexander, Note on Riemann spaces, B.A.M.S., May (1920) 370-372.
- [2] J. Cheeger, On the Hodge Theory of Riemannian Pseudomanifolds. A.M.S. Proc. Sym. Pure Math XXXVI, 1980.
- [3] J. Cheeger, On the spectral geometry of spaces with cone-like singularities, Proc. Nat. Acad. Sci. (U.S.A.) V. 76, 1979, 2103-2106.
- [4] J. Cheeger, Spectral geometry of spaces with cone-like singularities (Preprint 1978).
- [5] J. Cheeger, R. MacPherson, and M. Goresky,  $L^2$ -cohomology and intersection homology of algebraic varieties, Annals of Math. Studies, 102, Seminar on Differential Geometry, S. T. Yau, ed., Princeton Univ. Press.
- [6] P. Griffiths and J. Harris, Principles of algebraic geometry, J. Wiley, 1978.

J.CHEEGER

State University of New-York  
at Stony Brook

Department of Mathematics  
STONY BROOK, N.Y.11794

U.S.A.