

Astérisque

BOHUMIL CENKL

RICHARD PORTER

**Algebraic categories and the homotopy theory
of some C.W. complexes**

Astérisque, tome 113-114 (1984), p. 173-178

http://www.numdam.org/item?id=AST_1984__113-114__173_0

© Société mathématique de France, 1984, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ALGEBRAIC CATEGORIES AND THE HOMOTOPY THEORY
OF SOME C.W. COMPLEXES
by BOHUMIL CENKL and RICHARD PORTER^(*)

Here we present an equivalence between the homotopy category of certain C.W. complexes of finite type and algebraic homotopy categories of commutative differential graded algebras over the integers and differential graded Lie algebras over the integers.

This result adds to the number of homotopy categories of C.W. complexes of finite type for which there is a known algebraic equivalent [1], [10], [11].

Differential forms provide the bridge between algebra and geometry.

For graded rings, H^* , satisfying the C-r condition (see definition below), the result leads to a parameterization of the set of all homotopy types of 1-connected finite C.W. complexes with integer cohomology ring isomorphic to H^* . Here ; our techniques are motivated by analogous results in rational homotopy theory [5], [6], [7].

For the remainder of this paper, r denotes an integer, $r \geq 3$.

DEFINITION : A graded group H^* (with grading indicated by upper index) is said to satisfy the C-r condition if multiplication by p defines an automorphism on H^{r+k} for all pairs of positive integers (p,k) with $2p - 3 \leq k+1$.

DEFINITION : A graded group H_* (with grading indicated by lower index) is said to satisfy the H-r condition if H_r has no elements of order 2 and multiplication by p defines an automorphism on H_{r+k} for all pairs of positive integers (p,k) with $2p - 3 \leq k+2$.

(*) Richard Porter expresses his gratitude to the Université des Sciences et Techniques de Lille and to the University of Sussex for their hospitality while parts of the work reported here were being done.

REMARK : For X , an $r-1$ connected C.W. complex of finite type, $H^*(X;Z)$ satisfies the $C-r$ condition iff $H_*(X;Z)$ satisfies the $H-r$ condition.

DEFINITION : A space X is called a $C-r$ space if X is $(r-1)$ connected and is homotopy equivalent to a C.W. complex of finite type whose cohomology groups, $H^*(X;Z)$, satisfy the $C-r$ condition.

PROPOSITION : For each integer $r \geq 3$ the homotopy category of $C-r$ spaces is equivalent to

a) an algebraic homotopy category of $(r-1)$ connected commutative differential graded algebras of finite type over the integers whose cohomology groups satisfy the $C-r$ condition ;
and to.

b) an algebraic homotopy category of $(r-1)$ reduced differential graded Lie algebras, L , of finite type over the integers with the property that the homology groups of $L/[L,L]$ satisfy the $H-(r-1)$ condition.

The proof has two main parts. First, the filtered algebra of differential forms introduced in [2] is used to obtain equivalences between the tame homotopy theory of Dwyer [4] and algebraic homotopy theories of commutative differential graded algebras and differential graded Lie algebras [3].

Secondly, it is shown that tame localization restricted to the category of $C-r$ spaces is an equivalence onto its image.

EXAMPLE : For N a product of primes greater than 3 the set of homotopy types of simply connected C.W. complexes X with

$$H^*(X;Z) = \left\{ \begin{array}{ll} Z & * = 0,3 \\ Z/NZ & * = 4,8 \\ 0 & \text{otherwise} \end{array} \right\}$$

and trivial cup product is in 1-1 correspondence with the orbits of the group of

matrices $\begin{pmatrix} a_1 & 0 \\ a_2 & a_3 \end{pmatrix}$ with a_1, a_3 units in Z/NZ and a_2 an arbitrary element

in Z/NZ acting by multiplication on the left on the set of two by one matrices with entries in Z/NZ .

Proof : For k a nonnegative integer, S_k , denotes the smallest subring of the rationals containing $\frac{1}{p}$ for each prime p with $2p - 3 \leq k$.

For $r \geq 3$ an $(r-1)$ connected space X is called tame if π_{r+k} is an S_k

module for each $k \geq 0$. If each $\pi_{r+k}(X)$ is a finitely generated S_k module, we say that X is a tame space of finite type. Tame spaces were introduced by Dwyer in [4] who proved that the homotopy category of tame $(r-1)$ connected spaces ($r \geq 3$) is equivalent to an algebraic homotopy category of tame $(r-1)$ reduced differential graded Lie algebras over Z .

We list some properties of tame spaces :

- (i) Given an $(r-1)$ connected space X , there is a tame space X_T and a map $X \rightarrow X_T$ inducing isomorphisms $\pi_{r+k}(X) \otimes S_k \simeq \pi_{r+k}(X_T)$ for $k \geq 0$.
- (ii) Given a map $f : X \rightarrow Y$ of $(r-1)$ connected spaces there is a map $f_T : X_T \rightarrow Y_T$, unique up to homotopy, such that the diagram below homotopy commutes

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X_T & \xrightarrow{f_T} & Y_T \end{array}$$

We call X_T the tame localization of X .

- iii) The homotopy category of $(r-1)$ connected tame spaces of finite type is equivalent to an algebraic homotopy category of $(r-1)$ connected commutative differential graded algebras of finite type over the integers [3].

The results of [3] can be used to prove

- iv) The homotopy category of $(r-1)$ connected tame spaces of finite type is equivalent to an algebraic homotopy category of $(r-1)$ reduced differential graded Lie algebras of finite type over the integers.

The equivalences in iii) and iv) restricted to tame spaces of the form X_T with X a $C-r$ space give :

- v) The homotopy category of tame spaces of the form X_T for X a $C-r$ space is equivalent to

- a) an algebraic homotopy category of $(r-1)$ connected commutative differential graded algebras of finite type over the integers whose cohomology groups satisfy the $c-r$ condition ;

- b) an algebraic homotopy category of $(r-1)$ reduced differential graded Lie algebras, L , of finite type over the integers such that the homology groups of $L/[L,L]$ satisfy the $H(r-1)$ condition.

The proof of the Proposition is completed by proving :

LEMMA : For $r \geq 3$, tame localization induces an equivalence between the homotopy category of C - r spaces and the homotopy category of tame spaces of the form X_T for some C - r space X .

Proof : It suffices to show that for any two C - r spaces X and Y tame localization induces a bijection between $[X, Y]$ and $[X_T, Y_T]$. From ii) above it follows that $[X_T, Y_T]$ can be replaced by $[X, Y_T]$ and hence it is enough to show that the map $Y \rightarrow Y_T$ induces a bijection $[X, Y] \rightarrow [X, Y_T]$.

From the general theory of Moore-Postnikov systems and factorizations of maps [8] it follows that up to a homotopy equivalence Y is the inverse limit of a sequence of fibrations

$$\begin{array}{ccccccc}
 E_r = Y_T & \xleftarrow{P_{r+1}} & E_{r+1} & \xleftarrow{P_{r+2}} & E_{r+2} & \xleftarrow{\quad} & E_{N-1} \xleftarrow{P_N} E_N \xleftarrow{\quad} \\
 & \nearrow f_r & \nearrow f_{r+1} & \nearrow f_{r+2} & \nearrow f_{N-1} & \nearrow f_N & \\
 & & & & Y = \varprojlim_N E_N & &
 \end{array}$$

where $f_r : Y \rightarrow Y_T$ is tame localization and $E_{N-1} \xleftarrow{P_N} E_N$ is the pullback of the path space fibration over a map from E_{N-1} to the Eilenberg-MacLane space, $K(\pi_N(Y_T, Y), N)$.

Hence the obstructions to the existence of a map $\tilde{g}$ corresponding to the dotted arrow in the solid arrow diagram

$$\begin{array}{ccc}
 & & Y \\
 & \nearrow \tilde{g} & \downarrow f_r \\
 X & \xrightarrow{g} & Y_T
 \end{array}$$

lie in the groups $H^N(X : \pi_N(Y_T, Y))$.

The obstructions to the existence of a map $\tilde{H}$ corresponding to the dotted arrow in the solid arrow diagram

$$\begin{array}{ccc}
 X \times \{0,1\} & \xrightarrow{\quad} & Y \\
 \downarrow & \nearrow \tilde{H} & \downarrow f_r \\
 X \times I & \xrightarrow{H} & Y_T
 \end{array}$$

lie in the groups $H^N((X \times I, X \times \{0,1\}) : \pi_N(Y_T, Y)) \simeq H^{N-1}(X : \pi_N(Y_T, Y))$.

The proof of the lemma is completed by showing :

CLAIM : For any two C - r spaces X and Y , $H^\ell(X : \pi_N(Y_T, Y)) = 0$ for $\ell \geq N-1 \geq r-1$.

Proof of claim : For $k \geq 0$, the long exact homotopy sequence of the pair (Y_T, Y) remains exact when tensored with S_k . Since $\pi_{r+k}(Y) \otimes S_k \rightarrow \pi_{r+k}(Y_T) \otimes S_k$ is an isomorphism it follows that $\pi_{r+k}(Y_T, Y) \otimes S_k = 0$ for $k \geq 0$. For $k = 0$, $S_k = \mathbb{Z}$ and hence $\pi_r(Y_T, Y) = 0$.

To prove that $\pi_{r+1}(Y_T, Y) = 0$ note that since Y is a C - r space, Y has the same rational homotopy type as a (possibly empty) wedge of r -dimensional spheres. The minimal model [9] of a wedge of r -spheres does not have any generators in dimension $r+1$, (recall $r \geq 3$) and hence $\pi_{r+1}(Y)$ is a finite abelian group. It follows that $\pi_{r+1}(Y) \rightarrow \pi_{r+1}(Y_T) \simeq \pi_{r+1}(Y) \otimes S_1$ is an epimorphism and hence $\pi_{r+1}(Y_T, Y) = 0$.

At this point we have shown $H^\ell(X : \pi_N(Y_T, Y)) = 0$ for $N = r$, $N = r+1$.

Assume $\ell \geq N-1 > r$. Write $\ell = r+k$ and $N = r+j+1$ with $k \geq j \geq 1$.

$\pi_N(Y_T, Y) \otimes S_j = 0$ implies that $\pi_N(Y_T, Y)$ is a torsion group each of whose elements has order some product of primes $p_1 \times p_2 \dots \times p_t$ with $2p_i - 3 \leq j+1$ for $1 \leq i \leq t$.

On the other hand, the condition that X be a C - r space implies

that $H^\ell(X : \mathbb{Z}) = H^{r+k}(X : \mathbb{Z})$ is an S_{k+1} module and a finitely generated abelian group. Hence, $H^\ell(X : \mathbb{Z})$ is a finite abelian group each of whose elements has order some product of primes $p_1 \times \dots \times p_t$ with $2p_i - 3 > k+1$ for $1 \leq i \leq t$. Since $k \geq j$ it follows that $H^\ell(X : \mathbb{Z}) \otimes \pi_N(Y_T, Y) = 0$ and $\text{Tor}(H^\ell(X : \mathbb{Z}), \pi_N(Y_T, Y)) = 0$ for $\ell \geq N-1 > r$. Hence $H^\ell(X : \pi_N(Y_T, Y)) = 0$ for $\ell \geq N-1 > r$ by the universal coefficient theorem for cohomology.

REFERENCES

- [1] BAUES, H.J. - The homotopy theory in a cofibration category, preprint.
- [2] CENKL, B and PORTER, R. - Differential forms and torsion in the fundamental group. To appear in Advances in Math.
- [3] CENKL, B. and PORTER, R. - Tame Homotopy Theory. Proceedings of the summer conference on manifolds and homotopy theory, 1980.
- [4] DWYER, W.G. - Tame Homotopy Theory, Topology, Vol. 18, 321-338 (1979).
- [5] FELIX, Y. Dénombrement des Types de K-Homotopie Théorie de la Déformation, Mémoire de la Société Mathématique de France, tome 108/Fascicule 3.
- [6] HALPERIN, S. and STASHEFF, J. - Obstructions to Homotopy Equivalences, Advances in Math. 32, 233-279 (1979).
- [7] SCHLESSINGER, M. and STASHEFF, J. - Deformation theory and rational homotopy type, preprint.
- [8] SPANIER, E.H. - Algebraic Topology, MacGraw Hill, 1966.
- [9] SULLIVAN, D. - Infinitesimal computations in topology, Pub. Math. 47 (1978) I.H.E.S. 269-332.
- [10] WHITEHEAD, J.H.C. - Combinatorial Homotopy I., Bull. Amer. Math. Soc. 55, 213-245 (1949)
- [11] WHITEHEAD, J.H.C. - The homotopy type of a special kind of polyhedron. Ann. Soc. Polon. Math. 21 (1948) 176-186 (1949).
