

COMPOSITIO MATHEMATICA

G. FODOR

On a theorem in the theory of binary relations

Compositio Mathematica, tome 8 (1951), p. 250

http://www.numdam.org/item?id=CM_1951__8__250_0

© Foundation Compositio Mathematica, 1951, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

On a theorem in the theory of binary relations

by

G. Fodor

Szeged.

This paper was inspired by a theorem of Lázár¹⁾ which we shall state below.

Let M be a set of positive measure. To every $x \in M$ we adjoin a set of elements $y \neq x$ of M which may have the cardinal number of the continuum. This means that we define a function $y = \varphi(x)$ of $x \in M$, where $y \in M$, which may assume continuously many values for every x , whereas the equation $x = \varphi(x)$ cannot occur. If neither of the two equations $y = \varphi(x)$ and $x = \varphi(y)$ holds, the two elements x and y are called independent.

Let $\{\varphi(x)\}$ denote the set of values $\varphi(x)$ for a given x ; assume moreover that x is not a point of accumulation of $\{\varphi(x)\}$ and that $\{\varphi(x)\}$ is of measure zero.

THEOREM. We can find a set of positive exterior measure of elements of M so that any pair of its elements are independent.

Proof. As the set $\{\varphi(x)\}$ does not contain x and as x is no point of accumulation, its complement $C\{\varphi(x)\}$ with respect to M contains x and an interval I_x surrounding x . Now for every x let us choose in I_x a closed segment $S(x)$ with rational endpoints. Hence all values of $\varphi(x)$ are situated outside $S(x)$.

Now the segments $S(x)$ form an enumerable system S_1, S_2, \dots . To every S_n there belongs at least one $x \in M$, such that $S(x) = S_n$. Let N_n denote the set of all $x \in M$ with $S(x) = S_n$. We assert that there is at least one segment S_n for which N_n has positive exterior measure. For suppose that to every S_n the set N_n would be of measure zero. Then, as $M \subset N_1 + N_2 + \dots$, by the well-known theorem that the sum of enumerably many sets of measure zero is a set of measure zero, we must conclude that M was a set of measure zero.

The elements $x \in N_n$ all belong to S_n ; they are evidently independent as the adjoined values are all outside the segment S_n .

Lázár's theorem²⁾ only states that under the same assumptions we can find a set with the power of the continuum.

¹⁾ *Compositio Mathematica* 3 (1936), p. 304.

²⁾ Evidently Lázár uses the word "condensationpoint" where only a point of accumulation is meant.