

COMPOSITIO MATHEMATICA

B. H. NEUMANN

Corrigendum and addendum to “Ascending derived series”

Compositio Mathematica, tome 13 (1956-1958), p. 128

http://www.numdam.org/item?id=CM_1956-1958__13__128_0

© Foundation Compositio Mathematica, 1956-1958, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Corrigendum and addendum to “Ascending derived series”

by

B. H. Neumann*

Dr Graham Higman has pointed out to me that the construction of the example in § 9 is incorrect as it stands, because H_i'' is not, as claimed, a direct product with amalgamations of copies of H_{i-2} , but of copies of H_{i-1}' (p. 61, lines 12—11 from the bottom). I am indebted to Dr Higman for the following modification of the construction, to give an example with all the properties stated in the last 8 lines of § 9. The definition of the groups H_i remains unchanged, but we denote their inductive limit by H^* ; we denote by L_{i-1} the group generated by all the isomorphic copies that arise from H_i by forming the successive direct products with amalgamated Z , for all positive i : Thus L_{2i-1} is what was denoted by G_{2i} ; what was denoted by G_{2i-1} will again be denoted by G_{2i-1} . It follows from (8.3) that $L_i'' = L_{i-1}'$. Thus if we put $L_i' = G_i$ for all positive i , then $G_{i+1}' = G_i$, and

$$\{1\} = G_0 \subset G_1 \subset G_2 \subset \dots$$

is an infinite ascending derived series, with $G_1 = Z$ of order 2. The inductive limit of this series is again denoted by G^* . The construction in § 10 requires no modification.

Dr Higman has also remarked that a consequence of Theorem 7.3 is the following COROLLARY: *A finitely generated group is isomorphic to a term of its derived series only if it coincides with it.*

The University, Manchester, 13.

(Oblatum 10-2-57).

*) Compositio Math 13, 47—64 (1956).