

# COMPOSITIO MATHEMATICA

RAM SINGH

**Correction : “On a class of starlike functions”**

*Compositio Mathematica*, tome 21, n° 2 (1969), p. 230-231

[http://www.numdam.org/item?id=CM\\_1969\\_\\_21\\_2\\_230\\_0](http://www.numdam.org/item?id=CM_1969__21_2_230_0)

© Foundation Compositio Mathematica, 1969, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Correction

### On a class of starlike functions

by

Ram Singh

Compositio Mathematica 19 (1968), 78-82

### Introduction

In a recent paper (Comp. Math., 19 (1968), 78-82) with the above title we investigated the class  $\bar{S}$  of functions

$$f(z) = z + a_2 z^2 + \dots$$

that are analytic and starlike in  $|z| < 1$  and satisfy the condition  $|\{z f'(z)/f(z)\} - 1| \leq 1$  in  $|z| < 1$ . Theorem 4 of that paper, giving the radius of convexity of the class  $\bar{S}$ , contains an error and as such the result obtained is not sharp. Below we give the correct proof of that theorem.

**THEOREM.** *Each function  $f(z) \in \bar{S}$  maps*

$$|z| < \frac{(3 - \sqrt{5})}{2}$$

*onto a convex domain.*

**PROOF.** If  $f(z) \in \bar{S}$ , we can write

$$(1) \quad z \frac{f'(z)}{f(z)} = 1 + \psi(z),$$

when  $\psi(z)$  is analytic in  $|z| < 1$  and satisfies  $|\psi(z)| \leq 1$  and  $\psi(0) = 0$ . Logarithmic differentiation of (1) yields:

$$z \frac{f''(z)}{f'(z)} = \frac{z\psi'(z)}{1 + \psi(z)} + \psi(z).$$

Making use of the fact that  $|\psi(z)| \leq |z|$  and

$$|\psi'(z)| \leq \frac{1 - |\psi(z)|^2}{(1 - |z|^2)},$$

we find that

$$\begin{aligned} \left| z \frac{f''(z)}{f'(z)} \right| &\leq \frac{|z|(1-|\psi(z)|^2)}{(1-|\psi(z)|)(1-|z|^2)} + |\psi(z)| \\ &= \frac{|z|(1+|\psi(z)|)}{(1-|z|^2)} + |\psi(z)| \\ &\leq \frac{2|z|-|z|^2}{(1-|z|)}. \end{aligned}$$

Therefore,  $f(z)$  is convex if

$$\frac{2|z|-|z|^2}{1-|z|} < 1$$

or

$$|z| < \frac{(3-\sqrt{5})}{2}.$$

To show that  $(3-\sqrt{5})/2$  is the exact radius of convexity, we consider the function

$$f_0(z) = z \exp(z).$$

A little calculation shows that for this function

$$1 + \frac{zf_0''(z)}{f_0'(z)} = \frac{z^2+3z+1}{1+z}$$

vanishes when  $z = (\sqrt{5}-3)/2$ .