

COMPOSITIO MATHEMATICA

JANOS GALAMBOS

The absolute continuity of a limit law for Sylvester series

Compositio Mathematica, tome 35, n° 2 (1977), p. 125-128

<http://www.numdam.org/item?id=CM_1977__35_2_125_0>

© Foundation Compositio Mathematica, 1977, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

THE ABSOLUTE CONTINUITY OF A LIMIT LAW FOR SYLVESTER SERIES*

Janos Galambos

Every real number $x \in (0, 1]$ has a unique representation in the form

$$(1) \quad x = 1/d_1 + 1/d_2 + \cdots + 1/d_n + \cdots$$

where the $d_j = d_j(x)$ are positive integers satisfying

$$(2) \quad d_1 \geq 2 \quad \text{and} \quad d_{j+1} \geq d_j^2 - d_j + 1, \quad j \geq 1.$$

The representation (1) and (2) is called the Sylvester expansion of x . It was shown in [1] that, as $n \rightarrow +\infty$,

$$(3) \quad \lim 2^{-n} \log d_n(x) = \beta(x)$$

exists and is positive. The behaviour of $\beta(x)$ drew much attention in the literature; see, in particular, Vervaat [5], pp. 148–151, who particularly stresses the need for solving the following problem (correct the typing error and thus $\log d_n$ is to replace d_n in part (b) of his statement). Put

$$(4) \quad F(z) = P(\beta(x) < z),$$

where P stands for Lebesgue measure. The question is whether the function $F(z)$ is continuous for all real z . In the present note we settle this question by proving the following result.

THEOREM. *The distribution function $F(z)$ of (4) is absolutely continuous. Its derivative $f(z)$ is continuous for all z .*

In the proof, we need the following lemmas.

* This research was done while the author was on Research and Study Leave from Temple University and, as a Fellow of the Humboldt Foundation, he was at the Goethe University, Frankfurt am Main.

LEMMA 1: *Let $\varphi(t)$ be the characteristic function of $F(z)$. Assume that $|\varphi(t)|$ is integrable over the whole real line. Then the derivative $f(z)$ of $F(z)$ exists and is continuous for all z .*

Its proof can be found in [4], p. 267.

The following lemma was obtained in [2], p. 188.

LEMMA 2: $\beta(x)$ of (3) has the form

$$\beta(x) = \frac{1}{2} \log d_1(x) + \frac{1}{2} \sum_{n=1}^{+\infty} 2^{-n} \log \frac{d_{n+1}}{d_n^2}.$$

Our last lemma was independently discovered by the present author [3], p. 138 and by Vervaat [5], p. 111.

LEMMA 3: *Put in (1)*

$$(5) \quad x_n = 1/d_n + 1/d_{n+1} + \dots.$$

Then $y_n = d_n(d_n - 1)x_{n+1}$ is uniformly distributed on the interval $(0, 1)$. Furthermore, y_n is stochastically independent of $d_1, d_2, \dots, d_n$.

PROOF OF THE THEOREM: Put

$$\beta_N(x) = \frac{1}{2} \log d_1(x) + \frac{1}{2} \sum_{n=1}^N 2^{-n} \log \{d_{n+1}(x)/d_n^2(x)\}$$

and

$$\varphi_N(t) = \int_0^1 \exp \{it\beta_N(x)\} dx, \quad t \text{ real.}$$

By Lemma 2,

$$(6) \quad \varphi(t) = \lim \varphi_N(t) \quad (N \rightarrow +\infty).$$

Our aim now is to get a recursive formula for $\varphi_N(t)$. This is done by plugging in y_N into the formula for $\beta_N(x)$ and we then apply Lemma 3. We write

$$(7) \quad \frac{d_{N+1}}{d_N^2} = \frac{d_{N+1}x_{N+1}}{y_N} \cdot \frac{d_N - 1}{d_N}.$$

Since, by (2) and (5),

$$1 < d_{N+1}x_{N+1} \leq d_{N+1}/(d_{N+1} - 1),$$

(7) reduces to

$$d_{N+1}/d_N^2 = (1/y_N)(1 + \vartheta/d_N), \quad |\vartheta| \leq 1.$$

Hence,

$$(8) \quad \beta_N(x) = \beta_{N-1}(x) - 2^{-N-1} \log y_N + 2^{-N-1} \log(1 + \vartheta/d_N),$$

where $|\vartheta| \leq 1$. Before we turn to $\varphi_N(t)$, we estimate the last term of (8), in order to make it independent of x . First define the numbers D_n by

$$D_1 = 2, \quad D_{n+1} = D_n^2 - D_n + 1, \quad n \geq 1.$$

In view of (2), $d_n(x) \geq D_n$ for all n and all x . Thus

$$(9) \quad |\log(1 + \vartheta/d_N)| \leq 2/D_N.$$

We now appeal to Lemma 3. It says that

$$(10) \quad \int_0^1 \exp \{it \log(1/y_N(x))\} dx = 1/(1 - it).$$

Also, since $\beta_{N-1}(x)$ depends on $d_1, d_2, \dots, d_N$ only, $\beta_{N-1}(x)$ and $y_N(x)$ are independent. Therefore, as is well known from the elements of probability theory, (8)–(10) yield

$$(11) \quad \varphi_N(t) = \varphi_{N-1}(t)(1 - it/2^{N+1})^{-1}(1 + \vartheta_N t/2^{N+1} D_N),$$

where $|\vartheta_N| \leq 2$. We thus have from (6) and (11) that, for an arbitrary integer $M \geq 1$ and with a suitable sequence ϑ_N of numbers which satisfy $|\vartheta_N| \leq 2$,

$$\varphi(t) = \varphi_M(t) \prod_{N=M+2}^{+\infty} \{(1 - it/2^N)^{-1}(1 + \vartheta_{N-1} t/2^N D_{N-1})\}.$$

From the above formula, one can easily estimate the tails of $\varphi(t)$ by making use of the fact that $\log D_N > 2^{N-3}$ and that $|\varphi_M(t)| \leq 1$. Combining these estimates with the fact that, as any characteristic function, $\varphi(t)$ is continuous, we obtain that $|\varphi(t)|$ is integrable on the whole real line. Lemma 1 therefore completes the proof.

REFERENCES

- [1] P. ERDÖS, A. RÉNYI and P. SZÜSZ: On Engel's and Sylvester's series. *Annales Univ. Sci. Budapest, Sectio Math.* 1 (1958) 7–12.
- [2] J. GALAMBOS: The ergodic properties of the denominators in the Oppenheim expansion of real numbers into infinite series of rationals. *Quart. J. Math. Oxford Second Ser.* 21 (1970) 177–191.
- [3] J. GALAMBOS: Further ergodic results on the Oppenheim series. *Quart. J. Math. Oxford Second Ser.* 25 (1974) 135–141.
- [4] A. RÉNYI: *Wahrscheinlichkeitsrechnung mit einem Anhang über Informationstheorie*. VEB Deutscher Verlag der Wissenschaften, Berlin, 1962.

- [5] W. VERVAAT: Success epochs in Bernoulli trials with applications in number theory, *Mathematical Centre Tracts*, Vol. 42, Mathematisch Centrum Amsterdam 1972.

(Oblatum 8-IX-1975)

Department of Mathematics
Temple University
Philadelphia, Pa. 19122
USA.