

COMPOSITIO MATHEMATICA

JERZY JURKIEWICZ

Linearizing some $\mathbb{Z}/2\mathbb{Z}$ actions on affine space

Compositio Mathematica, tome 76, n° 1-2 (1990), p. 243-245

<http://www.numdam.org/item?id=CM_1990__76_1-2_243_0>

© Foundation Compositio Mathematica, 1990, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Linearizing some $\mathbb{Z}/2\mathbb{Z}$ actions on affine space

JERZY JURKIEWICZ

Institute of Mathematics, University of Warsaw, P.K.i. N. 9p., 00–901, Warsaw, Poland

Received 3 December 1988; accepted 20 July 1989

Let V be the affine space k^n over an algebraically closed field k , G a linearly reductive group and $A: G \times V \rightarrow V$ a group action with a fixed point, say the origin. Then for all $g \in G$ let me denote by $A(g)$ the corresponding automorphism of V . We have

$$A(g) = L(g) + D(g)$$

where $L(g), D(g) \in \text{End } V$, $L(g)$ linear and $D(g)$ the sum of terms of higher degrees. Let me recall the well known linearization problem: is the action A linearizable, i.e. conjugated to the linear action $L: G \times V \rightarrow V$ (see e.g. [B] and [K])? Recently counter-examples have been found, see [S] and [K + S], so it is reasonable to study additional assumptions on the action A . One of them is considered in the present paper.

First I want to define some morphism $\sigma_A: V \rightarrow V$ which turns out to be a conjugating automorphism for A , provided σ_A is invertible. It will be done using the Reynolds operator i.e. the equivariant projection $\rho: \mathcal{O}(G) \rightarrow k$. For a finite dimensional k -space W we have the unique linear map $\int_G: \text{Mor}(G, W) \rightarrow W$ such that for all linear maps $f: W \rightarrow k$ the induced diagram

$$\begin{array}{ccc} \int_G : \text{Mor}(G, W) & \longrightarrow & W \\ \downarrow f_* & & \downarrow f \\ \rho : \mathcal{O}(G) & \longrightarrow & k \end{array}$$

is commutative. Now let $\phi: G \rightarrow \text{End}(V)$ be such a map that the induced map $G \times V \rightarrow V$ is an algebraic morphism. Then $W := \text{lin hull}(\phi(G))$ is finite dimensional, hence $\int_G \phi$ is a well-defined element of $\text{End}(V)$. Let us apply the above to the map $\phi: G \ni g \mapsto L(g^{-1})A(g) \in \text{End}(V)$ and set $\sigma = \sigma_A = \int \phi$ (compare [J]). We have

$$L(h)\sigma = \int_{g \in G} L(h)L(g^{-1})A(g) = \left(\int_{g \in G} L(hg^{-1})A(gh^{-1}) \right) A(h) = \sigma A(h)$$

for all $h \in G$.

So σ invertible implies that $A(h) = \sigma^{-1}L(h)\sigma$. In particular the action A is linearizable. Later we will give an example of an action A which can be linearized but for which σ_A is not invertible.

As mentioned in [J], the morphism σ_A can be interpreted as an average deviation of A from being linear.

CONJECTURE (Kraft, Procesi). Assume for some $d \geq 2$

$$A(g) = L(g) + H_d(g) + H_{d+1}(g) + \cdots + H_{2d-2}(g), \quad \text{for all } g,$$

where $H_m(g)$ is a homogeneous endomorphism of V of degree m . Then σ_A is invertible. In particular the action A is linearizable.

THEOREM. *The above conjecture is true in the following cases*

1. G linearly reductive, $d = 2$ and $\text{char } k \neq 2$,
2. G diagonalizable, $d = 2$ and $\text{char } k$ arbitrary,
3. $G = \mathbb{Z}/2\mathbb{Z}$, d arbitrary and $\text{char } k = 0$.

Cases 1 and 2 are the objects of [J].

Proof for the case 3. Let I denote the identity map of V . We can write: $G = \{I, L + D\}$, where L and D are endomorphisms of V , L linear and $D = H_d + \cdots + H_{2d-2}$. We have $L^2 = (L + D)^2 = I$. It follows that

$$LD + D(L + D) = 0. \tag{1}$$

Let me denote by \tilde{H}_d the d -linear symmetric map from V^d to V corresponding to H_d . Then we have

$$D(L + D) = DL + d\tilde{H}_d(L, \dots, L, H_d) + \cdots$$

where the first summand consists of terms of degrees $d, \dots, 2d - 2$, the second is of degree $2d - 1$ and all further summands have higher degrees. Considering the possible cancellations in (1) we obtain:

$$-LD = DL = D(L + D). \tag{2}$$

By definition $\sigma = \frac{1}{2}(I + (I + LD)) = I - \frac{1}{2}DL$. We will prove that $I + \frac{1}{2}DL$ is the inverse of σ .

LEMMA. $D(I + mDL) = D$ for $m = 0, 1, 2, \dots$

Proof. Suppose the above holds for some $m - 1, m > 0$. By (2), $D = D(I + DL)$. Therefore

$$D = D(I + (m - 1)DL)(I + DL) = D(I + DL + (m - 1)DL(I + DL)).$$

On the other hand $DL(I + DL) = -LD(I + DL) = -LD = DL$, and we are done.

Since $\text{char}(k) = 0$ the Lemma implies that $D(I + rDL) = D$ for all $r \in k$. Then taking $r = \frac{1}{2}$ we have

$$(I - \frac{1}{2}DL)(I + \frac{1}{2}DL) = I + \frac{1}{2}DL + \frac{1}{2}LD(I + \frac{1}{2}DL) = I,$$

and the same applies if we interchange the order of factors at the left hand side. Q.E.D.

EXAMPLE OF A NON INVERTIBLE σ . Let the linear endomorphism L of k^2 be given by $L(x, y) = (x, -y)$ and an automorphism τ by $\tau(x, y) = (x - (x + y)^2, y + (x + y)^2)$ so that $\tau^{-1}(x, y) = (x + (x + y)^2, y - (x + y)^2)$. The automorphism $\tau^{-1}L\tau$ has order two, so it defines an action of the group of order two on k^2 . The corresponding endomorphism $\sigma = \frac{1}{2}(I + L\tau^{-1}L\tau)$ takes (x, y) to $(x - u + v, y + u + v)$, where $u = \frac{1}{2}(x + y)^2$, $v = \frac{1}{2}(x - y - 2(x + y)^2)^2$. Direct computation shows that the Jacobian determinant of σ is

$$J(\sigma) = 1 - 4(x^2 + y^2) + 8(x^3 + y^3) + 24(x^2y + xy^2).$$

Therefore the endomorphism σ is not invertible, while the considered group action can obviously be linearized.

References

[B] Bass, H., Algebraic group actions on affine spaces. Group actions on rings. Contemporary Mathematics, Vol. 43 (1985), AMS, 1–24.
 [J] Jurkiewicz, J., On the linearization of actions of linearly reductive groups; Commentari Mathematici Helvetici 64 (1989) 508–513.
 [K] Kraft, H., Algebraic group actions on affine spaces. Geometry today, Progress in Mathematics. Birkhäuser 1984, 251–266.
 [S] G.V. Schwarz, Exotic algebraic group actions, preprint.
 [K + F] Kraft, H. and Schwarz, G. V., Reductive group actions on affine spaces with one-dimensional quotient. To appear in Contemporary Mathematics, Proceedings of the Conference on Group Actions and Invariant Theory, Montreal 1988.