

COMPOSITIO MATHEMATICA

KATSURO SAKAI

**Connecting direct limit topologies with metrics
on infinite-dimensional manifolds**

Compositio Mathematica, tome 81, n° 2 (1992), p. 237-245

http://www.numdam.org/item?id=CM_1992__81_2_237_0

© Foundation Compositio Mathematica, 1992, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Connecting direct limit topologies with metrics on infinite-dimensional manifolds

KATSURO SAKAI

Institute of Mathematics, University of Tsukuba, Tsukuba-city 305, Japan

Received 20 November 1990; accepted 23 April 1991

Abstract. Let \mathbb{R}^∞ and Q^∞ be the direct limits of the towers $\mathbb{R}^1 \subset \mathbb{R}^2 \subset \mathbb{R}^3 \subset \dots$ and $Q^1 \subset Q^2 \subset Q^3 \subset \dots$, where $Q = [-1, 1]^\omega$ is the Hilbert cube. And let σ and Σ be the spaces $\bigcup_{n \in \mathbb{N}} \mathbb{R}^n$ and $\bigcup_{n \in \mathbb{N}} Q^n$ with relative topologies inherited from the product topologies of \mathbb{R}^ω and Q^ω , respectively. Let M_i ($i = 1, 2$) have two topologies $\tau_1^i \subset \tau_2^i$ such that (M_i, τ_1^i) is a σ -manifold and (M_i, τ_2^i) is an \mathbb{R}^∞ -manifold (or (M_i, τ_1^i) is a Σ -manifold and (M_i, τ_2^i) is a Q^∞ -manifold). If (M_1, τ_1^1) and (M_2, τ_1^2) have the same homotopy type, there exists $h: M_1 \rightarrow M_2$ which is a homeomorphism in both topologies. This gives the answers to the questions in [We, NLC15].

0. Introduction

A manifold modeled on a given space E is called an E -manifold. We naturally identify \mathbb{R}^n with the subset of the countable infinite product \mathbb{R}^ω . The set $\bigcup_{n \in \mathbb{N}} \mathbb{R}^n$ has two natural topologies,⁽¹⁾ that is, the weak topology with respect to the tower $(\mathbb{R}^n)_{n \in \mathbb{N}}$ and the relative topology inherited from the product topology of \mathbb{R}^ω . We denote these spaces by $\text{dir lim } \mathbb{R}^n$ and \mathbb{R}_f^ω or simply by \mathbb{R}^∞ and σ , respectively. The realization $|K|$ of a simplicial complex K has also two natural topologies, that is, the weak (or Whitehead) topology and the metric topology. These spaces are denoted by $|K|_w$ and $|K|_m$, respectively. A countable simplicial complex is called a *combinatorial ∞ -manifold* if the star of each vertex is combinatorially equivalent to the countable infinite full simplicial complex Δ^∞ [Sa4]. It is shown in [Sa5] (cf. [Sa6]) that for a combinatorial ∞ -manifold K , $|K|_w$ is an \mathbb{R}^∞ -manifold and $|K|_m$ is a σ -manifold. In [Pa, §3], R. S. Palais pointed out a Banach manifold of fiber bundle sections admits an atlas such that the transitions are also homeomorphisms with respect to the bounded weak topology of the model (cf. [He1]). Then it is natural to ask whether a combinatorial ∞ -manifold has the similar structure, that is, it admits an atlas such that the transitions are homeomorphisms in both topologies. Such a manifold may be called an $(\mathbb{R}^\infty, \sigma)$ -manifold. Concerning $(\mathbb{R}^\infty, \sigma)$ -manifolds, we can ask the same questions as [Ge, NLC9] (cf. [We, NLC15]).

Let $Q = [-1, 1]^\omega$ be the Hilbert cube. Similarly as the above, we can define $Q^\infty = \text{dir lim } Q^n$ and $\Sigma = \bigcup_{n \in \mathbb{N}} Q^n$ as the subspace of Q^ω . Let $\text{LIP}(X, Y)$ be the

⁽¹⁾According to [Ke], such a space with two topologies is called a *bitopological space*.

space of Lipschitz maps from a compact metric space X to a metric space Y with sup-metric. For $k > 0$, let $k\text{-LIP}(X, Y)$ be the subspace of $\text{LIP}(X, Y)$ consisting of all k -Lipschitz maps (i.e., maps of Lipschitz constant $\leq k$). Then $\text{LIP}(X, Y)$ has another topology, namely the weak topology with respect to the tower $(n\text{-LIP}(X, Y))_{n \in \mathbb{N}}$. We denote this space by $\text{LIP}(X, Y)_w$. It is proved in [Sa7] and [Sa8] that $\text{LIP}(X, Y)$ is a Σ -manifold and $\text{LIP}(X, Y)_w$ is a Q^∞ -manifold in case X is non-discrete and Y is a Euclidean polyhedron without isolated points or a Lipschitz manifold with $\dim Y > 0$. In this setting, we may also ask whether $\text{LIP}(X, Y)$ admits an atlas such that the transitions are homeomorphisms in both topologies. Such a manifold may be called a (Q^∞, Σ) -manifold. Concerning (Q^∞, Σ) -manifolds, we can ask similar questions (cf. [We, NLC15]).

In this paper, we give affirmative answers for these questions by proving the following theorem:

MAIN THEOREM. *Let M_i ($i=1, 2$) have two topologies $\tau_1^i \subset \tau_2^i$ such that (M_i, τ_1^i) is a σ -manifold and (M_i, τ_2^i) is an \mathbb{R}^∞ -manifold (or (M_i, τ_1^i) is a Σ -manifold and (M_i, τ_2^i) is a Q^∞ -manifold). If (M_1, τ_1^1) and (M_2, τ_1^2) have the same homotopy type, there exists $h: M_1 \rightarrow M_2$ which is a homeomorphism in both topologies.*

1. Corollaries and remarks on the model spaces

Combining the main theorem with [Sa5, Corollary 3 (resp. 3')] or the open embedding theorem for σ -manifolds (resp. Σ -manifolds) [Ch, Theorem 2.4], we have the following:

COROLLARY 1. *Any \mathbb{R}^∞ -manifold or any σ -manifold has the unique $(\mathbb{R}^\infty, \sigma)$ -manifold structure. And any Q^∞ -manifold or any Σ -manifold has the unique (Q^∞, Σ) -manifold structure.*

In the sequel corollaries, we assume the following:

(*) *M has two topologies $\tau_1 \subset \tau_2$ such that (M, τ_1) is a σ -manifold and (M, τ_2) is an \mathbb{R}^∞ -manifold (or (M, τ_1) is a Σ -manifold and (M, τ_2) is a Q^∞ -manifold).*

Combining the main theorem with the corresponding results for σ -manifolds (or Σ -manifolds) in [Ch] or by results in [Sa6], we have the following corollaries:

COROLLARY 2 (Open embedding theorem). *There exists an open embedding $h: (M, \tau_1) \rightarrow \sigma$ (or $h: (M, \tau_1) \rightarrow \Sigma$) such that $h: (M, \tau_2) \rightarrow \mathbb{R}^\infty$ (or $h: (M, \tau_2) \rightarrow Q^\infty$) is also an open embedding. Hence M is an $(\mathbb{R}^\infty, \sigma)$ -manifold (or a (Q^∞, Σ) -manifold).*

COROLLARY 3 (Triangulation theorem). *There exists a simplicial complex K and a homeomorphism $h: (M, \tau_1) \rightarrow |K|_m$ (or $h: (M, \tau_1) \rightarrow |K|_m \times Q$) such that $h: (M, \tau_2) \rightarrow |K|_w$ (or $h: (M, \tau_2) \rightarrow |K|_w \times Q$) is also a homeomorphism. (Then K must be a combinatorial ∞ -manifold [Sa6]).*

COROLLARY 4 (Stability theorem). *There exists a homeomorphism $h: (M, \tau_1) \times \sigma \rightarrow (M, \tau_1)$ (or $h: (M, \tau_1) \times \Sigma \rightarrow (M, \tau_1)$) such that $h: (M, \tau_2) \times \mathbb{R}^\infty \rightarrow (M, \tau_2)$ (or $h: (M, \tau_2) \times Q^\infty \rightarrow (M, \tau_2)$) is also a homeomorphism.*

COROLLARY 5 (Negligibility theorem). *If A is a Z -set in (M, τ_1) then there exists a homeomorphism $h: M \setminus A \rightarrow M$ in both topologies τ_1 and τ_2 .*

Here a closed set A in a space X is said to be a Z -set if the identity map of X is arbitrarily close to maps $f: X \rightarrow X \setminus A$.

The main theorem and Corollaries 2 and 4 answer affirmatively to the questions for $(\mathbb{R}^\infty, \sigma)$ -manifolds (or (Q^∞, Σ) -manifold) corresponding to [Ge, NLC9] (cf. [We, NLC15]). And Corollary 2 gives affirmative answers for the questions concerning a combinatorial ∞ -manifold and $LIP(X, Y)$.

Here we observe the topologies on the model spaces. One should note that $\bigcup_{n \in \mathbb{N}} \mathbb{R}^n$ has many different metric topologies. For instance, there are the relative topologies inherited from $l_p, p \in \mathbb{N} \cup \{\infty\}$. We denote these spaces by l_p^f . For each $n \in \mathbb{N}$, let $e_n \in \mathbb{R}^\omega$ be defined by $e_n(n) = 1$ and $e_n(i) = 0$ if $i \neq n$. Then e_n is convergent to 0 in σ but not in l_p^f for any p . If $p > q$, $\sum_{i=1}^n n^{-1} \cdot e_i$ is convergent to 0 in l_p^f but not in l_q^f . However it is well-known that all l_p^f are homeomorphic to σ . The main theorem asserts that these are homeomorphic to σ by a homeomorphism of \mathbb{R}^∞ onto itself. It is also known that $\text{dir lim } I^n$ and l_p^∞ are homeomorphic to \mathbb{R}^∞ and σ , respectively. But now it is shown that they are homeomorphic by a single homeomorphism. One should note that Q^∞ and Σ are represented by linear topological spaces. Let \mathbb{R}_Q^∞ denote the space $\bigcup_{n \in \mathbb{N}} n \cdot Q$ with the relative topology inherited from \mathbb{R}^ω . Then $\text{dir lim } n \cdot Q$ and \mathbb{R}_Q^∞ are linear topological spaces which are homeomorphic to Q^∞ and Σ respectively. Now it is also shown that they are homeomorphic by a single homeomorphism. Let $Q_0 = \prod_{i \in \mathbb{N}} [-2^{-i}, 2^{-i}]$ be another expression of the Hilbert cube. Then Q_0 is contained in all l_p as a subspace. Let $l_p^{Q_0}$ denote the space $\bigcup_{n \in \mathbb{N}} n \cdot Q_0$ with the relative topology inherited from l_p . Finally we note that $\text{dir lim } n \cdot Q_0$ and $l_p^{Q_0}$ are homeomorphic to Q^∞ and Σ by a single homeomorphism.

2. Fine homotopy equivalence

Let $f, g: X \rightarrow Y$ be maps and \mathcal{U} an open cover of Y . It is said that f is \mathcal{U} -homotopic to g and denoted by $f \simeq_{\mathcal{U}} g$ if there is a homotopy $h: X \times I \rightarrow Y$ such that $h_0 = f, h_1 = g$ and each $h(\{x\} \times I)$ is contained in some $U \in \mathcal{U}$. Such a homotopy h is called a \mathcal{U} -homotopy. In case Y is a metric space and $\text{mesh } \mathcal{U} < \varepsilon$, we say that f is ε -homotopic to g and write $f \simeq_\varepsilon g$. In this case, we call h an ε -homotopy. Observe that a homotopy h is an ε -homotopy if and only if $\text{diam } h(\{x\} \times I) < \varepsilon$ for each $x \in X$. Let $X_0 \subset X$. We denote $f \simeq g \text{ rel } X_0$ when there is a homotopy $h: X \times I \rightarrow Y$ such that $h_0 = f, h_1 = g$ and $h_t|X_0 = f|X_0$ for all $t \in I$.

A map $f: X \rightarrow Y$ is called a *fine homotopy equivalence* if for any open cover \mathcal{U} of Y there is a map $g: Y \rightarrow X$ such that

$$f \circ g \simeq_{\mathcal{U}} \text{id}_Y \quad \text{and} \quad g \circ f \simeq_{f^{-1}(\mathcal{U})} \text{id}_X.$$

Such a map g is called a \mathcal{U} -*homotopy inverse* of f .

Let $(X_n)_{n \in \mathbb{N}}$ be a tower of subsets of a metric space $X = (X, d)$. We say that $(X_n)_{n \in \mathbb{N}}$ has the *mapping absorption property for compacta* in X if for any compactum $A \subset X$, $\varepsilon > 0$ and $n \in \mathbb{N}$, there is a map $f: A \rightarrow X_m$ for some $m \geq n$ such that $f|_{A \cap X_n} = \text{id}$ and $d(f, \text{id}) < \varepsilon$. The proof of the following lemma is similar to [Sa3, Theorem 1].

LEMMA 1. *Let $(X_n)_{n \in \mathbb{N}}$ be a tower of compacta in an ANR⁽²⁾ X with the mapping absorption property for compacta in X . Then $\varphi = \text{id}_X: \text{dir lim } X_n \rightarrow X$ is a fine homotopy equivalence.*

Sketch of the proof. For any open cover \mathcal{U} of X , let \mathcal{V} be an open 2-nd star-refinement of \mathcal{U} . Then X is \mathcal{V} -homotopy dominated by a simplicial complex K , i.e., there are maps $f: X \rightarrow |K|_w$ and $g: |K|_w \rightarrow X$ such that $g \circ f \simeq_{\mathcal{V}} \text{id}$. Using the mapping absorption property, we have a map $g': |K|_w \rightarrow X$ which is \mathcal{V} -homotopic to g and maps each simplex of K into some X_n , whence $g': |K|_w \rightarrow \text{dir lim } X_n$ is a map. Thus we have a map $\psi = g' \circ f: X \rightarrow \text{dir lim } X_n$ such that $\varphi \circ \psi \simeq_{st \mathcal{V}} \text{id}_X$. Let $h: X \times I \rightarrow X$ be a $(st \mathcal{V})$ -homotopy such that $h_0 = \text{id}_X$ and $h_1 = \varphi \circ \psi$. Again using the mapping absorption property, we have maps $h'_n: X_n \times I \rightarrow X$, $n \in \mathbb{N}$, such that $h'_n|_{X_{n-1} \times I} = h'_{n-1}$, $h'_n|_{X_n \times \{0, 1\}} = h|_{X_n \times \{0, 1\}}$, $h'_n \simeq_{\mathcal{V}} h|_{X_n \times I}$ and h'_n maps $X_n \times I$ into some X_m . These induce a \mathcal{U} -homotopy $h': (\text{dir lim } X_n) \times I \rightarrow \text{dir lim } X_n$ such that $h'_0 = \text{id}$ and $h'_1 = \psi \circ \varphi$. \square

As a consequence of Lemma 1, we prove the following:

LEMMA 2. *Let $(X_n)_{n \in \mathbb{N}}$ be a tower of compact ANR's in an ANR X with $X = \bigcup_{n \in \mathbb{N}} X_n$. Then $\text{id}: \text{dir lim } X_n \rightarrow X$ is a fine homotopy equivalence.*

Proof. By Lemma 1, it suffices to show that $(X_n)_{n \in \mathbb{N}}$ has the mapping absorption property for compacta in X . To this end, let $A \subset X$ be a compactum, $\varepsilon > 0$ and $n \in \mathbb{N}$. Put $\varepsilon_{n-1} = \varepsilon$. Since each X_i is a compact ANR, we can inductively choose $\varepsilon_i > 0$, $i \geq n$, so that for any two maps $f, g: Z \rightarrow X_i$ defined on an arbitrary space Z if $d(f, g) < \varepsilon_i$ and $f|_{Z_0} = g|_{Z_0}$ for some $Z_0 \subset Z$ then $f \simeq_{\varepsilon_i/3} g$ rel. Z_0 [Du, Corollary 2.2, Theorem 2.5]. Let N_i be a closed neighborhood of X_i in X and $r_i: N_i \rightarrow X_i$ a retraction of N_i onto X_i such that $d(r_i, \text{id}) < \varepsilon_i/3$. From compactness, $A \subset \bigcup_{i=n}^m N_i$ for some $m \geq n$. For each $i = n, \dots, m$, let $A_i = A \cap N_i$ and $\tilde{A}_i = \bigcup_{j=1}^m A_j$, and let $A_0 = A \cap X_n$. Then

⁽²⁾ANR = absolute neighborhood retract for metrizable spaces.

$A_0 \subset A_i$ and $r_i|_{A_0} = \text{id}$ for each $i = n, \dots, m$. Observe

$$r_m|_{\tilde{A}_m \cap A_{m-1}} \simeq_{\varepsilon_{m-2}/3} r_{m-1}|_{\tilde{A}_m \cap A_{m-1}} \text{ rel } A_0.$$

By the homotopy extension theorem [vM, Theorem 5.1.3], $r_m|_{A_m \cap A_{m-1}}$ extends to a map $r'_{m-1}: A_{m-1} \rightarrow X_{m-1}$ such that

$$r'_{m-1} \simeq_{\varepsilon_{m-2}/3} r_{m-1} \text{ rel } A_0.$$

We extend r_m to a map $\tilde{r}_{m-1}: \tilde{A}_{m-1} \rightarrow X_m$ by $\tilde{r}_{m-1}|_{A_{m-1}} = r'_{m-1}$. Then $d(\tilde{r}_{m-1}, \text{id}) < 2\varepsilon_{m-2}/3$ and $\tilde{r}_{m-1}|_{A_0} = \text{id}$. Observe

$$\tilde{r}_{m-1}|_{\tilde{A}_{m-1} \cap A_{m-2}} \simeq_{\varepsilon_{m-3}/3} r_{m-2}|_{\tilde{A}_{m-1} \cap A_{m-2}} \text{ rel } A_0.$$

Similarly as above, \tilde{r}_{m-1} extends to a map $\tilde{r}_{m-2}: \tilde{A}_{m-2} \rightarrow X_m$ such that $d(\tilde{r}_{m-2}, \text{id}) < 2\varepsilon_{m-3}/3$ and $\tilde{r}_{m-2}|_{A_0} = \text{id}$. By induction, we obtain a map $\tilde{r}_n: A = \tilde{A}_n \rightarrow X_m$ such that $d(\tilde{r}_n, \text{id}) < 2\varepsilon_{n-1}/3 < \varepsilon$ and $\tilde{r}_n|_{A \cap X_n} = \tilde{r}_n|_{A_0} = \text{id}$. □

Under the same assumption (*) as corollaries in the Introduction we have the following:

COROLLARY 6. *The identity $\text{id}: (M, \tau_2) \rightarrow (M, \tau_1)$ is a fine homotopy equivalence.*

Proof. Since (M, τ_2) is an \mathbb{R}^∞ -manifold (or a Q^∞ -manifold), $(M, \tau_2) = \text{dir lim } X_n$ for some tower $(X_n)_{n \in \mathbb{N}}$ of compact ANR's [Sa2, Proposition 1.8] (or the proof of [He2, Theorem 4]). Since (M, τ_1) is an ANR, id is a fine homotopy equivalence by Lemma 2. □

A closed set A in a space X is said to be a *strong Z-set* if the identity map of X is arbitrarily close to maps $f: X \rightarrow X$ such that $A \cap \text{cl } f(X) = \emptyset$. It is well-known that each compact set in a σ -manifold (or a Σ -manifold) is a strong Z-set. Combining Corollary 6 with [Sa5, Proposition 1 (or 1')], we have the following:

PROPOSITION. *Let M be an \mathbb{R}^∞ -manifold (or a Q^∞ -manifold) and d a continuous metric on M . Then (M, d) is a σ -manifold (or a Σ -manifold) if and only if $\text{id}: M \rightarrow (M, d)$ is a fine homotopy equivalence and each compact set in (M, d) is a strong Z-set.*

3. The proof of main theorem

We prove the main theorem by the method employed in [Sa2]. To this end, we modify Corollary 6 as follows:

LEMMA 3. Under the assumption (*), let $A \subset B \subset M$ such that A is τ_2 -compact, B is τ_1 -compact and finite dimensional (there is no dimensional assumption in the (Σ, Q^∞) -case). Then for any τ_1 -open cover \mathcal{U} of M , $\text{id}: (M, \tau_2) \rightarrow (M, \tau_1)$ has a \mathcal{U} -homotopy inverse map $h: (M, \tau_1) \rightarrow (M, \tau_2)$ such that $h|_A = \text{id}$ and $h|_B$ is an embedding.

Proof. For simplicity, we write $X = (M, \tau_1)$ and $X_\infty = (M, \tau_2)$. By Corollary 6, $\varphi = \text{id}: X_\infty \rightarrow X$ is a fine homotopy equivalence. Let $f: X \rightarrow X_\infty$ be a \mathcal{V} -homotopy inverse of φ , where \mathcal{V} is an open star-refinement of \mathcal{U} . Then

$$f|_A = f \circ \varphi|_A \simeq_{\varphi^{-1}(\mathcal{V})} \text{id}.$$

Since X_∞ is an absolute neighborhood extensor for metrizable spaces, we can apply the homotopy extension theorem to obtain a map $g: X \rightarrow X_\infty$ such that $g|_A = \text{id}$ and $g \simeq_{\varphi^{-1}(\mathcal{V})} f$. By [Sa2, Lemma 1.5], we have an embedding $j: B \rightarrow X_\infty$ such that $j|_A = g|_A = \text{id}$ and $j \simeq_{\varphi^{-1}(\mathcal{V})} g|_B$. Again using the homotopy extension theorem, we extend j to a map $h: X \rightarrow X_\infty$ such that $h \simeq_{\varphi^{-1}(\mathcal{V})} g$. Then $h|_A = j|_A = \text{id}$, $h|_B = j$ is an embedding and

$$\varphi \circ h \simeq_{\mathcal{V}} \varphi \circ g \simeq_{\mathcal{V}} \varphi \circ f \simeq_{\mathcal{V}} \text{id}_X$$

and

$$h \circ \varphi \simeq_{\varphi^{-1}(\mathcal{V})} g \circ \varphi \simeq_{\varphi^{-1}(\mathcal{V})} f \circ \varphi \simeq_{\varphi^{-1}(\mathcal{V})} \text{id}_{X_\infty}.$$

In other words, h is a \mathcal{U} -homotopy inverse of φ . □

Proof of main theorem. For simplicity, we write $X = (M_1, \tau_1^1)$, $X_\infty = (M_1, \tau_2^1)$, $Y = (M_2, \tau_1^2)$, $Y_\infty = (M_2, \tau_2^2)$, $\varphi = \text{id}: X_\infty \rightarrow X$ and $\psi = \text{id}: Y_\infty \rightarrow Y$ and we denote both admissible metrics for X and Y by the same d . We can write $X_\infty = \text{dir lim } X_n$ and $Y_\infty = \text{dir lim } Y_n$, where each X_n and Y_n are compact and finite dimensional (there is no dimensional restriction in the (Q^∞, Σ) -case). Note that X_n and Y_n are compact in both topologies. We may assume that X_1 is a single point. Recall that any compact set in a σ -manifold (or a Σ -manifold) is a Z -set. Since X and Y are σ -manifolds (or Σ -manifolds) having the same homotopy type, there is a homeomorphism $f: X \rightarrow Y$ [Ch, Theorem 2.2].

Put $m(1) = 1$ and $f_1 = f$. Since $X_{m(1)}$ is a single point, $f_1(X_{m(1)}) \subset Y_{n(1)}$ for some $n(1) \in \mathbb{N}$. Let \mathcal{U}_1 be an open cover of M such that $\text{mesh } f_1(\mathcal{U}_1) < 2^{-2}$. By Lemma 3, φ has a \mathcal{U}_1 -homotopy inverse j_1 such that $j_1|_{X_{m(1)}} = \text{id}$ and $j_1|_{f_1^{-1}(Y_{n(1)})}$ is an embedding. Then $j_1 \circ f_1^{-1}(Y_{n(1)}) \subset X_{m(2)}$ for some $m(2) > m(1)$. Since $\varphi \circ j_1 \circ f_1^{-1}|_{Y_{n(1)}}$ and $f_1^{-1}|_{Y_{n(1)}}$ are \mathcal{U}_1 -homotopic Z -embeddings, we can apply the homeomorphism extension theorem [Ch, Theorem 2.25]⁽³⁾ to obtain

⁽³⁾The covering estimation of [Ch] was improved (cf. [Sa1]).

a homeomorphism $g_1: Y \rightarrow X$ such that $g_1|_{Y_{n(1)}} = \varphi \circ j_1 \circ f_1^{-1}|_{Y_{n(1)}}$ and $g_1 \simeq_{\mathcal{U}_1} f_1^{-1}$. Then $g_1(Y_{n(1)}) \subset X_{m(2)}$ and $g_1 \circ f_1|_{X_{m(1)}} = \varphi \circ j_1|_{X_{m(1)}} = \text{id}$. Since $g_1 \circ f_1$ is \mathcal{U}_1 -near to id_X , $d(f_1, g_1^{-1}) = d(f_1 \circ g_1, \text{id}_Y) < 2^{-2}$.

Let \mathcal{V}_2 be an open cover of Y such that $st \mathcal{V}_2 < \mathcal{V}_1$, $\text{mesh } \mathcal{V}_2 < 2^{-2}$ and $\text{mesh } g_1(\mathcal{V}_2) < 2^{-3}$. By Lemma 3, $\psi: Y_\infty \rightarrow Y$ has a \mathcal{V}_2 -homotopy inverse k_2 such that $k_2|_{Y_{n(1)}} = \text{id}$ and $k_2|_{g_1^{-1}(X_{m(2)})}$ is an embedding. Then $k_2 \circ g_1^{-1}(X_{m(2)}) \subset Y_{n(2)}$ for some $n(2) > n(1)$. Since $\psi \circ k_2 \circ g_1^{-1}|_{X_{m(2)}}$ and $g_1^{-1}|_{X_{m(2)}}$ are \mathcal{V}_2 -homotopic Z -embeddings, we can obtain a homeomorphism $f_2: X \rightarrow Y$ such that $f_2|_{X_{m(2)}} = \psi \circ k_2 \circ g_1^{-1}|_{X_{m(2)}}$ and $f_2 \simeq_{\mathcal{V}_2} g_1^{-1}$, so $d(f_2, g_1^{-1}) < 2^{-2}$. Then $f_2(X_{m(2)}) \subset Y_{n(2)}$, $f_2 \circ g_1|_{Y_{n(1)}} = \psi \circ k_1|_{X_{m(1)}} = \text{id}$ and $d(g_1, f_2^{-1}) = d(g_1 \circ f_2, \text{id}) < 2^{-3}$. Moreover

$$d(f_1, f_2) \leq d(f_1, g_1^{-1}) + d(f_2, g_1^{-1}) < 2^{-1}.$$

Let \mathcal{U}_2 be an open cover of X such that $\text{mesh } \mathcal{U}_2, \text{mesh } f_2(\mathcal{U}_2) < 2^{-3}$. By Lemma 3, $\varphi: X_\infty \rightarrow X$ has a \mathcal{U}_2 -homotopy inverse j_2 such that $j_2|_{X_{m(2)}} = \text{id}$ and $j_2|_{f_2^{-1}(Y_{n(2)})}$ is an embedding. Then $j_2 \circ f_2^{-1}(Y_{n(2)}) \subset X_{m(3)}$ for some $m(3) > m(2)$. Since $\varphi \circ j_2 \circ f_2^{-1}|_{Y_{n(2)}}$ and $f_2^{-1}|_{Y_{n(2)}}$ are \mathcal{U}_2 -homotopic Z -embeddings, we can obtain a homeomorphism $g_2: Y \rightarrow X$ such that $g_2|_{Y_{n(2)}} = \varphi \circ j_2 \circ f_2^{-1}|_{Y_{n(2)}}$ and $g_2 \simeq_{\mathcal{U}_2} f_2^{-1}$. Then $g_2(Y_{n(2)}) \subset X_{m(3)}$, $g_2 \circ f_2|_{X_{m(2)}} = \varphi \circ j_2|_{X_{m(2)}} = \text{id}$ and $d(f_2, g_2^{-1}) = d(f_2 \circ g_2, \text{id}_Y) < 2^{-3}$. Moreover

$$d(g_1, g_2) \leq d(g_1, f_2^{-1}) + d(g_2, f_2^{-1}) < 2^{-2}.$$

Thus by induction, we can obtain homeomorphisms $f_i: X \rightarrow Y$, $g_i: Y \rightarrow X$, $i \in \mathbb{N}$, and sequences $m(1) < m(2) < \dots, n(1) < n(2) < \dots$ such that

$$\begin{aligned} d(f_i, f_{i+1}) &< 2^{-i}, & g(g_i, g_{i+1}) &< 2^{-i}, \\ f_{i+1}|_{X_{m(i)}} &= f_i|_{X_{m(i)}}, & f_i(X_{m(i)}) &\subset Y_{n(i)}, \\ g_{i+1}|_{Y_{n(i)}} &= g_i|_{Y_{n(i)}}, & g_i(Y_{n(i)}) &\subset X_{m(i+1)}, \\ g_i \circ f_i|_{X_{m(i)}} &= \text{id} & \text{and } f_{i+1} \circ g_i|_{Y_{n(i)}} &= \text{id}. \end{aligned}$$

Define maps $f_\infty: X_\infty \rightarrow Y_\infty$ and $g_\infty: Y_\infty \rightarrow X_\infty$ by $f_\infty|_{X_{m(i)}} = f_i|_{X_{m(i)}}$ and $g_\infty|_{Y_{n(i)}} = g_i|_{Y_{n(i)}}$. Then $f_\infty \circ g_\infty = \text{id}$ and $g_\infty \circ f_\infty = \text{id}$. Hence, $f_\infty: X_\infty \rightarrow Y_\infty$ is a homeomorphism. Since

$$d(f_n, f_\infty) \leq \sum_{i \geq n} d(f_i, f_{i+1}) < \sum_{i \geq n} 2^{-i} = 2^{-n+1},$$

f_n is uniformly convergent to f_∞ . Hence, the same map $f_\infty: X \rightarrow Y$ is continuous in the metric topology. Similarly, $g_\infty: Y \rightarrow X$ is also continuous. In other words, $f_\infty: X \rightarrow Y$ is also a homeomorphism. □

REMARKS. In the proof, f_∞ can be taken arbitrarily close to f . In fact, for any open cover \mathcal{U} of Y , Y has an admissible metric d such that

$$\{\{y \in Y \mid d(x, y) < 1\} \mid x \in Y\} < \mathcal{U}.$$

Using this metric, f_∞ is \mathcal{U} -near to f since $d_Y(f, f_\infty) < 2^0 = 1$. As a consequence, the homeomorphism of the stability theorem (Corollary 4) can be taken arbitrarily close to the projection and the one of the negligibility theorem (Corollary 5) can be taken arbitrarily close to the inclusion with respect to τ_1 .

Furthermore, if $A \subset M_1$ is τ_1^1 -compact and $f(A) \subset M_2$ is τ_2^2 -compact, then f_∞ can be taken to satisfy $f_\infty|_A = f|_A$. In fact, we can assume $X_1 = A$ in the above argument, whence $f_\infty|_A = f|_A$. By the homeomorphism extension theorem for σ -manifolds (or Σ -manifolds), we have the following corollary under the same assumption (*) stated in the Introduction:

COROLLARY 7 (Homeomorphism extension theorem). *Let $f: A \rightarrow B$ be a homeomorphism between τ_2 -compact sets in M . If f is homotopic to id in (M, τ_1) , then f extends to $\tilde{f}: M \rightarrow M$ which is a homeomorphism in both topologies. If f is \mathcal{U} -homotopic to id for a τ_1 -open cover \mathcal{U} of M , \tilde{f} is \mathcal{U} -near to id .*

References

- [Du] Dugundji, J.: Locally equiconnected spaces and absolute neighborhood retracts, *Fund. Math* 57 (1965), 187–193.
- [Ch] Chapman, T.A.: Dense sigma-compact subsets of infinite-dimensional manifolds, *Trans. Amer. Math. Soc.* 154 (1971), 399–426.
- [Ge] Geoghegan, R. (ed.): Open problems in infinite-dimensional topology, *Topology Proc.* 4 (1979), 287–330.
- [He1] Heisey, R.E.: Partitions of unity and a closed embedding theorem for (C^b, b^*) -manifolds, *Trans. Amer. Math. Soc.* 206 (1975), 281–294.
- [He2] Heisey, R.E.: Manifolds modelled on the direct limit of Hilbert cubes, in J.C. Cantrell (ed.), *Geometric Topology*, Academic Press, New York (1979), 609–619.
- [Ke] Kelly, J.C.: Bitopological spaces, *Proc. London Math. Soc.* (3) 13 (1963), 71–89.
- [Pa] Palais, R.S.: Banach manifolds of fiber bundle sections, *Actes Congrès Intern. Math.*, Nice, 1970, vol. 2, Gauthier-Villars, Paris (1971), 243–249.
- [vM] van Mill, J.: Infinite-Dimensional Topology, Prerequisites and Introduction, *North-Holland Math. Library* 43, Elsevier Sci. Publ., Amsterdam (1989).
- [Sa1] Sakai, K.: Embeddings of infinite-dimensional manifold pairs and remarks stability and deficiency, *J. Math. Soc. Japan* 29 (1977), 262–280.
- [Sa2] Sakai, K.: On \mathbb{R}^∞ -manifolds and Q^∞ -manifolds, *Topology Appl.* 18 (1984), 69–79.
- [Sa3] Sakai, K.: Fine homotopy equivalences of metric simplicial complexes, *Bull. Polish Acad. Sci. Math.* 34 (1986), 89–97.
- [Sa4] Sakai, K.: Combinatorial infinite-dimensional manifolds and \mathbb{R}^∞ -manifolds, *Topology Appl.* 26 (1987), 287–300.
- [Sa5] Sakai, K.: On topologies of triangulated infinite-dimensional manifolds, *J. Math. Soc. Japan* 39 (1987), 287–300.
- [Sa6] Sakai, K.: Simplicial complexes triangulating infinite-dimensional manifolds, *Topology Appl.* 29 (1988), 167–183.

- [Sa7] Sakai, K.: The space of Lipschitz maps from a compactum to an absolute neighborhood LIP extensor, *Fund. Math.*, in press.
- [Sa8] Sakai, K.: A Q^∞ -manifold topology of the space of Lipschitz maps, preprint.
- [We] West, J.E. (ed.): Open problems in infinite-dimensional topology, in J. van Mill and G.M. Reed (eds), *Open Problems in Topology*, Elsevier Sci. Publ., Amsterdam (1990), 523–597.