

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

GRZEGORZ JARZEMBSKI

A new proof of Reiterman's theorem

Cahiers de topologie et géométrie différentielle catégoriques, tome 35, n° 3 (1994), p. 239-247

http://www.numdam.org/item?id=CTGDC_1994__35_3_239_0

© Andrée C. Ehresmann et les auteurs, 1994, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A NEW PROOF OF REITERMAN'S THEOREM

by Grzegorz JARZEMBSKI

In [3] J.Reiterman introduced a concept of an implicit operation in order to get a very elegant finite analog of the Birkhoff variety theorem. In this note we analyse implicit operations from a categorical point of view. As a result we obtain a simple, purely categorical proof of Reiterman theorem.

1 Basic concepts

For all unexplained notions of category theory we refer the reader to [2].

We shall identify classes of objects of a category considered with full subcategories they generate. For every natural number n , by $\tilde{n}$ we denote the set $\{1, 2, \dots, n\}$.

Let $(\mathcal{A}, U : \mathcal{A} \longrightarrow \text{Set})$ be an arbitrary concrete category over sets. For any natural number n , by an n -ary implicit operation in $\mathcal{A}$ we mean any natural transformation $\phi : U^n \longrightarrow U$ [3].

Roughly speaking, an implicit operation is a family of functions $(\phi_A : U^n A \longrightarrow UA : A \in \text{Ob}\mathcal{A})$ "compatible" with all $\mathcal{A}$ -morphisms.

By $IO(\mathcal{A})_n$ we shall denote the class of all n -ary implicit operations in $\mathcal{A}$.

Definition 1 Any pair (ϕ, ψ) of n -ary implicit operations in $\mathcal{A}$ is called here an "equation"; (notation $(\phi = \psi)$).

We say that an equation $(\phi = \psi)$ is satisfied in $B \in \text{Ob}\mathcal{A}$ iff $\phi_B = \psi_B$.

For a given set of equations E , by $\text{Mod}E$ we denote the full subcategory of $\mathcal{A}$ consisting of $\mathcal{A}$ -objects satisfying all equations in E .

We call a full subcategory $\mathcal{D} \subset \mathcal{A}$ equationally definable iff $\mathcal{D} = \text{Mod}E$ for some set of equations E .

The following observation needs only a routine verification:

Lemma 1 Every equationally definable full subcategory $\mathcal{D} \subset \mathcal{A}$ satisfies the following:

1. $\mathcal{D}$ is closed under formation of all existing concrete limits,
2. whenever $A \in \mathcal{D}$, $h : A \longrightarrow B$ and Uh is a surjection, then $B \in \mathcal{D}$ ($\mathcal{D}$ is closed under formation of homomorphic images),
3. whenever $A \in \mathcal{D}$, $m : B \longrightarrow A$ and Um is a monomorphism, then $B \in \mathcal{D}$ ($\mathcal{D}$ is closed under formation of subobjects).

J. Reiterman has proved the converse of Lemma 1 for equationally definable classes of finite algebras of an arbitrary finite type Ω ([3]). His result has been next generalized by Banaschewski ([1]) for arbitrary finitary types. In this paper we present a new proof of Reiterman theorem based on a categorical analysis of the concept of an implicit operation.

2 Two restrictions

Definition 2 A concrete category $(\mathcal{A}, U)$ is said to be small-based provided for every natural number n $IO(\mathcal{A})_n$ is a set.

Let $(\tilde{n} \downarrow U)$ denote the category of all U -morphisms with a domain $\tilde{n}$. Let

$$U_n : (\tilde{n} \downarrow U) \longrightarrow \text{Set}$$

be the "forgetful functor", i.e. $U_n(h : \tilde{n} \longrightarrow UA, A) = UA$, for every (h, A) in $(\tilde{n} \downarrow U)$.

Lemma 2 A concrete category $(\mathcal{A}, U)$ is small-based if and only if for every natural number n there exists a limit of the functor U_n and then

$$IO(\mathcal{A})_n \cong \lim U_n$$

Proof. A limit of U_n is formed by the set $IO(\mathcal{A})_n$ together with limit projections $\pi_{(h,A)} : IO(\mathcal{A})_n \longrightarrow U_n(h, A) = UA$ such that

$$\pi_{(h,A)}(\phi) = \phi_A(h)$$

for every $(h, A) \in (\tilde{n} \downarrow U)$ and $\phi \in IO(\mathcal{A})_n$.

Definition 3 We say that a concrete category $(\mathcal{A}, U)$ has enough implicit operations iff for every $A, B \in \mathcal{A}$ and $h : UA \longrightarrow UB$, $h = Ug$ for some $\mathcal{A}$ -morphism $g : A \longrightarrow B$ provided h preserves all implicit operations ($h \cdot \phi_A = \phi_B \cdot h^n$ for every $n \in N$ and $\phi \in IO(\mathcal{A})_n$).

Lemma 3 Assume that a $(\mathcal{A}, U)$ is a concrete, small-based with enough implicit operations. Then there exists a (finitary) monad T over Set such that the following conditions are satisfied:

1. $(\mathcal{A}, U)$ is concretely isomorphic to a full subcategory of the Eilenberg-Moore category Set^T ,
2. for every $n \in N$, $IO(\mathcal{A})_n \cong \lim U_n$ is a carrier of a finitely generated monadic T -algebra such that every limit projection is a T -morphism.

Proof. Since $(\mathcal{A}, U)$ is small-based, the family of all implicit operations form a finitary type: $IO = (IO(\mathcal{A})_n : n \in N)$.

Let T be a monad over Set such that its Eilenberg-Moore category Set^T is the category of all IO -algebras. Since the concrete category considered has enough implicit operations, it is concretely isomorphic to a full subcategory of T -algebras.

In the considered case the functor U_n factorizes through the forgetful functor $U^T : Set^T \longrightarrow Set$ in an obvious way. U^T creates limits, hence, by Lemma 2, every set $IO(\mathcal{A})_n$ carries a structure of a monadic T -algebra such that every limit-projection $\pi_{(h, A)} : IO(\mathcal{A})_n \longrightarrow A$ is a homomorphism.

Obviously, every $IO(\mathcal{A})_n$ is then a finitely generated algebra of the type IO - it is generated by the set of natural projections $\{\pi_i^n : U^n \longrightarrow U : i = 1, 2, \dots, n\}$.

It must be stressed however that a monad T satisfying the conditions of Lemma 3 is not uniquely determined. It need not to be finitary too. An embedding $\mathcal{A} \longrightarrow Set^T$ will be used only in order to support an investigation on equationally definable subcategories of $\mathcal{A}$. As we will show later, in particular cases a "good" choice of a monad T is crucial.

3 Characterization of equationally definable subcategories

Assume that a concrete category $(\mathcal{A}, U)$ is a small-based category with enough implicit operations. Let $T = (T, \mu, \eta)$ be an arbitrary but fixed monad over Set such that the conditions stated in Lemma 3 are satisfied. We shall assume that $\mathcal{A}$ is a full subcategory of the Eilenberg-Moore category Set^T . The T -algebra of n -ary implicit operations will be denoted by $IO(\mathcal{A})_n$.

Let $\pi^n : \tilde{n} \longrightarrow IO(\mathcal{A})_n$ be a function such that for every $i \in \tilde{n}$, $\pi^n(i) = \pi_i^n : U^n \longrightarrow U =$ the i -th natural projection.

We assume also that the extension of π^n to the T -morphism $(T\tilde{n}, \mu_{\tilde{n}}) \longrightarrow IO(\mathcal{A})_n$ is surjective for every natural number n .

Definition 4 We call a T -algebra D a $T_{\mathcal{A}}$ -algebra iff for every $n \in N$ and a function $f : \tilde{n} \longrightarrow U^T D$ there exists a (unique) T -morphism $f^* : IO(\mathcal{A})_n \longrightarrow D$ such that

$$f^* \cdot \pi^n = f$$

We shall call f^* an extension of f .

Clearly, $\mathcal{A}$ is a subclass of $T_{\mathcal{A}}$ -algebras: for every $B \in Ob\mathcal{A}$ and $h : \tilde{n} \longrightarrow UB$ its extension is the limit-projection $\pi_{(h, \mathcal{A})} : IO(\mathcal{A})_n \longrightarrow B$ (Lemma 2).

Notice also that every $IO(\mathcal{A})_n$ is a $T_{\mathcal{A}}$ -algebra. This is a consequence of the fact that the class of $T_{\mathcal{A}}$ -algebras is closed under formation of limits. It is easily checked that the class of $T_{\mathcal{A}}$ -algebras is closed within Set^T under formation of homomorphic images and subalgebras, i.e. the class of $T_{\mathcal{A}}$ -algebras is the "Birkhoff subcategory generated by $\mathcal{A}$ " within Set^T .

Definition 5 For each $\phi \in IO(\mathcal{A})_n$ and a $T_{\mathcal{A}}$ -algebra D we define an n -ary operation in D , $\phi_D : (U^T D)^n \longrightarrow U^T D$ as follows:

$$\phi_D(h) = h^*(\phi)$$

for every $h : \tilde{n} \longrightarrow U^T D$.

Thus h^* gives "values of all n -ary implicit operations in D at the valuation h ".

It is easily checked that T -morphisms between $T_{\mathcal{A}}$ -algebras preserve these operations.

Having all implicit operations extended to all $T_{\mathcal{A}}$ -algebras we extend also a notion of satisfaction for equations of implicit operations:

Definition 6 *Let $\phi, \psi \in IO(\mathcal{A})_n$. A $T_{\mathcal{A}}$ -algebra A satisfies the equation $(\phi = \psi)$ iff for every $h : \tilde{n} \longrightarrow U^T A$,*

$$h^*(\phi) = h^*(\psi)$$

If $A \in Ob\mathcal{A}$ and $h : \tilde{n} \longrightarrow UA$ then $h^*(\phi) = \pi_{(h,A)}(\phi) = \phi_A(h)$, i.e. the newly defined concept of satisfaction and that of Definition 1 coincide for $\mathcal{A}$ -objects.

For any class $\mathcal{B}$ of $T_{\mathcal{A}}$ -algebras we write $H(\mathcal{B})$, $S(\mathcal{B})$, $P(\mathcal{B})$ in order to denote the class of all homomorphic images, subalgebras and products of the algebras from $\mathcal{B}$, resp.

We call a class $\mathcal{B}$ an *HSP-class* iff $\mathcal{B} = HSP(\mathcal{B})$.

Since we deal with finitary implicit operations only, we shall use one more closure operator. In what follows by $C(\mathcal{B})$ we shall denote the class of all $T_{\mathcal{A}}$ -algebras with all finitely generated subalgebras in $\mathcal{B}$ (a monadic T -algebra A is said to be *finitely generated* provided it is isomorphic to a quotient of a free monadic T -algebra $(T\tilde{n}, \mu_{\tilde{n}})$ for some natural number n).

Theorem 4 *For an arbitrary class $\mathcal{B}$ of $T_{\mathcal{A}}$ -algebras the following conditions are equivalent:*

1. $\mathcal{B}$ is an HSP-class and $\mathcal{B} = C(\mathcal{B})$
2. There exists a set E of equations (of implicit operations) such that $\mathcal{B}$ consists of all $T_{\mathcal{A}}$ -algebras satisfying all equations in E .

Proof. 2. $\Rightarrow$ 1. The equations $\mathcal{B} = P(\mathcal{B}) = H(\mathcal{B}) = S(\mathcal{B})$ need only a routine verification.

For every $T_{\mathcal{A}}$ -algebra B and a function $h : \tilde{n} \longrightarrow U^T B$ its extension h^* factorizes as a surjective T -morphism followed by a monomorphism:

$$h^* = m \cdot e : IO(\mathcal{A}_n) \longrightarrow D_h \longrightarrow B$$

Clearly, D_h is a finitely generated $T_{\mathcal{A}}$ -algebra. Hence a $T_{\mathcal{A}}$ -algebra B satisfies an equation $(\phi = \psi)$ iff every finitely generated subalgebra of B satisfies it.

This proves $C(\mathcal{B}) \subset \mathcal{B}$.

1. $\Rightarrow$ 2. Since $\mathcal{B}$ is closed under products and subalgebras, for every natural number n there exists a "reflection map" - a surjective T -morphism $e_n : IO(\mathcal{A})_n \longrightarrow B_n$ such that $B_n \in \mathcal{B}$ and every morphism of T -algebras $h : IO(\mathcal{A})_n \longrightarrow A$ with $A \in \mathcal{B}$ factorizes through e_n . Let

$$E_n = \ker(e_n) = \{(\phi, \psi) : \phi, \psi \in IO(\mathcal{A})_n, e_n(\phi) = e_n(\psi)\}$$

and

$$E = \bigcup (E_n : n \in N)$$

We prove that E is a set of equations we are looking for.

Clearly, each $T_{\mathcal{A}}$ -algebra in $\mathcal{B}$ satisfies all equations in E .

Assume that A is a finitely generated $T_{\mathcal{A}}$ -algebra satisfying all equations in E . A is finitely generated hence there exists a surjective T -morphism $f : IO(\mathcal{A})_n \longrightarrow A$ for some $n \in N$. Since A satisfies all equations in E , we obtain $E_n \subset \ker f$. Hence $f = f^\circ \cdot e_n$ for some T -morphism f° . f is surjective, hence f° is a surjective morphism, too.

Thus $A \in H(\mathcal{B}) = \mathcal{B}$.

An arbitrary $T_{\mathcal{A}}$ -algebra B satisfies all equations from E , iff all its finitely generated subalgebras satisfy all these equations, i.e. if all those subalgebras are in $\mathcal{B}$. This means $B \in C(\mathcal{B}) = \mathcal{B}$.

The proof is complete.

Corollary 5 . *For any class $\mathcal{B}$ of $T_{\mathcal{A}}$ -algebras, the class $CHSP(\mathcal{B})$ is the smallest equationally definable class of $T_{\mathcal{A}}$ -algebras containing $\mathcal{B}$.*

Proof. Observe that $HC(\mathcal{B}) \subset CH(\mathcal{B})$ and $SPC(\mathcal{B}) \subset CSP(\mathcal{B})$ for any class of $T_{\mathcal{A}}$ -algebras $\mathcal{D}$. We omit a routine calculation.

The next Corollary summarizes investigation of this section.

Corollary 6 *For any class $\mathcal{D} \subset \mathcal{A}$, the following conditions are equivalent*

1. $\mathcal{D}$ is an equationally definable class,
2. $\mathcal{D} = CHSP(\mathcal{D}) \cap \mathcal{A}$

4 A new proof of Reiterman theorem

Consider now equationally definable classes in the concrete category $Alg_{fin}\Omega$ of all finite algebras of a given finitary type Ω . We are going to give a new proof of Reiterman characterization theorem based on the results of the previous section. Our proof also covers the generalization of Reiterman theorem given by Banaschewski ([1]).

Since $Alg_{fin}\Omega$ is a small category, it is small-based. It follows from Lemma 2 that in this case the set of n -ary implicit operations is represented as a limit of the poset of all finite quotients of a free Ω -algebra generated by the set $\tilde{n} = \{1, 2, \dots, n\}$. From this it easily follows that every implicit operation ϕ is "locally explicit" - i.e., for every finite algebra A there is a term t with $t_A = \phi_A$ ([3]).

Obviously, the concrete category $Alg_{fin}\Omega$ has enough implicit operations.

For the category $Alg_{fin}\Omega$ we may consider three (at least) concrete full embeddings of it into the following Eilenberg-Moore categories over sets:

- the category $Alg\Omega$ of all Ω -algebras,
- the category $AlgIO$, where IO is the type built of all implicit operations in $Alg_{fin}\Omega$,
- the category of all compact Hausdorff IO -algebras and continuous homomorphisms (by endowing each finite algebra with the discrete topology).

In our proof of Reiterman theorem we shall use the third embedding.

Recall that by a *pseudovariety* of finite Ω -algebras we mean each class of finite algebras closed under formation of finite products, subalgebras and homomorphic images.

Theorem 7 (*Reiterman theorem*)

For any class $\mathcal{D} \subset Alg_{fin}\Omega$ the following conditions are equivalent:

1. $\mathcal{D}$ is a pseudovariety ,
2. $\mathcal{D} = ModE$ for some set of equations of implicit operations of finite algebras.

Proof. Each $ModE$ is clearly a pseudovariety.

Conversely, let D be a pseudovariety. To find E with $D = ModE$ consider the monad T over Set whose Eilenberg-Moore category is the category of all compact Hausdorff IO -algebras (and continuous homomorphisms).

Clearly, $Alg_{fin}\Omega$ is a full concrete subcategory of Set^T and for every natural n , the set of n -ary implicit operations in $Alg_{fin}\Omega$ is a carrier of a finitely generated monadic T -algebra (Lemma 3).

By Corollary 6, it is enough to prove that $\mathcal{D}$ contains every finite algebra C lying in $HSP(\mathcal{D}) \subset Set^T$. That is, there is a continuous and surjective homomorphism $e : B \longrightarrow C$, where B is a compact Hausdorff IO -algebra, and B is a closed subalgebra of $\prod_{i \in I} A_i$ for a collection $(A_i : i \in I) \subset \mathcal{D}$.

We may assume that every A_i has the form $A_i = B/\rho_i$ for some congruence ρ_i on B . Then

$$\bigcap (\rho_i : i \in I) = \nabla_B$$

where ∇_B denotes the diagonal in $B \times B$.

Every congruence ρ_i has a finite index and it is a kernel of a continuous projection $pr_i : B \longrightarrow A_i$, hence it is a closed subset of the compact space $B \times B$.

The epimorphism $e : B \longrightarrow C$ is continuous, the diagonal ∇_C is open in the compact space $C \times C$, hence the inverse image $ker(e) = (e \times e)^{-1}(\nabla_C)$ is an open subset of $B \times B$ containig ∇_B . Hence

$$\bigcap (\rho_i : i \in I) = \nabla_B \subset ker(e)$$

Since $B \times B$ is compact, there exists a finite subset $I_1 \subset I$ such that

$$\bigcap (\rho_i : i \in I_1) \subset ker(e)$$

Clearly, the congruence $\bigcap (\rho_i : i \in I_1)$ has a finite index. Thus $B_0 = B / \bigcap (\rho_i : i \in I_1)$ is a finite algebra and C is a homomorphic image of B_0 .

Moreover, B_0 is a subalgebra of the product of the family of finite algebras $\{A_i : i \in I_1\}$.

Hence $C \in \mathcal{D}$.

The proof is complete.

References

- [1] B.Banaschewski, The Birkhoff theorem for varieties of finite algebras, *Algebra Universalis* **17**(1983), 360-368.

- [2] E.Manes, *Algebraic Theories, Graduate Texts in Math. 26*, Springer-Verlag, Berlin 1976 .
- [3] J.Reiterman, The Birkhoff theorem for finite algebras, *Algebra Universalis* **14**, no.1 (1982),1-10.

Institute of Mathematics
N.Copernicus University,
Toru ħ 87-100, ul.Chopina 12
POLAND