

# CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

M.-A. MOENS

E. M. VITALE

## **Groupoids and the Brauer group**

*Cahiers de topologie et géométrie différentielle catégoriques*, tome 41, n° 4 (2000), p. 305-313

[http://www.numdam.org/item?id=CTGDC\\_2000\\_\\_41\\_4\\_305\\_0](http://www.numdam.org/item?id=CTGDC_2000__41_4_305_0)

© Andrée C. Ehresmann et les auteurs, 2000, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## GROUPOIDS AND THE BRAUER GROUP

by *M.-A. MOENS\** and *E.M. VITALE*

**Résumé.** On utilise les bigroupoïdes pour analyser la suite exacte reliant le groupe de Picard et le groupe de Brauer, aussi bien que la description K-théorique des groupes de Picard et de Brauer.

MSC 2000 : 13A20, 13D15, 16D90, 18Dxx, 18F25, 20L05.

### 1 Preliminaries

The existence of an exact sequence between the Picard group and the Brauer group of a commutative unital ring is a well known fact in algebraic K-theory. Similar exact sequences have been obtained starting for example from a ringed space, a Krull domain, etc. [1, 9, 16, 20, 23, 24, 25]. All these constructions are particular cases of the exact sequence between the Picard group and the Brauer group of a symmetric monoidal category [9, 10, 26].

Nevertheless, there are generalizations/enrichements of the Brauer group and of the related exact sequences which do not fit into the general theory developed in [10, 26]. Two interesting (and, for us, motivating) examples are the Brauer-Taylor group and the categorical Brauer group [5, 6, 7, 17, 18, 19, 21, 22]. We look for a more general approach which contains these new examples.

What makes this possible is that it remains a deep analogy between the Brauer group and these generalizations: they can be described as a kind of Picard group associated to a convenient monoidal bicategory [5, 12, 18,

---

\*Aspirant du F.N.R.S.

19]. The aim of our note is to show that this fact suffices to obtain some relevant results: the above mentioned exact sequence and the K-theoretical description of Picard and Brauer groups.

Let us describe now the general situation using the formalism of bicategories and tricategories. For the basic definitions, the reader can consult [3, 8, 11]. The situation to be kept in mind is given in the following commutative diagram.

$$\begin{array}{ccccc}
 \text{Tricategories} & \xrightarrow{cl_2} & \text{Bicategories} & \xrightarrow{cl_1} & \text{Categories} \\
 P_3 \downarrow & & P_2 \downarrow & & P_1 \downarrow \\
 \text{Trigroupoids} & \xrightarrow{cl_2} & \text{Bigroupoids} & \xrightarrow{cl_1} & \text{Groupoids}
 \end{array}$$

where  $P_i$  takes invertibles at each level (for an  $n$ -cell invertible means invertible up to invertible  $n + 1$ -cell);  $cl_1$  is the classifying category of a bicategory as in [3] (take 2-isomorphism classes of 1-cells as arrows) and  $cl_2$  is the analogous construction performed taking 3-isomorphism classes of 2-cells. Given a monoidal category (i.e., a bicategory with a single 0-cell)  $\mathbb{C}$ , its Picard group is the group  $P_1(cl_1(\mathbb{C}))$ . If  $\mathbb{C}$  is symmetric and has stable coequalizers (cf. [8, 26]), we can build up the monoidal bicategory (i.e., tricategory with a single 0-cell)  $\text{Mon}\mathbb{C}$  of unital monoids and unital bimodules. Then the Brauer group of  $\mathbb{C}$  is the group  $P_1(cl_1(cl_2(\text{Mon}\mathbb{C})))$ .

In the final remark to section 2 in [26], it is suggested to take  $\text{Mon}\mathbb{C}$  as primitive notion. The point is that, despite of the previous definitions of Picard and Brauer groups, the majority part of the computations are performed passing through

$$\text{Bicategories} \xrightarrow{P_2} \text{Bigroupoids} \xrightarrow{cl_1} \text{Groupoids}$$

instead of

$$\text{Bicategories} \xrightarrow{cl_1} \text{Categories} \xrightarrow{P_1} \text{Groupoids}.$$

Once this is clearly recognized, it becomes reasonable to take as primitive the bigroupoid  $\mathbb{B} = P_2(cl_2(\text{Mon}\mathbb{C}))$ , called in [27] the Brauer cat-group of  $\mathbb{C}$ . We test this point of view in the next two sections.

## 2 The K-theoretical description

For the definition of the Grothendieck group  $K_0$  and of the Whitehead group  $K_1$  the reader can see [2]. Let  $\mathbb{B} = (\mathbb{B}, \otimes, I, ()^*, \dots)$  be a compact closed groupoid (or symmetric cat-group), i.e. a bigroupoid with a single 0-cell and symmetric (braided is enough) as monoidal category [13, 14, 15, 27].

**Proposition 1** (i)  $K_0(\mathbb{B})$  is isomorphic to  $cl_1(\mathbb{B})$

(ii)  $K_1(\mathbb{B})$  is isomorphic to  $\mathbb{B}(I, I)$  (the group of automorphisms of  $\mathbb{B}$  at  $I$ ).

*Proof* (i) is obvious from the universal property of  $K_0$ .

(ii) Consider the category  $\Omega\mathbb{B}$  whose objects are arrows  $f : A \rightarrow A$  in  $\mathbb{B}$  and whose arrows are commutative diagrams of the form

$$\begin{array}{ccc} A & \xrightarrow{f} & A \\ x \downarrow & & \downarrow x \\ B & \xrightarrow{g} & B. \end{array}$$

$\Omega\mathbb{B}$  is a monoidal category with a composition between objects. To have an isomorphism between  $K_1(\mathbb{B})$  and  $\mathbb{B}(I, I)$ , we need a surjective map  $\gamma$  from the objects of  $\Omega\mathbb{B}$  to  $\mathbb{B}(I, I)$ , constant on connected components, which sends tensor product and composition on the composition of  $\mathbb{B}(I, I)$ , and such that the following condition holds: if  $\gamma(f : A \rightarrow A) = (1_I : I \rightarrow I)$ , then  $f : A \rightarrow A$  is isomorphic to  $1_A : A \rightarrow A$  in  $\Omega\mathbb{B}$ . We define  $\gamma$  in the following way:

$$\gamma(f : A \rightarrow A) = I \xrightarrow{\eta_A} A \otimes A^* \xrightarrow{f \otimes 1_{A^*}} A \otimes A^* \xrightarrow{\eta_A^{-1}} I$$

where  $\eta_A$  is the unit of the duality  $A^* \dashv A$ . The proof that  $\gamma$  verifies the various conditions is a quite straightforward step-by-step transcription of the proof of proposition 2.4 in [26]. For this reason, we will only show that  $\gamma$  sends tensor product on composition. This probably is the simplest one between the five conditions, but it seems to us enough to give the flavour of the proof and to illustrate the role of the compact closed structure of  $\mathbb{B}$ .

Given two objects  $f : A \rightarrow A$  and  $g : B \rightarrow B$  in  $\mathbb{B}$ , we have to show that the following diagram commutes

$$\begin{array}{ccc}
 A \otimes B \otimes (A \otimes B)^* & \xrightarrow{f \otimes g \otimes 1} & A \otimes B \otimes (A \otimes B)^* \\
 \eta_{A \otimes B} \uparrow & & \downarrow \eta_{A \otimes B}^{-1} \\
 I & \xrightarrow{\eta_A \cdot (f \otimes 1) \cdot \eta_A^{-1}} & I \xrightarrow{\eta_B \cdot (g \otimes 1) \cdot \eta_B^{-1}} I
 \end{array}$$

But, as in every compact closed category, there is a natural isomorphism  $u_{AB} : (A \otimes B)^* \rightarrow B^* \otimes A^*$  such that

$$\begin{array}{ccc}
 I & \xrightarrow{\eta_{A \otimes B}} & A \otimes B \otimes (A \otimes B)^* \\
 \eta_A \downarrow & & \downarrow 1 \otimes 1 \otimes u_{AB} \\
 A \otimes A^* \cong A \otimes I \otimes A^* & \xrightarrow{1 \otimes \eta_B \otimes 1} & A \otimes B \otimes B^* \otimes A^*
 \end{array}$$

commutes. Then, the following diagram also commutes

$$\begin{array}{ccc}
 A \otimes B \otimes (A \otimes B)^* & \xrightarrow{f \otimes g \otimes 1} & A \otimes B \otimes (A \otimes B)^* \\
 \eta_{A \otimes B} \uparrow & & \downarrow \eta_{A \otimes B}^{-1} \\
 I & & I \\
 \eta_A \downarrow & & \uparrow \eta_A^{-1} \\
 A \otimes A^* \cong A \otimes I \otimes A^* & & A \otimes I \otimes A^* \cong A \otimes A^* \\
 1 \otimes \eta_B \otimes 1 \downarrow & & \uparrow 1 \otimes \eta_B^{-1} \otimes 1 \\
 A \otimes B \otimes B^* \otimes A^* & \xrightarrow{f \otimes g \otimes 1 \otimes 1} & A \otimes B \otimes B^* \otimes A^*
 \end{array}$$

Now the fact that

$$\begin{array}{ccc}
 I & \xrightarrow{\eta_A \cdot (f \otimes 1) \cdot \eta_A^{-1}} & I \xrightarrow{\eta_B \cdot (g \otimes 1) \cdot \eta_B^{-1}} I \\
 \eta_A \downarrow & & \uparrow \eta_A^{-1} \\
 A \otimes A^* \cong A \otimes I \otimes A^* & & A \otimes I \otimes A^* \cong A \otimes A^* \\
 1 \otimes \eta_B \otimes 1 \downarrow & & \uparrow 1 \otimes \eta_B^{-1} \otimes 1 \\
 A \otimes B \otimes B^* \otimes A^* & \xrightarrow{f \otimes g \otimes 1 \otimes 1} & A \otimes B \otimes B^* \otimes A^*
 \end{array}$$

commutes is a routine diagram argument using the functoriality of  $\otimes$  and the various natural and coherent isomorphisms of a symmetric monoidal category.  $\square$

### 3 The exact sequence

The K-theoretical description given in the previous proposition is the key result used in [2, 10, 9] to obtain a Picard-Brauer exact sequence. Here we sketch a more direct method, which follows the lines of section 2 in [26].

For this, consider two compact closed groupoids  $\mathbb{A}$  and  $\mathbb{B}$  and a monoidal functor  $F : \mathbb{A} \rightarrow \mathbb{B}$ . We can construct the abelian group  $\mathcal{F}$  having as elements classes of triples  $(A_1, b, A_2)$  with  $A_1, A_2$  in  $\mathbb{A}$  and  $b : FA_1 \rightarrow FA_2$  in  $\mathbb{B}$ . Two triples  $(A_1, b, A_2)$  and  $(A'_1, b', A'_2)$  are equivalent if there exist  $a_1 : A_1 \rightarrow A'_1$ ,  $a_2 : A_2 \rightarrow A'_2$  in  $\mathbb{A}$  such that

$$\begin{array}{ccc} FA_1 & \xrightarrow{b} & FA_2 \\ Fa_1 \downarrow & & \downarrow Fa_2 \\ FA'_1 & \xrightarrow{b'} & FA'_2 \end{array}$$

commutes. The operation in  $\mathcal{F}$  is induced by the tensor product in  $\mathbb{B}$ . Now consider the subgroup  $\mathcal{N}$  of  $\mathcal{F}$  spanned by the elements of the form  $[A, 1_{FA}, A]$  for  $A$  in  $\mathbb{A}$ , and take the quotient group  $\pi : \mathcal{F} \rightarrow \mathcal{F}/\mathcal{N} = \overline{\mathcal{F}}$ . There are also two morphisms:

- $F_1 : \mathbb{B}(I, I) \rightarrow \mathcal{F}, \quad (b : I \rightarrow I) \mapsto [I, FI \cong I \xrightarrow{b} I \cong FI, I]$
- $F_2 : \mathcal{F} \rightarrow cl_1(\mathbb{A}), \quad [A_1, b, A_2] \mapsto [A_1 \otimes A_2^*].$

Moreover, since  $\mathcal{N}$  is contained in the kernel of  $F_2$ , the morphism  $F_2$  factors through  $\pi$ ; let  $F'_2 : \overline{\mathcal{F}} \rightarrow cl_1(\mathbb{A})$  be the factorisation.

**Proposition 2** *With the previous notations, the sequence of abelian groups and morphisms*

$$\mathbb{A}(I, I) \longrightarrow \mathbb{B}(I, I) \xrightarrow{F_1 \cdot \pi} \overline{\mathcal{F}} \xrightarrow{F'_2} cl_1(\mathbb{A}) \longrightarrow cl_1(\mathbb{B})$$

*is exact.*

*Proof* A direct proof can be done following the proof of Proposition 2.2 in [26]. Alternatively, one can work in the following way: first observe that, even if  $\mathbb{B}$  in general does not have coequalizers, one can construct  $\text{Mon}\mathbb{B}$  because, up to isomorphism, the unique monoid in  $\mathbb{B}$  is  $I$  and then the needed coequalizers

are trivial. Now, following [27], we obtain a 2-exact sequence of symmetric cat-groups

$$\mathbb{A} \rightarrow \mathbb{B} \rightarrow \mathbb{F} \rightarrow cl_2(\text{Mon}\mathbb{A}) \rightarrow cl_2(\text{Mon}\mathbb{B}).$$

Taking, for each cat-group, the abelian group of automorphisms of the unit object, we have the requested exact sequence of abelian groups. (As far as  $cl_2(\text{Mon}\mathbb{B})$  is concerned, observe that it is the Brauer cat-group of  $\mathbb{B}$  [27], so that its group of automorphisms is the Picard group of  $\mathbb{B}$ . Since  $\mathbb{B}$  is already a compact closed groupoid, its Picard group is nothing but  $cl_1(\mathbb{B})$ .)  $\square$

## 4 Conclusion

The interest of our technique is that now we can apply Proposition 1 and 2 choosing as compact closed groupoids those defined by convenient monoidal bicategories.

1. Consider a unital commutative ring  $R$  and let  $\mathbb{C} = R\text{-mod}$  be the category of  $R$ -modules. If we put  $\mathbb{B} = P_2(cl_2(\text{Mon}\mathbb{C}))$ , Proposition 1 gives the well-known K-theoretical interpretation of the Picard and Brauer groups of  $R$ , and Proposition 2 gives the classical Picard-Brauer exact sequence.
2. With  $\mathbb{C} = R\text{-mod}$  as in the previous example, instead of  $\text{Mon}\mathbb{C}$  consider the monoidal bicategory  $\text{Mon}^{reg}\mathbb{C}$  of regular (but not necessarily unital)  $R$ -algebras and regular bimodules [12]. Then, taking  $\mathbb{B} = P_2(cl_2(\text{Mon}^{reg}\mathbb{C}))$ , Proposition 2 gives an exact sequence connecting the Picard and the Brauer-Taylor groups of  $R$ , and Proposition 1 provides a K-theoretical interpretation of the Brauer-Taylor group.
3. Again with  $\mathbb{C} = R\text{-mod}$ , consider the monoidal bicategory  $\text{Dist}\mathbb{C}$  of small  $\mathbb{C}$ -categories and distributors [5, 8]. Taking  $\mathbb{B} = P_2(cl_2(\text{Dist}\mathbb{C}))$ , Proposition 2 gives an exact sequence between the Picard and the categorical Brauer groups of  $R$ .
4. The three previous examples can be generalized taking as  $\mathbb{C}$  any symmetric monoidal category with stable coequalizers.

5. A curiosity: if we take  $\mathbb{B} = P_2(R\text{-mod})$ , then Proposition 2, which in the first example gives the Picard-Brauer sequence, gives now the Unit-Picard sequence, because  $\mathbb{B}(I, I)$  is isomorphic to the group of units of  $R$ .

## References

- [1] B. AUSLANDER: *The Brauer group of a ringed space*, J. Algebra 4 (1966) 220-273.
- [2] H. BASS: *Algebraic K-Theory*, W.A. Benjamin Inc. (1968).
- [3] J. BÉNABOU: *Introduction to bicategories*, Springer LNM 40 (1967) 1-77.
- [4] F. BORCEUX: *Handbook of categorical algebra*, Cambridge University Press (1994).
- [5] F. BORCEUX, E.M. VITALE: *Azumaya categories*, Applied Categorical Structures (to appear).
- [6] S. CAENEPEEL: *Brauer groups, Hopf algebras and Galois theory*, Kluwer Academic Publishers (1998).
- [7] S. CAENEPEEL, F. GRANDJEAN: *A note on Taylor's Brauer group*, Pacific J. Math. 186 (1998) 13-27.
- [8] B. DAY, R. STREET: *Monoidal bicategories and Hopf algebroids*, Adv. Math. 129 (1997) 99-157.
- [9] J.M. FERNANDEZ VILABOA, R. GONZALEZ RODRIGUEZ, E. VILLANUEVA NOVOA: *The Picard-Brauer five-term exact sequence for a cocommutative finite Hopf algebra*, J. Algebra 186 (1996) 384-400.
- [10] R. GONZALEZ RODRIGUEZ: *La sucesion exacta ...*, Ph. D. Thesis, Santiago de Compostela (1994).

- [11] R. GORDON, A.J. POWER, R. STREET: *Coherence for tricategories*, Mem. Am. Math. Soc. 558 (1995).
- [12] F. GRANDJEAN, E.M. VITALE: *Morita equivalence for regular algebras*, Cahier de Topologie Géométrie Différentielle Catégoriques 39(1998) 137-153.
- [13] S. KASANGIAN, E.M. VITALE: *Factorization systems for symmetric cat-groups*, Theory and Applications of Categories 7 (2000) 47-70 (available at <http://www.tac.mta.ca/tac/>).
- [14] G.M. KELLY, M.L. LAPLAZA: *Coherence for compact closed categories*, J. Pure Applied Algebra 19 (1980) 193-213.
- [15] M.L. LAPLAZA: *Coherence for categories with group structure: an alternative approach*, J. Algebra 84 (1983) 305-323.
- [16] F.W. LONG: *The Brauer group of dimodule algebras*, J. Algebra 30 (1974) 559-601.
- [17] B. MITCHELL: *Separable algebroids*, Mem. Am. Math. Soc. 333 (1985).
- [18] M.-A. MOENS, U. BERNI-CANANI, F. BORCEUX: *On regular presheaves and regular graphs*, preprint (1999).
- [19] M.-A. MOENS, F. BORCEUX: *On Azumaya graphs*, preprint (2000).
- [20] M. ORZECZ: *Brauer groups and class groups for a Krull domain*, Springer LNM 917 (1982) 66-90.
- [21] I. RAEBURN, J.L. TAYLOR: *The bigger Brauer group and étale cohomology*, Pacific J. Math. 119 (1985) 445-463.
- [22] J.L. TAYLOR: *A bigger Brauer group*, Pacific J. Math. 103 (1982) 162-203.
- [23] A. VERSCHOREN: *Exact sequences for relative Brauer groups and Picard groups*, J. Pure Applied Algebra 28 (1983) 93-108.

- [24] A. VERSCHOREN: *Brauer groups and class groups for Krull domains: A K-theoretic approach*, J. Pure Applied Algebra 33 (1984) 219-224.
- [25] A. VERSCHOREN: *Relative invariants of sheaves*, Marcel Dekker (1987).
- [26] E.M. VITALE: *The Brauer and Brauer-Taylor groups of a symmetric monoidal category*, Cahier Topologie Géométrie Différentielle Catégoriques 37 (1996) 91-122.
- [27] E.M. VITALE: *A Picard-Brauer exact sequence of categorical groups*, preprint (2000).

Marie-Anne Moens

Département de Mathématique, Université Catholique de Louvain,  
Chemin du Cyclotron 2, 1348 Louvain-la-Neuve, Belgium  
m.moens@agel.ucl.ac.be

Enrico M. Vitale

Département de Mathématique, Université catholique de Louvain,  
Chemin du Cyclotron 2, 1348 Louvain-la-Neuve, Belgium  
vitale@agel.ucl.ac.be