

# JOURNÉES ÉQUATIONS AUX DÉRIVÉES PARTIELLES

GERD GRUBB

## **Singular perturbations and parameter-dependent pseudo-differential boundary problems**

*Journées Équations aux dérivées partielles* (1986), p. 1-7

[http://www.numdam.org/item?id=JEDP\\_1986\\_\\_\\_A9\\_0](http://www.numdam.org/item?id=JEDP_1986___A9_0)

© Journées Équations aux dérivées partielles, 1986, tous droits réservés.

L'accès aux archives de la revue « Journées Équations aux dérivées partielles » (<http://www.math.sciences.univ-nantes.fr/edpa/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

SINGULAR PERTURBATIONS AND  
PARAMETER-DEPENDENT PSEUDO-DIFFERENTIAL BOUNDARY PROBLEMS

GERD GRUBB  
MATEMATISK INSTITUT, UNIVERSITETSPARKEN 5  
2100 COPENHAGUE, DANEMARK

The purpose of this talk is to show how the theory of parameter-dependent pseudo-differential boundary problems developed in Grubb [7-12] can be used to treat some important problems in singular perturbation theory, for differential operators as well as pseudo-differential operators.

The parameter-dependent ps.d.o. calculus was developed in particular for the study of resolvents of elliptic boundary value problems. Let  $\bar{\Omega}$  be a compact  $n$ -dimensional  $C^\infty$  manifold with boundary  $\partial\Omega$  and interior  $\Omega$ ; let  $E$  be an  $N$ -dimensional  $C^\infty$  vector bundle over  $\bar{\Omega}$  and  $F$  an  $M$ -dimensional  $C^\infty$  vector bundle over  $\partial\Omega$ . For a positive integer  $d$  we consider the system of operators defined within the Boutet de Monvel calculus [2]

$$(1) \quad \begin{pmatrix} P_\Omega + G \\ T \end{pmatrix} : C^\infty(E) \rightarrow \begin{matrix} C^\infty(E) \\ C^\infty(F) \end{matrix},$$

where  $P$  is a pseudo-differential operator in a bundle  $\tilde{E}$  extending  $E$  to an open neighborhood of  $\bar{\Omega}$  and  $P_\Omega$  is its restriction to  $\Omega$ ,  $P$  has the transmission property at  $\partial\Omega$ ,  $G$  is a singular Green operator in  $E$  of order  $d$  and class  $\leq d$ , and  $T = \{T_0, \dots, T_{d-1}\}$  is a system of trace operators from  $E$  to  $F_k$  of order  $k$  and class  $\leq k+1$ ; here  $F = F_0 \oplus \dots \oplus F_{d-1}$  and  $\dim F_k = M_k \geq 0$ , so that  $M = M_0 + \dots + M_{d-1}$ . (When  $M_k=0$ ,  $T_k$  is void.) Let  $B$  denote the  $L^2$ -realization defined by  $\{P_\Omega + G, T\}$ , that is,  $B$  acts like  $P_\Omega + G$  in  $L^2(E)$ , with domain

$$(2) \quad D(B) = \{u \in H^d(E) \mid Tu = 0\}.$$

The resolvent  $R_\lambda = (B - \lambda)^{-1}$  is constructed in our work under the following hypothesis (expressed in local coordinates, with  $x = (x', x_n) \in \bar{\mathbb{R}}_+^n$ ,  $\xi \in \mathbb{R}^n$ ,  $\mu \geq 0$ ):

Hypothesis of parameter-ellipticity on the ray  $\lambda = re^{i\theta_0}$ . Let  $\theta_0 \in [0, 2\pi]$ , and let  $\omega = \exp[i(\theta_0 - \pi)]$ .

(I) The principal, strictly homogeneous, interior symbol  $p^h(x, \xi) + \omega\mu^d$  is bijective in  $\mathbb{C}^N$  for all  $x \in \overline{\mathbb{R}}_+^n$ ,  $\xi \in \mathbb{R}^n \setminus 0$ ,  $\mu \geq 0$ .

(II) The principal, strictly homogeneous, boundary symbol operator

$$(3) \quad a^h(x', \xi', \mu, D_n) = \begin{pmatrix} p^h(x', 0, \xi', D_n)_\Omega + g^h(x', \xi', D_n) + \omega\mu^d \\ t^h(x', \xi', D_n) \end{pmatrix} : H^d(\overline{\mathbb{R}}_+)^N \rightarrow \begin{matrix} L^2(\mathbb{R}_+) \\ \times \\ \mathbb{C}^M \end{matrix}$$

is bijective for all  $x' \in \mathbb{R}^{n-1}$ ,  $\xi' \in \mathbb{R}^{n-1} \setminus 0$ ,  $\mu \geq 0$ .

(III) For  $\xi' \rightarrow 0$ ,  $a^h(x', \xi', \mu, D_n)$  converges in symbol norm to an operator  $a^h(x', 0, \mu, D_n)$  that is bijective from  $H^d(\overline{\mathbb{R}}_+)^N$  to  $L^2(\mathbb{R}_+)^N \times \mathbb{C}^M$  for all  $x'$ , all  $\mu > 0$ .

Conditions (I) and (II) are straightforward ellipticity hypotheses, whereas (III) plays a more special rôle. Condition (III) implies that the boundary condition is normal; i.e. the trace operator is of the form

$$(4) \quad T = \begin{pmatrix} s_{00}(x')\gamma_0 + T'_0 \\ s_{11}(x')\gamma_1 + s_{10}(x', D')\gamma_0 + T'_1 \\ \vdots \\ s_{d-1, d-1}(x')\gamma_{d-1} + \sum_{j < d-1} s_{d-1, j}(x', D')\gamma_j + T'_{d-1} \end{pmatrix},$$

where the  $T'_j$  are of class 0, and the coefficient matrix

$$(5) \quad \begin{pmatrix} s_{00}(x') \\ s_{11}(x') \\ \vdots \\ s_{d-1, d-1}(x') \end{pmatrix} : \mathbb{C}^N \rightarrow \mathbb{C}^M$$

is surjective at each  $x'$ . In fact, if  $N = 1$  (and in certain cases where  $N > 1$ , e.g. when  $\dim F_k = N$  or  $0$  for all  $k$ ), condition (III) is equivalent with normality. (In the most general cases, normality plus invertibility of a certain matrix function imply (III).)

(III) is usually naturally assumed (more or less tacitly) in the resolvent

constructions for differential operator boundary problems, cf. Agmon [1], Seeley [17]. It is of interest to discuss cases where (III) does not hold (cf. Rempel and Schulze [16] and Grubb [10-12]), but our main positive results are shown under this assumption.

We show that when (I)-(III) hold for a  $\theta_0$ , then the resolvent  $R_\lambda = (B-\lambda)^{-1}$  exists, as a bounded operator in  $L^2(E)$ , for  $\lambda = \mu^d \exp(i\theta_0)$ ,  $\mu \geq \mu_0$  (with a sufficiently large  $\mu_0$ ). It satisfies the estimates

$$(6) \quad \langle \mu \rangle^{s+d} \|R_\lambda f\|_0 + \|R_\lambda f\|_{d+s} \leq c_s (\langle \mu \rangle^s \|f\|_0 + \|f\|_s) \quad \text{for any } s \geq 0,$$

and the precise structure is described in a generalized Boutet de Monvel calculus. This is used in [12] to discuss: complex powers  $B^Z$  (when the Hypothesis holds for  $\theta_0 = \pi$ ); the "heat operator"  $\exp(-tB)$  and its trace, cf. also [10] (when the Hypothesis holds for all  $\theta_0 \in [\pi/2, 3\pi/2]$ , i.e. parabolicity holds); the index of any normal elliptic problem; spectral asymptotics; and singular perturbations (see also [11]). We describe here the last application.

Consider a boundary problem (all orders in the following are integers)

$$(7) \quad \begin{aligned} A_0 u &= f \quad \text{in } E, \\ T_{0,j} u &= \varphi_{0,j} \quad \text{at } \partial\Omega, \quad 0 \leq j < k'_0, \end{aligned}$$

where  $A_0 = P_{0,\Omega} + G_0$  is of order  $r \geq 0$  and class 0, and  $T_{0,j}: C^\infty(E) \rightarrow C^\infty(F_j^0)$  is of order  $j$  and class  $\leq j+1$  for each  $j$ ; here we denote  $\{T_{0,j}\}_{0 \leq j < k'_0} = T_0$ ,  $\oplus_{0 \leq j < k'_0} F_j^0 = F^0$  and  $\dim F^0 = M^0$ . Consider also an associated "perturbed problem"

$$(8) \quad \begin{aligned} (\varepsilon^d A_1 + A_0) u_\varepsilon &= f \quad \text{in } E, \\ T_{0,j} u_\varepsilon &= \varphi_{0,j} \quad \text{at } \partial\Omega \quad \text{for } 0 \leq j < k'_0, \\ T_{1,j} u_\varepsilon &= \varphi_{1,j} \quad \text{at } \partial\Omega \quad \text{for } k_1 \leq j < k_1, \end{aligned}$$

where  $A_1 = P_{1,\Omega} + G_1$  is of order  $r+d$  and class 0,  $d > 0$ , and  $T_{1,j}: C^\infty(E) \rightarrow C^\infty(F_j^1)$  is of order  $j$  and class  $\leq j+1$  for each  $j$ . Here we denote  $\{T_{1,j}\}_{k_1 \leq j < k'_1} = T_1$ ,  $\oplus_{k_1 \leq j < k'_1} F_j^1 = F^1$  and  $\dim F^1 = M^1$ . We assume:

$$1^0 \quad 0 \leq k'_0 < r, \quad \text{and} \quad \begin{pmatrix} A_0 \\ T_0 \end{pmatrix} \quad \text{is } \underline{\text{elliptic}} \text{ and bijective, with inverse}$$

$$(R_0 \quad K_0).$$

2<sup>0</sup>  $k'_0 < k_1 \leq r$  and  $0 \leq k'_1 - k_1 < d$ ,  $T_1$  is normal, and  $\{A_1, T_0, T_1\}$  is elliptic.

3<sup>o</sup> The parameter-dependent principal symbol  $\varepsilon^d p_1^h(x, \xi) + p_0^h(x, \xi)$  is bijective in  $\mathbb{C}^N$  for all  $x$ , all  $\xi \neq 0$  and all  $\varepsilon > 0$ ; and the parameter-dependent principal boundary symbol operator

$$(9) \quad \begin{pmatrix} \varepsilon^d a_1^h(x', \xi', D_n) + a_0^h(x', \xi', D_n) \\ t_0^h(x', \xi', D_n) \\ t_1^h(x', \xi', D_n) \end{pmatrix} : H^{r+d}(\overline{\mathbb{R}_+})^N \rightarrow \begin{matrix} L^2(\mathbb{R}_+)^N \\ \mathbb{C}^{M^0} \\ \mathbb{C}^{M^1} \end{matrix}$$

is bijective for all  $x'$ , all  $\xi' \neq 0$  and all  $\varepsilon > 0$ .

(An extra hypothesis is added if  $N > 1$  and not all  $\dim F_j^1$  are either  $N$  or  $0$ ; otherwise the above conditions suffice for our purposes.)

The problem (8) will be solvable for sufficiently small  $\varepsilon$ , and one is interested in the behavior of  $u_\varepsilon$  for  $\varepsilon \rightarrow 0$ . In fact,  $u_\varepsilon$  converges to  $u$  in certain spaces, and one wants to determine this behavior, and more precisely to get explicit formulas for the operator mapping  $\{f, \varphi_0, \varphi_1\}$  into  $u_\varepsilon - u$ .

The problem has been studied for differential operators by numerous authors; let us in particular mention Vishik and Lyusternik [18], Huet [13-15] and Frank [3], and the generalizations of Frank and Wendt [4-6] to certain pseudo-differential cases with rational symbols. See also their references.

Example. Consider the problems

$$(10) \quad \begin{aligned} \varepsilon^2 \Delta^2 u_\varepsilon - \Delta u_\varepsilon &= f & \text{in } \Omega, \\ \gamma_0 u_\varepsilon &= 0 & \text{at } \partial\Omega, \\ \gamma_1 u_\varepsilon &= 0 & \text{at } \partial\Omega, \end{aligned}$$

and

$$(11) \quad \begin{aligned} \varepsilon^2 \Delta^2 u'_\varepsilon - \Delta u'_\varepsilon &= f & \text{in } \Omega, \\ \gamma_0 u'_\varepsilon &= 0 & \text{at } \partial\Omega, \\ \gamma_2 u'_\varepsilon &= 0 & \text{at } \partial\Omega, \end{aligned}$$

that are singular perturbations of the Dirichlet problem

$$(12) \quad -\Delta u = f \quad \text{in } \Omega, \quad \gamma_0 u = 0 \quad \text{at } \partial\Omega.$$

An obvious reduction of (10) and (11) is to insert  $v_\varepsilon = -\Delta u_\varepsilon$  (so that  $u_\varepsilon = R_0 v_\varepsilon$ ). Multiplying by  $\mu^2 = \varepsilon^{-2}$  we reduce (10) to the problem

$$(13) \quad (-\Delta + \mu^2)v_\varepsilon = g \quad \text{in } \Omega, \quad \gamma_1 R_0 v_\varepsilon = 0 \quad \text{at } \partial\Omega,$$

and we reduce (11) to the problem

$$(14) \quad (-\Delta + \mu^2)v'_\varepsilon = g \quad \text{in } \Omega, \quad \gamma_2 R_0 v'_\varepsilon = 0 \quad \text{at } \partial\Omega,$$

where  $g = \mu^2 f$ . Here

$$(15) \quad \gamma_2 R_0 = s(x')\gamma_0 + T'_0$$

is a normal trace operator with  $s(x') \neq 0$  for all  $x'$ , whereas  $\gamma_1 R_0$  is of order  $-1$  and not normal. Our resolvent theory applies easily to (14), but not to (13) where condition (III) is violated.

Completely similar considerations hold in the general case: If we insert  $u_\varepsilon = R_0 v_\varepsilon + K_0 \varphi_0$ , we can reduce problem (8) to a resolvent problem, where the boundary condition is normal if the lowest order in  $T_1$  is  $\geq r$ ; in which case the resolvent construction is directly applicable. But if  $T_1$  contains nontrivial entries of order  $< r$ , the boundary condition resulting from this reduction is not normal, so condition (III) is violated.

The violation of condition (III) need not be prohibitive in constant coefficient cases on  $\overline{\mathbb{R}}_+^n$ , where it is sometimes possible to get solvability results and resolvent formulas e.g. by working in other (parameter-dependent) Sobolev spaces than the most usual ones; see Frank and Wendt [4-6] for a treatment of certain types of ps.d.o. singular perturbations, and see Rempel and Schulze [16] Part I for some further observations.

However, for variable coefficient cases, the failure of condition (III) is a severe handicap if one wants to use pseudo-differential techniques, for it means (in the symbol classes we consider) that the terms in the inversion formulas arising from differentiation and lower order symbols do not have a better  $\mu$ -behavior than the principal part (in fact they generally blow up in  $\mu$  for  $\mu \rightarrow \infty$ ), so the usual series expansion of the resolvent does not converge.\*) (Indeed, in Part II of Rempel-Schulze [16] on variable coefficient cases, there are made hypotheses in order to avoid this phenomenon, which imply condition (III) for Boutet de Monvel type operators. Frank and Wendt [4-6] give very few details concerning this point for boundary problems, and we have not been able to (re)construct their argument.)

Now it turns out that the problem of the violation of condition (III) (the so-called negative regularity) can be avoided altogether if we use a little more

\*) -----  
When (I)-(III) hold, the  $\xi$ -derivatives and lower order terms are at least  $\mu^{-\frac{1}{2}}$  better than the principal term.

of the Boutet de Monvel calculus before going on to the resolvent. By the hypotheses on the various orders, there exists an integer  $\ell$  such that

$$(16) \quad k_0' < r - \ell \leq k_1$$

(for example,  $\ell = 1$  in (10)). Moreover, there exists an elliptic ps.d.o.  $\Lambda_{-\ell}$  (restricted to  $\Omega$ ) such that  $\Lambda_{-\ell}$  maps  $H^s(E)$  isomorphically onto  $H^{s+\ell}(E)$  for all  $s \geq 0$ . (The symbol of  $\Lambda_{-\ell}$  is a modified version of  $\langle \xi' \rangle - i\xi_n^{-\ell}$ .) Instead of inserting  $u_\varepsilon = R_0 v_\varepsilon + K_0 \varphi_0$ , we now insert  $u_\varepsilon = R_0 (\Lambda_{-\ell})^{-1} w_\varepsilon + K_0 \varphi_0$  in (8), and arrive by composition with  $\Lambda_{-\ell}$  and multiplication by  $\mu^d = \varepsilon^{-d}$  at the problem

$$(17) \quad \Lambda_{-\ell} A_1 R_0 (\Lambda_{-\ell})^{-1} w_\varepsilon + \mu^d w_\varepsilon = g \text{ in } E, \quad T_1 R_0 (\Lambda_{-\ell})^{-1} w_\varepsilon = \varphi_1 \text{ at } \partial\Omega,$$

$g = \Lambda_{-\ell} \mu^d f$ , where the new interior operator

$$(18) \quad A = \Lambda_{-\ell} A_1 R_0 (\Lambda_{-\ell})^{-1} = [\Lambda_{-\ell} P_1 P_0^{-1} (\Lambda_{-\ell})^{-1}]_\Omega + G$$

is of order  $d$  and class 0, and the new trace operator

$$(19) \quad T = T_1 R_0 (\Lambda_{-\ell})^{-1}$$

is normal. The hypotheses  $1^0-3^0$  then assure that (17) satisfies conditions (I)-(III) if  $N = 1$ , or  $N > 1$  and  $\dim F_j^1 = N$  or 0 for all  $j$  (in the remaining cases, and extra matrix condition is added).

Now the theorems of Grubb [7-12] can be applied to (17), which gives an explicit solution operator for  $\mu \geq \mu_0$  ( $\mu_0$  sufficiently large), i.e. for  $\varepsilon \in ]0, \mu_0^{-1}]$ . This is then carried back to give the solution operator for (8), and the various resolvent and operator estimates can be used to obtain estimates of  $u_\varepsilon - u$  for  $\varepsilon \rightarrow 0$ , as well as expansions in powers of  $\varepsilon^{\frac{1}{2}}$ .

Further details are given in Grubb [11] and [12].

Also problems with a more complicated  $\varepsilon$ -dependence can be treated by use of the general  $\mu$ -dependent calculus in Grubb [12].

We are grateful to Denise Huet for having called our attention to singular perturbation problems and their relation to resolvents, which not only resulted in the present work, but also at an earlier stage led to improvements in [12].

#### REFERENCES:

- [1] S. Agmon: On the eigenfunctions and on the eigenvalues of general elliptic boundary value problems, Comm. Pure Appl. Math. 15 (1962), 119-147.

- [2] L. Boutet de Monvel: Boundary problems for pseudo-differential operators, *Acta Math.* 126 (1971), 11-51.
- [3] L.S. Frank: Coercive singular perturbations I: A priori estimates, *Ann. Mat. pura appl.* 119 (1979), 41-113.
- [4] L.S. Frank and W.D. Wendt: Coercive singular perturbations II: Reduction to regular perturbations and applications, *Comm. Part. Diff. Equ.* 7 (1982), 469-535.
- [5] L.S. Frank and W.D. Wendt: Coercive singular perturbations III: Wiener-Hopf operators, *J. d'Analyse Math.* 43 (1984), 88-135.
- [6] L.S. Frank and W.D. Wendt: Erratum, *Comm. Part. Diff. Equ.* 10 (1985), 1227.
- [7] G. Grubb: On Pseudo-Differential Boundary Problems. Reports no. 2 (1979), 1, 2, 7 and 8 (1980), Copenhagen University Math. Dept. Publication Series.
- [8] G. Grubb: Problèmes aux limites dépendant d'un paramètre, *C.R. Acad. Sci. Paris (Sér. I)* 292 (1981), 581-583.
- [9] G. Grubb: La résolvante d'un problème aux limites pseudo-différentiel elliptique, *C.R. Acad. Sci. Paris (Sér. I)* 292 (1981), 625-627.
- [10] G. Grubb: The heat equation associated with a pseudo-differential boundary problem, *Seminar Analysis 1981-82*, Akad. Wiss. Berlin, 27-41. [Also available as Preprint no. 2, 1982, Copenhagen Univ. Math. Dept.]
- [11] G. Grubb: Une méthode pseudo-différentielle pour les perturbations singulières elliptiques, *C.R. Acad. Sci. Paris (Sér. I)* 301 (1985), 427-430.
- [12] G. Grubb: *Functional Calculus of Pseudo-Differential Boundary Problems*, Progress in Mathematics, vol. 65, Birkhäuser Boston 1986.
- [13] D. Huet: *Décomposition spectrale et opérateurs*. Presses universitaires de France, Paris 1977.
- [14] D. Huet: Proper approximation of a normed space and singular perturbations, *Analytical and Numerical Approaches to Asymptotic Problems in Analysis*, North Holland Publ. Co. 1981, 87-98.
- [15] D. Huet: Convergence propre et perturbations singulières coercives, *C.R. Acad. Sci. Paris* 301 (Sér. I) (1985), 435-438; Stabilité inverse d'une famille d'opérateurs opérant dans des espaces normés et dépendant d'un paramètre, *ibid.*, 675-677.
- [16] S. Rempel and B.-W. Schulze: Complex powers for pseudo-differential boundary problems I and II, *Math. Nachr.* 111 (1983), 41-109, and *Math. Nachr.* 116 (1984), 269-314.
- [17] R. Seeley: The resolvent of an elliptic boundary problem, *Amer. J. Math.* 91 (1969), 889-920.
- [18] M.I. Vishik and L. Lyusternik: Regular degeneration and boundary layer for linear differential equations with small parameters, *Uspekhi Mat. Nauk* 12 (1957), 3-122 = *Amer. Math. Soc. Transl. (2)*, 20 (1962), 239-264.