

# JOURNÉES ÉQUATIONS AUX DÉRIVÉES PARTIELLES

ANDERS MELIN

## **Some problems in inverse scattering theory**

*Journées Équations aux dérivées partielles* (1987), p. 1-3

[http://www.numdam.org/item?id=JEDP\\_1987\\_\\_\\_\\_A3\\_0](http://www.numdam.org/item?id=JEDP_1987____A3_0)

© Journées Équations aux dérivées partielles, 1987, tous droits réservés.

L'accès aux archives de la revue « Journées Équations aux dérivées partielles » (<http://www.math.sciences.univ-nantes.fr/edpa/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## Some problems in inverse scattering theory.

ANDERS MELIN

Department of Mathematics  
University of Lund

We shall consider the Schrödinger operator  $H_v = -\Delta + v(x)$  in  $\mathbf{R}^n$ , where  $n = 3, 5, \dots$ . We assume that  $v \in \mathcal{V}$ , i.e.

$$(1) \quad \int (1 + |x|)^{|\alpha| - (n-2)} |v^{(\alpha)}(x)| dx < \infty$$

for any  $\alpha$ .

Some of the main problems we consider are the following:

- (a) Analysis of bound states and poles of the scattering matrix.
- (b) Backward scattering.
- (c) The characterization problem for scattering matrices.

This talk will be a continuation of the authors lecture at École Polytechnique [6], and we shall mainly give some comments to (a).

We shall study families of intertwining operators  $A$  such that

$$(2) \quad H_v A = A H_0$$

or equivalently

$$(3) \quad (\Delta_x - \Delta_y - v(x))A(x, y) = 0.$$

(We shall always identify operators with their distribution kernels.) Let  $\mathcal{M}$  be the set of all  $U(x, y) \in L^1_{loc}$  such that

$$\|U\|_{\mathcal{M}} = \max \left\{ \sup_x \int |U(x, y)| dy, \sup_y \int |U(x, y)| dx \right\} < \infty.$$

Then  $\|U\|_{L^p \rightarrow L^p} \leq \|U\|_{\mathcal{M}}$  for  $1 \leq p \leq \infty$  if  $U \in \mathcal{M}$ . We let  $\mathcal{M}_\theta$  be the subspace of  $\mathcal{M}$  consisting of  $U$  such that  $\langle y - x, \theta \rangle \geq 0$  in its support. Here  $\theta \in S^{n-1}$  and  $\mathcal{M}_{\theta, \lambda}$  is the set of  $U$  in  $\mathcal{M}_\theta$  such that

$$e^{-\lambda \langle y - x, \theta \rangle} U(x, y) \in \mathcal{M}_\theta.$$

The spaces  $\mathcal{M}$ ,  $\mathcal{M}_\theta$  and  $\mathcal{M}_{\theta, \lambda}$  are Banach algebras. Finally  $\tilde{\mathcal{M}}_{\theta, \lambda}$  is defined by the following conditions:

$$\int |U(x, y)| dy \rightarrow 0 \text{ as } |x| \rightarrow \infty, x/|x| \rightarrow \theta$$

and

$$\int |U(x, y)| dx \rightarrow 0 \text{ as } |y| \rightarrow \infty, y/|y| \rightarrow -\theta.$$

**Example.** If  $q \in L^1(\mathbf{R}^n)$  we let  $[q]$  be the convolution operator with kernel  $q(x - y)$ . If  $\langle x, \theta \rangle \leq 0$  in the support of  $q$ , then  $(I - [q])^{-1}$  exists in  $I + \mathcal{M}_{\theta, \lambda}$  when  $\lambda$  is large.

**THEOREM 1.** Let  $v \in \mathcal{V}$  be real valued and  $\theta \in S^{n-1}$ . Then there is a unique  $A_\theta \in \cup_{\lambda \geq 0} I + \tilde{\mathcal{M}}_{\theta, \lambda}$  such that  $H_\nu A = AH_0$ . Moreover,  $A_{-\theta}^* \circ A_\theta = I$ .

The distribution  $A_\theta$  is constructed as the infinite sum  $\sum_0^\infty U_N$ , where  $U_0(x, y) = \delta(x - y)$ , and

$$U_{N+1} = E_\theta * (vU_N),$$

Here  $(vU_N)(x, y) = v(x)U_N(x, y)$ , and  $E_\theta$  is the fundamental solution for  $\Delta_x - \Delta_y$ , which is uniquely determined from the following conditions:

- (i)  $\langle y - x, \theta \rangle \geq 0$  in the support of  $E_\theta$ ,
- (ii)  $E_\theta(x + t\theta, y + t\theta) \rightarrow 0$  in  $\mathcal{D}'(\mathbb{R}^n \times \mathbb{R}^n)$  as  $|t| \rightarrow \infty$ .
- (iii)  $E_\theta = \sum c_{\alpha, \beta} \partial_x^\alpha \partial_y^\beta h_{\alpha, \beta}$ , where  $\phi(x - y)h_{\alpha, \beta}(x, y) \in \mathcal{M}$  for any  $\phi \in C_0$ .

**THEOREM 2.** There exists a family of  $L^1$  functions  $q_\theta$  in  $\mathbb{R}^n$  which depend continuously on  $\theta$  and are supported in the set where  $\langle x, \theta \rangle \leq 0$ , such that

$$A_\theta(I - [q_\theta]) \in I + \mathcal{M}_\theta.$$

**COROLLARY 3.** Assume that  $v \in C_0^\infty$ . Then the scattering matrix  $S_k(\theta, \theta')$  is analytic in the upper half-plane  $\Im k \geq 0$  after multiplication by  $1 - \widehat{q_\theta}(-k)$ .

**Sketch of proof .** One first constructs  $B_\theta \in I + \tilde{\mathcal{M}}_{\theta, 0}$  so that

$$B_\theta^{-1} H_\nu B_\theta = H_0 + \sum_1^N f_j \otimes g_j,$$

where  $f_j$  and  $g_j$  are in  $L^1$  together with all their derivatives.

Next one defines the  $L^1$  functions  $q_{jk}$  by the formula

$$(4) \quad q_{jk}(y) = \int (\check{f}_j * g_k)(x) E_\theta(x, y) dx.$$

Set  $[Q] = [q_{jk}]$ , where the right-hand side is considered as a  $N \times N$  matrix of convolution operators, and define the vector valued function  $\vec{h} = (h_1, \dots, h_N)$  by the equation

$$\vec{h}{}^{\circ\circ}(I - [Q])\vec{g},$$

where  ${}^{\circ\circ}(I - [Q])$  denotes the co-factor matrix of  $I - [Q]$ . We can now define the  $L^1$  function  $q = q_\theta$  by the equation

$$\det(I - [Q]) = I - [q].$$

It is easy to see that  $\langle x, \theta \rangle \leq 0$  in the support of  $q_\theta$ . Set

$$C_\theta = I - [q_\theta] + F_\theta,$$

where  $F_\theta = \sum_1^N E_\theta * (f_j \otimes h_j)$ . Then  $H_\nu(B_\theta C_\theta) = (B_\theta C_\theta)H_0$ . Therefore, if we set

$$R(x, y) = A_\theta^{-1} B_\theta C_\theta - \delta(x - y),$$

then  $(\Delta_x - \Delta_y)R = 0$  and  $\langle y - x, \theta \rangle \geq 0$  in its support. From a uniqueness result for  $\Delta_x - \Delta_y$  one then finds that  $R$  is constant in the direction of  $(\theta, \theta)$ , i.e.  $R(x + t\theta, y + t\theta) = R(x, y)$  when  $t$  is any real number. Since  $R + [q] \in \tilde{\mathcal{M}}_{\theta, \lambda}$  we conclude that  $R + [q] = 0$ . Hence

$$A_\theta(I - [q_\theta]) = B_\theta C_\theta \in I + \tilde{\mathcal{M}}_{\theta, 0}$$

and the proof is complete.

## REFERENCES

1. M.Cheney, *Inverse scattering in dimension 2*, J. Math. Phys. **25** (1984), 94-107.
2. L.D. Faddeev, *The inverse problem in quantum theory of scattering*, J. Math. Phys. **4** (1963), 72-104.
3. L.D. Faddeev, *Inverse problem of quantum scattering theory, II*, J. Sov. Math. **5** (1976), 334-396.
4. A. Melin, *Operator methods for inverse scattering on the real line*, Comm. in Partial Differential Eqs. **10** (1985), 677-766.
5. A.Melin, *Intertwining methods in the theory of inverse scattering*, to appear in Int. J. of Quantum Chemistry.
6. A.Melin, *Sem.Eq. Der. Part. 1986-1987, École Polytechnique*.
7. R.G. Newton, *Inverse scattering II. Three dimensions*, J. Math. Phys. **21** (1980), 1698-1715.
8. R. G. Newton, *An inverse spectral problem in three dimensions*, SIAM-AMS Proceedings **14**, 81-90.