

JOURNAL  
DE  
MATHÉMATIQUES

PURES ET APPLIQUÉES

FONDÉ EN 1836 ET PUBLIÉ JUSQU'EN 1874

PAR JOSEPH LIOUVILLE

---

LAWRENCE MARKUS

**On completeness of invariant measures defined by differential equations**

*Journal de mathématiques pures et appliquées* 9<sup>e</sup> série, tome 31 (1952), p. 341-353.

[http://www.numdam.org/item?id=JMPA\\_1952\\_9\\_31\\_\\_341\\_0](http://www.numdam.org/item?id=JMPA_1952_9_31__341_0)



NUMDAM

Article numérisé dans le cadre du programme  
Gallica de la Bibliothèque nationale de France  
<http://gallica.bnf.fr/>

et catalogué par Mathdoc  
dans le cadre du pôle associé BnF/Mathdoc  
<http://www.numdam.org/journals/JMPA>

---

---

*On completeness of invariant measures defined  
by differential equations;*

**PAR LAWRENCE MARKUS.**

Harvard University.

---

**I. — Introduction.**

Let the real, first order, non-singular, ordinary differential system

$$(1) \quad \frac{dx}{dt} = f(x, y), \quad \frac{dy}{dt} = g(x, y) \quad (f^2 + g^2 > 0),$$

with

$$f, g \in C^{(\alpha)} \quad (\alpha = 1, 2, 3, \dots, \infty, A) \quad (1)$$

be defined in an open plane set  $R$ . Then through each point  $P_0 : (x_0, y_0) \in R$  there exists an unique solution curve

$$x = x(t; x_0, y_0), \quad y = y(t; x_0, y_0),$$

initiating at  $P_0$  for  $t=0$ , and defined for some maximal « time » interval

$$\tau_-(P_0) < t < \tau_+(P_0).$$


---

(1)  $f(x_1, x_2, \dots, x_n) \in C^{(0)}$  in  $R$ , an open set of Euclidean  $n$ -space, in case  $f(x)$  is continuous, in  $n$  variables in  $R : f(x_1, x_2, \dots, x_n) = f(x) \in C^{(K)}$  ( $K=1, 2, \dots$ ) in  $R$  in case all the partial derivatives of  $f(x)$  up to and including those of order  $K$  exist (are finite) and are continuous in  $R$ .  $f(x) \in C^{(\infty)}$  in case all partial derivatives of  $f(x)$  exist and are continuous in  $R$ .  $f(x) \in C^{(A)}$  in case  $f(x)$  is analytic in  $R$ , that is, near each point in  $R$ ,  $f(x)$  has an absolutely convergent, real,  $n$ -variable power series representation. We define  $0 < 1 < 2 < \dots < \infty < A$  and also  $\infty \pm K = \infty$  and  $A \pm K = A$ .

The subset of Euclidean 3-space in which  $x(t; x_0, y_0)$  and  $y(t; x_0, y_0)$  are defined is open and therein these functions, as well as  $\frac{\partial x}{\partial t}$  and  $\frac{\partial y}{\partial t}$ , are in class  $C^{(\alpha)}$ . The transformations  $T_t$

$$(x_0, y_0) \rightarrow x(t; x_0, y_0), y(t; x_0, y_0),$$

from an open subset  $R_0 \subset R$  onto open subsets  $R_t \subset R$ , from a one-parameter local group, or stream, of  $C^{(\alpha)}$ -homeomorphisms <sup>(2)</sup>.

Suppose the ordinary differential system (1) were exact, that is,  $\frac{\partial f}{\partial x} + \frac{\partial g}{\partial y} \equiv 0$  and  $R$  were a simply-connected region (open, connected set <sup>(3)</sup>). Then  $\{T_t\}$  forms a measure true stream, that is,  $T_t$  is measurability preserving ( $T_t$  and  $T^{-1}$  preserve Lebesgue measurability of sets) and also  $T_t$  preserves the magnitude of the measure. In this case, there exists a stream function  $\psi(x, y) \in C^{(\alpha+1)}$  in  $R$  such that  $f = \frac{\partial \psi}{\partial y}$ ,  $g = -\frac{\partial \psi}{\partial x}$  and  $\psi(x, y)$  is a principal integral <sup>(4)</sup> of the first order, partial differential equation

$$(2) \quad f(x, y) \frac{\partial \psi}{\partial x} + g(x, y) \frac{\partial \psi}{\partial y} = 0.$$

That is,  $\psi(x, y)$  is a solution of (2) which is constant on no open set and therefore the class of all solutions of (2) are precisely those functions in  $C^{(1)}$  in  $R$  which are functionally dependent on  $\psi(x, y)$ .

Even if the ordinary differential system (1) is not exact, there may exist a principal integral  $\psi(x, y) \in C^{(2)}$  of the corresponding partial differential equation (2). Then

$$\mu(x, y) = \left[ \frac{\psi_x^2 + \psi_y^2}{f^2 + g^2} \right]^{\frac{1}{2}} \in C^{(1)}$$

<sup>(2)</sup> A homeomorphism  $T$  of an open set  $R_1$  of Euclidean  $n$ -space onto a second open set  $R_2$  of the same space is called a  $C^{(\beta)}$ -homeomorphism,  $\beta = 0, 1, 2, \dots, \infty$ , if in case both  $T$  and  $T^{-1}$  are expressed by functions of class  $C^{(\beta)}$ .

<sup>(3)</sup> For notation, see E. HOPF, *Ergodentheorie* (Berlin, 1937).

<sup>(4)</sup> E. KAMKE, *Differentialgleichungen Reeller Funktionen* (Chelsea, 1947), p. 323.

is a non-negative integrating factor for (1), that is,

$$\mu f = \psi_y, \quad \mu g = -\psi_x.$$

In case (1) is exact then  $\mu(x, y)$  is a constant.

Because of the relation (5).

$$(3) \quad \frac{d}{dt} \left[ \iint_{R_0} \mu(x, y) \frac{\partial(x, y)}{\partial(x_0, y_0)} dx_0 dy_0 \right] = \iint_{R_0} [(\mu f)_x + (\mu g)_y] dx_0 dy_0 = 0$$

we can define a new « invariant » measure  $m_\mu$  on the  $\sigma$ -ring <sup>(6)</sup> L of Lebesgue measurable sets of R by

$$m_\mu(\Lambda) = \iint_{\Lambda} \mu(x, y) dx, dy$$

for  $\Lambda \in L$ . Then the local group of  $C^{(\alpha)}$ -homeomorphisms  $\{T_t\}$  is a  $m_\mu$ -measure true stream. Here  $m_\mu$  is a non-negative, completely additive set function on L. For any compact set  $K \subset R$ ,  $m_\mu(K) < \infty$ .  $m_\mu$  is an absolutely continuous measure in terms of the Lebesgue measure  $m_L$ , that is, if  $m_L(\Lambda) = 0$ , then  $m_\mu(\Lambda) = 0$ . The purpose of this note is to point out that if  $\mu(x, y) \geq 0$  and vanishes on no open set, even if  $\mu(x, y) \in C^{(\infty)}$ , there can exist a set  $Z \in L$  such that  $m_\mu(Z) = 0$  but  $m_L(Z) > 0$ . In this case the invariant measure  $m_\mu$  is not complete since there exist non-measurable subsets of Z. However in case  $\mu(x, y)$  is analytic in R,  $m_\mu(\Lambda) = 0$  if and only if  $m_L(\Lambda) = 0$  and  $m_\mu$  is a complete measure.

## II. — Homeomorphisms which preserve measurability.

Any simply-connected plane region R can be mapped by a  $C^{(\alpha)}$ -homeomorphism onto the entire plane. Thus we can replace R by the entire plane in any measure-theoretic considerations which are invariant under such a map.

Let  $\theta$  be a Borel set of Euclidean  $n$ -space. Let S be the  $\sigma$ -ring of

(\*) H. POINCARÉ, *Méthodes nouvelles de la Mécanique céleste*, t. III, chap. 26.

(6) Sometimes called  $\sigma$ -field. For notation, see P. HALMOS, *Measure Theory* (New-York, 1950).

all subsets of  $\theta$ .  $S$  actually forms an algebraic commutative Boolean (idempotent) ring with unity element under the operations of symmetric difference  $\Delta$  for « addition » and intersection  $\cap$  for « multiplication ». Because of the simple, well-known formulae for union, difference, and complementation

$$(4) \quad \begin{cases} \varphi \cup \psi = (\varphi \Delta \psi) \Delta (\varphi \cap \psi), \\ \varphi - \psi = \varphi \Delta (\varphi \cap \psi), \\ \varphi' = \varphi \Delta \theta \end{cases}$$

all of the set-theoretic properties of  $S$  are determined by its structure as an algebraic ring under  $\Delta$  and  $\cap$ . Let  $B$  be the  $\sigma$ -ring of Borel sets of  $\theta$  (smallest  $\sigma$ -ring containing the open sets of  $\theta$ ), with the Borel measure  $m_b$ .  $B$  is an algebraic subring of  $S$ . The subset  $b \subset B$  of Borel sets with measure zero forms an ideal in the ring  $B$ . The  $\sigma$ -ring  $L$  of Lebesgue measurable subsets of  $\theta$  (all sets of the form  $\eta \Delta \nu$  where  $\eta \in B$  and  $\nu$  is a subset of some set in  $b$ ) with Lebesgue measure  $m_{(L)}$  is an algebraic subring of  $S$  and a superring of  $B$ . The ideal  $l \subset L$  of sets with Lebesgue measure zero is a superring of  $b$ , and consists of all sets of the form  $\beta \Delta \nu$  where  $\beta \in b$  and  $\nu$  is a subset of some set of  $b$ .

Let  $\theta_1$  and  $\theta_2$  be two Borel sets of Euclidean spaces and let  $S_1, B_1, b_1, L_1, l_1$  and  $S_2, B_2, b_2, L_2, l_2$  be the corresponding  $\sigma$ -rings described above. Let  $T$  be a homeomorphism of  $\theta_1$  onto  $\theta_2$ . Since  $T$  is a one-to-one point transformation,  $T$  induces a set mapping  $\tau$  from  $S_1$  onto  $S_2$ . Clearly  $\tau$  is an algebraic isomorphism of  $S_1$  onto  $S_2$ .

**THEOREM I.** — *Let  $T$  be a homeomorphism of  $\theta_1$  onto  $\theta_2$ , Borel sets of Euclidean spaces. Then the induced set-map  $\tau$  is an isomorphism of  $B_1$  onto  $B_2$ . Furthermore the following four conditions are equivalent :*

1.  $T$  is a  $m_L$ -measurability preserving transformation;
2.  $\tau$  induces an isomorphism of  $L_1$  onto  $L_2$ ;
3.  $\tau$  induces an isomorphism of  $b_1$  onto  $b_2$ ;
4.  $\tau$  induces an isomorphism of  $l_1$  onto  $l_2$ .

*Proof.* — Since  $T$  is a homeomorphism, open sets of  $\theta_1$  correspond to open sets of  $\theta_2$  under  $\tau$  and thus there is a one-to-one corres-

pendence between the sets of  $B_1$  and  $B_2$ . Since  $\tau$  is an isomorphism of  $S_1$  onto  $S_2$ ,  $\tau$  is therefore an isomorphism of  $B_1$  onto  $B_2$ .

Since  $\tau$  is an isomorphism of  $S_1$  onto  $S_2$ , conditions 1 and 2 are equivalent by the definition of measurability preserving transformations.

Suppose condition 3 holds; we shall deduce condition 2. If  $\Lambda_1 \in L_1$ , then  $\Lambda_1 = \eta_1 \triangle \nu_1$  where  $\eta_1 \in B_1$  and  $\nu_1 \subset \beta_1 \in b_1$ . Then

$$\tau(\Lambda_1) = \Lambda_2 = \tau(\eta_1) \triangle \tau(\nu_1) = \eta_2 \triangle \nu_2,$$

with

$$\tau(\eta_1) = \eta_2 \in B_2 \quad \text{and} \quad \tau(\nu_1) = \nu_2 \subset \tau(\beta_1) = \beta_2 \in b_2.$$

Thus  $\tau(\Lambda_1) = \Lambda_2 \in L_2$  and  $\tau$  maps  $L_1$  into  $L_2$ . Using the inverse homeomorphism  $T^{-1}$  we see that  $\tau^{-1}$  maps  $L_2$  into  $L_1$  and since  $\tau$  is one-to-one on  $S_1$ ,  $\tau$  is an isomorphism of  $L_1$  onto  $L_2$ .

Conversely suppose  $\tau$  is an isomorphism of  $L_1$  onto  $L_2$ . Suppose  $\tau(\beta_1) = \beta_2 \in B_2 - b_2$  for some  $\beta_1 \in b_1$ . Then  $m_b(\beta_2) > 0$  and there exists a set  $(^7)$   $\chi_2 \subset \beta_2$  such that  $\chi_2 \in S_2 - L_2$ . But then  $\tau^{-1}(\chi_2) = \chi_1 \subset \beta_1$  and thus  $\chi_1 \in L_1$ . But  $\tau(\chi_1) \notin L_2$  and this contradicts the hypothesis that  $\tau$  maps  $L_1$  onto  $L_2$ . Thus  $\tau$  maps  $b_1$  into  $b_2$  and, as above,  $\tau$  is an isomorphism of  $b_1$  onto  $b_2$ .

Since  $b_i = B_i \cap l_i$  ( $i=1, 2$ ), condition 4 clearly implies 3. Because  $l_i$  consists of those sets of  $L_i$  which are contained in sets of  $b_i$  ( $i=1, 2$ ), conditions 2 and 3 imply 4.

Q. E. D.

**COROLLARY.** — *If  $T$  is a  $C^{(1)}$ -homeomorphism of  $\theta_1$  onto  $\theta_2$ , open sets of an Euclidean  $n$ -space, then  $T$  is  $m_L$ -measurability preserving.*

*Proof.* — We need show only that  $\tau(b_1) \subset b_2$ . If  $\beta_1 \in b_1$ , then  $\beta_1$  has a covering by a countable number of compact closed  $n$ -balls  $\bar{R}_i$  with

$$\sum_{i=1}^{\infty} m_B(\bar{R}_i) = 1.$$

In  $\bar{R}_i$  the continuous Jacobian of  $T$  is bounded,  $|J(T)| < K_i$ . Then

$$m_B(\beta_1 \cap \bar{R}_i) = 0,$$

(<sup>7</sup>) See P. HALMOS, *op. cit.*, p. 70. The existence of a non-measurable linear set combined with an application of Fubini's theorem gives this result.

and we cover  $\beta_1 \cap \bar{R}_i$  by an open set  $o_i$  with  $m_B(o_i) = \frac{\delta}{2^i K_i}$ . Then

$$m_B(\tau(o_i)) \leq \frac{\delta}{2^i}$$

and thus

$$\sum_{i=1}^{\infty} m_B(\tau(o_i)) \leq \delta.$$

Thus  $m_B(\tau(\beta_1)) \leq \delta$  and therefore, since  $\delta$  is an arbitrary positive number,  $\tau(\beta_1) = \beta_2 \in I_2$ . But certainly  $\beta_2 \in B_2$ . Thus  $\beta_2' \in b_2$ . Therefore  $\tau(b_1) \subset b_2$  and using  $T^{-1}$ , which also has a continuous Jacobian,  $\tau^{-1}(b_2) \subset b_1$  and therefore  $\tau$  maps  $b_1$  onto  $b_2$ .

Q. E. D.

An example of a  $C^{(0)}$ -homeomorphism of the linear interval  $[0, 1]$  onto itself which is not measurability preserving is given by  $T$

$$x \rightarrow f(x) = \frac{1}{2} [x + \psi(x)]$$

where  $\psi(x)$  is the continuous, but not absolutely continuous, non-decreasing, Cantor function <sup>(8)</sup>. If  $K$  is the compact Cantor set, then  $m_B(K) = 0$  but  $m_B(f(K)) = \frac{1}{2}$ .

### III. — Completeness of invariant measures.

We shall construct a function  $\mu(x, y)$  such that :

1.  $\mu(x, y) \in C^{(\infty)}$  in the entire plane;
2.  $\mu(x, y) \geq 0$  and the plane set  $Z$  defined by  $\mu(x, y) = 0$  contains no interior points;
3.  $m_B(Z) = \infty$ .

Let  $z$  be a compact, nowhere dense point set of the real line such that the linear Borel measure  $m'_B(z) > 0$ . A closed point set is nowhere dense if and only if it contains no interior point. Such a

---

<sup>(8)</sup> P. HALMOS, *op. cit.*, p. 83.

set  $z$  is given in Hobson <sup>(9)</sup> and can be described briefly as follows.

Let  $[0, 1]$  be divided into  $m > 2$  equal parts and the last exempted from further division. Then let the remaining  $m - 1$  parts each be divided into  $m^2$  equal parts, the last of each being exempted from further division. Let the remaining parts be then divided into  $m^3$  equal parts, the last of these in each case being exempted from further division. If this process is carried out a countable number of times, the endpoints of the divisions, together with their accumulation points, form a nowhere dense, compact set  $z$ .

This set  $z$  has measure

$$0 < \prod_{i=1}^{\infty} \left(1 - \frac{1}{m^i}\right) < 1$$

for after  $i$  operations, the measure of the union of the exempted segments is

$$\frac{1}{m} + \frac{m-1}{m^3} + \frac{(m-1)(m^2-1)}{m^6} + \dots + \frac{(m-1)(m^2-1)\dots(m^{i-1}-1)}{m^{\frac{i(i+1)}{2}}}$$

or

$$1 - \left(1 - \frac{1}{m}\right) \left(1 - \frac{1}{m^2}\right) \dots \left(1 - \frac{1}{m^i}\right).$$

Thus  $m'_n(z) > 0$  and indeed, may be chosen arbitrarily close to one if the dividing ratio  $m$  be large enough. The complement of  $z$  in  $[0, 1]$  is a countable union of disjoint open intervals  $I_n$  and  $z$  consists of the closure of the endpoints of the  $I_n$ .

The distance from a real number  $x$  to the compact set  $z$  is  $d(x) \geq 0$  and  $d(x) = 0$  if and only if  $x \in z$ .

Then for any point  $(x, y)$  of the plane we define

$$(5) \quad \begin{cases} \mu(x, y) = 0 & \text{in case } x \in z. \\ \mu(x, y) = e^{-\frac{1}{x^2}} & \text{in case } x < 0. \\ \mu(x, y) = e^{-\frac{1}{(x-1)^2}} & \text{in case } x > 1. \\ \mu(x, y) = e^{-\frac{\Delta_n^2}{4} - \left(\frac{\Delta_n}{2} - d(x)\right)^2} & \text{in case } x \in I_n \end{cases}$$

<sup>(9)</sup> A. E. HOBSON, *The theory of functions a real variable* (Cambridge, 1921), p. 119 and also p. 164.

where  $\Delta_n = b_n - a_n$  is the length of  $I_n = (a_n, b_n)$ . Then  $\mu(x, y) \geq 0$  and, defining the plane set  $(x, y) \in Z$  in case  $x \in I_n$ ,  $\mu(x, y) = 0$  if and only if  $(x, y) \in Z$ . The set  $Z$  is closed and contains no interior points of the plane. Moreover  $m_n^A(Z) = \infty$ . We shall show that  $\mu(x, y) \in C^{(\infty)}$  and then clearly

$$m_\mu(Z) = \iint_Z \mu(x, y) dx dy = 0.$$

If  $x < 0$  or  $x > 1$ , then  $\mu(x, y) \in C^{(\infty)}$ . If  $x \in I_n$ , that is,  $a_n < x < b_n$ , then

$$\frac{\Delta_n^2}{4} - \left( \frac{\Delta_n}{2} - d(x) \right)^2 = (x - a_n)(b_n - x)$$

and thus  $\mu(x, y) \in C^{(\infty)}$  for these regions.

For  $x \in I_n$ ,

$$\frac{\Delta_n^2}{4} - \left( \frac{\Delta_n}{2} - d(x) \right)^2 = 2\Delta_n d - d^2 < d - d^2 < d.$$

Thus

$$|\mu(x, y)| \leq e^{-\frac{1}{d(x)}} < \frac{1}{d(x)^2} e^{-\frac{1}{d(x)}}.$$

By induction it is easy to show that

$$\frac{\partial^k \mu}{\partial x^k} = \frac{P_k((x - a_n), (b_n - x))}{[(x - a_n)(b_n - x)]^{2k}} e^{-\frac{1}{(x - a_n)(b_n - x)}}$$

where  $P_k$  is a polynomial and since

$$d(x) \leq |x - a_n| < 1, \quad d(x) \leq |b_n - x| < 1,$$

we have

$$|P_k((x - a_n)(b_n - x))| < \bar{P}_k$$

an upper bound independent of  $I_n$  and thus

$$(6) \quad \left| \frac{\partial^k \mu}{\partial x^k} \right| < \frac{\bar{P}_k}{d(x)^{2k+1}} e^{-\frac{1}{d(x)}} \quad \text{for } k = 0, 1, 2, \dots$$

For sufficiently small positive  $d$ , the function  $\frac{1}{d^{2k+1}} e^{-\frac{1}{d}}$  is strictly increasing.

Finally consider a point  $(x_0, y_0) \in Z$ . Then  $\mu(x_0, y_0) = 0$  and

for  $|x - x_0| < \delta$ , a sufficiently small positive number,  $d(x) < \delta$  and

$$0 \leq |\mu(x, y) - \mu(x_0, y_0)| \leq \frac{\bar{P}_0}{d(x)^2} e^{-\frac{1}{d(x)}} < \frac{\bar{P}_0}{\delta^2} e^{-\frac{1}{\delta}}.$$

Thus  $\mu(x, y)$  is continuous at each point of  $Z$  and  $\mu(x, y) \in C^{(0)}$  in the plane. We prove  $\mu(x, y) \in C^{(\infty)}$  by induction. Suppose

$$\mu(x, y) \in C^{(k)} \quad \text{and} \quad \frac{\partial^k \mu}{\partial x^k}(x_0, y_0) = 0.$$

Then for  $0 < |x - x_0| < \delta$ , a sufficiently small positive number,  $d(x) < \delta$  and

$$\left| \frac{\frac{\partial^k \mu(x, y)}{\partial x^k} - \frac{\partial^k \mu(x_0, y_0)}{\partial x^k}}{x - x_0} \right| < \frac{\bar{P}_k e^{-\frac{1}{\delta}}}{\delta \delta^{2k+1}}.$$

Thus  $\frac{\partial^{k+1} \mu(x_0, y_0)}{\partial x^{k+1}} = 0$ . From the inequality

$$\left| \frac{\partial^{k+1} \mu}{\partial x^{k+1}} \right| < \frac{\bar{P}_{k+1}}{d(x)^{2k+2}} e^{-\frac{1}{d(x)}}$$

it is clear that

$$\lim_{\substack{x=x_0 \\ y=y_0}} \frac{\partial^{k+1} \mu(x, y)}{\partial x^{k+1}} = 0$$

and thus  $\frac{\partial^{k+1} \mu(x, y)}{\partial x^{k+1}}$  exists and is continuous at each point of  $Z$  and

therefore at every point of the plane. Since  $\frac{\partial \mu}{\partial y} \equiv 0$  we have

$$\mu(x, y) \in C^{(k+1)} \quad \text{and} \quad \frac{\partial^{k+1} \mu(x_0, y_0)}{\partial x^{k+1}} = 0.$$

Therefore the induction is complete and  $\mu(x, y) \in C^{(\infty)}$  in the plane.

**THEOREM II.** — *Let  $R$  be simply-connected plane region. Then there exists a real, first order non-singular ordinary differential equation.*

$$(1) \quad \frac{dx}{dt} = f(x, y), \quad \frac{dy}{dt} = g(x, y) \quad (f^2 + g^2 > 0)$$

with  $f, g \in C^{(\infty)}$  in  $R$ , with an integrating factor  $\mu(x, y) \in C^{(\infty)}$ , defining an invariant measure

$$m_\mu(\Lambda) = \iint_{\Lambda} \mu(x, y) dx dy$$

on the plane Lebesgue  $m_L$ -measurable sets, such that

1.  $\mu(x, y) \in C^{(\infty)}$  in  $R$ ;
2.  $\mu(x, y) \geq 0$ ;
3.  $\mu(x, y)$  vanishes on no open set;
4.  $\frac{\partial}{\partial x}(\mu f) + \frac{\partial}{\partial y}(\mu g) \equiv 0$  in  $R$

and the  $m_\mu$ -measure is not complete.

If  $f, g$  are only in  $C^{(1)}$  with  $f^2 + g^2 > 0$  in  $R$ , and if  $\mu(x, y) \in C^{(A)}$  in  $R$  is an integrating factor, that is, satisfies conditions 2, 3, and 4, then the  $m_\mu$ -measure is complete, that is,  $m_\mu(\Lambda) = 0$  if and only if  $m_L(\Lambda) = 0$ .

*Proof.* — We first prove the theorem in the case  $R$  is the entire plane and then use theorem I for the general simply-connected region.

Consider the differential system

$$(7) \quad \frac{dx}{dt} = \mu(x, y), \quad \frac{dy}{dt} = -2 \frac{\partial \mu}{\partial x} y + 1$$

where  $\mu(x, y)$  is the function described above. This system is  $C^{(\infty)}$  and also non-singular since  $\mu(x_0, y_0) = 0$  if and only if  $(x_0, y_0) \in Z$  in which case  $\frac{\partial \mu}{\partial x} = 0$  and  $(-2\mu_x y + 1) = 1$ . Moreover, it is not exact but has an integrating factor of  $\mu(x, y)$  since

$$\frac{\partial}{\partial x}(\mu^2) + \frac{\partial}{\partial y}(-2\mu_x \mu_y + \mu) \equiv 0.$$

To show the  $m_\mu$ -measure is not complete let  $\chi \in Z$  be a non- $m_L$ -measurable set. This is possible since  $m_L(Z) = \infty$ . But

$$m_\mu(Z) = \iint_Z \mu(x, y) dx dy = 0$$

and thus  $\chi$  is a non- $m_\mu$ -measurable subset of a set  $Z$  of  $m_\mu$ -measure zero. Therefore the  $m_\mu$ -measure is not complete.

Next let  $\mu(x, y) \in C^{(A)}$  be any integrating factor of 1 in the plane. Clearly if

$$m_L(\Lambda) = 0$$

then

$$m_{\mu}(\Lambda) = \iint_{\Lambda} \mu \, dx \, dy = 0.$$

Conversely let  $m_{\mu}(\Lambda) = 0$ . Then there exist two sets  $\Lambda_1, \Lambda_2 \in \mathcal{L}$  such that  $m_L(\Lambda_1) = 0$ , and

$$\Lambda_1 \cap \Lambda_2 = 0, \quad \Lambda_1 \cup \Lambda_2 = \Lambda, \quad \text{and} \quad \Lambda_2 \subset Z,$$

the set of zeros of  $\mu(x, y)$ . We shall show that  $m_L(Z) = 0$ .

Since  $Z$  is closed,  $Z \in \mathcal{L}$ . Then for almost all horizontal lines  $h_c: y = y_c$ , the linear measure of  $Z \cap h_c$  exists. If  $Z \cap h_c$  contains only a countable number of points, it has linear measure zero. If  $Z \cap h_c$  contains a non-countable number of points, then since  $\mu(x, y_c)$  is an analytic function of one argument,  $Z \cap h_c = h_c$  and thus has infinite linear measure. However there are only a countable number of lines  $h_c$  on which  $m_L(Z \cap h_c) = \infty$ . For if otherwise, there would be a finite accumulation point  $\bar{y}$  of the corresponding ordinates and then, for each fixed  $x_0$ ,  $\mu(x_0, y)$  is an analytic function of  $y$  and must vanish. Thus  $\mu(x, y) \equiv 0$  which contradicts the hypothesis of the theorem that  $\mu(x, y)$  vanishes on no open set. Therefore on almost all horizontal lines  $m_L(Z \cap h_c) = 0$ . Therefore by Fubini's theorem  $m_L(Z) = 0$ . Since  $\Lambda \subset \Lambda_1 \cup Z$  and  $m_L(\Lambda_1) = 0$ , we have  $m_L(\Lambda) = 0$ .

Next consider any subset  $\Lambda' \subset \Lambda$ . Then  $\Lambda' \in \mathcal{L}$  and  $m_L(\Lambda') = 0$ . Thus  $m_{\mu}(\Lambda') = 0$  and the  $m_{\mu}$ -measure is complete. The theorem is proved in case R is the entire plane.

Again return to the case of a general simply-connected plane region  $R$ . Let  $T: (x, y) \rightarrow (u(x, y), v(x, y))$  be a  $C^{(A)}$ -homeomorphism <sup>(10)</sup> of the plane onto the  $(u, v)$ -region  $R$ . By  $T$  the

<sup>(10)</sup> First map the plane onto the square  $|x^1| < 1, |y^1| < 1$  by

$$x^1 = \frac{2}{\pi} \operatorname{tg}^{-1} x, \quad y^1 = \frac{2}{\pi} \operatorname{tg}^{-1} y.$$

Then use the Riemann conformal mapping theorem.

differential system (7) is carried into a non-singular  $C^{(\infty)}$ -differential system defined in  $R$

$$(8) \quad \frac{du}{dt} = u_x f + u_y g = F(u, v), \quad \frac{dv}{dt} = v_x f + v_y g = G(u, v)$$

where  $F, G \in C^{(\infty)}$  in  $R$  and  $F^2 + G^2 > 0$ . The corresponding integrating factor is

$$M(u, v) = \mu \left| \frac{\partial(x, y)}{\partial(u, v)} \right|.$$

with the corresponding integral  $\psi(x(u, v), y(u, v))$ , as is seen from the following matrix equation

$$(9) \quad \begin{pmatrix} \psi_u \\ \psi_v \end{pmatrix} = \begin{pmatrix} x_u & y_u \\ x_v & y_v \end{pmatrix} \begin{pmatrix} \psi_x \\ \psi_y \end{pmatrix} = \begin{pmatrix} x_u & y_u \\ x_v & y_v \end{pmatrix} \mu \begin{pmatrix} -g \\ f \end{pmatrix}$$

and

$$\begin{pmatrix} -g \\ f \end{pmatrix} = \begin{pmatrix} y_v & -y_u \\ x_v & x_u \end{pmatrix} \begin{pmatrix} -G \\ F \end{pmatrix}.$$

Thus

$$(10) \quad \begin{pmatrix} \psi_u \\ \psi_v \end{pmatrix} = \mu(x(u, v), y(u, v)) \frac{\partial(x, y)}{\partial(u, v)} \begin{pmatrix} -G \\ F \end{pmatrix} = \pm M \begin{pmatrix} -G \\ F \end{pmatrix}.$$

Clearly  $M(u, v) \geq 0$ ,  $M \in C^{(\infty)}$  and vanishes only on the set  $T(Z)$  which contains no interior points. Since  $T$  is a  $C^{(1)}$ -homeomorphism it preserves the set properties measurability and also of having zero measure. Thus  $m_L(T(Z)) > 0$  and thus the  $m_M$ -measure is not complete.

A similar argument proves that, in case  $\mu(x, y) \in C^{(A)}$  is an integrating factor of (1) in  $R$ , a simply-connected region, the resulting invariant measure is complete.

Q. E. D.

COROLLARY. — *Let*

$$(1) \quad \frac{dx}{dt} = f(x, y) \quad \frac{dy}{dt} = g(x, y) \quad (f^2 + g^2 > 0)$$

with  $f, g \in C^{(A)}$  in a simply-connected plane region  $R$  have a principal integral  $\psi(x, y)$  such that

1.  $\psi(x, y) \in C^{(A)}$  in  $R$ ;

2.  $\psi(x, y)$  is constant on no open set of  $R$ ;
3.  $\psi(x, y)$  is constant along each solution curve of (1).

Then the differential system (1) has a complete invariant measure.

*Proof.* — Let

$$\mu(x, y) = \left[ \frac{\psi_x^2 + \psi_y^2}{f^2 + g^2} \right]^{\frac{1}{2}}$$

be the integrating factor corresponding to the integral  $\psi(x, y) \in C^{(A)}$ . Then  $\mu(x, y) \in C^{(A)}$  and, by the theorem, the invariant measure

$$m_\mu(\Lambda) = \iint_\Lambda \mu(x, y) \, dx \, dy$$

is complete.

Q. E. D.

We shall discuss criteria that (1) should have an analytic integral in a later paper.

