

TAKAO KOMATSU

The fractional part of $n\theta + \alpha$ and Beatty sequences

Journal de Théorie des Nombres de Bordeaux, tome 7, n° 2 (1995),
p. 387-406

http://www.numdam.org/item?id=JTNB_1995__7_2_387_0

© Université Bordeaux 1, 1995, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

The fractional part of $n\theta + \phi$ and Beatty sequences

par TAKAO KOMATSU

1. Introduction

Let θ be real and q_n be the denominator of the n -th convergent of the continued fraction expansion of θ . Thus when θ is rational, $\theta = p_N/q_N$ for some integer N . Denote the fractional part of θ by $\{\theta\}$, the floor, that is integer part, of θ by $[\theta]$, and its distance from the nearest integer by $\|\theta\|$.

It is well known that the continued fraction convergents are best approximations to θ in the sense that

$$\min_{0 < x < q_n} \|x\theta\| = \|q_{n-1}\theta\| \quad \text{for } n \geq 1.$$

The corresponding results for one-sided best approximations are readily derived, see for example van Ravenstein [9], and are

$$\min_{0 < x < q_n} \{x\theta\} = \begin{cases} \{q_{n-1}\theta\} & \text{if } n \text{ is odd;} \\ \{(q_n - q_{n-1})\theta\} & \text{if } n \text{ is even,} \end{cases} \quad \text{for } n \geq 2,$$

and

$$\max_{0 < x < q_n} \{x\theta\} = \begin{cases} \{(q_n - q_{n-1})\theta\} & \text{if } n \text{ is odd;} \\ \{q_{n-1}\theta\} & \text{if } n \text{ is even,} \end{cases} \quad \text{for } n \geq 2.$$

The purpose of this paper is to give similar results in the inhomogeneous case $x\theta + \phi$, where θ and ϕ are real numbers, not necessarily irrational, and to apply these results to find the characteristic word

$$f(n; \theta, \phi) = \lfloor (n+1)\theta + \phi \rfloor - \lfloor n\theta + \phi \rfloor \quad \text{for } n = 1, 2, \dots$$

of the Beatty sequence. A Beatty sequence is a sequence of the form $\lfloor n\theta + \phi \rfloor$ for fixed real numbers θ and ϕ . The homogeneous case $\phi = 0$ has been dealt

1991 *Mathematics Subject Classification.* 11B83..

Mots clés : continued fraction, Beatty sequence.

Manuscrit reçu le 23 mai 1994

with extensively (see for example [3], [4], [5], [8], [10], [11], [12], and [2] for additional references). The inhomogeneous case $\phi \neq 0$ is also discussed from several viewpoints (see [5], [7], [8]), but these sources do not necessarily make it easy to get the actual Beatty sequences. The paper [5] is complete on the theory, but only the limited periodic case yields Beatty sequences. The paper [8] provides a very effective description of Beatty sequences, but regrettably, it is now known that the results do not match the facts (see the corrected version, [6]).

2. The minimum of the fractional part of $x\theta + \phi$

We first consider how to find an integer κ satisfying

$$\min_{x \in \mathbb{Z}} \{x\theta + \phi\} = \{\kappa\theta + \phi\}.$$

When both θ and ϕ are rational, the problem is not too difficult. But how should we treat the irrational case? The answer is given in the following theorem. Using this result, we can find κ directly in every case — though two values must be compared when θ is irrational.

THEOREM 1. *Let θ, ϕ be real numbers, and let p_n/q_n denote the n -th convergent of θ . Then for $n \geq 2$,*

$$\min_{0 \leq x < q_n} \{x\theta + \phi\} = \{\kappa_n\theta + \phi\} < \frac{2}{q_n},$$

where $\kappa_n \equiv (-1)^n q_{n-1}(\lfloor q_n \phi \rfloor + t) \pmod{q_n}$ and $t = 0$ or $(-1)^n$.

When $\theta = p_n/q_n$ ($n \geq 2$), $\{x\theta + \phi\}$ has period q_n in x and

$$\min_{x \in \mathbb{Z}} \{x\theta + \phi\} = \min_{0 \leq x < q_n} \{x\theta + \phi\} = \{\kappa_n\theta + \phi\} < \frac{1}{q_n};$$

and $t = 0$.

Proof. Set $\alpha_n = \lfloor q_n \phi \rfloor$. Consider the case $\theta \neq p_n/q_n$. We recall that

$$\frac{p_2}{q_2} < \frac{p_4}{q_4} < \dots < \theta < \dots < \frac{p_3}{q_3} < \frac{p_1}{q_1} \quad \text{and} \quad \left| \theta - \frac{p_n}{q_n} \right| < \frac{1}{q_n q_{n+1}} < \frac{1}{q_n^2}.$$

When n is odd,

$$\frac{p_n x + \alpha_n - 1}{q_n} < -\frac{x}{q_n^2} + x \frac{p_n}{q_n} + \phi < x\theta + \phi < \frac{p_n x + \alpha_n + 1}{q_n}.$$

To minimize $\{x\theta + \phi\}$, we need

$$\begin{aligned} \frac{xp_n + \alpha_n}{q_n} \in \mathbb{Z} & \quad \text{if} \quad x\theta + \phi \geq \frac{xp_n + \alpha_n}{q_n}, \\ \frac{xp_n + \alpha_n - 1}{q_n} \in \mathbb{Z} & \quad \text{if} \quad x\theta + \phi < \frac{xp_n + \alpha_n}{q_n}. \end{aligned}$$

And this is possible since $(p_n, q_n) = 1$ and $0 \leq x < q_n$.

Now, since (q_{n-1}, p_{n-1}) is a solution of

$$p_n x - q_n y = (-1)^{n-1},$$

$x = (-1)^n q_{n-1} \alpha_n$, $y = (-1)^n p_{n-1} \alpha_n$ is a solution of the diophantine equation

$$p_n x - q_n y = -\alpha_n.$$

Thus, the solutions of this equation are

$$\begin{aligned} x &\equiv (-1)^n q_{n-1} [q_n \phi] \pmod{q_n}, \\ y &\equiv (-1)^n p_{n-1} [q_n \phi] \pmod{p_n}. \end{aligned}$$

If we find x with $0 \leq x < q_n$, we get

$$\begin{aligned} \kappa_n &= (-1)^n q_{n-1} [q_n \phi] - (-1)^n q_n [q_{n-1} \phi] \quad \text{or} \\ \kappa_n &= (-1)^n q_{n-1} [q_n \phi] - (-1)^n q_n [q_{n-1} \phi] + q_n. \end{aligned}$$

Therefore, when n is odd and $(xp_n + \alpha_n)/q_n \in \mathbb{Z}$,

$$\kappa_n \equiv -\alpha_n q_{n-1} \pmod{q_n}.$$

In like manner, when n is odd and $(xp_n + \alpha_n - 1)/q_n \in \mathbb{Z}$,

$$\kappa_n \equiv -(\alpha_n - 1)q_{n-1} \pmod{q_n}.$$

Similarly, when n is even,

$$\frac{p_n x + \alpha_n}{q_n} < x\theta + \phi < \frac{x}{q_n^2} + x \frac{p_n}{q_n} + \phi < \frac{p_n x + \alpha_n + 2}{q_n}.$$

Thus,

$$[\kappa_n \theta + \phi] = \frac{p_n \kappa_n + \alpha_n}{q_n} \quad \text{or} \quad \frac{p_n \kappa_n + \alpha_n + 1}{q_n},$$

that is

$$\kappa_n \equiv \alpha_n q_{n-1} \quad \text{or} \quad (\alpha_n + 1) q_{n-1} \pmod{q_n}.$$

When $\theta = p_n/q_n$ for some $n \geq 2$,

$$\frac{xp_n + \alpha_n}{q_n} \leq x\theta + \phi < \frac{xp_n + \alpha_n}{q_n} + \frac{1}{q_n}.$$

Now

$$\kappa_n \equiv (-1)^n \alpha_n q_{n-1} \pmod{q_n}$$

gives

$$(xp_n + \alpha_n)/q_n \in \mathbb{Z}.$$

Remark. If we disallow $x = 0$ (in order that van Ravenstein's result [9] is a particular case), Theorem 1 must be modified. In this case

$$\kappa_n = \left\{ (-1)^{n-1} \frac{q_{n-1}}{q_n} \right\} q_n$$

when $0 \leq \{\phi\} < 1/q_n$.

Corresponding results for the maximum can be obtained symmetrically: One uses the ceiling instead of the floor in the proof above.

COROLLARY. Let θ, ϕ be any real numbers and let p_n/q_n be the n -th convergent of θ . Then for $n \geq 2$

$$\max_{0 \leq x < q_n} \{x\theta + \phi\} = \{\rho_n \theta + \phi\} > 1 - \frac{2}{q_n},$$

where $\rho_n \equiv (-1)^n q_{n-1}([q_n \phi] + t + 1) \pmod{q_n}$ and $t = 0$ or $(-1)^n$.

When $\theta = p_n/q_n$ ($n \geq 2$), $\{x\theta + \phi\}$ has period q_n in x and

$$\max_{x \in \mathbb{Z}} \{x\theta + \phi\} = \max_{0 \leq x < q_n} \{x\theta + \phi\} = \{\rho_n \theta + \phi\} > 1 - \frac{1}{q_n};$$

and $t = 0$.

Remark. In the range $0 < x < q_n$ in which $x = 0$ is disallowed,

$$\rho_n = \left\{ (-1)^n \frac{q_{n-1}}{q_n} \right\} q_n$$

when $1 - 1/q_n \leq \{\phi\} < 1$.

3. THE MINIMUM VALUE OF $\|x\theta + \phi\|$

Combining the results for minimum and maximum, we get the following theorem:

THEOREM 2. *Let θ, ϕ be any real numbers and let p_n/q_n be the n -th convergent of θ . Then for $n \geq 2$*

$$\min_{0 \leq x < q_n} \|x\theta + \phi\| = \|\xi_n\theta + \phi\|,$$

where $\xi_n \equiv (-1)^n q_{n-1}(\lfloor q_n \phi \rfloor + t) \pmod{q_n}$ and $t = -1, 0$ or 1 if n is odd; and $t = 0, 1$ or 2 if n is even.

When $\theta = p_n/q_n$ ($n \geq 2$), $\|x\theta + \phi\|$ has period q_n in x and

$$\min_{x \in \mathbb{Z}} \|x\theta + \phi\| = \min_{0 \leq x < q_n} \|x\theta + \phi\| = \|\xi_n\theta + \phi\|;$$

and $t = 0$ or 1 .

Furthermore, if $a_{n+1} \geq 2$, then for any real θ and ϕ we have $t = 0$ or 1 . And for any real θ and ϕ and for infinitely many n with $n \geq 2$ we have $t = 0$ or 1 .

Proof. The first two assertions are obvious. It suffices to deal with the case when n is odd. Then from the first assertion there are three cases, say

$$\begin{aligned}\xi_n^{(0)} &= \left\{ -\frac{q_{n-1}}{q_n} \alpha_n \right\} q_n, \\ \xi_n^{(-1)} &= \left\{ -\frac{q_{n-1}}{q_n} (\alpha_n - 1) \right\} q_n, \\ \xi_n^{(1)} &= \left\{ -\frac{q_{n-1}}{q_n} (\alpha_n + 1) \right\} q_n.\end{aligned}$$

We will show that $\|\xi_n^{(-1)}\theta + \phi\| \geq \|\xi_n^{(0)}\theta + \phi\|$ if $a_{n+1} \geq 2$. Now

$$\begin{aligned}\|\xi_n^{(-1)}\theta + \phi\| &= \{\xi_n^{(-1)}\theta + \phi\} = (\xi_n^{(-1)}\theta + \phi) - \frac{p_n \xi_n^{(-1)} + \alpha_n - 1}{q_n} \\ &= -\left(\frac{p_n}{q_n} - \theta\right) \xi_n^{(-1)} + \left(\phi - \frac{\alpha_n}{q_n}\right) + \frac{1}{q_n}.\end{aligned}$$

Since

$$-\frac{1}{q_{n+1}} < -\left(\frac{p_n}{q_n} - \theta\right) \xi_n^{(-1)} < 0 \quad \text{and} \quad 0 \leq \phi - \frac{\alpha_n}{q_n} < \frac{1}{q_n},$$

it follows that

$$0 < \frac{1}{q_n} - \frac{1}{q_{n+1}} < \|\xi_n^{(-1)}\theta + \phi\| < \frac{2}{q_n}. \quad (1)$$

On the other hand

$$\begin{aligned} \|\xi_n^{(0)}\theta + \phi\| &= \min[\{\xi_n^{(0)}\theta + \phi\}, 1 - \{\xi_n^{(0)}\theta + \phi\}] \\ &= \begin{cases} -(p_n/q_n - \theta)\xi_n^{(0)} + (\phi - \alpha_n/q_n), & \text{if this value is positive,} \\ (p_n/q_n - \theta)\xi_n^{(0)} - (\phi - \alpha_n/q_n), & \text{otherwise.} \end{cases} \end{aligned}$$

Since

$$-\frac{1}{q_n} < -\left(\frac{p_n}{q_n} - \theta\right)\xi_n^{(0)} + \left(\phi - \frac{\alpha_n}{q_n}\right) < \frac{1}{q_{n+1}},$$

we have

$$0 < \|\xi_n^{(0)}\theta + \phi\| < \frac{1}{q_{n+1}}. \quad (2)$$

From (1) and (2) the fact $q_{n+1} = a_{n+1}q_n + q_{n-1} \geq 2q_n$ now yields the conclusion.

Using Hurwitz's theorem for infinitely many p_n/q_n , that is for infinitely many n

$$\frac{1}{q_n} - \frac{1}{\sqrt{5}q_n} < \|\xi_n^{(-1)}\theta + \phi\| < \frac{2}{q_n} \quad \text{and} \quad 0 < \|\xi_n^{(0)}\theta + \phi\| < \frac{1}{\sqrt{5}q_n},$$

we obtain the last part of Theorem 2.

Remark. In the range $0 < x < q_n$, in which $x = 0$ is disallowed,

$$\xi_n = \left\{ \frac{q_{n-1}}{q_n} \right\} q_n \quad \text{or} \quad \left\{ -\frac{q_{n-1}}{q_n} \right\} q_n$$

when $0 \leq \{\phi\} < 1/q_n$ or $1 - 1/q_n \leq \{\phi\} < 1$.

4. The third possibility in Theorem 2

If θ is irrational and $a_{n+1} = 1$ for some $n(\geq 2)$, there may be the third possibility, namely $t = -1$ when n is odd; $t = 2$ when n is even. In this section we describe θ and ϕ which give this third possibility.

Once again it suffices to consider just odd n . Again set $\alpha_n = \lfloor q_n \phi \rfloor$ and $\xi_n^{(t)} \equiv -q_{n-1}(\alpha_n + t) \pmod{q_n}$, where $t = -1, 0, 1$. Then, the third possibility occurs if

$$\|\xi_n^{(-1)}\theta + \phi\| < \|\xi_n^{(0)}\theta + \phi\|, \|\xi_n^{(1)}\theta + \phi\|. \quad (3)$$

But (3) is equivalent to

$$x\theta + \phi - \frac{p_n x + \alpha_n - 1}{q_n} < \frac{1}{2q_n}$$

or

$$\left(\frac{p_n}{q_n} - \theta\right)x > \frac{1}{2q_n} + \left(\phi - \frac{\alpha_n}{q_n}\right), \quad (4)$$

where $x = \xi_n^{(-1)}$. Since $\phi - \alpha_n/q_n \geq 0$ and $0 \leq x \leq q_n - 1$, θ must satisfy

$$\frac{p_n}{q_n} - \theta > \frac{c}{2q_n(q_n - 1)}, \quad (5)$$

where c is a constant with $1 < c < 2$, which we determine later. Let $a_{n+1} = 1$, a_n and a_{n+2} be large positive integers, say L_n and L_{n+2} , respectively for odd n in the continued fraction expansion of θ . Then

$$p_n - q_n\theta = \frac{1}{\theta_{n+1}q_n + q_{n-1}} = \frac{1}{(1 + 1/(L_{n+2} + \epsilon))q_n + q_{n-1}} > \frac{1}{2(q_n - 1)},$$

where $\theta_{n+1} = [a_{n+1}, a_{n+2}, \dots]$ and $0 < \epsilon < 1$. Thus, this θ satisfies (5). By comparing with (5), we get

$$1 < c < 2 \left(\frac{q_n - 1}{q_{n+1} + q_n/(L_{n+2} + \epsilon)} \right) < 2.$$

Since we may take L_{n+2} as large as we like, the condition on c is

$$1 < c < 2 \left(\frac{q_n - 1}{q_{n+1}} \right) < 2.$$

This condition will be sharpened in (8).

We still have to make $x \equiv -q_{n-1}(\alpha_n - 1) \pmod{q_n}$ large enough and have to make $\phi - \alpha_n/q_n \geq 0$ small enough to satisfy (4). First, we select x , and then may choose $\alpha_n = k_n p_n + 1$, $x = q_n - k_n$. To satisfy the conditions of (4), the value of k_n must be in the range

$$k_n < q_n - \frac{(1 + 2\{q_n \phi\})(q_n - 1)}{c} \leq q_n - \frac{q_n - 1}{c}.$$

Together with the condition for c we get

$$1 \leq k_n < q_n - \frac{q_{n+1}}{2} = \frac{q_n - q_{n-1}}{2} < \frac{q_n}{2}.$$

This condition will be sharpened in (7). We conclude that x is as large as

$$\frac{q_{n+1}}{2} = \frac{q_n + q_{n-1}}{2} < x \leq q_n - 1.$$

Next, we select ϕ . We can say that the condition (3) is

$$\xi_n^{(-1)}\theta + \phi - \frac{p_n\xi_n^{(-1)} + \alpha_n - 1}{q_n} < \frac{p_n\xi_n^{(0)} + \alpha_n}{q_n} - (\xi_n^{(0)}\theta + \phi),$$

where the two fractions are integers. When $\xi_n^{(-1)} = q_n - k_n$ we have $\xi_n^{(0)} = q_n - q_{n-1} - k_n$ since $q_n > L_n q_{n-1}$ and given the conditions on k_n . Recalling the definition of α_n , we get

$$\frac{\alpha_n}{q_n} < \phi < \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta + p_n.$$

This is equivalent to

$$\begin{aligned} \left\{ \frac{\alpha_n}{q_n} \right\} &< \{\phi\} < \left\{ \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta \right\} \\ &= \frac{q_{n-1}\theta - p_{n-1}}{2} + (p_n - q_n\theta) + \{k_n\theta\}, \end{aligned}$$

whence

$$\{k_n\theta\} < 1 - \frac{q_{n-1}\theta - p_{n-1}}{2} - (p_n - q_n\theta) \left(< 1 - \frac{1}{2q_{n+1}} - \frac{1}{q_{n+1} + q_n} \right). \quad (6)$$

To see that, we note

$$\frac{1}{q_n} \leq k_n \left(\frac{p_n}{q_n} - \theta \right) + \frac{1}{q_n} < \frac{q_n - q_{n+1}/2}{q_n q_{n+1}} + \frac{1}{q_n} < \frac{3}{2q_n},$$

whence we have

$$\frac{q_{n-1}\theta - p_{n-1}}{2} + (p_n - q_n\theta) + \{k_n\theta\} = \left\{ \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta + p_n \right\}$$

and

$$\left\{ \frac{\alpha_n}{q_n} \right\} = k_n \left(\frac{p_n}{q_n} - \theta \right) + \frac{1}{q_n} + \{k_n \theta\}.$$

Among these k_n satisfying (6) we choose k_n such that the left hand side of the inequality defining the range of ϕ is less than the right hand side, so

$$k_n \left(\frac{p_n}{q_n} - \theta \right) + \frac{1}{q_n} < \frac{q_{n-1}\theta - p_{n-1}}{2} + (p_n - q_n\theta),$$

that is,

$$1 \leq k_n < q_n - \frac{q_{n-1}}{2} - \frac{1}{2(p_n - q_n\theta)} = \left(1 - \frac{\theta_{n+1}}{2} \right) q_n - q_{n-1} < \frac{q_n}{2} - q_{n-1}. \quad (7)$$

When $n \geq 3$, it follows from (5) that such k_n exist if we take c as

$$1 < \frac{2(q_n - 1)}{2q_n - q_{n-1}} < c < \frac{2(q_n - 1)}{q_{n+1}} < 2. \quad (8)$$

We remark that $2(q_n - 1)/(2q_n - q_{n-1})$ is near to 1 and $2(q_n - 1)/q_{n+1}$ is near to 2 because L_n is large.

When there is a k_n satisfying (7) and

$$\left\{ \frac{k_n p_n + 1}{q_n} \right\} < \left\{ \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta \right\},$$

we can select ϕ as

$$\left\{ \frac{k_n p_n + 1}{q_n} \right\} < \{\phi\} < \left\{ \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta \right\}. \quad (9)$$

The θ and ϕ so selected give the third possibility for odd n . When n is even, instead of (9) we can choose ϕ as

$$\left\{ \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta \right\} < \{\phi\} < \left\{ \frac{k_n p_n - 1}{q_n} \right\}, \quad (9')$$

By these steps, we can always find θ and ϕ to give the third possibility for any odd n , or any even n when $a_{n+1} = 1$, by taking L_n and L_{n+2} large enough so that there is an integer k_n satisfying the conditions (7) and (9) (or (9')). Moreover, it is possible to obtain ϕ giving the third possibility for infinitely many odd (or even) n 's.

We now show the way to find the next interval such that $I_{n+2} \subset I_n$:

Take an increasing sequence of constants $\{\delta_n\}$ with $0 < \delta_n < 1/3$ for all positive integers n . For each odd n (≥ 3) we shall always choose

$$L_{n+2} \geq L_n \geq \max \left(L, \frac{1}{\delta_{n+1} - \delta_n}, \frac{1}{\delta_n - \delta_{n-1}} \right) \quad \text{and} \quad k_n < \delta_n q_n,$$

where L is an integer with $L \geq 8$ and k_n satisfying condition (7). Put

$$\alpha_n = \left\{ \frac{k_n p_n}{q_n} \right\} + \frac{1}{q_n} \quad \text{and} \quad \beta_n = \alpha_n + \frac{1}{mq_n}, \quad (10)$$

where using the notation of the ceiling

$$m = \left\lceil \frac{6L^2 + 8L + 2}{L^2 - 7L - 2} \right\rceil,$$

and we assume that

$$\alpha \geq 2/q_n \quad \text{and} \quad \beta_n \leq 1 - 1/2q_n. \quad (11)$$

Then, comparing with (7),

$$\left\{ \frac{k_n p_n + 1}{q_n} \right\} = \alpha_n < \beta_n < \left\{ \frac{q_{n-1}\theta - p_{n-1}}{2} - (q_n - k_n)\theta \right\}$$

and $\beta_n \leq 1 - 1/2q_n$ yields (6). Let I_n be the interval of (10). Note that if $I_{n+2} \subset I_n$ then I_{n+2} will automatically satisfy condition (11).

We examine the fractional parts of various integer multiples of p_{n+2}/q_{n+2} with an eye to combining them to find an integer k_{n+2} such that

$$\left\{ k_{n+2} \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{1}{q_{n+2}}$$

is in a subinterval of I_n . First,

$$0 < \frac{p_n}{q_n} - \frac{p_{n+2}}{q_{n+2}} < \frac{p_n}{q_n} - \frac{p_{n+1}}{q_{n+1}} = \frac{1}{q_n q_{n+1}},$$

so

$$0 < \left\{ k_n \frac{p_n}{q_n} \right\} - \left\{ k_n \frac{p_{n+2}}{q_{n+2}} \right\} < \frac{\delta_n}{q_{n+1}}. \quad (12)$$

We also have

$$\left\{ q_{n+1} \frac{p_{n+2}}{q_{n+2}} \right\} = q_{n+1} \frac{p_{n+2}}{q_{n+2}} - p_{n+1} = \frac{1}{q_{n+2}}, \quad (13)$$

$$\begin{aligned} \left\{ -q_n \frac{p_{n+2}}{q_{n+2}} \right\} &= \left\{ \left(-q_n \frac{p_{n+2}}{q_{n+2}} + q_n \frac{p_{n+1}}{q_{n+1}} \right) - \left(q_n \frac{p_{n+2}}{q_{n+2}} - p_n \right) \right\} \\ &= -\frac{q_n}{q_{n+1}q_{n+2}} + \frac{1}{q_{n+1}}, \end{aligned}$$

and

$$\left\{ q_{n-1} \frac{p_{n+2}}{q_{n+2}} \right\} = \frac{1}{q_{n+1}} + \frac{1}{q_{n+2}} - \frac{q_n}{q_{n+1}q_{n+2}}, \quad (14)$$

where the third equality can be obtained by adding the first two. From (12) and (14) we have

$$\begin{aligned} -\frac{1}{q_{n+2}} &< \left\{ k_n \frac{p_n}{q_n} \right\} + \frac{1}{q_n} - \left(\left\{ (k_n + q_{n-1}) \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{1}{q_{n+2}} \right) \\ &< \frac{\delta_n}{q_{n+1}} + \frac{1}{q_n} - \frac{1}{q_{n+1}} < \frac{\delta_{n+1}}{q_{n+1}}. \end{aligned}$$

Now let l be the non-negative integer satisfying

$$\frac{l}{q_{n+2}} < \alpha_n - \left\{ (k_n + q_{n-1}) \frac{p_{n+2}}{q_{n+2}} \right\} < \frac{l+1}{q_{n+2}}.$$

Using (13) we get

$$\begin{aligned} \alpha_n &< \left\{ (k_n + q_{n-1}) \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{l}{q_{n+2}} + \frac{1}{q_{n+2}} \\ &= \left\{ (k_n + q_{n-1}) \frac{p_{n+2}}{q_{n+2}} \right\} + \left\{ l q_{n+1} \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{1}{q_{n+2}} \\ &= \left\{ k_{n+2} \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{1}{q_{n+2}} \quad (15) \end{aligned}$$

and

$$\begin{aligned} &\left\{ k_{n+2} \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{1}{q_{n+2}} + \frac{1}{mq_{n+2}} \\ &= \left\{ (k_n + q_{n-1}) \frac{p_{n+2}}{q_{n+2}} \right\} + \frac{l}{q_{n+2}} + \frac{1}{q_{n+2}} + \frac{1}{mq_{n+2}} \\ &< \alpha_n + \frac{1}{q_{n+2}} + \frac{1}{mq_{n+2}} < \beta_n, \quad (16) \end{aligned}$$

where $k_{n+2} = k_n + q_{n-1} + lq_{n+1}$ for some l with

$$0 \leq l \leq \frac{\delta_{n+1}q_{n+2}}{q_{n+1}}.$$

Now

$$k_{n+2} < (l+1)q_{n+1} \leq \delta_{n+2}q_{n+2}.$$

Having chosen L_{n+2} and k_{n+2} we define α_{n+2} and β_{n+2} analogously to α_n and β_n in (10). In view of (15) and (16) we have $I_{n+2} \subset I_n$.

Therefore,

$$\begin{aligned}\theta &= [0, 1, 1, L_3, 1, L_5, \dots, 1, L_{2i-1}, 1, L_{2i+1}, \dots] \\ \phi &\in \dots I_{2i+1} \subset I_{2i-1} \subset \dots \subset I_5 \subset I_3 \subset [0, 1]\end{aligned}$$

give the third possibility for each odd $n = 3, 5, \dots, 2i-1, 2i+1, \dots$.

EXAMPLES.

$$\begin{aligned}\theta &= \frac{\sqrt{35}+4}{19} = [0, 1, 1, 10, 1, 10, \dots] \quad (= 0.52189893\dots) \\ \phi &= 0.573777\dots\end{aligned}$$

give the third possibility for $n = 3, 5, 7, 9, \dots$.

$$\begin{aligned}\theta &= \sqrt{35} - 5 = [0, 1, 10, 1, 10, \dots] \quad (= 5.91607978\dots) \\ \phi &= 0.81678\dots\end{aligned}$$

give the third possibility for $n = 2, 4, 6, 8, \dots$.

5. The sorting of $\{x\theta + \phi\}$

We can solve the sorting problem by using the former results. When θ is irrational, let q_n be the denominator of the n -th convergent of θ , as above. When θ is rational, let $\theta = p_n/q_n$, where p_n and q_n (> 0) are coprime. Let $(\{u_j\theta + \phi\})$, $j = 1, 2, \dots, N$, be the ordered sequence of fractional parts. That is, $\{u_1, u_2, \dots, u_N\} = \{0, 1, 2, \dots, N-1\}$, and $\{u_j\theta + \phi\} < \{u_{j+1}\theta + \phi\}$.

When θ is irrational and $\phi = 0$, the following result is well known:

LEMMA. When $N = q_n$, for $j = 1, 2, \dots, q_n$

$$u_j = \left\{ \frac{(j-1)u_2}{q_n} \right\} q_n,$$

where $u_2 = q_{n-1}$ if n is odd, $u_2 = q_n - q_{n-1}$ if n is even.

Proof. This follows immediately from Lemma 2.1 and Theorem 3.3 of [9].

We will sort the fractional part of $x\theta + \phi$, which includes the result above.

THEOREM 3. When $N = q_n$, for $j = 1, 2, \dots, q_n$

$$u_j = \left\{ (-1)^n \frac{q_{n-1}}{q_n} (\lfloor q_n \phi \rfloor + t - j + 1) \right\} q_n,$$

where $t = 0$ or $(-1)^n$ and is independent of j .

Proof. Set $\alpha_n = \lfloor q_n \phi \rfloor$. When $x\theta + \phi$ is not integer, for each x there exists an integer t satisfying

$$\begin{aligned} \dots < \frac{xp_n + \alpha_n + t - 1}{q_n} < \frac{xp_n + \alpha_n + t}{q_n} < x\theta + \phi \\ < \frac{xp_n + \alpha_n + t + 1}{q_n} < \frac{xp_n + \alpha_n + t + 2}{q_n} < \dots \end{aligned}$$

Let $x^{(t)} = x_n^{(t)}$ with $0 \leq x^{(t)} < q_n$ satisfy

$$x^{(t)} p_n + \alpha_n + t \equiv 0 \pmod{q_n}.$$

Then

$$\{x^{(t)}\theta + \phi\} < \{x^{(t-1)}\theta + \phi\} < \{x^{(t-2)}\theta + \phi\} < \dots$$

From Theorem 1 t must be 0 or $(-1)^n$.

When $x\theta + \phi$ is integer, there exists x satisfying

$$\frac{xp_n + \alpha_n}{q_n} = x\theta + \phi.$$

In particular, $t = 0$ when θ is rational, or if $0 \leq \phi < 1/q_n$.

6. Application to the inhomogeneous Beatty sequence

In this section, we introduce the easiest method for getting the inhomogeneous Beatty sequences in every case. Indeed, one easily obtain the sequence by the use our theorems once one knows the corresponding homogeneous Beatty sequences; see the examples in §7.

Consider the sequence of differences, that is the characteristic sequence

$$f(n; \theta, \phi) = \lfloor (n+1)\theta + \phi \rfloor - \lfloor n\theta + \phi \rfloor \quad n = 1, 2, \dots$$

First, let θ be rational. Then, the following theorem is essential.

THEOREM 4. *Let θ be a rational number, ϕ a real number, and let κ be an integer satisfying*

$$\min_{x \in \mathbb{Z}} \{x\theta + \phi\} = \{\kappa\theta + \phi\}.$$

Then for all integer n we have $f(n + \kappa; \theta, \phi) = f(n; \theta, 0)$.

Proof. Plainly

$$\lfloor (n + \kappa)\theta + \phi \rfloor + \{(n + \kappa)\theta + \phi\} = (n + \kappa)\theta + \phi = n\theta + \lfloor \kappa\theta + \phi \rfloor + \{\kappa\theta + \phi\}.$$

Because $0 \leq \{(n + \kappa)\theta + \phi\} - \{\kappa\theta + \phi\} < 1$, we have

$$\lfloor n\theta \rfloor = \lfloor (n + \kappa)\theta + \phi \rfloor - \lfloor \kappa\theta + \phi \rfloor.$$

Since also $\lfloor (n + 1)\theta \rfloor = \lfloor (n + \kappa + 1)\theta + \phi \rfloor - \lfloor \kappa\theta + \phi \rfloor$, the result follows.

Next, we consider the case θ irrational. The main theorem is Theorem 5, which will follow from Proposition 1 and Proposition 2. The method of proof is similar to that of the rational case. As seen in the assertions, we need not necessarily take an integer x which minimizes the fractional part of $x\theta + \phi$, but it is surely the best way to use such an integer.

PROPOSITION 1. *Let θ be an irrational number with continued fraction expansion*

$$\theta = [a_0, a_1, a_2, \dots],$$

and let ϕ be a real number. For $i = 0, 1, 2, \dots$ denote the convergents of θ by p_i/q_i .

If $a_{i+1}/q_{i+1} \geq \{k\theta + \phi\}$ for a non-negative integer k , then for all integer n with $-q_i + 1 \leq n \leq q_i - 2$ we have $f(n + k; \theta, \phi) = f(n; \theta, 0)$.

This proof depends on the following lemmata:

LEMMA 1. Let θ be an irrational number and ϕ any real number. If $a_{i+1}/q_{i+1} \geq \{k\theta + \phi\}$ for a non-negative integer k , then for every positive odd integer i

$$\{(q_i - q_{i-1})\theta + (k\theta + \phi)\} \geq \{(q_i - q_{i-1})\theta\},$$

$$\{q_{i-1}\theta - (k\theta + \phi)\} \leq \{q_{i-1}\theta\};$$

and for every positive even integer i

$$\{q_{i-1}\theta + (k\theta + \phi)\} \geq \{q_{i-1}\theta\},$$

$$\{(q_i - q_{i-1})\theta - (k\theta + \phi)\} \leq \{(q_i - q_{i-1})\theta\}.$$

Proof. When i is odd,

$$(q_i - q_{i-1})\frac{p_i}{q_i} = p_i - p_{i-1} - \frac{1}{q_i} \quad \text{and} \quad 0 < \frac{p_i}{q_i} - \theta < \frac{1}{q_i q_{i+1}}.$$

Then,

$$\{(q_i - q_{i-1})\theta + (k\theta + \phi)\} = \left\{ -(q_i - q_{i-1}) \left(\frac{p_i}{q_i} - \theta \right) - \frac{1}{q_i} + (k\theta + \phi) \right\},$$

$$\{(q_i - q_{i-1})\theta\} = \left\{ -(q_i - q_{i-1}) \left(\frac{p_i}{q_i} - \theta \right) - \frac{1}{q_i} \right\}.$$

Since

$$1 - \frac{1}{q_i} > \{(q_i - q_{i-1})\theta\} > 1 - \frac{q_i - q_{i-1}}{q_i q_{i+1}} - \frac{1}{q_i},$$

together with

$$(q_i - q_{i-1}) \left(\frac{p_i}{q_i} - \theta \right) + \frac{1}{q_i} > \frac{1}{q_i} \frac{a_{i+1}}{q_{i+1}} \geq \{k\theta + \phi\}$$

the first inequality follows.

When i is even,

$$q_{i-1}\frac{p_i}{q_i} = p_{i-1} - \frac{1}{q_i} \quad \text{and} \quad 0 < \theta - \frac{p_i}{q_i} < \frac{1}{q_i q_{i+1}}.$$

Then,

$$\{q_{i-1}\theta + (k\theta + \phi)\} = \left\{ q_{i-1} \left(\theta - \frac{p_i}{q_i} \right) - \frac{1}{q_i} + (k\theta + \phi) \right\},$$

$$\{q_{i-1}\theta\} = \left\{ q_{i-1} \left(\theta - \frac{p_i}{q_i} \right) - \frac{1}{q_i} \right\}.$$

Since

$$1 - \frac{1}{q_i} < \{q_{i-1}\theta\} < 1 + \frac{q_{i-1}}{q_i q_{i+1}} - \frac{1}{q_i} < 1,$$

together with

$$\frac{1}{q_i} = \frac{a_{i+1}}{q_{i+1}} + \frac{q_{i-1}}{q_i q_{i+1}} \{k\theta + \phi\} + q_{i-1} \left(\theta - \frac{p_i}{q_i} \right),$$

the third inequality is proved.

The second and the fourth inequalities are proved similarly.

LEMMA 2. *If $a_{i+1}/q_{i+1} \geq \{k\theta + \phi\}$ for a non-negative integer k , then for every integer n with $|n| < q_i$,*

$$\{n\theta + (k\theta + \phi)\} \geq \{n\theta\}.$$

Proof. First let $n > 0$. By the result of [9] and the first and the third inequalities in Lemma 1, Lemma 2 holds for n which maximizes the fractional part when $0 < n < q_i$ for integers $i \geq 2$, whence together with the condition $a_{i+1}/q_{i+1} \geq \{k\theta + \phi\}$ the lemma holds for the other positive n .

When $n = 0$, the result is trivial.

Finally, if $n < 0$, by [9] and the second and the fourth inequalities in Lemma 1, for $0 < n' < q_i$,

$$\{n'\theta - (k\theta + \phi)\} \leq \{n'\theta\}.$$

Hence, together with

$$\{-n\theta + (k\theta + \phi)\} = 1 - \{n\theta - (k\theta + \phi)\} \quad \text{and} \quad \{-n\theta\} = 1 - \{n\theta\},$$

we have the claim.

Proof of Proposition 1. If $\{k\theta + \phi\} = 0$, then for any integer n

$$f(n+k; \theta, \phi) = \lfloor (n+1)\theta + \{k\theta + \phi\} \rfloor - \lfloor n\theta + \{k\theta + \phi\} \rfloor = f(n; \theta, 0).$$

By Lemma 2, $\{n\theta + (k\theta + \phi)\} = \{n\theta + \{k\theta + \phi\}\}$ is never located on the the number line to the left of $\{n\theta\}$. Therefore, if $\{k\theta + \phi\} \neq 0$, for each integer n with $|n| < q_i$

$$\{n\theta + (k\theta + \phi)\} - \{n\theta\} = \{k\theta + \phi\}.$$

This gives

$$f(n+k; \theta, \phi) = f(n; \theta, 0).$$

As the inequality of Lemma 2 may not hold for $n = q_i$, there is the possibility that

$$f(q_i - 1 + k; \theta, \phi) \neq f(q_i - 1; \theta, 0).$$

PROPOSITION 2. *Let θ be an irrational number with continued fraction expansion*

$$\theta = [a_0, a_1, a_2, \dots],$$

and let ϕ be a real number. Denote by p_i/q_i ($i = 1, 2, \dots$) the i -th convergent of the continued fraction expansion of θ . Let κ_i be a non-negative integer satisfying

$$\min_{0 \leq x < q_i} \{x\theta + \phi\} = \{\kappa_i\theta + \phi\}.$$

If $l \geq i$ for an integer i with $i \geq 2$, then for all integers n with

$$-\kappa_l \leq n \leq q_i - \kappa_l - 2$$

we have $f(n + \kappa_l; \theta, \phi) = f(n; \theta, 0)$.

Proof. This is similar to the proof of Theorem 4.

Now, we combine Propositions 1 and 2 to get the main theorem for an irrational θ .

THEOREM 5. *Let θ be an irrational number with continued fraction expansion*

$$\theta = [a_0, a_1, a_2, \dots],$$

and ϕ a real number. Denote by p_i/q_i ($i = 1, 2, \dots$) the i -th convergent of the continued fraction expansion of θ . Let κ_i be a non-negative integer satisfying

$$\min_{0 \leq x < q_i} \{x\theta + \phi\} = \{\kappa_i\theta + \phi\}.$$

If $a_{i+1}/q_{i+1} \geq \{\kappa_l\theta + \phi\}$ for $l \geq i \geq 2$, then for all integers n with $0 \leq n \leq q_i - 2 + \kappa_l$,

$$f(n; \theta, \phi) = f(n - \kappa_l; \theta, 0).$$

Remark. We don't have to care for the mysterious condition $a_{i+1}/q_{i+1} \geq \{\kappa_l \theta + \phi\}$ as long as $a_{i+1} \geq 2$ ($i = l-1, l-2$) or $i \leq l-3$. For, always $a_{i+1}/q_{i+1} \geq 2/q_l > \{\kappa_l \theta + \phi\}$ in these cases.

7. Examples

Let $\theta = \sqrt{2}$, $\phi = \sqrt{3}$. Since

$$\sqrt{2} = [1, 2, 2, \dots] \quad \text{and} \quad q_1, q_2, \dots = 2, 5, 12, 29, 70, 169, \dots,$$

by Theorem 1

$$\begin{aligned} \kappa_4 &= \left\{ (-1)^4 \frac{12}{29} [29\sqrt{3}] \right\} \cdot 29 = 20 \quad \text{or} \\ \kappa_4 &= \left\{ (-1)^4 \frac{12}{29} ([29\sqrt{3}] + (-1)^4) \right\} \cdot 29 = 3. \end{aligned}$$

From

$$\{20 \cdot \sqrt{2} + \sqrt{3}\} = 0.016322 \dots \quad \text{and} \quad \{3 \cdot \sqrt{2} + \sqrt{3}\} = 0.9746914 \dots,$$

we get $\kappa_4 = 20$.

Next, since by Theorem 5

$$\frac{2}{70} = 0.0285714 \dots > \left\{ 20 \cdot \sqrt{2} + \sqrt{3} \right\},$$

we have

$$f(n; \sqrt{2}, \sqrt{3}) = f(n-20; \sqrt{2}, 0) \quad \text{for } 0 \leq n \leq 47.$$

For example, according to the method of Fraenkel et. al. [4], as

$$f(1; \sqrt{2}, 0) = 1, \quad f(2; \sqrt{2}, 0) = 2 \quad \text{and} \quad T = (2, 5, 12, 29, 70, \dots),$$

we get

$$\begin{aligned} T_1 &= (12)^\infty, \\ T_2 &= (T_1, 5)^\infty = (12121)^\infty, \\ T_3 &= (T_1, 12)^\infty = (121211212112)^\infty, \\ T_4 &= (T_1, 29)^\infty = (121211212112121211212112121)^\infty. \end{aligned}$$

Furthermore,

$$f(0; \sqrt{2}, 0) = 1, \quad f(-1; \sqrt{2}, 0) = 2 \quad \text{and} \quad f(-n; \theta, 0) = f(n-1; \theta, 0).$$

Thus, $f(n; \sqrt{2}, \sqrt{3})$ for $0 \leq n \leq 47$, that is $f(n; \sqrt{2}, 0)$ for $-20 \leq n \leq 27$, is

$$211212121121211212121212121211212112121121211212112121,$$

where the underline indicates the place $f(0; \sqrt{2}, 0)$.

Let $\theta = 36/25$, $\phi = \sqrt{5}$. Since

$$36/25 = [1, 2, 3, 1, 2] \quad \text{and} \quad q_1, q_2, q_3, q_4 = 2, 7, 9, 25,$$

by Theorem 1

$$\kappa_4 = \left\{ (-1)^4 \frac{9}{25} \lfloor 25\sqrt{5} \rfloor \right\} \cdot 25 = 20.$$

Next, by Theorem 4 we have

$$f(n; 36/25, \sqrt{5}) = f(n-20; 36/25, 0) \quad \text{for all integer } n.$$

For example, according to the method of Fraenkel et. al. [4], as

$$f(1; 36/25, \sqrt{5}) = 1, \quad f(2; 36/25, \sqrt{5}) = 2 \quad \text{and} \quad T = (2, 7, 9, 25),$$

we get

$$\begin{aligned} T_1 &= (12)^\infty, \\ T_2 &= (T_1, 7)^\infty = (1212121)^\infty, \\ T_3 &= (T_1, 9)^\infty = (121212112)^\infty, \\ T_4 &= (T_1, 25)^\infty = (1212121121212121121212121)^\infty. \end{aligned}$$

Therefore, $f(n; 36/25, \sqrt{5})$ for $n \in \mathbb{Z}$ is

$$\dots 12121211212121211212121212121212121211212121212121212121 \dots,$$

where the underline indicates the place $f(0; 36/25, 0)$ and the double-underline indicates the place $f(0; 36/25, \sqrt{5})$.

Of course, we can use the form $36/25 = [1, 2, 3, 1, 1]$ to have the same characteristic sequence.

8. Acknowledgment

I wish to thank Dr P. A. B. Pleasants and Professor A. J. van der Poorten for their advice and assistance in my preparation of the present paper.

REFERENCES

- [1] J. M. BORWEIN and P. B. BORWEIN, *On the generating function of the integer part: $[n\alpha + \gamma]$* , J. Number Theory **43** (1993), 293–318.
- [2] T. C. BROWN, *Descriptions of the characteristic sequence of an irrational*, Canad. Math. Bull. **36** (1993), 15–21.
- [3] L. V. DANILOV, *Some class of transcendental numbers*, Mat. Zametki **12** (1972), 149–154= Math. Notes **12** (1972), 524–527.
- [4] A. S. FRAENKEL, M. MUSHKIN and U. TASSA, *Determination of $[n\theta]$ by its sequence of differences*, Canad. Math. Bull. **21** (1978), 441–446.
- [5] Sh. ITO and S. YASUTOMI, *On continued fractions, substitutions and characteristic sequences $[nx + y] - [(n-1)x + y]$* , Japan. J. Math. **16** (1990), 287–306.
- [6] T. KOMATSU, *A certain power series associated with Beatty sequences*, manuscript.
- [7] T. KOMATSU, *On the characteristic word of the inhomogeneous Beatty sequence*, Bull. Austral. Math. Soc. **51** (1995), 337–351.
- [8] K. NISHIOKA, I. SHIOKAWA and J. TAMURA, *Arithmetical properties of certain power series*, J. Number Theory **42** (1992), 61–87.
- [9] T. VAN RAVENSTEIN, *The three gap theorem (Steinhaus conjecture)*, J. Austral. Math. Soc. (Series A) **45** (1988), 360–370.
- [10] T. VAN RAVENSTEIN, G. WINLEY and K. TOGNETTI, *Characteristics and the three gap theorem*, Fibonacci Quarterly **28** (1990), 204–214.
- [11] K. B. STOLARSKY, *Beatty sequences, continued fractions, and certain shift operators*, Canad. Math. Bull. **19** (1976), 473–482.
- [12] B. A. VENKOV, *Elementary Number Theory*, Wolters-Noordhoff, Groningen, 1970.

Takao KOMATSU
 School of Mathematics, Physics,
 Computing and Electronics
 Macquarie University NSW 2109
 Australia
 e-mail : komatsu@mpce.mq.edu.au