

XIAODONG CAO

WENGUANG ZHAI

**The distribution of square-free numbers of the form  $[n^c]$**

*Journal de Théorie des Nombres de Bordeaux*, tome 10, n° 2 (1998),  
p. 287-299

[http://www.numdam.org/item?id=JTNB\\_1998\\_\\_10\\_2\\_287\\_0](http://www.numdam.org/item?id=JTNB_1998__10_2_287_0)

© Université Bordeaux 1, 1998, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## The distribution of square-free numbers of the form $[n^c]$

par XIAODONG CAO et WENGUANG ZHAI

RÉSUMÉ. Nous montrons que pour  $1 < c < \frac{61}{36} = 1.6944\dots$ , la suite  $[n^c]$  ( $n = 1, 2, \dots$ ) contient une infinité d'entiers sans facteur carré ; cela améliore un résultat antérieur dû à Rieger qui obtenait l'infinitude de ces entiers pour  $1 < c < 1.5$ .

ABSTRACT. It is proved that the sequence  $[n^c]$  ( $n = 1, 2, \dots$ ) contains infinite squarefree integers whenever  $1 < c < \frac{61}{36} = 1.6944\dots$ , which improves Rieger's earlier range  $1 < c < 1.5$ .

### 1. INTRODUCTION

A positive integer  $n$  is called squarefree if it is a product of different primes. Following a paper of Stux [15], Rieger showed in [11] that for all real  $c$  with  $1 < c < 1.5$ , the equation

$$(1.1) \quad S_c(x) = \sum_{\substack{n \leq x \\ [n^c] \text{ squarefree}}} 1 = \frac{6}{\pi^2}x + O(x^{\frac{2c+1}{4}+\varepsilon})$$

holds, which is an immediate consequence of Deshouillers [4]. Here  $[t]$  denotes the fractional part of  $t$  and  $\varepsilon$  is a positive constant small enough. It is an easy exercise to prove that

$$(1.2) \quad S_c(x) = \frac{6}{\pi^2}x + o(x)$$

for  $0 < c \leq 1$ . When  $1 < c < 2$ , one still expects (1.2) to hold, but if  $c = 2$ ,  $[n^c]$  is always a square, so that  $S_c(x) = 0$ .

It is worth remarking that Stux [15] has shown that  $S_c(x)$  tends to infinity for almost all positive real  $1 < c < 2$  (in the sense of Lebesgue measure), however this result provides no specific value of  $c$ .

---

Manuscrit reçu le 26 février 1998.

*Mots-clés.* Square-free number, exponential sum, exponent pair.

This work is supported by National Natural Science Foundation of China (Grant N° 19801021).

The aim of this paper is to further improve Rieger’s range  $1 < c < 1.5$  by the method of exponential sums.

**Basic Proposition.** *Let  $1 < c < 2, \gamma = \frac{1}{c}$ , and  $x > 1$ . Then we have*

$$(1.3) \quad S_c(x) = \frac{6}{\pi^2}x + \Delta_c(x),$$

with

$$(1.4) \quad \Delta_c(x) = \sum_{n \leq x^c} |\mu(n)| (\psi(-(n+1)^\gamma) - \psi(-n^\gamma)) + O(x^{1-c/2}),$$

where  $\psi(t) = t - [t] - 1/2$  and  $\mu(n)$  is the well-known Möbius function.

Using the simple one-dimensional exponent pair, we can prove immediately from the Basic Proposition that

**Corollary.** *Let  $1 < c < 1.625$ , then for  $\varepsilon > 0$*

$$(1.5) \quad S_c(x) = \frac{6}{\pi^2}x + O(x^{\frac{8c+3}{16}+\varepsilon}).$$

Combining Fouvry and Iwaniec’s new method in [5] and Heath-Brown’s new idea in [6], we can prove the following better Theorem .

**Theorem.** *Let  $c$  be a real constant such that  $1 < c < 61/36$ , then*

$$(1.6) \quad S_c(x) = \frac{6}{\pi^2}x + O(x^{\frac{36(c+1)}{97}+\varepsilon}).$$

**Notations.**  $f(x) \ll g(x)$  means that  $f(x) = O(g(x))$ ,  $m \sim M$  means  $c_1M < m \leq c_2M$  for some constants  $c_1, c_2 > 0$ . We also use notations  $L = \log(x), e(x) = \exp(2\pi ix)$  and  $\psi(\theta) = \theta - [\theta] - 1/2$ . To simplify writing logarithms, we will assume that all parameters are bounded by a power of  $x$ . Throughout the paper we allow the constants implied by ‘ $O$ ’ or ‘ $\ll$ ’ to depend on only arbitrarily small positive number  $\varepsilon$  and  $c$  when it occurs.

## 2. PROOFS OF BASIC PROPOSITION AND COROLLARY

**Proof of Basic Proposition.** It is well-known that (see [9])

$$(2.1) \quad \sum_{n \leq x} |\mu(n)| = \frac{6}{\pi^2}x + O(x^{\frac{1}{2}})$$

and

$$(2.2) \quad |\mu(n)| = \sum_{d^2|n} \mu(d).$$

Obviously,  $[n^c]$  is square-free if and only if  $m^\gamma \leq n < (m+1)^\gamma, m$  square-free. Therefore

$$\begin{aligned}
 (2.3) \quad S_c(x) &= \sum_{\substack{n \leq x \\ [n^c] \text{ squarefree}}} 1 = \sum_{\substack{m \leq x^c \\ m \text{ squarefree}}} ([-m^\gamma] - [-(m+1)^\gamma]) + O(1) \\
 &= \sum_{m \leq x^c} |\mu(m)| ((m+1)^\gamma - m^\gamma) + E_{1c}(x) \\
 &= \gamma \sum_{m \leq x^c} |\mu(m)| m^{\gamma-1} + E_{2c}(x),
 \end{aligned}$$

where

$$E_{jc}(x) = \sum_{m \leq x^c} |\mu(m)| (\psi(-(m+1)^\gamma) - \psi(-m^\gamma)) + O(1), \quad j = 1, 2.$$

From (2.3), (2.1) and partial summation we can get the Basic Proposition at once.

The proof of Corollary will need the following two lemmas. Lemma 1 is well-known (see [1]), Lemma 2 is contained in Theorem 18 of Vaaler [16].

**Lemma 1.** *Let  $|g^{(m)}(x)| \sim YX^{1-m}$  for  $1 < X < x \leq 2X$  and  $m = 1, 2, \dots$ . Then*

$$(2.4) \quad \sum_{X < n \leq 2X} e(g(n)) \ll Y^\kappa X^\lambda + Y^{-1}$$

where  $(\kappa, \lambda)$  is any exponent pair.

**Lemma 2.** *Suppose  $J > 1$ . There is a function  $\psi^*(x)$  such that*

- (1)  $\psi^*(x) = \sum_{1 \leq |h| \leq J} \gamma(h)e(hx),$
- (2)  $\gamma(h) \ll \frac{1}{|h|} \quad \text{and} \quad (\gamma(h))' \ll \frac{1}{h^2},$
- (3)  $|\psi^*(x) - \psi(x)| \leq \frac{1}{2(J+1)} \sum_{|h| \leq J} (1 - \frac{|h|}{J})e(hx).$

By Lemma 1 and Lemma 2 we immediately obtain

**Lemma 3.** *Let  $y > 0, X > 1, 0 \leq \sigma < 1, g(n) = (n + \sigma)^\gamma$ . Then*

$$(2.5) \quad \sum_{n \sim X} \psi(yg(n)) \ll y^{\frac{\kappa}{1+\kappa}} X^{\frac{\lambda+\gamma\kappa}{1+\kappa}} + y^{-1} X^{1-\gamma}.$$

*Proof of Corollary.* Taking  $M = x^{\frac{8c-5}{16}}$ , by (2.2) we have

$$(2.6) \quad \sum_{n \leq x^c} |\mu(n)| (\psi(-(n+1)^\gamma) - \psi(-n^\gamma))$$

$$\begin{aligned}
 &= \sum_{d \leq x^{\frac{5}{2}}} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2 n + 1)^\gamma) - \psi(-(d^2 n)^\gamma)) \\
 &= \sum_{d \leq M} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2 n + 1)^\gamma) - \psi(-(d^2 n)^\gamma)) \\
 &\quad + \sum_{M < d \leq x^{\frac{5}{2}}} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2 n + 1)^\gamma) - \psi(-(d^2 n)^\gamma)).
 \end{aligned}$$

□

By Lemma 3 with  $(\kappa, \lambda) = (\frac{2}{7}, \frac{4}{7})$  and simple splitting argument we have

$$\begin{aligned}
 (2.7) \quad &\sum_{d \leq M} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2 n + 1)^\gamma) - \psi(-(d^2 n)^\gamma)) \\
 &\ll L \sum_{d \leq M} \left( (d^{2\gamma})^{\frac{\kappa}{1+\kappa}} (x^c d^{-2})^{\frac{\lambda+\gamma\kappa}{1+\kappa}} + (d^{2\gamma})^{-1} (x^c d^{-2})^{1-\gamma} \right) \\
 &\ll L \sum_{d \leq M} x^{\frac{4c+2}{9}} d^{-\frac{8}{9}} + x^{c-1} L \\
 &\ll x^{\frac{4c+2}{9}} M^{\frac{1}{9}} L + x^{c-1} L \\
 &\ll x^{\frac{8c+3}{16} + \varepsilon}.
 \end{aligned}$$

By Lemma 3 with  $(\kappa, \lambda) = (\frac{1}{6}, \frac{2}{3})$  we have

$$\begin{aligned}
 (2.8) \quad &\sum_{M < d \leq x^{\frac{5}{2}}} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2 n + 1)^\gamma) - \psi(-(d^2 n)^\gamma)) \\
 &\ll L \sum_{M < d \leq x^{\frac{5}{2}}} \left( (d^{2\gamma})^{\frac{\kappa}{1+\kappa}} (x^c d^{-2})^{\frac{\lambda+\gamma\kappa}{1+\kappa}} + (d^{2\gamma})^{-1} (x^c d^{-2})^{1-\gamma} \right) \\
 &\ll L \sum_{M < d \leq x^{\frac{5}{2}}} x^{\frac{4c+1}{7}} d^{-\frac{8}{7}} + x^{c-1} L \\
 &\ll x^{\frac{4c+1}{7}} M^{-\frac{1}{7}} L + x^{c-1} L \\
 &\ll x^{\frac{8c+3}{16} + \varepsilon}.
 \end{aligned}$$

Now the Corollary follows from (2.6), (2.7), (2.8) and the Basic Proposition.

3. SOME LEMMAS

**Lemma 4.** Let  $0 < a < b \leq 2a$ . Let  $f(\frac{z}{a})$  be holomorphic on an open convex set  $\mathbf{I}$  containing the real line segment  $[1, b/a]$ . Assume also that (1)  $|f''(\frac{z}{a})| \leq M$  on  $\mathbf{I}$ , (2)  $f(x)$  is real when  $x$  is real, (3)  $f''(x) \leq -cM$  for some  $c > 0$ . Let  $f'(b) = \alpha$ ,  $f'(a) = \beta$ . For each integer  $v$  in the range  $\alpha \leq v \leq \beta$ , define  $x_v$  by  $f'(x_v) = v$ . Then

$$\sum_{a < n \leq b} e(f(n)) = \sum_{\alpha < v \leq \beta} \frac{e(f(x_v) - vx_v - \frac{1}{8})}{\sqrt{|f''(x_v)|}} + O\left(M^{-\frac{1}{2}} + \log(2 + M(b - a))\right).$$

For the proof of Lemma 4, see Heath-Brown [6], Lemma 6.

**Lemma 5.** Suppose  $A_i, B_j, a_i$  and  $b_j$  are all positive numbers. If  $Q_1$  and  $Q_2$  are real with  $0 < Q_1 \leq Q_2$ , then there exists some  $q$  such that  $Q_1 \leq q \leq Q_2$  and

$$\sum_{i=1}^m A_i q^{a_i} + \sum_{j=1}^n B_j q^{-b_j} \leq 2^{m+n} \left( \sum_{i=1}^m \sum_{j=1}^n (A_i^{b_j} B_j^{a_i})^{\frac{1}{a_i+b_j}} + \sum_{i=1}^m A_i Q_1^{a_i} + \sum_{j=1}^n B_j Q_2^{-b_j} \right).$$

This is Lemma 3 of Srinivasan [14].

**Lemma 6.** Let  $0 < M \leq N < \mu N \leq \lambda M$ , and let  $a_m$  be complex numbers with  $|a_m| \leq 1$ . Then we have

$$\sum_{N < m \leq \mu N} a_m = \frac{1}{2\pi} \int_{-M}^M \left( \sum_{M < m \leq \lambda M} a_m m^{-it} \right) N^{it} (\mu^{it} - 1) t^{-1} dt + O(\log(2 + M)).$$

See Lemma 6 of Fouvry and Iwaniec [5].

**Lemma 7.** Let  $\alpha, \beta_1, \beta_2$  be given real numbers with  $\alpha(\beta_1 - 1)\beta_2 \neq 0$  and  $\alpha \notin N$ . Let  $|a_m| \leq 1, |b_{m_1 m_2}| \leq 1, y \neq 0$  and

$$S = S(M, M_1, M_2) = \sum_{m \sim M} \sum_{m_1 \sim M_1} \sum_{m_2 \sim M_2} a_m b_{m_1 m_2} e(y m^\alpha m_1^{\beta_1} m_2^{\beta_2}).$$

Let  $F = |y| M^\alpha M_1^{\beta_1} M_2^{\beta_2}$ . Then we have

$$SL^{-3} \ll F^{\frac{\kappa}{2(1+\kappa)}} M^{\frac{1+\kappa+\lambda}{2(1+\kappa)}} (M_1 M_2)^{\frac{2+\kappa}{2(1+\kappa)}} + M(M_1 M_2)^{\frac{1}{2}} + M^{\frac{1}{2}} M_1 M_2 + F^{-\frac{1}{2}} M M_1 M_2$$

where  $(\kappa, \lambda)$  is any exponent pair.

Lemma 7 can be proved in the same way as the proof of Theorem 2 of Baker [1]. The idea of the proof is due to Heath-Brown [6].

**Lemma 8.** *Under the conditions of Lemma 7, if we further suppose that  $F \gg M$ , then*

$$SL^{-3} \ll \frac{^{22}\sqrt{(M_1M_2)^{19}M^{13}F^3} + M_1M_2M^{\frac{5}{8}} \sqrt[16]{1 + M^7F^{-4}}}{+ \sqrt[4]{(M_1M_2)^3M^4(1 + FM^{-2})}}.$$

*Proof.* This is Theorem 3 of Liu [8], which is proved essentially by the large sieve inequality developed by Bombieri and Iwaniec [3]. But the term

$$^{32}\sqrt{(M_1M_2)^{29}M^{28}F^{-2} \max(1, F^5M^{-10})}$$

in Liu’s result is superfluous, since

$$\begin{aligned} ^{32}\sqrt{(M_1M_2)^{29}M^{28}F^{-2}} &= \left( \sqrt[16]{F^{-4}M^{17}M_1M_2} \right)^{1/4} \left( M^{5/8}M_1M_2 \right)^{3/8} \\ &\quad \times \left( M^4\sqrt[4]{M_1^3M_2^3} \right)^{3/8}, \\ ^{32}\sqrt{(M_1M_2)^{29}M^{18}F^3} &= \left( \sqrt[22]{F^3M^{13}M_1^{19}M_2^{19}} \right)^{11/16} \\ &\quad \times \left( M^{5/8}M_1M_2 \right)^{5/16} M^{-5/128}. \end{aligned}$$

□

**Lemma 9.** *Let  $\frac{1}{2} < \gamma < 1, M > 1, H > 1, |c(h)| \leq 1, |b(d)| \leq 1$ . Let  $N = N_j = x^cM^{-2}2^{-j}, (j = 1, 2, \dots), F = H(M^2N)^\gamma$  and*

$$S(H, M, N) = \sum_{h \sim H} \sum_{d \sim M} \sum_{x^c d^{-2} 2^{-j} < n \leq x^c d^{-2} 2^{-j+1}} c(h)b(d)e(hd^2n)^\gamma,$$

then

$$\begin{aligned} (3.1) \quad S(H, M, N)L^{-4} &\ll F^{\frac{1}{2}} \left( M^{1+\lambda+\kappa} N^\kappa H^{2+\kappa} \right)^{\frac{1}{2(1+\kappa)}} + (FM)^{\frac{1}{2}} H \\ &\quad + M(NH)^{\frac{1}{2}} + MH + F^{-\frac{1}{2}} MNH. \end{aligned}$$

*Proof.* By Lemma 4 and partial summation we get

$$\begin{aligned} (3.2) \quad S(H, M, N) &\ll F^{-\frac{1}{2}} N \left| \sum_{h \sim H} \sum_{d \sim M} \sum_{\tilde{n} \in I} a(d)b(h, \tilde{n})e\left( (hd^{2\gamma})^{\frac{1}{1-\gamma}} \tilde{n}^{\frac{\gamma}{\gamma-1}} \right) \right| \\ &\quad + F^{-\frac{1}{2}} MNH + LMH, \end{aligned}$$

where  $|a(d)| \leq 1, |b(h, \tilde{n})| \leq 1$  and

$$I = (\gamma 2^{(j-1)(1-\gamma)} x^{1-c} h d^2, \gamma 2^{j(1-\gamma)} x^{1-c} h d^2] \subset [c_1 \frac{F}{N}, c_2 \frac{F}{N}]$$

for some  $c_1, c_2 > 0$ .

Applying Lemma 6 to the variable  $\tilde{n}$ , we obtain that for some  $|b_1(h, \tilde{n})| \leq 1$

(3.3)

$$S(H, M, N)L^{-1} \ll F^{-\frac{1}{2}}N \left| \sum_{h \sim H} \sum_{d \sim M} \sum_{\tilde{n} \sim F/N} a(d)b_1(h, \tilde{n})e\left((hd^{2\gamma})^{\frac{1}{1-\gamma}} \tilde{n}^{\frac{\gamma}{\gamma-1}}\right) \right| + F^{-\frac{1}{2}}MNH + LMH.$$

Now we use Lemma 7 to estimate the above sum, with  $M, H, F/N$  in place of  $M, M_1, M_2$ . This completes the proof of Lemma 9.  $\square$

**Lemma 10.** *Under the conditions of Lemma 9, we have*

$$(3.4) \quad S(H, M, N)L^{-4} \ll \sqrt[22]{F^{11}M^{13}N^3H^{19}} + \sqrt[8]{F^4M^5H^8} + \sqrt[16]{F^4M^{17}H^{16}} + \sqrt[4]{FM^4NH^3} + \sqrt[4]{F^2M^2NH^3} + F^{-\frac{1}{2}}MNH.$$

*Proof.* Applying Lemma 8 to estimate the sum in (3.3), with  $M, H, F/N$  in place of  $M, M_1, M_2$ , we can obtain the bound (3.4) if we notice

$$MH = \left(\sqrt[16]{F^4M^{17}H^{16}}\right)^{2/3} \left(F^{-1/2}MNH\right)^{1/3} (MN^8)^{-1/24}.$$

$\square$

#### 4. PROOF OF THEOREM

Taking  $Y = x^{\frac{c}{3}}$ , we have

$$\begin{aligned} & \sum_{n \leq x^c} |\mu(n)| (\psi(-(n+1)^\gamma) - \psi(-n^\gamma)) \\ &= \sum_{d \leq x^{\frac{c}{2}}} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2n+1)^\gamma) - \psi(-(d^2n)^\gamma)) \\ (4.1) \quad &= \sum_{d \leq Y} \mu(d) \sum_{n \leq x^c d^{-2}} (\psi(-(d^2n+1)^\gamma) - \psi(-(d^2n)^\gamma)) \\ &+ \sum_{n \leq Y} \sum_{Y < d \leq x^{\frac{c}{2}} n^{-\frac{1}{2}}} \mu(d) (\psi(-(d^2n+1)^\gamma) - \psi(-(d^2n)^\gamma)) \\ &= \sum_1 + \sum_2, \text{ say.} \end{aligned}$$

Let  $M > 1$  and  $N = N_j = x^c M^{-2} 2^{-j}$  for  $j = 1, 2, \dots$ . Define

(4.2)

$$T(M, N) = \sum_{M < d \leq 2M} \mu(d) \sum_{\frac{x^c}{d^2 2^j} < n \leq \frac{x^c}{d^2 2^{j-1}}} (\psi(-(d^2n+1)^\gamma) - \psi(-(d^2n)^\gamma)).$$



**Lemma 11.** *We have*

$$(4.3) \quad T(M, N)L^{-5} \ll \left( (M^2N)^{(1+\gamma)(1+\kappa)} M^{\lambda-\kappa} \right)^{\frac{1}{3+2\kappa}} + (M^2N)^{\frac{1+\gamma}{3}} \\ + MN^{\frac{1}{2}} + M^{1-\gamma}N^{1-\frac{\gamma}{2}} + ((M^2N)^{9\gamma}M^2N^4)^{\frac{1}{16}},$$

where  $(\kappa, \lambda)$  is any exponent pair.

*Proof.* By Lemma 2 we get for any  $J > 0$

$$(4.4) \quad T(M, N) \ll MNJ^{-1} + \sum_H \frac{1}{H} |S(H, M, N)| + \sum_H \frac{1}{H} |S_1(H, M, N)|,$$

where  $H = J, \frac{J}{2}, \frac{J}{2^2}, \dots$  and

$$S_1(H, M, N) = \sum_{h \sim H} c(h) \sum_{M < d \leq 2M} b_1(d) \sum_{\frac{x^c}{d^{2^{2^j}}} < n \leq \frac{x^c}{d^{2^{2^j-1}}}} e(-h(d^2n + 1)^\gamma),$$

$|c(h)| \ll 1, b_1(d) \ll 1, S(H, M, N)$  is defined in Lemma 9. □

Write  $\Phi_h(y) = e(h(y^\gamma - (y + 1)^\gamma)) - 1$ . By partial summation and Lemma 1, we have

$$(4.5) \quad \sum_{X < n \leq 2X} e(-h(d^2n + 1)^\gamma) - e(-h(d^2n)^\gamma) \\ = \sum_{X < n \leq 2X} \Phi_h(d^2n)e(-h(d^2n)^\gamma) \\ \ll \max_{X \leq t \leq 2X} |\Phi_h(d^2t)| \left| \sum_{X < n \leq t} e(-h(d^2n)^\gamma) \right| \\ + \int_X^{2X} \left| \frac{\partial \Phi_h(d^2t)}{\partial t} \right| \left| \sum_{X < n \leq t} e(-h(d^2n)^\gamma) \right| dt \\ \ll h(d^2X)^{\gamma-1} \max_{X \leq t \leq 2X} \left| \sum_{X < n \leq t} e(h(d^2n)^\gamma) \right| \\ \ll h(d^2X)^{\gamma-1} \{ (hd^{2\gamma}X^{\gamma-1})^\kappa X^\lambda + (hd^{2\gamma}X^{\gamma-1})^{-1} \} \\ \ll h^{1+\kappa}(d^2X)^{\gamma-1+\kappa\gamma} X^{\lambda-\kappa} + d^{-2}.$$

Here we used  $\Phi_h(d^2t) \ll h(d^2X)^{\gamma-1}$  and  $\frac{\partial \Phi_h(d^2t)}{\partial t} \ll hd^2(d^2X)^{\gamma-2}$  for  $X \leq t \leq 2X$ .

Now take  $(\kappa, \lambda) = (2/7, 4/7)$  in (4.5) we have

$$\begin{aligned}
 (4.6) \quad S_1(H, M, N) &= \sum_{h \sim H} c(h) \sum_{M < d \leq 2M} b_1(d) \sum_{\frac{x^c}{d^{2j}} < n \leq \frac{x^c}{d^{2j-1}}} e(-hd^{2\gamma}n^\gamma) \\
 &+ \sum_{h \sim H} c(h) \sum_{M < d \leq 2M} b_1(d) \sum_{\frac{x^c}{d^{2j}} < n \leq \frac{x^c}{d^{2j-1}}} e(-h(d^2n + 1)^\gamma) - e(-hd^{2\gamma}n^\gamma) \\
 &= \sum_{h \sim H} c(h) \sum_{M < d \leq 2M} b_1(d) \sum_{\frac{x^c}{d^{2j}} < n \leq \frac{x^c}{d^{2j-1}}} e(-hd^{2\gamma}n^\gamma) \\
 &\quad + \sum_h \sum_d O(h^{9/7}(d^2N)^{9\gamma/7-1}N^{2/7} + d^{-2}) \\
 &= \sum_{h \sim H} c(h) \sum_{M < d \leq 2M} b_1(d) \sum_{\frac{x^c}{d^{2j}} < n \leq \frac{x^c}{d^{2j-1}}} e(-hd^{2\gamma}n^\gamma) \\
 &\quad + O\left(H^{\frac{16}{7}}(M^2N)^{\frac{9\gamma}{7}}M^{-1}N^{-\frac{5}{7}}\right).
 \end{aligned}$$

From (4.4)–(4.6) we get

$$(4.7) \quad T(M, N) \ll \frac{MN}{J} + J^{\frac{9}{7}}(M^2N)^{\frac{9\gamma}{7}}M^{-1}N^{-\frac{5}{7}} + \sum_H \frac{1}{H} |S(H, M, N)|,$$

where in  $S(H, M, N)$  the coefficient of  $d$  is  $b(d)$  or  $b_1(d)$ .

Now use Lemma 9 to estimate the above sum and then choose a best  $J \in (0, +\infty)$  by Lemma 5, we obtain the bound (4.3).

**Lemma 12.** *We have*

$$\begin{aligned}
 (4.8) \quad T(M, N)L^{-5} &\ll \sqrt[30]{(M^2N)^{11\gamma}M^{21}N^{11}} + \sqrt[12]{(M^2N)^{4\gamma}M^9N^4} \\
 &+ \sqrt[20]{(M^2N)^{4\gamma}M^{21}N^4} + \sqrt[5]{(M^2N)^{2\gamma}M^3N^2} \\
 &+ \sqrt[4]{(M^2N)^\gamma M^4N} + \sqrt[16]{(M^2N)^{9\gamma}M^2N^4} + M^{1-\gamma}N^{1-\frac{7}{2}}.
 \end{aligned}$$

*Proof.* In the proof of Lemma 11, using Lemma 10 in place of Lemma 9 to estimate the sum in (4.8), we can get Lemma 12. □

**Lemma 13.** *We have*

$$(4.9) \quad \sum_1 \ll x^{\frac{36(c+1)}{97}}L^8.$$

*Proof.* In the proof of Lemma 13, we will use  $M^2N \ll x^c$ .

To estimate  $\sum_1$ , we consider the following cases:

Case (i)  $M \leq x^{\frac{32c+323}{1164}}$ .

Similar to (2.6), by Lemma 3 with  $(\kappa, \lambda) = (\frac{13}{31}, \frac{16}{31})$  we get

$$\begin{aligned}
 (4.10) \quad & \sum_{M < d \leq 2M} \mu(d) \sum_{n \leq \frac{x^c}{d^2}} \psi(-(d^2n + 1)^\gamma) - \psi(-(d^2n)^\gamma) \\
 & \ll Lx^{\frac{16c+13}{44}} \sum_{M < d \leq 2M} d^{-\frac{32}{44}} + x^{c-1}L \\
 & \ll x^{\frac{16c+13}{44}} \left(x^{\frac{32c+323}{1164}}\right)^{\frac{12}{44}} L \\
 & \ll x^{\frac{36(c+1)}{97}} L.
 \end{aligned}$$

Case (ii)  $Z_1 = x^{\frac{32c+323}{1164}} < M \leq x^{\frac{14(c+1)}{97}} = Z$ .

By Lemma 11 with  $(\kappa, \lambda) = (\frac{11}{53}, \frac{33}{53})$ (see [6, pp. 265]) we obtain

$$\begin{aligned}
 (4.11) \quad & \sum_{M < d \leq 2M} \mu(d) \sum_{n \leq \frac{x^c}{d^2}} (\psi(-(d^2n + 1)^\gamma) - \psi(-(d^2n)^\gamma)) \\
 & \ll \left\{ \left(x^c\right)^{\frac{64(1+\gamma)}{53}} M^{\frac{22}{53}} \right\}^{\frac{53}{181}} + (x^c)^{\frac{1+\gamma}{3}} \\
 & \quad + x^{\frac{c}{2}} + (x^c)^{1-\frac{7}{2}} M^{-1} + \left\{ (x^c)^{9\gamma+4} M^{-6} \right\}^{\frac{1}{16}} \} L^7 \\
 & \ll \left\{ \left(x^{64(1+c)} Z^{22}\right)^{\frac{1}{181}} + x^{\frac{1+c}{3}} + x^{\frac{c}{2}} + x^{c-\frac{1}{2}} Z_1^{-1} + \left(x^{9+4c} Z_1^{-6}\right)^{\frac{1}{16}} \right\} L^7 \\
 & \ll x^{\frac{36(c+1)}{97}} L^7.
 \end{aligned}$$

Case (iii)  $Z = x^{\frac{14(c+1)}{97}} < M \leq Y$ .

By Lemma 12 we have

$$\begin{aligned}
 (4.12) \quad & \sum_{M < d \leq 2M} \mu(d) \sum_{n \leq \frac{x^c}{d^2}} (\psi(-(d^2n + 1)^\gamma) - \psi(-(d^2n)^\gamma)) \\
 & \ll \left\{ \sqrt[30]{(x^c)^{11+11\gamma} M^{-1}} + \sqrt[12]{(x^c)^{4+4\gamma} M} + \sqrt[20]{(x^c)^{4+4\gamma} M^{13}} \right. \\
 & \quad \left. + \sqrt[5]{(x^c)^{2+2\gamma} M^{-1}} + \sqrt[4]{(x^c)^{1+\gamma} M^2} + \sqrt[16]{(x^c)^{4+9\gamma} M^{-6}} \right. \\
 & \quad \left. + (x^c)^{1-\frac{7}{2}} M^{-1} \right\} L^7 \\
 & \ll \left\{ \sqrt[30]{x^{11(1+c)} Z^{-1}} + \sqrt[12]{x^{4(1+c)} Y} + \sqrt[20]{x^{4(1+c)} Y^{13}} \right. \\
 & \quad \left. + \sqrt[5]{x^{2(1+c)} Z^{-1}} + \sqrt[4]{x^{1+c} Y^2} + \sqrt[16]{x^{4c+9} Z^{-6}} + x^{c-\frac{1}{2}} Z^{-1} \right\} L^7 \\
 & \ll x^{\frac{36(c+1)}{97}} L^7.
 \end{aligned}$$

Combining (4.10)—(4.12) completes the proof of Lemma 13. □

**Lemma 14.** *We have*

$$(4.13) \quad \sum_2 \ll x^{\max(\frac{7c}{12}, \frac{32c+9}{66}, \frac{5c+3}{12})} L^8.$$

*Proof.* Let  $N \geq 1$ ,  $M = M_j = \frac{x^{\frac{c}{2}}}{N^{\frac{1}{2}2^j}}$  for  $j = 1, 2, \dots$ , and  $F = H(M^2N)^\gamma$ , take  $J = (M^2N)^{\frac{1-\gamma}{2}}$ . By lemma 2 we have

$$(4.14) \quad \begin{aligned} T_1(N, M) &= \sum_{n \sim N} \sum_{\frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^j}} < d \leq \frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^{j-1}}} r(d) (\psi(-(d^2n + 1)^\gamma) - \psi(-(d^2n)^\gamma)) \\ &\ll MNJ^{-1} + \sum_H \frac{1}{H} |T_1(H, N, M)| + \sum_H \frac{1}{H} |T_2(H, N, M)|, \end{aligned}$$

where  $H$  runs through  $J, \frac{J}{2}, \frac{J}{2^2}, \dots$ , and

$$(4.15) \quad T_1(H, N, M) = \sum_{h \sim H} \sum_{n \sim N} \sum_{\frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^j}} < d \leq \frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^{j-1}}} c(h)r(d)e(-h(d^2n)^\gamma),$$

$$T_2(H, N, M) = \sum_{h \sim H} \sum_{n \sim N} \sum_{\frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^j}} < d \leq \frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^{j-1}}} c_1(h)r_1(d)e(-h(d^2n + 1)^\gamma),$$

for some  $c(h) \ll 1, c_1(h) \ll 1, r(d) \ll 1, r_1(d) \ll 1$ .

Since  $e(-h(d^2n + 1)^\gamma) - e(-h(d^2n)^\gamma) \ll |h|(d^2n)^{\gamma-1}$ , we have

$$\begin{aligned} T_2(H, N, M) &= \sum_{h \sim H} \sum_{n \sim N} \sum_{\frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^j}} < d \leq \frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^{j-1}}} c_1(h)r_1(d)e(-h(d^2n)^\gamma) \\ &+ \sum_{h \sim H} \sum_{n \sim N} \sum_{\frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^j}} < d \leq \frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^{j-1}}} c_1(h)r_1(d)\Phi_h(d^2n)e(-h(d^2n)^\gamma) \\ &\ll \sum_{h \sim H} \sum_{n \sim N} \sum_{\frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^j}} < d \leq \frac{x^{\frac{c}{2}}}{n^{\frac{1}{2}2^{j-1}}} c_1(h)r_1(d)e(-h(d^2n)^\gamma) \\ &\quad + MN(M^2N)^{\gamma-1}LH. \end{aligned}$$

So it suffices to bound  $T_1(H, M, N)$ .

Now first applying Lemma 6 to the variable  $d$  and then using Lemma 8 to estimate the sum directly with  $(\alpha, \beta_1, \beta_2) = (2\gamma, \gamma, 1)$ , we get

$$L^{-4}T_1(N, M) \ll \sqrt{(MN)^2(M^2N)^{\gamma-1}} + \sum_H \frac{1}{H} \left\{ {}^{22}\sqrt{(HN)^{19}M^{13}F^3} \right. \\ \left. + HNM^{\frac{5}{8}} \sqrt[16]{1 + M^7F^{-4}} + \sqrt[4]{(HN)^3M^4(1 + FM^{-2})} \right\}.$$

Using the bound  $M^2N \ll x^c$ , one has

$$(4.16) \quad L^{-6}T_1(N, M) \ll \sqrt{(x^c)^\gamma N} + \sqrt[44]{(x^c)^{13+6\gamma}N^{25}} + \sqrt[16]{x^{5c}N^{11}} \\ + \sqrt[32]{x^{17c-8}N^{15}} + \sqrt[4]{x^{2c}N} + \sqrt[4]{x^{1+c}N^2}.$$

Now we note that  $N \leq x^{\frac{5}{3}}$ , by (4.16) and simple splitting argument, the bound (4.13) could be obtained at once.  $\square$

Finally, Combining Lemma 13, Lemma 14, (4.1) and the basic Proposition completes the proof of Theorem .

**Acknowledgement.** The authors would like to thank the referee for his kind and helpful suggestions.

#### REFERENCES

- [1] R.C. Baker, *The square-free divisor problem*. Quart. J. Math. Oxford Ser.(2) **45** (1994), no. 179, 269–277.
- [2] R.C. Baker, G. Harman, J. Rivat, *Primes of the form  $[n^c]$* . J. Number Theory **50** (1995), 261–277.
- [3] E. Bombieri, H. Iwaniec, *On the order of  $\zeta(\frac{1}{2} + it)$* . Ann. Scuola Norm. Sup. Pisa Cl. Sci.(4) **13** (1986), no. 3, 449–472.
- [4] J.-M. Deshouillers, *Sur la répartition des nombres  $[n^c]$  dans les progressions arithmétiques*. C. R. Acad. Sci. Paris Sér. A **277** (1973), 647–650.
- [5] E. Fouvry, H. Iwaniec, *Exponential sums with monomials*. J. Number Theory **33** (1989), 311–333.
- [6] D.R. Heath-Brown, *The Pjateckiĭ-Šapiro prime number theorem*. J. Number Theory **16** (1983), 242–266.
- [7] M.N. Huxley, M. Nair, *Power free values of polynomials III*. Proc. London Math. Soc.(3) **41** (1980), 66–82.
- [8] H.-Q. Liu, *On the number of abelian groups of given order*. Acta Arith. **64** (1993), 285–296.
- [9] H.L. Montgomery, R.C. Vaughan, *The distribution of square-free numbers*. in Recent Progress in Number Theory vol 1, pp. 247–256, Academic Press, London-New York, 1981.
- [10] M. Nair, *Power free values of polynomials*. Mathematika **23** (1976), 159–183.
- [11] G.J. Rieger, *Remark on a paper of Stux concerning squarefree numbers in non-linear sequences*. Pacific J. Math. **78** (1978), 241–242.
- [12] F. Roesler, *Über die Verteilung der Primzahlen in Folgen der Form  $[f(n+x)]$* . Acta Arith. **35** (1979), 117–174.
- [13] F. Roesler, *Squarefree integers in non-linear sequences*. Pacific J. math. **123** (1986), 223–225.
- [14] B.R. Srinivasan, *Lattice problems of many-dimensional hyperboloids II*. Acta Arith. **37** (1963), 173–204.
- [15] I.E. Stux, *Distribution of squarefree integers in non-linear sequences*. Pacific J. Math. **59** (1975), 577–584.

- [16] J.D. Vaaler, *Some extremal problems in Fourier analysis*. Bull. Amer. Math. Soc. **12** (1985), 183–216.

Xiaodong CAO  
Beijing Institute of Petrochemical Technology,  
Daxing, Beijing 102600  
P. R. China

Wenguang ZHAI  
Department of Mathematics  
Shandong Normal University  
Jinan, Shandong 250014  
P.R. China