

JEAN-LOUP MAUCLAIRE

**An almost-sure estimate for the mean of generalized
 Q -multiplicative functions of modulus 1**

Journal de Théorie des Nombres de Bordeaux, tome 12, n° 1 (2000),
p. 1-12

http://www.numdam.org/item?id=JTNB_2000__12_1_1_0

© Université Bordeaux 1, 2000, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

An almost-sure estimate for the mean of generalized Q -multiplicative functions of modulus 1

par JEAN-LOUP MAUCLAIRE

RÉSUMÉ. Soit $Q = (Q_k)_{k \geq 0}$, $Q_0 = 1$, $Q_{k+1} = q_k Q_k$, $q_k \geq 2$, $k \geq 0$, une échelle de Cantor, Z_Q le groupe compact $\prod_{0 \leq j} \mathbb{Z}/q_j \mathbb{Z}$, et μ sa mesure de Haar normalisée. À un élément x of Z_Q écrit $x = \{a_0, a_1, a_2, \dots\}$, $0 \leq a_k \leq q_{k+1} - 1$, $k \geq 0$, on associe la suite $x_k = \sum_{0 \leq j \leq k} a_j Q_j$. On montre que si g est une fonction Q -multiplicative unimodulaire, alors

$$\lim_{x_k \rightarrow x} \left(\frac{1}{x_k} \sum_{n \leq x_k - 1} g(n) - \prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a \leq q_j - 1} g(a Q_j) \right) = 0 \quad \mu\text{-p.s.}$$

ABSTRACT Let $Q = (Q_k)_{k \geq 0}$, $Q_0 = 1$, $Q_{k+1} = q_k Q_k$, $q_k \geq 2$, be a Cantor scale, Z_Q the compact projective limit group of the groups $\mathbb{Z}/Q_k \mathbb{Z}$, identified to $\prod_{0 \leq j \leq k-1} \mathbb{Z}/q_j \mathbb{Z}$, and let μ be its normalized Haar measure. To an element $x = \{a_0, a_1, a_2, \dots\}$, $0 \leq a_k \leq q_{k+1} - 1$, of Z_Q we associate the sequence of integral valued random variables $x_k = \sum_{0 \leq j \leq k} a_j Q_j$. The main result of this article is that, given a complex Q -multiplicative function g of modulus 1, we have

$$\lim_{x_k \rightarrow x} \left(\frac{1}{x_k} \sum_{n \leq x_k - 1} g(n) - \prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g(a Q_j) \right) = 0 \quad \mu\text{-a.e.}$$

1. INTRODUCTION

Let \mathbb{N} be the set of non-negative integers, and let $Q = (Q_k)_{k \geq 0}$, $Q_0 = 1$, be an increasing sequence of positive integers. Using the greedy algorithm, to every element n of \mathbb{N} , one can associate a representation

$$n = \sum_{k=0}^{+\infty} \varepsilon_k(n) Q_k$$

which is unique if for every K ,

$$\sum_{k=0}^{K-1} \varepsilon_k(n) Q_k < Q_K.$$

The simplest examples are the q -adic scale, q integer, $q \geq 2$, and its generalization, the Cantor scale $Q_{k+1} = q_k Q_k$, $Q_0 = 1$, $q_k \geq 2$, $k \geq 0$. In this article, we are concerned with the Cantor scale. For a given integer $n \geq 1$, we denote by $k(n)$ the maximal index k for which $\varepsilon_k(n)$ is different from zero. The integers $\varepsilon_k(n)$ are the digits from n in the basis Q . We recall that if G is an abelian group, a G -valued arithmetical function f such that

$$f(n) = \sum_{k=0}^{k(n)} f(\varepsilon_k(n) Q_k) \quad \text{for } n \geq 1 \quad \text{and} \quad f(0) = 0_G,$$

is called a Q -additive function, an extension of the notion of q -additive function introduced by A. O. Gelfond in the q -adic case [4]. We recall that a real-valued sequence $f(n)$ has an asymptotic distribution if there exists a distribution function F such that for all continuity points x of F , the probability measures defined by $\mu_N(x) = N^{-1} \text{card}\{n \leq N; f(n) \leq x\}$ tends to $F(x)$ as N tends to infinity. In the case of the q -adic scale, necessary and sufficient conditions for the existence of an asymptotic distribution for a real-valued q -additive function have been given by H. Delange in 1972 [3]. J. Coquet [2] considered in 1975 the same kind of problem in cases of Cantor scales and obtained mainly sufficient conditions. In both cases, it appears essential to have information on the difference

$$\left(\frac{1}{x} \sum_{0 \leq n < x} g(n) - \prod_{0 \leq j \leq k(x)} \frac{1}{q_j} \sum_{0 \leq a < q_j} g(a Q_j) \right),$$

where $g(\cdot)$ is any Q -multiplicative function of modulus 1, and more precisely, to get a characterization of

$$(1) \quad \lim_{x \rightarrow +\infty} \left(\frac{1}{x} \sum_{0 \leq n < x} g(n) - \prod_{0 \leq j \leq k(x)} \frac{1}{q_j} \sum_{0 \leq a < q_j} g(a Q_j) \right) = 0.$$

In fact, if the sequence $\{q_j\}_{j \geq 0}$ is bounded, the relation 1 is always true. But if $\{q_j\}_{j \geq 0}$ is unbounded, the situation is quite different. In [1], G. Barat constructs a Q -multiplicative function h with values 1 or -1 such that

$$\lim_{x \rightarrow +\infty} \prod_{0 \leq j \leq k(x)} \frac{1}{q_j} \sum_{0 \leq a < q_j} h(a Q_j)$$

exists and is a positive number while

$$\liminf_{x \rightarrow +\infty} \frac{1}{x} \sum_{n < x} h(n)$$

is less than or equal to zero. This difference is due to the existence of a *first digit phenomenon*, unavoidable for unbounded sequence $\{q_j\}_{0 \leq j}$, as remarked by E. Manstavičius in a recent article [6].

Let \mathbf{Z}_Q denote the group of Q -adic integers, considered as the compact projective limit group of $\mathbf{Z}/Q_k\mathbf{Z}$ and identified to $\prod_{0 \leq k} \mathbf{Z}/q_k\mathbf{Z}$ (see [5], p. 109). The products

$$\prod_{0 \leq j \leq k(n)} \frac{1}{q_j} \sum_{0 \leq a < q_j} g(aQ_j)$$

are clearly related to this group in the following way: an element a of \mathbf{Z}_Q can be written $a = (a_0, a_1, \dots)$, $0 \leq a_k \leq q_k - 1$, $0 \leq k$, and we may identify an element of \mathbf{N} with an element of \mathbf{Z}_Q which has only a finite number of digits different from zero. For all $a = (a_0, a_1, \dots)$ belonging to \mathbf{Z}_Q , we define on \mathbf{Z}_Q the sequence of \mathbf{N} -valued random variables $x_k(\cdot)$ given by $x_k(a) = \sum_{j=0}^k a_j Q_j$, the compact group \mathbf{Z}_Q being endowed with its normalized Haar measure μ , and clearly

$$\prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g(aQ_j) = \int_{\mathbf{Z}_Q} g(x_k) d\mu.$$

In this article, we show roughly speaking that although the relation 1 is not always true according to the example of G. Barat (for unbounded sequence $\{q_j\}_{j \geq 0}$), it is almost surely true for a path chosen at random.

2. RESULTS

2.1. Main theorem.

Theorem 1. *Let g be a unimodular Q -multiplicative function and set*

$$m_j(g) = \frac{1}{q_j} \sum_{0 \leq a < q_j} g(aQ_j).$$

Then, the relation

$$\lim_{k \rightarrow \infty} \left(\frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} g(n) - \prod_{0 \leq j \leq k} m_j(g) \right) = 0$$

holds μ -a.e.

2.2. Consequence of Theorem 1.

Theorem 2. *Let G be a metrizable locally compact abelian group with group law denoted by $+$. Γ denotes the dual group of G endowed with its Haar measure m , and let $f(n)$ be a G -valued Q -additive function. Given a sequence $A(k)$ in G , we denote by F_k^A the distribution of the G -valued*

function defined on \mathbf{Z}_Q by $t \mapsto (f(x_k(t)) - A(k))$, and by $\delta_{(a)}$ the measure consisting in a unit mass at the point a .

The following assertions are equivalent:

i) there exists a sequence $A(k)$ in G and a probability measure ν on G such that the sequence of distributions F_k^A converges vaguely to ν (i.e., $\lim_k \int_G \varphi dF_k^A = \int_G \varphi d\nu$ for all continuous maps $\varphi : G \rightarrow \mathbf{C}$ with compact support);

ii) there exists a sequence $A(k)$ in G and a probability measure ν on G such that μ -a.e., the sequence of random measures $\frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} \delta_{(f(n) - A(k))}$ converges vaguely to ν as k tends to infinity;

iii) the set X of characters g of Γ for which there exists an integer $N(g)$ such that $\prod_{j \geq N(g)} m_j(g) \neq 0$ is not m -negligible.

Remarks. 1) Assertion iii) is always satisfied if G is a compact metrizable group, for X is not empty (it contains the trivial character), and consequently is not m -negligible.

2) Necessary and sufficient conditions for the continuity of ν can be easily found, since ν appears as a convolution of measures on \mathbf{Z}_Q : in fact, the same method as in [7] (p. 84–87), gives that X is a closed and open subgroup. Denoting by H the orthogonal of X and by T_H the canonical projection $G \mapsto G/H$, the measure ν is not continuous if and only if H is finite and

$$\lim_{k \rightarrow \infty} \sum_{0 \leq j \leq k} \frac{1}{q_j} \sum_{\substack{0 \leq a < q_j \\ T_H(f(aQ_j)) \neq 0}} 1 < +\infty.$$

2.3. Proof of Theorem 2. A straightforward adaptation of the argument given in [7] (p 84–87) leads to, *primo* if one of the assumptions i), ii), iii), holds, then, X is a closed and open subgroup of Γ ; and *secundo*, there exists a probability measure ν on G and a G -valued sequence $\{A(k)\}_k$ such that for all g in Γ , the sequence

$$\{\bar{g}(A(k)) \prod_{0 \leq j \leq k} m_j(g \circ f)\}_k$$

tends to $\hat{\nu}(g)$ where $\hat{\nu}$ is the Fourier transform of ν . This is due to the fact that for g in X there exists an $N(g)$ for which the relation $\prod_{j \geq N(g)} m_j(g) \neq 0$,

holds. Hence we get by Theorem 1 that for all g , the sequence

$$\left\{ \frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} g(f(n) - A(k)) \right\}_k$$

converges to $\hat{\nu}(g)$ μ -a.e. Next, we use the Fubini theorem on the measured space $(\Gamma \times \mathbf{Z}_Q, m \otimes \mu)$ in an essential way, by saying that since Γ is countable at infinity and \mathbf{Z}_Q is compact, both of the measures m and μ are σ -finite

and so, μ -a.e., the sequence $\{\frac{1}{x_k} \sum_{n < x_k(\cdot)} g(f(n) - A(k))\}_k$ converges to $\hat{\nu}(g)$ m -a.e.. In order to prove that μ -a.e., the sequence

$$\left\{ \frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} \delta_{(f(n) - A(k))} \right\}_k$$

converges vaguely to ν , it suffices to show that for any real-valued continuous function F defined on G whose support is compact, the sequence

$$\left\{ \frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} F(f(n) - A(k)) \right\}_k$$

converges to $\nu(F)$. This can be done as follows. Take any $\varepsilon > 0$; by assumption on F , there exists V , a symmetric neighborhood of the origin in G , such that for all t in G and all u in V , one has $|F(t+u) - F(t)| \leq \varepsilon$. Denoting by M the Haar measure on G normalized with respect to m , we have

$$\begin{aligned} \left| F(t) - \frac{1}{M(V)} \int_V F(t+u) dM(u) \right| &= \left| \frac{1}{M(V)} \int_V (F(t+u) - F(t)) dM(u) \right| \\ &\leq \frac{1}{M(V)} \int_V |F(t+u) - F(t)| dM(u) \\ &\leq \frac{1}{M(V)} \int_V \varepsilon dM(u) \leq \varepsilon. \end{aligned}$$

The function $F_V(t)$ defined by

$$F_V(t) = \frac{1}{M(V)} \int_V F(t+u) dM(u)$$

is the convolution product of F with the characteristic function of V normalized by the constant $M(V)^{-1}$. Therefore, the Fourier transform \hat{F}_V is integrable and we get

$$\begin{aligned} \frac{1}{x_k} \sum_{n < x_k} F(f(n) - A(k)) &= \frac{1}{x_k} \sum_{n < x_k} \int_{\Gamma} \hat{F}_V(g) \bar{g}(f(n) - A(k)) dm(g) \\ &= \int_{\Gamma} \left(\frac{1}{x_k} \sum_{n < x_k} \hat{F}_V(g) \bar{g}(f(n) - A(k)) \right) dm(g). \end{aligned}$$

By the Lebesgue dominated convergence theorem,

$$\begin{aligned}
& \lim_{k \rightarrow +\infty} \frac{1}{x_k} \sum_{n \leq x_k - 1} F(f(n) - A(k)) \\
&= \lim_{k \rightarrow +\infty} \int_{\Gamma} \left(\frac{1}{x_k} \sum_{n \leq x_k - 1} \hat{F}_V(g) \bar{g}(f(n) - A(k)) \right) dm(g) \\
&= \int_{\Gamma} \hat{F}_V(g) \left(\lim_{k \rightarrow +\infty} \frac{1}{x_k} \sum_{n \leq x_k - 1} \bar{g}(f(n) - A(k)) \right) dm(g) \\
&= \nu(F_V) \mu\text{-a.e.}
\end{aligned}$$

Now, since ν is a probability measure and $|F - F_V| \leq \varepsilon$, the sequence

$$\left\{ \frac{1}{x_k} \sum_{n \leq x_k - 1} F(f(n) - A(k)) - \nu(F) \right\}_k$$

is bounded in modulus by 2ε ; this implies

$$\lim_{k \rightarrow +\infty} \frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} F(f(n) - A(k)) = \nu(F) \mu\text{-a.e.}$$

Therefore, the sequence $\left\{ \frac{1}{x_k(\cdot)} \sum_{n < x_k(\cdot)} \delta_{(f(n) - A(k))} \right\}_k$ converges vaguely μ -a.e. to ν .

3. PROOF OF THEOREM 1

Notation and conventions

Given an arbitrary arithmetical function f , we set

$$S_N(f) = \sum_{0 \leq n \leq N} f(n), \quad M_{N-1}(f) = \sum_{0 \leq n < Q_N} f(n), \quad \widetilde{M}_N(f) = Q^{-1} M_N(f).$$

Notice that we have the identity $M_{N-1}(f) = S_{Q_N-1}(f)$ and for any Q -multiplicative function f ,

$$M_{N-1}(f) = \prod_{0 \leq k < N} m_k(f).$$

By convention, the result of a summation (resp. a product) on an empty set will be 0 (resp. 1).

A - Toolbox.

Proposition 1. *Let g be a Q -multiplicative function of modulus 1 and assume that the sequence $\{\widetilde{M}_k(g)\}_k$ does not tend to 0. Then, there exists a*

sequence $\{\alpha_k\}_{k \geq 0}$ of complex numbers of modulus 1 such that

$$\sum_{k=0}^{+\infty} \frac{1}{q_k} \sum_{a=0}^{q_k-1} |1 - g(aQ_k)\alpha_k|^2 < +\infty.$$

Proof. By our assumption, all the complex numbers $m_j(g)$ are different from zero. Put $\alpha_j = m_j(\bar{g}(\cdot))|m_j(g(\cdot))|^{-1}$ where $\bar{g}(\cdot)$ is the complex conjugate of $g(\cdot)$. The product $\alpha_j m_j(g)$ is equal to $|m_j(g)|$ and the sequence $\{|\widetilde{M}_{k+1}|\}_k$ is convergent. Therefore,

$$\sum_{k=0}^{+\infty} (1 - \alpha_k m_k(g)) < +\infty.$$

From

$$\sum_{k=0}^{+\infty} (1 - \alpha_k m_k(g)) = \sum_{k=0}^{+\infty} \frac{1}{q_k} \sum_{0 \leq a < q_k} (1 - g(aQ_k)\alpha_k)$$

we get a fortiori that the series $\sum_{k=0}^{+\infty} \frac{1}{q_k} \sum_{0 \leq a < q_k} \operatorname{Re}(1 - g(aQ_k)\alpha_k)$ converges and since $|g(aQ_k)\alpha_k| = 1$, we deduce

$$\sum_{k=0}^{+\infty} \frac{1}{q_k} \sum_{0 \leq a < q_k} |1 - g(aQ_k)\alpha_k|^2 < +\infty.$$

□

According to Proposition 1, we introduce the sequence of arithmetical functions $g_k^*(n)$ defined by

$$g_k^*(n) = \prod_{0 \leq j \leq k} g(a_j Q_j) \alpha_j$$

where n is written in base Q as $n = \sum_{j=0}^k a_j Q_j$. This means that if $k(n)$ is the index of the last digit of n which is different from zero, $g_k^*(n)$ is equal to

$$\left(\prod_{0 \leq j \leq k(n)} g(a_j Q_j) \alpha_j \right) \cdot \left(\prod_{k(n) < j \leq k} \alpha_j \right).$$

We extend g_k^* by $g_k^*(x) = g_k^* \circ x_k$ which we also denote g_k^* . Moreover, for simplification, we shall use the notation $g^*(aQ_j) = g(aQ_j)\alpha_j$.

Proposition 2. *If the sequence $\{\widetilde{M}_{k+1}(f)\}_{k \geq 0}$ does not converge to 0, there exists a subset E_∞ of \mathbf{Z}_Q such that $\mu(E_\infty) = 1$ and for every $a = (a_0, a_1, \dots)$ in E_∞ , the sequence $k \mapsto g_k^*(a)$ converges.*

Proof. The sequence of finite groups $\mathbf{Z}/Q_k\mathbf{Z}$, $k \geq 0$, induces a filtration on the μ -measured space \mathbf{Z}_Q , and the complex-valued sequence of adapted functions for this filtration defined by

$$g_k^*(\cdot) \left(\prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g^*(aQ_j) \right)^{-1}$$

is a martingale. Since we have

$$\prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g^*(aQ_j) = \left| \prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g^*(aQ_j) \right|$$

and

$$\left| \prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g^*(aQ_j) \right|^{-1}$$

is bounded, this martingale is bounded and so, it converges μ -a.e. But the sequence

$$\left\{ \prod_{0 \leq j \leq k} \frac{1}{q_j} \sum_{0 \leq a < q_j} g^*(aQ_j) \right\}_k$$

is convergent. Hence we obtain that the sequence $\{g_k^*(\cdot)\}$ converges μ -a.e. \square

Proposition 3. *If the sequence $\{\widetilde{M}_{k+1}(f)\}_k$ does not tend to 0, there exists a subset F_∞ of \mathbf{Z}_Q such that $\mu(F_\infty) = 1$ and for every $x = (a_0(x), a_1(x), \dots)$ in F_∞ , one has*

$$\lim_{\substack{k \rightarrow +\infty \\ a_k(x) \neq 0}} \frac{1}{a_k(x)} \sum_{0 \leq a < a_k(x)} |1 - g^*(aQ_k)|^2 = 0.$$

Proof. Assume that the sequence $\{\widetilde{M}_{k+1}(f)\}_k$ does not tend to 0. Using the same notations as in Proposition 2, we have by Proposition 1

$$\sum_{k=0}^{+\infty} \frac{1}{q_k} \sum_{a=0}^{q_k-1} |1 - g(aQ_k)\alpha_k|^2 < +\infty.$$

Let σ_k be defined by $\sigma_k = \frac{1}{q_k} \sum_{a=0}^{q_k-1} |1 - g(aQ_k)\alpha_k|^2$. For x in \mathbf{Z}_Q , we write $x = (a_0(x), a_1(x), \dots)$, $0 \leq a_k(x) \leq q_k - 1$, $0 \leq k$ and we remark that, on the sequence of the $a_k(x)$ different from 0, one has

$$\begin{aligned} \frac{1}{a_k(x)} \sum_{0 \leq a < a_k(x)} |1 - g^*(aQ_k)|^2 &\leq \frac{1}{a_k(x)} \sum_{0 \leq a < q_k} |1 - g^*(aQ_k)|^2 \\ &\leq \frac{q_k}{a_k(x)} \left(\frac{1}{q_k} \sum_{0 \leq a < q_k} |1 - g^*(aQ_k)|^2 \right) \end{aligned}$$

$$\leq \frac{q_k}{a_k(x)} \sigma_k.$$

Since $\sum_k \sigma_k < +\infty$, it is known that there exists an increasing positive function h tending to infinity when k tends to infinity such that $\sum_k \sigma_k h(k) < +\infty$ and $\prod_{k=0}^{+\infty} (1 - \sigma_k h(k)) > 0$. We consider the set $F(h)$ of points x in \mathbf{Z}_Q such that for all k , the inequality

$$[q_k \sigma_k h(k)] \leq a_k(x) \leq q_k - 1$$

holds, where $[\cdot]$ denotes the integral part function. This set $F(h)$ is closed, and its measure $\mu(F(h))$ is equal to

$$\prod_{k=0}^{+\infty} \frac{1}{q_k} (q_k - [q_k \sigma_k h(k)]),$$

and we have

$$\mu F(h) \geq \prod_{k=0}^{+\infty} \frac{1}{q_k} (q_k - q_k \sigma_k h(k)).$$

Now, we remark that this last product can be written $\prod_{k=0}^{+\infty} (1 - \sigma_k h(k))$ and so, $\mu F(h) \neq 0$. For an x in $F(h)$, we consider the condition $[q_k \sigma_k h(k)] \leq a_k(x) \leq q_k - 1$, when $a_k(x) \neq 0$. If $[q_k \sigma_k h(k)]$ is not 0, then we have

$$\begin{aligned} \frac{q_k}{a_k(x)} \sigma_k &\leq \frac{q_k}{[q_k \sigma_k h(k)]} \sigma_k \leq \frac{q_k \sigma_k h(k)}{[q_k \sigma_k h(k)]} \cdot \frac{q_k}{q_k \sigma_k h(k)} \sigma_k \\ &\leq \frac{q_k \sigma_k h(k)}{[q_k \sigma_k h(k)]} \frac{1}{h(k)} \leq \frac{2}{h(k)} \end{aligned}$$

and in this case, we get $\lim_{k \rightarrow +\infty} \frac{q_k}{a_k(x)} \sigma_k = 0$. The case where $[q_k \sigma_k h(k)] = 0$ remains. We have $0 \leq q_k \sigma_k h(k) < 1$, i.e. $q_k \sigma_k < 1/h(k)$. Hence

$$\frac{q_k}{a_k(x)} \sigma_k \leq \frac{q_k}{1} \sigma_k \leq q_k \sigma_k \leq \frac{1}{h(k)} = o(1), \quad k \rightarrow +\infty.$$

To obtain our result, we remark that the sequence of functions h_r indexed by positive integers r and defined by $h_r(n) = h(n)$ if $n > r$ and $h(n)r^{-1}$ otherwise, satisfies the same requirements as h . Now, the sequence of closed sets $F(h_r)$ is increasing with r and $\lim_{r \rightarrow +\infty} \mu(F(h_r)) = 1$. This gives immediately that F_∞ , the union of the $F(h_r)$, is a measurable set of measure 1. Now, if x belongs to F_∞ , it belongs to some $F(h_r)$ and as a consequence, along the sequence k such that $a_k(x) \neq 0$, we have

$$\begin{aligned} \frac{1}{a_k(x)} \sum_{0 \leq a < a_k(x)} |1 - g^*(aQ_k)|^2 &\leq \frac{q_k}{a_k(x)} \sigma_k \leq q_k \sigma_k \\ &\leq \frac{2}{h_r(k)} = o(1), \quad k \rightarrow +\infty. \end{aligned}$$

□

Proposition 4. *If the sequence $\{\widetilde{M}_{k+1}(f)\}_{k \geq 0}$ converges to zero, then*

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \sum_{0 \leq n < N} g(n) = 0.$$

Proof. This Proposition is due to J.Coquet [2]. □

B- End of the proof

1- First case: *the sequence $\{\widetilde{M}_{k+1}(f)\}_k$ tends to zero.*

From Proposition 4, $\lim_{N \rightarrow +\infty} \sum_{0 \leq n < N-1} f(n) = 0$, and $(x_k)_k$ tends to infinity μ -a.e. due to the fact that $x_k(a)$ is bounded if and only if a has only a finite number of nonzero digits. This means exactly that a is an integer; but $\mu(\mathbb{N}) = 0$.

2- Second case: *the sequence $\{\widetilde{M}_{k+1}(f)\}_k$ does not tend to zero.*

We consider the intersection of the sets E_∞ and F_∞ given in Proposition 2 and Proposition 3 respectively. Notice that $\mu(E_\infty \cap F_\infty) = 1$. Our aim is to prove that for every ξ in $E_\infty \cap F_\infty$

$$\lim_{k \rightarrow +\infty} \left(\frac{1}{x_k(\xi)} \sum_{n < x_k(\xi)} g(n) - \widetilde{M}_{k+1}(g) \right) = 0.$$

The sequence of functions $k \mapsto g_k^*(n)$ and the constants α_j are defined as in Proposition 2. Let ξ be an element of $E_\infty \cap F_\infty$ and denote $x_k(\xi)$ by x_k for short. We have:

$$\begin{aligned} S_{x_k}(g_k^*) &= \left(\sum_{0 \leq a < a_k} g(aQ_k)\alpha_k \right) M_{k-1}(g_{k-1}^*) + (g(a_kQ_k)\alpha_k) S_{x_{k-1}}(g_{k-1}^*) \\ &= \left(\sum_{0 \leq a < a_k} g(aQ_k)\alpha_k \right) M_{k-1}(g_{k-1}^*) + (g_k^*(\xi))(\overline{g_{k-1}^*(\xi)}) S_{x_{k-1}}(g_{k-1}^*), \end{aligned}$$

and by iteration

$$\begin{aligned} S_{x_k}(g_k^*) &= \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} \left(\sum_{0 \leq a < a_j(\xi)} g(aQ_j)\alpha_j \right) \left(\prod_{r=0}^{j-1} \sum_{a=0}^{q_r-1} g_{j-1}^*(aQ_r) \right) \\ &= \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} \left(\sum_{0 \leq a < a_j(\xi)} g(aQ_j)\alpha_j \right) (M_{j-1}(g_{j-1}^*)) \end{aligned}$$

If $a_j(\xi) \neq 0$, the choice of ξ in F_∞ implies

$$\sum_{0 \leq a < a_j(\xi)} g^*(aQ_j) = a_j(\xi)(1 + \varepsilon_j),$$

with $\varepsilon_j = o(1)$ when j tends to infinity. Since g is of modulus 1 and $Q_j^{-1}M_{j-1}(g_{j-1}^*)$ is bounded by 1,

$$\begin{aligned} & \left| S_{x_k}(g_k^*) - \sum_{j=0}^k (g_k^*(\xi)) \overline{g_j^*(\xi)} (a_j(\xi)) (M_{j-1}(g_{j-1}^*)) \right| \\ &= \left| S_{x_k}(g_k^*) - \sum_{j=0}^k (g_k^*(\xi)) \overline{g_j^*(\xi)} \cdot (a_j(\xi)) \cdot ((Q_j^{-1}M_{j-1}(g_{j-1}^*)) Q_j) \right| \\ &\leq \sum_{j=0}^k \varepsilon_j a_j(\xi) Q_j. \end{aligned}$$

Consequently

$$\left| S_{x_k}(g_k^*) - \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} a_j(\xi) M_{j-1}(g_{j-1}^*) \right| = o(x_k), \quad (k \rightarrow +\infty).$$

Since ξ belongs to E_∞ , $\{g_k^*(\xi)\}_k$ converges, and as a consequence, the sequence $\eta_{j,k} = |g_k^*(\xi) \cdot \overline{g_j^*(\xi)} - 1|$ tends to 0 when k and j , $j \leq k$, tend to infinity independently.

This implies

$$\left| \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} a_j(\xi) M_{j-1}(g_{j-1}^*) - \sum_{j=0}^k a_j(\xi) M_{j-1}(g_{j-1}^*) \right| \leq \sum_{j=0}^k \eta_{j,k} a_j(\xi) Q_j,$$

and so, when $k \rightarrow +\infty$,

$$\left| \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} a_j(\xi) M_{j-1}(g_{j-1}^*) - \sum_{j=0}^k a_j(\xi) M_{j-1}(g_{j-1}^*) \right| = o(x_k).$$

Moreover, $Q_j^{-1}M_{j-1}(g_{j-1}^*)$ tends to a limit, say $\widetilde{M}_\infty(g_\infty^*)$. Hence

$$\begin{aligned} & \left| \sum_{j=0}^k a_j(\xi) \cdot M_{j-1}(g_{j-1}^*) - \widetilde{M}_\infty(g_\infty^*) \sum_{j=0}^k a_j(\xi) Q_j \right| \\ &\leq \sum_{j=0}^k |Q_j^{-1}M_{j-1}(g_{j-1}^*) - \widetilde{M}_\infty(g_\infty^*)| \cdot a_j(\xi) Q_j = o(x_k), \quad (k \rightarrow +\infty). \end{aligned}$$

Finally

$$\begin{aligned}
& \left| S_{x_k}(g_k^*) - \widetilde{M}_\infty(g_\infty^*) \sum_{j=0}^k a_j(\xi) Q_j \right| \\
& \leq \left| S_{x_k}(g_k^*) - \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} \cdot a_j(\xi) \cdot M_{j-1}(g_{j-1}^*) \right| \\
& + \left| \sum_{j=0}^k g_k^*(\xi) \overline{g_j^*(\xi)} \cdot a_j(\xi) \cdot M_{j-1}(g_{j-1}^*) - \sum_{j=0}^k a_j(\xi) \cdot M_{j-1}(g_{j-1}^*) \right| \\
& + \left| \sum_{j=0}^k a_j(\xi) \cdot M_{j-1}(g_{j-1}^*) - \widetilde{M}_\infty(g_\infty^*) \sum_{j=0}^k a_j(\xi) Q_j \right|.
\end{aligned}$$

It then follows that

$$S_{x_k}(g_k^*) = \widetilde{M}_\infty(g_\infty^*) \cdot x_k + o(x_k), (k \rightarrow +\infty).$$

To obtain the result, it is enough to notice that from $Q_{k+1}^{-1} M_k(g_k^*) - \widetilde{M}_\infty(g_\infty^*) = o(1)$ we obtain $S_{x_k}(g_k^*) = Q_{k+1}^{-1} M_k(g_k^*) \cdot x_k + o(x_k)$, and replacing g_k^* by its value, we get

$$S_{x_k}(g_k^*) = S_{x_k}(g) \prod_{j=0}^k \alpha_j, \quad M_k(g_k^*) = M_k(g) \prod_{j=0}^k \alpha_j.$$

and this leads to $S_{x_k}(g) - (M_k(g) Q_{k+1}^{-1}) \cdot x_k = o(x_k)$.

REFERENCES

- [1] G. Barat, *Echelles de numération et fonctions arithmétiques associées*. Thèse de doctorat, Université de Provence, Marseille, 1995.
- [2] J. Coquet, *Sur les fonctions S-multiplicatives et S-additives*. Thèse de doctorat de Troisième Cycle, Université Paris-Sud, Orsay, 1975.
- [3] H. Delange, *Sur les fonctions q-additives ou q-multiplicatives*. Acta Arithmetica **21** (1972), 285–298.
- [4] A.O. Gelfond, *Sur les nombres qui ont des propriétés additives ou multiplicatives données*. Acta Arithmetica **13** (1968), 259–265.
- [5] E. Hewit, K.A. Ross, *Abstract harmonic analysis*. Springer-Verlag, 1963.
- [6] E. Manstavičius, *Probabilistic theory of additive functions related to systems of numeration*. New trends in Probability and Statistics Vol.4 (1997), VSP BV & TEV, 412–429.
- [7] J.-L. Maucclair, *Sur la répartition des fonctions q-additives*. J. Théorie des Nombres de Bordeaux **5** (1993), 79–91.

Jean-Loup MAUCLAIRE
 Théorie des Nombres (UMR 7586)
 15 rue du Chevaleret
 F-75013 Paris cedex
 E-mail : mauclai@ccr.jussieu.fr