MORDECHAY B. LEVIN

**On normal lattice configurations and simultaneously
normal numbers**

Journal de Théorie des Nombres de Bordeaux, tome 13, n° 2 (2001),
p. 483-527

[<http://www.numdam.org/item?id=JTNB_2001__13_2_483_0>](http://www.numdam.org/item?id=JTNB_2001__13_2_483_0)

© Université Bordeaux 1, 2001, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

On normal lattice configurations and simultaneously normal numbers

par MORDECHAY B. LEVIN

*Dedicated to Professor Michel Mendès France
in the occasion of his 65th birthday*

RÉSUMÉ. Soient $q, q_1, \dots, q_s \geq 2$ des entiers et $\alpha_1, \alpha_2, \dots$ des nombres réels. Dans cet article, on montre que la borne inférieure de la discrédance de la suite double

$$(\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\})_{m,n=1}^{M N}$$

coïncide (à un facteur logarithmique près) avec la borne inférieure de la discrédance des suites ordinaires $(x_n)_{n=1}^{MN}$ dans un cube de dimension s ($s, M, N = 1, 2, \dots$). Nous calculons aussi une borne inférieure de la discrédance (à un facteur logarithmique près) de la suite $(\{\alpha_1 q_1^n\}, \dots, \{\alpha_s q_s^n\})_{n=1}^N$ (problème de Korobov).

ABSTRACT. Let $q, q_1, \dots, q_s \geq 2$ be integers, and let $\alpha_1, \alpha_2, \dots$ be a sequence of real numbers. In this paper we prove that the lower bound of the discrepancy of the double sequence

$$(\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\})_{m,n=1}^{M N}$$

coincides (up to a logarithmic factor) with the lower bound of the discrepancy of ordinary sequences $(x_n)_{n=1}^{MN}$ in s -dimensional unit cube ($s, M, N = 1, 2, \dots$). We also find a lower bound of the discrepancy (up to a logarithmic factor) of the sequence $(\{\alpha_1 q_1^n\}, \dots, \{\alpha_s q_s^n\})_{n=1}^N$ (Korobov's problem).

1. Introduction.

1.1. A number $\alpha \in (0, 1)$ is said to be *normal* to the base q , if in a q -ary expansion of α , $\alpha = d_1 d_2 \dots$ ($d_i \in \{0, 1, \dots, q-1\}, i = 1, 2, \dots$), each fixed finite block of digits of length k appears with an asymptotic frequency of q^{-k} along the sequence $(d_i)_{i \geq 1}$. Normal numbers were introduced by

Borel (1909). Champernowne (1935) gave an explicit construction of such a number, namely

$$\theta = .1\ 2\ 3\ 4\ 5\ 6\ 7\ 8\ 9\ 10\ 11\ 12\ \dots,$$

obtained by successively concatenating all the natural numbers.

1.1.1. We denote by $\mathbb{N}$ the set of non-negative integers. Let $d, q \geq 2$ be two integers, $\mathbb{N}^d = \{(n_1, \dots, n_d) \mid n_i \in \mathbb{N}, i = 1, \dots, d\}$, $\Delta = \{0, 1, \dots, q-1\}$, $\Omega = \Delta^{\mathbb{N}^d}$.

We shall call $\omega \in \Omega$ a *configuration (lattice configuration)*. A configuration is a function $\omega : \mathbb{N}^d \rightarrow \Delta$. Let $\mathbf{h}, \mathbf{N} \in \mathbb{N}^d$, $\mathbf{h} = (h_1, \dots, h_d)$, $\mathbf{N} = (N_1, \dots, N_d)$. We denote a *rectangular block* by

$$F_{\mathbf{N}} = \{(f_1, \dots, f_d) \in \mathbb{N}^d \mid 0 \leq f_i < N_i, i = 1, \dots, d\},$$

$G = G_{\mathbf{h}}$ is a fixed block of digits $G = \{g_{i_1, \dots, i_d} \in \Delta \mid i_j \in [0, h_j), j = 1, \dots, d\}$.

1.1.2. Definition. A lattice configuration, $\omega \in \Omega$, is said to be *normal (rectangular normal)* if for any $\mathbf{h} \in \mathbb{N}^d$ with $h_1 \cdots h_d \geq 1$ and block of digits $G_{\mathbf{h}}$,

$$(1) \quad \#\{\mathbf{n} \in F_{\mathbf{N}} \mid \omega(\mathbf{n} + \mathbf{i}) = g_{i_1, \dots, i_d} \ \forall \mathbf{i} \in F_{\mathbf{h}}\} - q^{-h_1 \cdots h_d} N_1 \cdots N_d = o(N_1 \cdots N_d),$$

where $\mathbf{i} = (i_1, \dots, i_d)$, and $\max(N_1, \dots, N_d) \rightarrow \infty$.

It is evident that almost every $\omega \in \Omega$ is normal. The constructive proof of the existence of the normal lattice configuration is given in [LS1], [LS2].

Below, to simplify the calculations we consider only the case of $d = 2$.

1.1.3. Let $(\mathbf{x}_n)$ be an infinite sequence of points in an s -dimensional unit cube $[0, 1]^s$; $v = [0, \gamma_1] \times \cdots \times [0, \gamma_s]$ be a box in $[0, 1]^s$; and $A_v(N)$ be a number of indexes $n \in [1, N]$ such that $\mathbf{x}_n$ lies in v . The sequence $(\mathbf{x}_n)$ is said to be *uniformly distributed* in $[0, 1]^s$ if for every box v , $A_v(N)/N \rightarrow \gamma_1 \cdots \gamma_s$. The quantity

$$(2) \quad D(N) = D((\mathbf{x}_n)_{n=1}^N) = D^{(s)}((\mathbf{x}_n)_{n=1}^N) = \sup_{v \in [0, 1]^s} \left| \frac{1}{N} A_v(N) - \gamma_1 \cdots \gamma_s \right|$$

is called the *discrepancy* of $(\mathbf{x}_n)_{n=1}^N$.

It is known (Roth, [Ro]) that for any sequence in $[0, 1]^s$,

$$\overline{\lim}_{N \rightarrow \infty} ND(N) / \log^{s/2} N > 0,$$

and according to the well-known conjecture (see for example [Ni, p. 32, 33]),

$$(3) \quad \overline{\lim}_{N \rightarrow \infty} ND(N) / \log^s N > 0.$$

1.1.4. The double sequence $(\mathbf{u}_{n,m}) \in [0, 1]^s$, $(n, m = 1, 2, \dots)$ is said to be uniformly distributed (Cigler, [Ci]) if

$$D^{(s)}(\{\mathbf{u}_{n,m}\}_{n=1}^N \{m=1\}^M) = o(1), \quad \text{with} \quad \max(N, M) \rightarrow \infty.$$

Kirschenhofer and Tichy [KiTi] investigated double sequences over finite sets (see also references in [DrTi, p.364], [KN, p. 18]).

1.2. It is known (Wall, 1949) that a number α is normal to the base q if and only if the sequence $\{\alpha q^n\}_{n \geq 1}$ is uniformly distributed in $[0, 1)$ (see [KN, p. 70]). It is easy to prove similarly (see Appendix of this paper) the following statement:

Proposition 1. *Let $q \geq 2$ be integer, $d_{m,n} \in \{0, 1, \dots, q-1\}$, $m, n = 1, 2, \dots$. The lattice configuration $(d_{m,n})_{m,n \geq 1}$ is normal if and only if for all $s \geq 1$ the double sequence*

$$(4) \quad (\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\})_{m,n \geq 1}$$

is uniformly distributed in $[0, 1)^s$, where

$$(5) \quad \alpha_m = \sum_{n=1}^{\infty} d_{m,n} / q^n.$$

1.2.1. In [Le3] it was proved explicitly that there exists a normal number α with

$$(6) \quad D(\{\alpha q^n\}_{n=1}^N) = O(N^{-1} \log^2 N), \quad N \rightarrow \infty.$$

The estimate of discrepancy was previously known $O(N^{-2/3} \log^{4/3} N)$ (see [Ko2],[Le2]). According to (3), the estimate (6) cannot be improved essentially.

Our goal is to find a lower bound of discrepancy of the double sequence (4). The main idea of the paper is the using of small discrepancy sequences on the multidimensional unit cube to construct the sequence of reals $(\alpha_m)_{m \geq 1}$ (see (5) and (9)). Here we use a variant of Korobov's s -dimensional sequences ($s = 1, 2, \dots$) with optimal coefficients (see [Ko3]). We provide the following construction of a normal lattice configuration:

1.2.2 Construction. Let p_1, p_2 be distinct primes; $(q, p_1 p_2) = 1$,

$$(7) \quad k_0 = 0, \quad k_1 = [\log_q(p_1 p_2) + 1] p_1, \quad k_i = k_1 p_1^{[\log_{p_1} i]}, \quad i = 2, 3, \dots,$$

$$(8) \quad \begin{aligned} t_0 &= 1, \quad t_i = p_2^{[\log_{p_2} \log_2(i+1)]}, \quad i = 1, 2, \dots, \\ r(j) &= \min_{j \leq t_i p_1^i, \quad i=1,2,\dots} i, \quad j = 1, 2, \dots, \end{aligned}$$

$$(9) \quad \alpha_j = \sum_{i=r(j)}^{\infty} \sum_{n=\delta_i^{(j)} p_1^{i-1} k_{i-1}/k_i}^{p_1^i-1} \sum_{\nu=0}^1 \frac{1}{q^{k_i(2n+\nu)}} \left\{ \frac{a_{j_2, \nu}^{(i)} (p_2^i n + p_1^i j_1)}{p_1^i p_2^i} \right\}_{k_i},$$

where

$$(10) \quad \delta_i^{(j)} = 1 \text{ if } i > r(j); \text{ otherwise } \delta_i^{(j)} = 0, \quad \{x\}_k = [q^k \{x\}]/q^k, \\ j_2 \in \{0, 1, \dots, t_i - 1\}, \quad j_2 \equiv j \pmod{t_i}, \quad j_1 = (j - j_2)/t_i,$$

$$(11) \quad a_{j_2, \nu}^{(i)} \in \{0, 1, \dots, p_1^i p_2^i - 1\}, \quad \nu = 0, 1, \quad j = 1, 2, \dots.$$

Theorem 1. *There exist integers $a_{r, \nu}^{(m)}$ ($m, r = 1, 2, \dots, \nu = 0, 1$) satisfying (11), such that for all $s, N, M \geq 1$ we have*

$$(12) \quad D\left(\left(\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\}\right)_{1 \leq n \leq N, 0 \leq m < M}\right) \\ = O((MN)^{-1} (\log MN)^{2s+4} \log^2 \log MN)$$

with $\max(M, N) \rightarrow \infty$, and the constant implied by O only depends on s .

We note that according to (3), the estimate (12) cannot be improved by more than the power of the logarithmic multiplier.

Corollary. *Let $s, q \geq 2$. There exist numbers $\alpha_1, \dots, \alpha_s$ (simultaneously normal to the base q) such that*

$$D\left(\left(\{\alpha_1 q^n\}, \dots, \{\alpha_s q^n\}\right)_{n=1}^N\right) = O(N^{-1} \log^{2s+4+\epsilon} N).$$

The discrepancy estimate was previously known as $O(N^{-1/s})$ [Ko1] and $O(N^{-2/3} \log^{s+2} N)$ [Le2].

1.3. Let $s, q_1, \dots, q_s \geq 2$ be integers. Numbers $\alpha_1, \dots, \alpha_s$ are said to be *simultaneously normal to the base $(q_1, \dots, q_s)$* [Ko1], [Ko3] if the sequence

$$(13) \quad (\{\alpha_1 q_1^n\}, \dots, \{\alpha_s q_s^n\})_{n \geq 1}$$

is uniformly distributed in $[0, 1)^s$.

In [Ko1], Korobov obtained the first examples of simultaneously normal numbers using normal periodic systems, completely uniformly distributed sequences, and estimates of trigonometric sums with exponential functions (see also [Ko3]). In [Ko1], Korobov constructed simultaneously normal numbers with

$$D\left(\left(\{\alpha_1 q_1^n\}, \dots, \{\alpha_s q_s^n\}\right)_{n=1}^N\right) = O(N^{-1/s})$$

and posed the problem of finding simultaneously normal numbers with a maximum decay of the discrepancy of the sequence (13). In [Le1], simultaneously normal numbers with $D_N = O(N^{-1/2} \log^{s+3/2} N)$ were constructed. Here we find simultaneously normal numbers with the discrepancy estimate $O(N^{-1} \log^{2s+2} N)$. We note that according to (3), this estimate cannot be improved by more than the power of the logarithmic multiplier.

1.3.1 Construction. Let p be prime; $(q_i, p) = 1$, $i = 1, \dots, s$;

$$(14) \quad \begin{aligned} k_1 &= p \max_{1 \leq i \leq s} [\log_{q_i} p + 1], \quad k_m = k_1 p^{\lfloor \log_p m \rfloor}, \quad n_1 = 0, \\ n_m &= n_{m-1} + 2k_{m-1} p^{m-1}, \quad m = 2, 3, \dots \end{aligned}$$

$$(15) \quad \alpha_i = \sum_{m=1}^{\infty} \sum_{n=0}^{p^m-1} \sum_{\nu=0}^1 \frac{1}{q_i^{n_m+k_m(2n+\nu)}} \left\{ \frac{a_{i,\nu}^{(m)} n}{p^m} \right\}_{k_m, i},$$

where $\{x\}_{k,i} = [\{x\}q_i^k]/q_i^k$, $i = 1, \dots, s$, $k = 1, 2, \dots$.

Theorem 2. Let $s \geq 2$. There exist integers $a_{i,\nu}^{(m)} \in \{0, 1, \dots, p^m - 1\}$ ($i = 1, \dots, s$; $\nu = 0, 1$; $m = 1, 2, \dots$) such that

$$D((\{\alpha_1 q_1^n\}, \dots, \{\alpha_s q_s^n\})_{n=1}^N) = O(N^{-1} \log^{2s+2} N), \quad N \rightarrow \infty.$$

We prove this theorem in Section 4. Theorem 1 is proved in Section 3. Section 2 contains auxiliary results.

2. Auxiliary results

First, some further notation is necessary. For integers $d \geq 1$ and $l \geq 2$, let $C_d(l)$ be the set of all nonzero lattice points $(h_1, \dots, h_d) \in \mathbb{Z}^d$ with $-l/2 < h_j \leq l/2$ for $1 \leq j \leq d$; $C(l) = \mathbb{Z} \cap (-l/2, l/2]$. Define

$$r(h, l) = \begin{cases} l \sin(\pi|h|/l) & \text{for } h \in C_1(l), \\ 1 & \text{for } h = 0, \end{cases}$$

and

$$(16) \quad r(\mathbf{h}, l) = \prod_{j=1}^d r(h_j, l)$$

for $\mathbf{h} = (h_1, \dots, h_d) \in C_d(l)$. For real t , the abbreviation $e(t) = e^{2\pi\sqrt{-1}t}$ is used. Subsequently, four known results are stated, which follow from [Ko3, Lemma 2], [Ei, Lemma 3], [Ko3, p.13, Ni, p. 35] and [Ni, Theorem 3.10], respectively.

Lemma 1. Let $p \geq 2, a$ be integers,

$$\delta_p(a) = \begin{cases} 1 & \text{if } a \equiv 0 \pmod{p}, \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\delta_p(a) = \frac{1}{p} \sum_{n=0}^{p-1} e(an/p).$$

Lemma 2. Let $q \geq 2$ be an integer. Then

$$\sum_{\substack{\mathbf{h} \in C_d(q) \\ \mathbf{h} \equiv \mathbf{0} \pmod{v}}} \frac{1}{r(\mathbf{h}, q)} < \frac{1}{v} \left(\frac{2}{\pi} \log q + \frac{7}{5} \right)^d$$

for any divisor v of q with $1 \leq v < q$.

Lemma 3. Let A, B, T be integers, $1 \leq B \leq T$. Then

$$\left| \sum_{n \in [A, A+B)} e(t_n) \right| \leq \sum_{h_0 \in (-T/2, T/2]} \frac{1}{r(h_0, T)} \left| \sum_{n=A}^{A+T-1} e\left(t_n + \frac{nh_0}{T}\right) \right|.$$

According to [Ni, p. 35] $|1/T \sum_{n \in [A, A+B)} e(nh_0/T)| \leq 1/r(h_0, T)$. Now the proof of Lemma 3 repeats that of [Ko3, p.13]. $\square$

Applying this lemma twice, we get

Corollary 1. Let $r \geq 1, M, M_1, N, N_1$ be integers, $M \in [1, p_2^r]$, $N \in [1, p_1^r]$. Then

$$\begin{aligned} \left| \sum_{n=N_1}^{N_1+N-1} \sum_{m=M_1}^{M_1+M-1} e(t_{nm}) \right| &\leq \sum_{h_{-1} \in C(p_1^r)} \sum_{h_{-2} \in C(p_2^r)} \frac{1}{r(h_{-1}, p_1^r) r(h_{-2}, p_2^r)} \\ &\times \left| \sum_{n=N_1}^{N_1+p_1^r-1} \sum_{m=M_1}^{M_1+p_2^r-1} e\left(t_{nm} + \frac{nh_{-1}}{p_1^r} + \frac{mh_{-2}}{p_2^r}\right) \right|. \end{aligned}$$

Lemma 4. Let $N \geq 1$ and $P \geq 2$ be integers. Let $\mathbf{t}_n = \mathbf{y}_n/P \in [0, 1)^d$ with $\mathbf{y}_n \in \{0, 1, \dots, P-1\}^d$ for $0 \leq n < N$. Then the discrepancy of the points $\mathbf{t}_0, \mathbf{t}_1, \dots, \mathbf{t}_{N-1}$ satisfies

$$D(\mathbf{t}_0, \mathbf{t}_1, \dots, \mathbf{t}_{N-1}) \leq \frac{d}{P} + \frac{1}{N} \sum_{\mathbf{h} \in C_d(P)} \frac{1}{r(\mathbf{h}, P)} \left| \sum_{n=0}^{N-1} e(\mathbf{h} \cdot \mathbf{t}_n) \right|.$$

Corollary 2. Let $T \geq N \geq 1$ and $P \geq 2$ be integers, $\mathbf{t}_n = \mathbf{y}_n/P \in [0, 1)^d$ with $\mathbf{y}_n \in \{0, 1, \dots, P-1\}^d$ for $0 \leq n < N$. Then

$$(17) \quad D(\mathbf{t}_0, \mathbf{t}_1, \dots, \mathbf{t}_{N-1}) \leq \frac{d}{P} + \frac{T}{N} \tilde{D}_T((\mathbf{t}_n)_{n \geq 0}),$$

where

$$\tilde{D}_T((\mathbf{t}_n)_{n \geq 0}) = \frac{1}{T} \sum_{\mathbf{h} \in C_d(P)} \sum_{h_0 \in (-T/2, T/2]} \frac{1}{r(\mathbf{h}, P)r(h_0, T)} \left| \sum_{n=0}^{T-1} e(\mathbf{h} \cdot \mathbf{t}_n + \frac{nh_0}{T}) \right|.$$

Corollary 3. Let $p_1^r \geq N \geq 1$, $p_2^r \geq M \geq 1$, and $p_1, p_2 \geq 2$ be integers, $\mathbf{t}_{n,m} = \mathbf{y}_{n,m}/p_1^r p_2^r \in [0, 1]^d$, with $\mathbf{y}_{n,m} \in \{0, 1, \dots, p_1^r p_2^r - 1\}^d$ for $n, m = 0, 1, \dots$. Then

$$(18) \quad MND^{(d)}((\mathbf{t}_{nm})_{M_1 \leq m < M_1+M, N_1 \leq n < N_1+N}) \leq \frac{dMN}{p_1^r p_2^r} + \hat{D}_r^{(d+2)}(\mathbf{t}_{nm}),$$

where

$$\begin{aligned} \hat{D}_r^{(d+2)}(\mathbf{t}_{nm}) = & \sum_{\mathbf{h} \in C_d(p_1^r p_2^r)} \sum_{h_{-1} \in C(p_1^r)} \sum_{h_{-2} \in C(p_2^r)} \frac{1}{r(\mathbf{h}, p_1^r p_2^r) r(h_{-1}, p_1^r) r(h_{-2}, p_2^r)} \\ & \times \left| \sum_{n=0}^{p_1^r-1} \sum_{m=0}^{p_2^r-1} e(\mathbf{h} \cdot \mathbf{t}_{nm} + \frac{nh_{-1}}{p_1^r} + \frac{mh_{-2}}{p_2^r}) \right|. \end{aligned}$$

Lemma 5. Let $\mathbf{x}_n \in [0, 1]^s$, $n = 1, 2, \dots$, $q_1, \dots, q_s \geq 2$ be integers, $q = \min(q_1, \dots, q_s)$, $k^{(1)}, \dots, k^{(s)} \geq 1$ be integers, and put $\mathbf{k} = (k^{(1)}, \dots, k^{(s)})$, $k = \min(k^{(1)}, \dots, k^{(s)})$. Then

$$(19) \quad D((\mathbf{x}_n)_{n=1}^N) \leq D((\{\mathbf{x}_n\}_{\mathbf{k}})_{n=1}^N) + \frac{s}{q^k},$$

$$(20) \quad D((\{\mathbf{x}_n\}_{\mathbf{k}})_{n=1}^N) \leq \frac{s}{q^k} + \max_{c_i \in [1, q_i^{k^{(i)}}], i=1, \dots, s} \left| \frac{1}{N} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in \prod_{i=1}^s [0, \gamma(c_i))\} - \prod_{i=1}^s \gamma(c_i) \right| \leq \frac{s}{q^k} + D((\mathbf{x}_n)_{n=1}^N),$$

where $\gamma(c_i) = c_i/q_i^{k^{(i)}}$, $\{\mathbf{x}_n\}_{\mathbf{k}} = (\{x_{n,1}\}_{k^{(1)},1}, \dots, \{x_{n,s}\}_{k^{(s)},s})$, $\{y\}_{m,i} = [q_i^m \{y\}]/q_i^m$, $m = 1, 2, \dots$.

Proof. Let $v = [0, \gamma_1) \times \dots \times [0, \gamma_s)$; $v' = \prod_{i=1}^s [0, \{\gamma_i\}_{k^{(i)},i})$. It is easy to see that

$$\begin{aligned} \#\{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v'\} & \leq \#\{1 \leq n \leq N \mid \{\mathbf{x}_n\} \in v\} \\ & \leq \#\{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v\}. \end{aligned}$$

Using (2), we obtain

$$\frac{1}{N} \#\{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v\} \leq \gamma_1 \cdots \gamma_s + D((\{\mathbf{x}_n\}_{\mathbf{k}})_{n=1}^N),$$

and

$$\frac{1}{N} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v'\} \geq \gamma_1 \cdots \gamma_s - D\left(\left(\{\mathbf{x}_n\}_{\mathbf{k}}\right)_{n=1}^N\right) - \left| \prod_{i=1}^s \gamma_i - \prod_{i=1}^s \{\gamma_i\}_{k^{(i)}, i} \right|.$$

Hence

$$D\left(\left(\mathbf{x}_n\right)_{n=1}^N\right) \leq D\left(\left(\{\mathbf{x}_n\}_{\mathbf{k}}\right)_{n=1}^N\right) + \sup_{\gamma_1, \dots, \gamma_s \in [0,1]^s} \left| \prod_{i=1}^s \gamma_i - \prod_{i=1}^s \{\gamma_i\}_{k^{(i)}, i} \right|.$$

Similarly to [Ni, Lemma 3.9], the second sum is not more than

$$1 - \prod_{i=1}^s \left(1 - \frac{1}{q_i^{k^{(i)}}}\right) \leq 1 - \prod_{i=1}^s \left(1 - \frac{1}{q^k}\right) \leq \frac{s}{q^k},$$

and we obtain the first part of the lemma.

Let $v'' = \prod_{i=1}^s [0, \{\gamma_i\}_{k^{(i)}, i} + 1/q_i^{k^{(i)}})$. It is easy to see that

$$\begin{aligned} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\} \in v'\} &\leq \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v\} \\ &\leq \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\} \in v''\} \end{aligned}$$

and

$$\begin{aligned} &\left| \frac{1}{N} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\} \in v'\} - \text{mes } v' \right| - |\text{mes } v - \text{mes } v'| \\ &\leq \left| \frac{1}{N} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v\} - \text{mes } v \right| \\ &\leq \left| \frac{1}{N} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\} \in v''\} - \text{mes } v'' \right| + |\text{mes } v'' - \text{mes } v|. \end{aligned}$$

Bearing in mind that $\max((\text{mes } v - \text{mes } v'), (\text{mes } v'' - \text{mes } v))$

$$\begin{aligned} &\leq \max_{c_i \in [0, q_i^{k^{(i)}}), i=1, \dots, s} \prod_{i=1}^s \frac{c_i + 1}{q_i^{k^{(i)}}} - \prod_{i=1}^s \frac{c_i}{q_i^{k^{(i)}}} \\ &\leq 1 - \prod_{i=1}^s \left(1 - \frac{1}{q_i^{k^{(i)}}}\right) \leq \frac{s}{q^k}, \end{aligned}$$

we find that

$$\begin{aligned} &\left| \frac{1}{N} \# \{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in v\} \right| \leq \frac{s}{q^k} \\ &+ \max_{c_i \in [1, q_i^{k^{(i)}}], i=1, \dots, s} \left| \frac{1}{N} \# \left\{1 \leq n \leq N \mid \{\mathbf{x}_n\}_{\mathbf{k}} \in \prod_{i=1}^s \left[0, \frac{c_i}{q_i^{k^{(i)}}}\right)\right\} - \prod_{i=1}^s \frac{c_i}{q_i^{k^{(i)}}} \right|. \end{aligned}$$

Now we obtain from (2) the second part of the lemma. $\square$

Lemma 6. Let $q_1, \dots, q_s \geq 2$ be integers, $q = \min(q_1, \dots, q_s)$, $k^{(1)}, \dots, k^{(s)} \geq 1$ be integers, $\ell^{(1)}, \dots, \ell^{(s)}$; put $k = \min(k^{(1)}, \dots, k^{(s)})$, $\ell = \min(\ell^{(1)}, \dots, \ell^{(s)})$; $\mathbf{x}_n = (x_{n,1}, \dots, x_{n,s})$, $\mathbf{y}_n = (y_{n,1}, \dots, y_{n,s})$, $\mathbf{z}_n = (z_{n,1}, \dots, z_{n,s})$, and $y_{n,i} = \{x_{n,i}\}_{k,i}^{(i)} + \frac{1}{q_i^{k(i)}} \{z_{n,i}\}_{\ell,i}^{(i)}$, $i = 1, \dots, s$, $n = 1, 2, \dots$. Then

$$D^{(s)}((\mathbf{y}_n)_{n=1}^N) \leq 2^s D^{(2s)}((\mathbf{x}_n, \mathbf{z}_n)_{n=1}^N) + \frac{s}{q^{k+\ell}}.$$

Proof. Let $S = \{1, 2, \dots, s\}$, $I \subset S$, $\gamma_i = c_i / q_i^{k(i)+\ell(i)}$, $c_i \in \{1, \dots, q_i^{k(i)+\ell(i)}\}$, $\gamma'_i = q_i^{k(i)} (\gamma_i - \{\gamma_i\}_{k(i),i}) = \{\gamma'_i\}_{\ell(i),i}$, $i = 1, \dots, s$,

$$\gamma_I = \prod_{i \in I} \{\gamma_i\}_{k(i),i} \prod_{i \in S \setminus I} q_i^{-k(i)} \gamma'_i,$$

$$\begin{aligned} B_I &= \left\{ 1 \leq n \leq N \mid \left(\{x_{n,i}\}_{k(i),i} \in [0, \{\gamma_i\}_{k(i),i}), \quad \forall i \in I \right), \quad \text{and} \right. \\ &\quad \left. \left(\{x_{n,i}\}_{k(i),i} = \{\gamma_i\}_{k(i),i}, \quad \text{and} \quad \{z_{n,i}\}_{\ell(i),i} \in [0, \gamma'_i] \quad \forall i \in S \setminus I \right) \right\}, \\ B &= \{1 \leq n \leq N \mid \{y_{n,i}\} \in [0, \gamma_i], \quad i = 1, \dots, s\}. \end{aligned}$$

It is easy to see that

$$\begin{aligned} \prod_{i \in [1,s]} \gamma_i &= \sum_{I \subset S} \gamma_I, \quad B = \bigcup_{I \subset S} B_I, \\ (21) \quad \left| \frac{1}{N} \# B - \prod_{i \in [1,s]} \gamma_i \right| &\leq \sum_{I \subset S} \left| \frac{1}{N} \# B_I - \gamma_I \right|, \end{aligned}$$

and

$$\begin{aligned} \{x_{n,i}\}_{k(i),i} \in [0, \{\gamma_i\}_{k(i),i}] &\iff \{x_{n,i}\} \in [0, \{\gamma_i\}_{k(i),i}); \\ \{x_{n,i}\}_{k(i),i} = \{\gamma_i\}_{k(i),i} &\iff \{x_{n,i}\} \in [\{\gamma_i\}_{k(i),i}, \{\gamma_i\}_{k(i),i} + 1/q_i^{k(i)}); \\ \{z_{n,i}\}_{\ell(i),i} \in [0, \gamma'_i] &\iff \{z_{n,i}\} \in [0, \gamma'_i] \quad (\gamma'_i = \{\gamma'_i\}_{\ell(i),i}). \end{aligned}$$

Applying (2), we obtain:

$$\begin{aligned} \left| \gamma_I - \frac{1}{N} \# B_I \right| &\leq \left| \gamma_I - \frac{1}{N} \# \left\{ 1 \leq n \leq N \mid \left(\{x_{n,i}\} \in [0, \{\gamma_i\}_{k(i),i}] \quad \forall i \in I \right), \right. \right. \\ &\quad \text{and} \quad \left(\{x_{n,i}\} \in [\{\gamma_i\}_{k(i),i}, \{\gamma_i\}_{k(i),i} + 1/q_i^{k(i)}] \right. \\ &\quad \left. \left. \text{and} \quad \{z_{n,i}\} \in [0, \gamma'_i] \quad \forall i \in S \setminus I \right) \right\} \right| \\ &\leq D^{(2s-\#I)} \left(((\{x_{n,i}\})_{i=1}^s, (\{z_{n,i}\})_{i \in S \setminus I})_{n=1}^N \right) \leq D^{(2s)}((\mathbf{x}_n, \mathbf{z}_n)_{n=1}^N). \end{aligned}$$

From (20) and (21) we get :

$$\begin{aligned}
 D\left(\left(\{\mathbf{y}_n\}\right)_{n=1}^N\right) &\leq \frac{s}{q^{k+\ell}} \\
 &+ \max_{\substack{c_i \in [1, q_i^{k(i)+\ell(i)}] \\ i=1, \dots, s}} \left| \frac{1}{N} \# \left\{ 1 \leq n \leq N \mid \{\mathbf{y}_n\} \in \prod_{i=1}^s [0, \gamma(c_i)) \right\} - \prod_{i=1}^s \gamma(c_i) \right| \\
 &\leq \frac{s}{q^{k+\ell}} + \max_{\substack{c_i \in [1, q_i^{k(i)+\ell(i)}] \\ i=1, \dots, s}} \sum_{I \subset S} \left| \frac{1}{N} \# B_I - \gamma_I \right| \leq \frac{s}{q^{k+\ell}} + 2^s D^{(2s)}((\mathbf{x}_n, \mathbf{z}_n)_{n=1}^N),
 \end{aligned}$$

where $\gamma(c_i) = c_i / q_i^{k(i)+\ell(i)}$, $i = 1, \dots, s$. □

Let $\varphi_0 = \varphi(p_1 p_2) / p_1 p_2$, where $\varphi(x)$ is a Euler function,

$$\begin{aligned}
 (22) \quad \Delta_m(b) &= \{0, 1, \dots, b^m - 1\}, \\
 \Delta_m^*(b) &= \{n \in \Delta_m(b) \mid (n, b) = 1\}.
 \end{aligned}$$

It is evident that

$$(23) \quad \# \Delta_m^*(p_1 p_2) = \varphi(p_1 p_2) p_1^{m-1} p_2^{m-1} = \varphi_0 p_1^m p_2^m, \quad m = 1, 2, \dots.$$

Now let

$$\begin{aligned}
 (24) \quad A(i, m, s, c_0, \dots, c_{2s-1}) &= p_1^m p_2^m \\
 &\times \sum_{h_{-1} \in C(p_1^m)} \sum_{h_{-2} \in C(p_2^m)} \sum_{\mathbf{h} \in C_{2s}(p_1^m p_2^m)} r^{-1}(h_{-1}, p_1^m) r^{-1}(h_{-2}, p_2^m) r^{-1}(\mathbf{h}, p_1^m p_2^m) \\
 &\times \delta_{p_1^m p_2^m} \left(h_{-1} M_2 p_2^m + h_{-2} M_1 p_1^m + \sum_{j=0}^{s-1} (q^i h_j c_j + h_{j+s} c_{j+s}) \right),
 \end{aligned}$$

where $M_1 p_1^m \equiv 1 \pmod{p_2^m}$ and $M_2 p_2^m \equiv 1 \pmod{p_1^m}$.

Lemma 7. *Let $i \geq 0$; $m, s \geq 1$ be integers. Then*

$$\begin{aligned}
 (25) \quad \frac{1}{(\varphi_0 p_1^m p_2^m)^{2s}} \sum_{(c_0, \dots, c_{2s-1}) \in (\Delta_m^*(p_1 p_2))^{2s}} A(i, m, s, c_0, \dots, c_{2s-1}) \\
 \leq K_1(s) m^{2s+2},
 \end{aligned}$$

with $K_1(s) = 12\varphi_0^{-2s} (\frac{2}{\pi} \log p_1 p_2 + \frac{7}{5})^{2s+2} p_1^2 p_2^2$.

Proof. We follow [Ko3, p. 191]. We denote the left side of (25) by σ_1 . Changing the order of the summation, from (24) we get

$$\begin{aligned}
 (26) \quad \sigma_1 &\leq \sum_{h_{-1} \in C(p_1^m)} \sum_{h_{-2} \in C(p_2^m)} \sum_{\mathbf{h} \in C_{2s}(p_1^m p_2^m)} r^{-1}(h_{-1}, p_1^m) \\
 &\times r^{-1}(h_{-2}, p_2^m) r^{-1}(\mathbf{h}, p_1^m p_2^m) E(h_{-1}, h_{-2}, \mathbf{h}),
 \end{aligned}$$

where

$$(27) \quad E(h_{-1}, h_{-2}, \mathbf{h}) = \frac{1}{(\varphi_0 p_1^m p_2^m)^{2s}} \sum_{(c_0, \dots, c_{2s-1}) \in (\Delta_m(p_1 p_2))^{2s}} p_1^m p_2^m \\ \times \delta_{p_1^m p_2^m} \left(h_{-1} M_2 p_2^m + h_{-2} M_1 p_1^m + \sum_{j=0}^{s-1} (q^i h_j c_j + h_{j+s} c_{j+s}) \right).$$

Let $(h_0, \dots, h_{2s-1}, p_1^m p_2^m) = p_1^{\alpha_1} p_2^{\alpha_2}$, and let

$$(28) \quad h_i = p_1^{\alpha_1 + \beta_{1,i}} p_2^{\alpha_2 + \beta_{2,i}} h'_i, \quad (h'_i, p_1 p_2) = 1, \quad \beta_{1,i}, \beta_{2,i} \geq 0, \\ i = 0, \dots, 2s-1.$$

This yields that there exist $\mu, \nu \in [0, 2s-1]$ such that $\beta_{1,\mu} = 0$ and $\beta_{2,\nu} = 0$. If $\mu = \nu$, then

$$(29) \quad \sum_{c_\mu \in [0, p_1^m p_2^m)} \delta_{p_1^{m-\alpha_1} p_2^{m-\alpha_2}} (c_\mu h'_\mu v_1 + v_3) = p_1^{\alpha_1} p_2^{\alpha_2} \quad \text{for } (v_1, p_1 p_2) = 1.$$

Now let $\mu \neq \nu$.

We find for integers v_1, v_2 and v_3 (with $(v_i, p_1 p_2) = 1$, $i = 1, 2$), that

$$(30) \quad \sum_{c_\mu, c_\nu \in [0, p_1^m p_2^m)} \delta_{p_1^{m-\alpha_1} p_2^{m-\alpha_2}} (c_\mu h'_\mu p_2^{\beta_{2,\mu}} v_1 + c_\nu h'_\nu p_1^{\beta_{1,\nu}} v_2 + v_3) \\ = \sum_{c_{\mu,1}, c_{\nu,1} \in [0, p_1^m)} \sum_{c_{\mu,2}, c_{\nu,2} \in [0, p_2^m)} \delta_{p_1^{m-\alpha_1} p_2^{m-\alpha_2}} \left((c_{\mu,1} M_2 p_2^m + c_{\mu,2} M_1 p_1^m) h'_\mu p_2^{\beta_{2,\mu}} v_1 \right. \\ \left. + (c_{\nu,1} M_2 p_2^m + c_{\nu,2} M_1 p_1^m) h'_\nu p_1^{\beta_{1,\nu}} v_2 + v_3 \right) = \sigma' \sigma'',$$

where

$$\sigma' = \sum_{c_{\mu,1}, c_{\nu,1} \in [0, p_1^m)} \delta_{p_1^{m-\alpha_1}} (c_{\mu,1} M_2 p_2^m h'_\mu p_2^{\beta_{2,\mu}} v_1 + c_{\nu,1} M_2 p_2^m h'_\nu p_1^{\beta_{1,\nu}} v_2 + v_3),$$

and

$$\sigma'' = \sum_{c_{\mu,2}, c_{\nu,2} \in [0, p_2^m)} \delta_{p_2^{m-\alpha_2}} (c_{\mu,2} M_1 p_1^m h'_\mu p_2^{\beta_{2,\mu}} v_1 + c_{\nu,2} M_1 p_1^m h'_\nu p_1^{\beta_{1,\nu}} v_2 + v_3).$$

Observe that $(M_2 p_2^m h'_\mu p_2^{\beta_{2,\mu}} v_1, p_1) = 1$, $(M_1 p_1^m h'_\nu p_1^{\beta_{1,\nu}} v_2, p_2) = 1$,

$$\sum_{c_{\mu,1} \in [0, p_1^m)} \delta_{p_1^{m-\alpha_1}} (c_{\mu,1} M_2 p_2^m h'_\mu p_2^{\beta_{2,\mu}} v_1 + v_4) = p_1^{\alpha_1},$$

and

$$\sum_{c_{\nu,2} \in [0, p_2^m)} \delta_{p_2^{m-\alpha_2}} (c_{\nu,2} M_1 p_1^m h'_\nu p_1^{\beta_{1,\nu}} v_2 + v_5) = p_2^{\alpha_2}.$$

Hence

$$(31) \quad \sigma' = p_1^{m+\alpha_1} \quad \text{and} \quad \sigma'' = p_2^{m+\alpha_2}.$$

If $h_{-1} \not\equiv 0 \pmod{p_1^{\alpha_1}}$ or $h_{-2} \not\equiv 0 \pmod{p_2^{\alpha_2}}$, then $E(h_{-1}, h_{-2}, \mathbf{h}) = 0$.
Now let

$$h_{-1} = p_1^{\alpha_1} h'_{-1}, \quad h_{-2} = p_2^{\alpha_2} h'_{-2}.$$

From (27) we see, that

$$(32) \quad E(p_1^{\alpha_1} h'_{-1}, p_2^{\alpha_2} h'_{-2}, p_1^{\alpha_1} p_2^{\alpha_2} \mathbf{h}') = \frac{1}{(\varphi_0 p_1^m p_2^m)^{2s}} \\ \times \sum_{(c_0, \dots, c_{2s-1}) \in (\Delta_r(p_1^m p_2^m))^{2s}} p_1^m p_2^m \times \delta_{p_1^{m-\alpha_1} p_2^{m-\alpha_2}} \left(h'_{-1} M_2 p_2^{m-\alpha_2} \right. \\ \left. + h'_{-2} M_1 p_1^{m-\alpha_1} + \sum_{j=0}^{s-1} (q^i h'_j c_j p_1^{\beta_{1,j}} p_2^{\beta_{2,j}} + h'_{j+s} c_{j+s} p_1^{\beta_{1,j+s}} p_2^{\beta_{2,j+s}}) \right).$$

Now, applying (29) for the case of $\mu=\nu$ and (30), (31) for the case of $\mu \neq \nu$, we get

$$E(p_1^{\alpha_1} h'_{-1}, p_2^{\alpha_2} h'_{-2}, p_1^{\alpha_1} p_2^{\alpha_2} \mathbf{h}') = \varphi_0^{-2s} p_1^{\alpha_1} p_2^{\alpha_2}.$$

We obtain from (26) that

$$\sigma_1 \leq \varphi_0^{-2s} \sum_{0 \leq \alpha_1, \alpha_2 \leq m} \sum_{h'_{-1} \in C(p_1^{m-\alpha_1})} \sum_{h'_{-2} \in C(p_2^{m-\alpha_2})} \sum_{\mathbf{h}' \in C_{2s}(p_1^{m-\alpha_1} p_2^{m-\alpha_2})} p_1^{\alpha_1} p_2^{\alpha_2} \times \\ (33) \quad \times r^{-1}(h'_{-1} p_1^{\alpha_1}, p_1^m) r^{-1}(h'_{-2} p_2^{\alpha_2}, p_2^m) r^{-1}(\mathbf{h}' p_1^{\alpha_1} p_2^{\alpha_2}, p_1^m p_2^m).$$

Applying Lemma 2, we get

$$\sum_{\mathbf{h}' \in C_{2s}(p_1^{m-\alpha_1} p_2^{m-\alpha_2})} r^{-1}(p_1^{\alpha_1} p_2^{\alpha_2} \mathbf{h}', p_1^m p_2^m) \leq p_1^{-\alpha_1} p_2^{-\alpha_2} \left(\frac{2m}{\pi} \log p_1 p_2 + \frac{7}{5} \right)^{2s},$$

and

$$(34) \quad \sum_{h'_{-1} \in C(p_1^{m-\alpha_1})} r^{-1}(p_1^{\alpha_1} h_{-1}, p_1^m) \leq 1 + p_1^{-\alpha_1} \left(\frac{2m}{\pi} \log p_1 + \frac{7}{5} \right).$$

Now, from (33) we obtain :

$$\begin{aligned}
 \sigma_1 &\leq \varphi_0^{-2s} \sum_{0 \leq \alpha_1, \alpha_2 \leq m} \left(\frac{2}{\pi} m \log p_1 p_2 + \frac{7}{5} \right)^{2s} \left(1 + p_1^{-\alpha_1} \left(\frac{2}{\pi} m \log p_1 + \frac{7}{5} \right) \right) \\
 &\quad \times \left(1 + p_2^{-\alpha_2} \left(\frac{2}{\pi} m \log p_2 + \frac{7}{5} \right) \right) \\
 &\leq \varphi_0^{-2s} \left(\frac{2}{\pi} m \log p_1 p_2 + \frac{7}{5} \right)^{2s} \left((m+1)^2 + (m+1) \left(\frac{2}{\pi} m \log p_1 p_2 + \frac{7}{5} \right) \right) \\
 &\quad \times \left(\frac{p_1}{p_1-1} + \frac{p_2}{p_2-1} \right) + \left(\frac{2}{\pi} m \log p_1 p_2 + \frac{7}{5} \right)^2 \frac{p_1}{p_1-1} \frac{p_2}{p_2-1} \\
 &\leq K_1(s) m^{2s+2}.
 \end{aligned}$$

□

Let

$$\mathbf{b}^{(m)} = (b_{0,0}^{(m)}, \dots, b_{t_m-1,0}^{(m)}, b_{0,1}^{(m)}, \dots, b_{t_m-1,1}^{(m)}),$$

$$\begin{aligned}
 (35) \quad B(\nu, i, j, s, \mathbf{b}^{(m)}) &= A(i, m, s, b_{j,\nu}^{(m)}, \dots, b_{j+s-1,\nu}^{(m)}, b_{j,\nu+1}^{(m)}, \\
 &\quad \dots, b_{j+s-1,\nu+1}^{(m)}),
 \end{aligned}$$

where $b_{j+t_m,\nu+2}^{(m)} = b_{j,\nu}^{(m)}$, $\nu \in \{0, 1\}$, $j \in \{0, 1, \dots, t_m - 1\}$,

$$\begin{aligned}
 (36) \quad \Omega_m &= \left\{ \mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m} \mid \sum_{s \in [1, t_m]} \sum_{j \in [0, t_m]} \right. \\
 &\quad \times \sum_{i \in [0, k_m-1]} \sum_{\nu \in \{0, 1\}} B(\nu, i, j, s, \mathbf{b}^{(m)}) (24K_1(s) t_m^2 k_m m^{2s+2})^{-1} > 1 \Big\}.
 \end{aligned}$$

Lemma 8. *With the notation defined above we have:*

$$\# \Omega_m \leq \frac{1}{12} (\# \Delta_m^*(p_1 p_2))^{2t_m}, \quad m = 1, 2, \dots$$

Proof. It follows from Lemma 7 and (35) that

$$\begin{aligned}
 &\frac{1}{(\# \Delta_m^*(p_1 p_2))^{2t_m}} \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} B(\nu, i, j, s, \mathbf{b}^{(m)}) = \frac{1}{(\# \Delta_m^*(p_1 p_2))^{2t_m}} \\
 &\times \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} A(i, m, s, b_{j,\nu}^{(m)}, \dots, b_{j+s-1,\nu}^{(m)}, b_{j,\nu+1}^{(m)}, \dots, b_{j+s-1,\nu+1}^{(m)}) \\
 &\leq K_1(s) m^{2s+2},
 \end{aligned}$$

where $\nu \in \{0, 1\}$, $j \in [0, t_m]$, and $i \in [0, k_m - 1]$.

Hence

$$\sum_{j \in [0, t_m]} \sum_{i \in [0, k_m-1]} \sum_{\nu \in \{0, 1\}} \frac{1}{(\# \Delta_m^*(p_1 p_2))^{2t_m}} \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} B(\nu, i, j, s, \mathbf{b}^{(m)})$$

$$\leq 2K_1(s)t_mk_m m^{2s+2},$$

and

$$\sum_{s \in [1, t_m]} \sum_{j \in [0, t_m]} \sum_{i \in [0, k_m - 1]} \sum_{\nu \in \{0, 1\}} \frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_m}} \\ \times \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} B(\nu, i, j, s, \mathbf{b}^{(m)}) \left(2K_1(s)t_m^2 k_m m^{2s+2} \right)^{-1} \leq 1.$$

Changing the order of the summation, we find that

$$(37) \quad \frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_m}} \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} \left(\sum_{s \in [1, t_m]} \left(24K_1(s)t_m^2 k_m m^{2s+2} \right)^{-1} \right. \\ \left. \times \sum_{j \in [0, t_m]} \sum_{i \in [0, k_m - 1]} \sum_{\nu \in \{0, 1\}} B(\nu, i, j, s, \mathbf{b}^{(m)}) \right) \leq 1/12.$$

Now, from (36), we obtain the assertion of the lemma. $\square$

Put

$$(38) \quad A_1(\alpha, i, m, s, c_0, \dots, c_{s-1}) = \sum_{h_{-2} \in C(p_2^{m+1})} \sum_{\substack{\mathbf{h} \in C_{2s}(p_2^{m+1}) \\ (h_s, \dots, h_{2s-1}, p_2^{m+1}) = p_2^\alpha}} \\ \times r^{-1}(h_{-2}, p_2^{m+1}) r^{-1}(\mathbf{h}, p_2^{m+1}) p_2^\alpha \delta_{p_2^\alpha} \left(h_{-2} + q^i p_2 \sum_{j=0}^{s-1} h_j c_j t_{m+1}/t_m \right),$$

and for $v \in [1, s-1]$

$$(39) \quad A_2(\alpha, i, m, v, c_0, \dots, c_{2s-1}) = \sum_{h_{-1} \in C(p_1^{m+1})} \sum_{\substack{\mathbf{h} \in C_{2s}(p_1^{m+1}) \\ (h_v, \dots, h_{s-1}, h_{v+s}, \dots, h_{2s-1}, p_1^{m+1}) = p_1^\alpha}} \\ \times r^{-1}(h_{-1}, p_1^m) r^{-1}(\mathbf{h}, p_1^m) p_1^\alpha \delta_{p_1^\alpha} \left(h_{-1} + p_1 k_{m+1}/k_m \sum_{j=0}^{v-1} (q^i h_j c_j + h_{j+s} c_{j+s}) \right).$$

Lemma 9. *Let $t_m \geq s \geq 1, \alpha \in [0, m+1], v \in [1, s-1], i \in [0, k_{m-1}]$ be integers. Then*

$$(40) \quad \frac{1}{(\#\Delta_m^*(p_2))^s} \sum_{(c_0, \dots, c_{s-1}) \in (\Delta_m^*(p_2))^s} A_1(\alpha, i, m, s, c_0, \dots, c_{s-1}) \\ \leq K_1(s)p_2(m+1)^{2s+1}$$

and

$$(41) \quad \frac{1}{(\#\Delta_m^*(p_1))^{2s}} \sum_{(c_0, \dots, c_{2s-1}) \in (\Delta_m^*(p_1))^{2s}} A_2(\alpha, i, m, v, c_0, \dots, c_{2s-1}) \leq K_1(s) p_1 (m+1)^{2s+1}.$$

Proof. We will prove the statement (41). The proof of (40) repeats that of (41). We denote the left side of (41) by σ_1 . Changing the order of the summation, from (39) we get

$$(42) \quad \sigma_1 = \sum_{h_{-1} \in C(p_1^{m+1})} \sum_{\substack{\mathbf{h} \in C_{2s}(p_1^{m+1}) \\ (h_v, \dots, h_{s-1}, h_{v+s}, \dots, h_{2s-1}, p_1^{m+1}) = p_1^\alpha}} r^{-1}(h_{-1}, p_1^{m+1}) \times r^{-1}(\mathbf{h}, p_1^{m+1}) E(h_{-1}, \mathbf{h}),$$

where

$$E(h_{-1}, \mathbf{h}) = \frac{1}{(\varphi(p_1) p_1^{m-1})^{2s}} \sum_{(c_0, \dots, c_{2s-1}) \in (\Delta_m^*(p_1))^{2s}} p_1^\alpha \times \delta_{p_1^\alpha} \left(h_{-1} + p_1 k_{m+1} / k_m \sum_{j=0}^{v-1} (q^i h_j c_j + h_{j+s} c_{j+s}) \right).$$

Let $(h_0, \dots, h_{v-1}, h_s, \dots, h_{v+s-1}, p_1^{m+1}) = p_1^{\alpha_1}$, and let

$$h_{\nu s+i} = p_1^{\alpha_1} h'_{\nu s+i}, \quad \nu = 0, 1, \quad i = 0, \dots, v-1.$$

Then there is $(\nu_0, i_0) \in \{0, 1\} \times \{0, \dots, v-1\}$ with $(h'_{\nu_0 s+i_0}, p_1) = 1$.

It is easy to see that

$$(43) \quad E(h_{-1}, \mathbf{h}) \leq \max_{\mathbf{c} \in (\Delta_m^*(p_1))^{2s}} \sigma(\mathbf{c}, i_0, \nu_0),$$

where

$$(44) \quad \sigma(\mathbf{c}, i_0, \nu_0) = \frac{1}{\varphi(p_1) p_1^{m-1}} \sum_{c_{\nu_0 s+i_0} \in \Delta_{m+1}^*(p_1)} p_1^\alpha \times \delta_{p_1^\alpha} \left(h_{-1} + p_1^{\alpha_1+1} k_{m+1} / k_m \sum_{j=0}^{v-1} (q^i h'_j c_j + h'_{j+s} c_{j+s}) \right).$$

We find that if $h_{-1} \not\equiv 0 \pmod{p_1^{\min(\alpha, \alpha_1)}}$, then $E(h_{-1}, \mathbf{h}) = 0$.

Now let $h_{-1} \equiv 0 \pmod{p_1^{\min(\alpha, \alpha_1)}}$, and let $h_{-1} = p_1^{\min(\alpha, \alpha_1)} h'_{-1}$.

From (44) we find that

$$(45) \quad \sigma(\mathbf{c}, i_0, \nu_0) = p_1^\alpha \quad \text{for } \alpha_1 \geq \alpha,$$

and

$$\begin{aligned}
 (46) \quad \sigma(\mathbf{c}, i_0, \nu_0) &\leq \frac{1}{\varphi(p_1)p_1^{m-1}} \sum_{c_{\nu_0 s+i_0} \in [0, p_1^m)} p_1^\alpha \\
 &\quad \times \delta_{p_1^{\alpha-\alpha_1}} \left(h'_{-1} + p_1 k_{m+1}/k_m \sum_{j=0}^{v-1} (q^i h'_j c_j + h'_{j+s} c_{j+s}) \right) \\
 &< \frac{1}{\varphi_0 p_1^m} \sum_{c_{\nu_0 s+i_0} \in [0, p_1^m)} p_1^\alpha \delta_{p_1^{\alpha-\alpha_1}} \left(f + p_1 k_{m+1}/k_m q^{(1-\nu_0)i} h'_{\nu_0 s+j_0} c_{\nu_0 s+j_0} \right) \\
 &\quad \text{for } \alpha_1 < \alpha,
 \end{aligned}$$

where

$$f = h'_{-1} + p_1 k_{m+1}/k_m \sum_{\substack{\nu \in \{0,1\}, j \in [0,v] \\ (\nu,j) \neq (\nu_0,j_0)}} q^{(1-\nu)i} h'_{\nu s+j} c_{\nu s+j}.$$

It is easy to verify that

$$\log_{p_1}(m+1) - \log_{p_1}(m) \leq 1, \quad m = 1, 2, \dots,$$

and

$$(47) \quad [\log_{p_1}(m+1)] - [\log_{p_1}(m)] \leq 1, \quad \text{for } m = 1, 2, \dots.$$

According to (7), we have

$$(48) \quad k_{m+1}/k_m = p_1^{[\log_{p_1}(m+1)] - [\log_{p_1}(m)]} \in \{1, p_1\} \quad \text{for } m = 1, 2, \dots.$$

Bearing in mind that $(q^{(1-\nu_0)i} h'_{\nu_0 s+j_0}, p_1) = 1$, we obtain from (46) that

$$\sigma(\mathbf{c}, i_0, \nu_0) \leq \varphi_0^{-1} p_1^{2+\alpha_1} \quad \text{for } \alpha_1 < \alpha.$$

Now, (45) and (43) imply that

$$E(h_{-1}, \mathbf{h}) \leq \varphi_0^{-1} p_1^{2+\min(\alpha, \alpha_1)} \delta_{p_1^{\min(\alpha, \alpha_1)}}(h_{-1}) \quad \text{for } \alpha_1 \in [0, m+1].$$

From (42) we obtain

$$\begin{aligned}
 \sigma_1 &\leq \sum_{\alpha_1 \in [0, m+1]} \sum_{\substack{h_{-1} \in C(p_1^{m+1}) \\ h_{-1} \equiv 0 \pmod{p_1^{\min(\alpha, \alpha_1)}}}} \sum_{\substack{\mathbf{h} \in C_{2s}(p_1^{m+1}) \\ (h_v, \dots, h_{s-1}, h_{v+s}, \dots, h_{2s-1}, p_1^{m+1}) = p_1^\alpha, \\ (h_0, \dots, h_{v-1}, h_s, \dots, h_{s+v-1}, p_1^{m+1}) = p_1^{\alpha_1}}} \\
 &\quad \times r^{-1}(h_{-1}, p_1^{m+1}) r^{-1}(\mathbf{h}, p_1^{m+1}) \varphi_0^{-1} p_1^{2+\min(\alpha, \alpha_1)}.
 \end{aligned}$$

Applying Lemma 2, we get

$$\begin{aligned}
 \sigma_1 &\leq \varphi_0^{-1} \sum_{\alpha_1 \in [0, m+1]} p_1^{2+\min(\alpha, \alpha_1)} \left(1 + p_1^{-\min(\alpha, \alpha_1)} \left(\frac{2(m+1)}{\pi} \log p_1 + \frac{7}{5} \right) \right) \\
 &\quad \times \left(\frac{2(m+1)}{\pi} \log p_1 + \frac{7}{5} \right)^{2s} p_1^{-\min(\alpha, \alpha_1)} \\
 &\leq \varphi_0^{-1} p_1^2 \left(\frac{2(m+1)}{\pi} \log p_1 + \frac{7}{5} \right)^{2s} \left(m+2 + 2 \left(\frac{2(m+1)}{\pi} \log p_1 + \frac{7}{5} \right) \right) \\
 &\leq \varphi_0^{-1} p_1^2 (m+1)^{2s+1} \left(\frac{2}{\pi} \log p_1 + \frac{7}{5} \right)^{2s} \left(\frac{m+2}{m+1} + 2 \left(\frac{2}{\pi} \log p_1 + \frac{7}{5} \right) \right) \\
 &\leq K_1(s) (m+1)^{2s+1}.
 \end{aligned}$$

□

Lemma 10. *Let*

$$\begin{aligned}
 (49) \quad \Omega_{m,1} &= \left\{ \mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m} \mid \sum_{s=1}^{t_m} \sum_{\alpha=0}^{m+1} \sum_{\mu=0}^{t_m-1} \sum_{i=0}^{k_m-1} \right. \\
 &\quad \times \left. \frac{A_1(\alpha, i, m, s, b_{\mu,1}^{(m)}, \dots, b_{\mu+s-1,1}^{(m)})}{12K_1(s)t_m^2 k_m (m+2)(m+1)^{2s+1}} > 1 \right\},
 \end{aligned}$$

and

$$\begin{aligned}
 (50) \quad \Omega_{m,2} &= \left\{ \mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m} \mid \sum_{s=1}^{t_m} \sum_{\alpha=0}^{m+1} \sum_{\mu \in (t_m-s, t_m)} \sum_{i=0}^{k_m-1} \sum_{\nu=0}^1 \right. \\
 &\quad \times \left. \frac{A_2(\alpha, i, m, t_m - \mu, b_{\mu,\nu}^{(m)}, \dots, b_{\mu+s-1,\nu}^{(m)}, b_{\mu,\nu+1}^{(m)}, \dots, b_{\mu+s-1,\nu+1}^{(m)})}{24K_1(s)t_m^2 k_m (m+2)(m+1)^{2s+1}} > 1 \right\}.
 \end{aligned}$$

Then

$$(51) \quad \#\Omega_{m,\nu} \leq \frac{1}{12} (\#\Delta_m^*(p_1 p_2))^{2t_m}, \quad \nu = 1, 2.$$

Proof. Let $\nu = 2$. It follows from Lemma 9, and (39) that

$$\begin{aligned}
 &\frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_m}} \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} \\
 &\quad A_2(\alpha, i, m, t_m - \mu, b_{\mu,\nu}^{(m)}, \dots, b_{\mu+s-1,\nu}^{(m)}, b_{\mu,\nu+1}^{(m)}, \dots, b_{\mu+s-1,\nu+1}^{(m)})
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(\#\Delta_m^*(p_1))^{2s}} \sum_{(b_{\mu,\nu}^{(m)}, \dots, b_{\mu+s-1,\nu+1}^{(m)}) \in (\Delta_m^*(p_1))^{2s}} \\
&\quad A_2(\alpha, i, m, t_m - \mu, b_{\mu,\nu}^{(m)}, \dots, b_{\mu+s-1,\nu}^{(m)}, b_{\mu,\nu+1}^{(m)}, \dots, b_{\mu+s-1,\nu+1}^{(m)}) \\
&\leq K_1(s) p_1 (m+1)^{2s+1},
\end{aligned}$$

for $s \in [1, t_m]$, $\alpha \in [0, m+1]$, $\mu \in (t_m - s, t_m)$, $i \in [0, k_m - 1]$, and $\nu \in \{0, 1\}$. Hence

$$\begin{aligned}
&\sum_{s=1}^{t_m} \sum_{\alpha=0}^{m+1} \sum_{\mu \in (t_m-s, t_m)} \sum_{i=0}^{k_m-1} \sum_{\nu=0}^1 \frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_m}} \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} \\
&\quad \times \frac{A_2(\alpha, i, m, t_m - \mu, b_{\mu,\nu}^{(m)}, \dots, b_{\mu+s-1,\nu}^{(m)}, b_{\mu,\nu+1}^{(m)}, \dots, b_{\mu+s-1,\nu+1}^{(m)})}{2K_1(s) p_1 t_m^2 k_m (m+2)(m+1)^{2s+1}} \leq 1.
\end{aligned}$$

Changing the order of the summation we obtain that

$$\begin{aligned}
&\frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_m}} \sum_{\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}} \sum_{s=1}^{t_m} \sum_{\alpha=0}^{m+1} \sum_{\mu \in (t_m-s, t_m)} \sum_{i=0}^{k_m-1} \sum_{\nu=0}^1 \\
&\quad \times \frac{A_2(\alpha, i, m, t_m - \mu, b_{\mu,\nu}^{(m)}, \dots, b_{\mu+s-1,\nu}^{(m)}, b_{\mu,\nu+1}^{(m)}, \dots, b_{\mu+s-1,\nu+1}^{(m)})}{24K_1(s) p_1 t_m^2 k_m (m+2)(m+1)^{2s+1}} \leq \frac{1}{12}.
\end{aligned}$$

Now, from (50) we obtain the desired result. Using (49), we similarly obtain (51) for the case of $\nu = 1$. $\square$

Now, from Lemmas 8 and 10 we get:

Corollary 4. *Let*

$$(52) \quad \Omega_{m,3} = \Omega_m \cup \Omega_{m,1} \cup \Omega_{m,2}.$$

Then

$$(53) \quad \#\Omega_{m,3} \leq \frac{1}{4} (\#\Delta_m^*(p_1 p_2))^{2t_m}, \quad m = 1, 2, \dots$$

Put

$$\begin{aligned}
(54) \quad B_1(i, \mu, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)}) &= \sum_{h_{-2} \in C(p_2^{m+1})} \sum_{\mathbf{h} \in C_{2s}(p_2^{m+1})} r^{-1}(h_{-2}, p_2^{m+1}) \\
&\quad \times r^{-1}(\mathbf{h}, p_2^{m+1}) p_2^{m+1} \delta_{p_2^{m+1}} \left(h_{-2} + \sum_{j=0}^{s-1} \left(q^i h_j b_{\mu+j,1}^{(m)} p_2 \frac{t_{m+1}}{t_m} + h_{s+j} c_{\mu+j,0}^{(m+1)} \right) \right),
\end{aligned}$$

where $c_{j+t_{m+1}, \nu+2}^{(m+1)} = c_{j,\nu}^{(m+1)}$, $\nu \in \{0, 1\}$, $j \in \{0, 1, \dots, t_{m+1} - 1\}$.

Lemma 11. *With the notation defined above, we have:*

$$(55) \quad \frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} B_1(i, \mu, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)}) \\ \leq p_2 \sum_{\alpha=0}^{m+1} A_1(\alpha, i, m, s, b_{\mu,1}^{(m)}, \dots, b_{\mu+s-1,1}^{(m)}) .$$

Proof. Let $(h_s, \dots, h_{2s-1}, p_2^{m+1}) = p_2^\alpha$, $h_i = h'_i p_2^\alpha$, for $i = s, \dots, 2s-1$, and let $(h'_{s+i_0}, p_2) = 1$ for some $i_0 \in [0, s-1]$. We denote the left side of (55) by σ . Changing the order of the summation we get from (54)

$$(56) \quad \sigma = \sum_{\alpha=0}^{m+1} \sum_{h_{-2} \in C(p_2^{m+1})} \sum_{\substack{\mathbf{h} \in C_{2s}(p_2^{m+1}) \\ (h_s, \dots, h_{2s-1}, p_2^{m+1}) = p_2^\alpha}} r^{-1}(h_{-2}, p_2^{m+1}) \\ \times r^{-1}(\mathbf{h}, p_2^{m+1}) W(h_{-2}, \mathbf{h}),$$

where

$$W(h_{-2}, \mathbf{h}) = \frac{1}{(\#\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} p_2^{m+1} \\ \times \delta_{p_2^{m+1}} \left(h_{-2} + \sum_{j=0}^{s-1} \left(q^i h_j b_{\mu+j,1}^{(m)} p_2 t_{m+1} / t_m + h_{s+j} c_{\mu+j,0}^{(m+1)} \right) \right).$$

It is easy to see that

$$W(h_{-2}, \mathbf{h}) = \frac{1}{(\#\Delta_{m+1}^*(p_1 p_2))^s} \sum_{(c_0, \dots, c_{s-1}) \in (\Delta_{m+1}^*(p_1 p_2))^s} p_2^{m+1} \\ \times \delta_{p_2^{m+1}} \left(f + \sum_{j=0}^{s-1} h_{s+j} c_j \right),$$

where

$$f = h_{-2} + \sum_{j=0}^{s-1} q^i h_j b_{\mu+j,1}^{(m)} p_2 t_{m+1} / t_m.$$

Hence

$$(57) \quad W(h_{-2}, \mathbf{h}) = \frac{1}{(\#\Delta_{m+1}^*(p_2))^s} \sum_{(c_0, \dots, c_{s-1}) \in (\Delta_{m+1}^*(p_2))^s} p_2^{m+1} \\ \times \delta_{p_2^{m+1}} \left(f + \sum_{j=0}^{s-1} p_2^\alpha h'_{s+j} c_j \right).$$

It is evident that if $f \not\equiv 0 \pmod{p_2^\alpha}$, then $W(h_{-2}, \mathbf{h}) = 0$.

Let $f \equiv 0 \pmod{p_2^\alpha}$, $f = f' p_2^\alpha$. Bearing in mind that $(h'_{s+i_0}, p_2) = 1$, we obtain

$$\sum_{c_{i_0} \in [0, p_2^{m+1})} \delta_{p_2^{m+1-\alpha}} \left(f' + \sum_{i \in [0, s-1], i \neq i_0} h'_{s+i} c_i + h'_{s+i_0} c_{i_0} \right) = p_2^\alpha.$$

Equation (57) shows that

$$\begin{aligned} W(h_{-2}, \mathbf{h}) &\leq \max_{\mathbf{c} \in (\Delta_{m+1}^*(p_2))^s} \sum_{c_{i_0} \in [0, p_2^{m+1})} (p_2 / \varphi(p_2)) \delta_{p_2^{m+1-\alpha}} \left(f' + \sum_{j=0}^{s-1} h'_{s+j} c_j \right) \\ &\leq (p_2 / \varphi(p_2)) p_2^\alpha \delta_{p_2^\alpha}(f). \end{aligned}$$

Then equation (56) implies that

$$\begin{aligned} \sigma &\leq \sum_{\alpha=0}^{m+1} \sum_{h_{-2} \in C(p_2^{m+1})} \sum_{\substack{\mathbf{h} \in C_{2s}(p_2^{m+1}) \\ (h_s, \dots, h_{2s-1}, p_2^{m+1}) = p_2^\alpha}} r^{-1}(h_{-2}, p_2^{m+1}) r^{-1}(\mathbf{h}, p_2^{m+1}) \\ &\quad \times p_2^{\alpha+1} \delta_{p_2^\alpha}(h_{-2} + \sum_{j=0}^{s-1} q^j h_j b_{\mu+j,1}^{(m)} p_2 t_{m+1} / t_m). \end{aligned}$$

Now from (38) we obtain the assertion of the lemma. $\square$

Corollary 5. Let $\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}$, $\mathbf{b}^{(m)} \notin \Omega_{m,3}$,

$$(58) \quad F_{m+1,1} = \left\{ \mathbf{b}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}} \mid \sum_{s=1}^{t_m} \sum_{\mu=0}^{t_{m+1}-1} \sum_{i=0}^{k_m-1} \times \frac{B_1(i, \mu, s, \mathbf{b}^{(m)}, \mathbf{b}^{(m+1)})}{96 K_1(s) p_2 t_{m+1}^2 k_m (m+1)^{2s+2}} > 1 \right\}.$$

Then

$$(59) \quad \# F_{m+1,1} \leq \frac{1}{4} (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}.$$

Proof. Changing the order of the summation, from Lemma 11 we have

$$\begin{aligned}
 (60) \quad & \frac{1}{(\#\Delta_m^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{s=1}^{t_m} \sum_{\mu \in [0, t_{m+1}]} \sum_{i \in [0, k_m - 1]} \\
 & \times \frac{B_1(i, \mu, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)})}{96K_1(s)p_2 t_{m+1}^2 k_m (m+1)^{2s+2}} \\
 & \leq \sum_{s=1}^{t_m} p_2 \sum_{\alpha \in [0, m+1]} \sum_{\mu \in [0, t_{m+1}]} \sum_{i \in [0, k_m - 1]} \frac{A_1(\alpha, i, m, s, b_{\mu,1}^{(m)}, \dots, b_{\mu+s-1,1}^{(m)})}{96K_1(s)p_2 t_{m+1}^2 k_m (m+1)^{2s+2}}.
 \end{aligned}$$

We denote the right side of (60) by σ . Bearing in mind (8) and that $b_{\mu+t_m,1}^{(m)} = b_{\mu,1}^{(m)}$, $\mu = 0, 1, \dots$, we find that

$$\begin{aligned}
 \sigma & \leq \frac{m+2}{8(m+1)} \sum_{s=1}^{t_m} \sum_{\alpha \in [0, m+1]} \sum_{\mu \in [0, t_m]} \sum_{i \in [0, k_m - 1]} \\
 & \times \frac{A_1(\alpha, i, m, s, b_{\mu,1}^{(m)}, \dots, b_{\mu+s-1,1}^{(m)})}{12K_1(s)p_2 t_m^2 k_m (m+2)(m+1)^{2s+1}}.
 \end{aligned}$$

Taking into account (49), (52), and that $\mathbf{b}^{(m)} \notin \Omega_{m,3}$, we deduce that $\sigma \leq (m+2)/8(m+1) \leq 1/4$.

Now, from (58) and (60), we obtain the desired result. $\square$

Let $x_2, x_3, x_5, x_6, y_2, y_4$ be integers, $y_2 \in (t_m - s, t_m)$,

$$\begin{aligned}
 (61) \quad B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)}) & = \sum_{h_{-1} \in C(p_1^{m+1})} \sum_{\mathbf{h} \in C_{2s}(p_1^{m+1})} \\
 & \times r^{-1}(h_{-1}, p_1^{m+1}) r^{-1}(\mathbf{h}, p_1^{m+1}) p_1^{m+1} \\
 & \times \delta_{p_1^{m+1}} \left(h_{-1} + p_1 k_{m+1} / k_m \sum_{0 \leq j < t_m - y_2} (q^{x_3} h_j b_{y_2+j, x_2}^{(m)} + h_{s+j} b_{y_2+j, x_2+1}^{(m)}) \right. \\
 & \quad \left. + \sum_{t_m - y_2 \leq j < s} (q^{x_6} h_j c_{y_4+j, x_5}^{(m+1)} + h_{s+j} c_{y_4+j, x_5+1}^{(m+1)}) \right).
 \end{aligned}$$

Lemma 12. Let $s \leq t_m$, $y_2 \in (t_m - s, t_m)$, $\mathbf{b}^{(m)} \notin \Omega_{m,3}$. Then

$$(62) \quad \frac{1}{(\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)}) \\ \leq p_1 \sum_{\alpha=0}^{m+1} A_2(\alpha, x_3, m, t_m - y_2, b_{y_2, x_2}^{(m)}, \dots, b_{y_2+s-1, x_2}^{(m)}, b_{y_2, x_2+1}^{(m)}, \\ \dots, b_{y_2+s-1, x_2+1}^{(m)}) .$$

Proof. Let $(h_{t_m-y_2}, \dots, h_{s-1}, h_{t_m-y_2+s}, \dots, h_{2s-1}, p_1^{m+1}) = p_1^\alpha$, and let $h_i = h'_i p_1^\alpha$, $i \in [t_m - y_2, s) \cup [t_m - y_2 + s, 2s)$.

Then $(h_{j_0}, p_1) = 1$ for some $j_0 = j_1 + s j_2$, with $j_1 \in [t_m - y_2, s)$, and $j_2 \in \{0, 1\}$.

We denote the left side of (62) by σ . Changing the order of the summation, we obtain from (61) that

$$(63) \quad \sigma = \sum_{\alpha=0}^{m+1} \sum_{h_{-1} \in C(p_1^{m+1})} \sum_{\substack{\mathbf{h} \in C_{2s}(p_1^{m+1}), \\ (h_{t_m-y_2}, \dots, h_{s-1}, h_{t_m-y_2+s}, \dots, h_{2s-1}, p_1^{m+1}) = p_1^\alpha}} \\ \times r^{-1}(h_{-1}, p_1^{m+1}) r^{-1}(\mathbf{h}, p_1^{m+1}) W(h_{-1}, \mathbf{h}),$$

where

$$W(h_{-1}, \mathbf{h}) = \frac{1}{(\#\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} p_1^{m+1} \\ \times \delta_{p_1^{m+1}} \left(h_{-1} + p_1 k_{m+1}/k_m \sum_{0 \leq j < t_m - y_2} (q^{x_3} h_j b_{y_2+j, x_2}^{(m)} + h_{s+j} b_{y_2+j, x_2+1}^{(m)}) \right. \\ \left. + \sum_{t_m - y_2 \leq j < s} (q^{x_6} h_j c_{y_4+j, x_5}^{(m+1)} + h_{s+j} c_{y_4+j, x_5+1}^{(m+1)}) \right).$$

We now obtain

$$W(h_{-1}, \mathbf{h}) \leq \max_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \frac{1}{\#\Delta_{m+1}^*(p_1)} \sum_{c_{y_4+j_1, x_5+j_2} \in \Delta_{m+1}^*(p_1)} \\ \times p_1^{m+1} \delta_{p_1^{m+1}} \left(f + p_1^\alpha \sum_{t_m - y_2 \leq j < s} (q^{x_6} h'_j c_{y_4+j, x_5}^{(m+1)} + h'_{s+j} c_{y_4+j, x_5+1}^{(m+1)}) \right),$$

where

$$f = h_{-1} + p_1 k_{m+1}/k_m \sum_{0 \leq j < t_m - y_2} (q^{x_3} h_j b_{y_2+j, x_2}^{(m)} + h_{s+j} b_{y_2+j, x_2+1}^{(m)}) .$$

It is easy to see that if $f \not\equiv 0 \pmod{p_1^\alpha}$, then $W(h_{-1}, \mathbf{h}) = 0$. Let $f \equiv 0 \pmod{p_1^\alpha}$, $f = f' p_1^\alpha$. Bearing in mind that $(q^{x_6(1-j_2)} h'_{j_2 s+j_1}, p_1) = 1$, we have

$$\sum_{c_{y_4+j_1, x_5+j_2} \in [0, p_1^{m+1})} \delta_{p_1^{m+1-\alpha}} \left(f' + q^{x_6(1-j_2)} h'_{j_2 s+j_1} c_{y_4+j_1, x_5+j_2} \right. \\ \left. + \sum_{\substack{(\nu, j) \in \{0,1\} \times [t_m - y_2, s) \\ (\nu, j) \neq (j_2, j_1)}} q^{x_6(1-\nu)} h'_{\nu s+j} c_{y_4+j, x_5+\nu} \right) = p_1^\alpha.$$

Hence

$$W(h_{-1}, \mathbf{h}) \leq (p_1/\varphi(p_1)) p_1^\alpha \delta_{p_1^\alpha}(f) \\ \leq p_1^{\alpha+1} \delta_{p_1^\alpha} \left(h_{-1} + p_1 k_{m+1}/k_m \sum_{0 \leq j < t_m - y_2} (q^{x_3} h_j b_{y_2+j, x_2}^{(m)} + h_{s+j} b_{y_2+j, x_2+1}^{(m)}) \right).$$

Now, from (63) and (39), we obtain the assertion of the lemma. $\square$

Corollary 6. Let $\mathbf{b}^{(m)} \in (\Delta_m^*(p_1 p_2))^{2t_m}$, $\mathbf{b}^{(m)} \notin \Omega_{m,3}$, $x_2 = 2\{([x_6/k_m] + x_5 k_{m+1}/k_m)/2\}$, $x_3 = k_m\{x_6/k_m\}$, $y_4 = t_{m+1}\{(t_m(p_1^m - 1) + y_2)/t_{m+1}\}$, $m = 1, 2, \dots$,

$$(64) \quad F_{m+1,2} = \left\{ \mathbf{b}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}} \mid \sum_{s=1}^{t_m} \sum_{x_5=0}^1 \sum_{x_6=0}^{k_{m+1}-1} \right. \\ \left. \times \sum_{y_2 \in (t_m-s, t_m)} \frac{B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{b}^{(m)}, \mathbf{b}^{(m+1)})}{192 K_1(s) p_1 t_m^2 k_{m+1} (m+1)^{2s+2}} > 1 \right\}.$$

Then

$$(65) \quad \#F_{m+1,2} \leq \frac{1}{4} (\#\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}.$$

Proof. Changing the order of the summation, from Lemma 12 we have

$$(66) \quad \frac{1}{(\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{x_5=0}^1 \sum_{x_6=0}^{k_{m+1}-1} \\ \times \sum_{y_2 \in (t_m-s, t_m)} B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)}) \\ \leq p_1 \sum_{\alpha=0}^{m+1} \sum_{x_5=0}^1 \sum_{x_6=0}^{k_{m+1}-1} \sum_{y_2 \in (t_m-s, t_m)} \\ \times A_2(\alpha, x_3, m, t_m - y_2, b_{y_2, x_2}^{(m)}, \dots, b_{y_2+s-1, x_2}^{(m)}, b_{y_2, x_2+1}^{(m)}, \dots, b_{y_2+s-1, x_2+1}^{(m)}).$$

According to (7), $k_{m+1}/k_m = p_1^{[\log_{p_1}(m+1)] - [\log_{p_1} m]}$.

Let $p_1 = 2$ and $k_{m+1}/k_m > 1$. Then $x_2 = 2\{[x_6/k_m]/2\}$, $x_3 = k_m\{x_6/k_m\}$. If x_6 passes the set $\{0, 1, \dots, k_{m+1} - 1\}$, then (x_2, x_3) passes $k_{m+1}/(2k_m)$ times the set $\{0, 1\} \times \{0, 1, \dots, k_m - 1\}$.

Now let $p_1 \geq 3$ or $p_1 = 2$ and $k_{m+1}/k_m = 1$. We find that $x_2 = 2\{([x_6/k_m] + x_5)/2\}$, $x_3 = k_m\{x_6/k_m\}$. If (x_5, x_6) passes the set $\{0, 1\} \times \{0, 1, \dots, k_{m+1} - 1\}$, then (x_2, x_3) passes k_{m+1}/k_m times the set $\{0, 1\} \times \{0, 1, \dots, k_m - 1\}$. Therefore, for both of the cases, if (x_5, x_6) passes the set $\{0, 1\} \times \{0, 1, \dots, k_{m+1} - 1\}$, then (x_2, x_3) passes k_{m+1}/k_m times the set $\{0, 1\} \times \{0, 1, \dots, k_m - 1\}$. Hence

$$\begin{aligned}
 (67) \quad & \frac{1}{(\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{\mathbf{c}^{(m+1)} \in (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}}} \sum_{s=1}^{t_m} \sum_{x_5=0}^1 \sum_{x_6=0}^{k_{m+1}-1} \\
 & \times \sum_{y_2 \in (t_m - s, t_m)} \frac{B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{b}^{(m)}, \mathbf{c}^{(m+1)})}{192K_1(s)p_1 t_m^2 k_{m+1} (m+1)^{2s+2}} \\
 & \leq \frac{(m+2)}{8(m+1)} \sum_{s=1}^{t_m} \sum_{\alpha=0}^{m+1} \sum_{x_2=0}^1 \sum_{x_3=0}^{k_m-1} \sum_{y_2 \in (t_m - s, t_m)} \\
 & \times \frac{p_1 A_2(\alpha, x_3, m, t_m - y_2, b_{y_2, x_2}^{(m)}, \dots, b_{y_2 + s - 1, x_2}^{(m)}, b_{y_2, x_2 + 1}^{(m)}, \dots, b_{y_2 + s - 1, x_2 + 1}^{(m)})}{24K_1(s)p_1 t_m^2 k_m (m+2)(m+1)^{2s+1}}.
 \end{aligned}$$

Bearing in mind (50), (52), and that $\mathbf{b}^{(m)} \notin \Omega_{m,3}$, we deduce that the left side of (67) is less than $(m+2)/(8(m+1)) \leq 1/4$. Then from (64) we obtain the desired result. $\square$

Now we choose vectors $\mathbf{a}^{(m)}$ ($m = 1, 2, \dots$) for the construction in (9) in the following way:

According to (53),

$$\# \left((\Delta_m^*(p_1 p_2))^{2t_m} \setminus \Omega_{m,3} \right) > 0.$$

We take $\mathbf{a}^{(1)}$ arbitrarily from the set $(\Delta_1^*(p_1 p_2))^{2t_1} \setminus \Omega_{1,3}$.

Taking into account (53), (59), and (65), we obtain that the set

$$(68) \quad F_{m+1,3} = (\Delta_{m+1}^*(p_1 p_2))^{2t_{m+1}} \setminus \left(\Omega_{m+1,3} \cup F_{m+1,1} \cup F_{m+1,2} \right).$$

is not empty. Let $\mathbf{a}^{(1)}, \dots, \mathbf{a}^{(m)}$ be chosen. Then we choose $\mathbf{a}^{(m+1)}$ arbitrarily so that

$$(69) \quad \mathbf{a}^{(m+1)} \in F_{m+1,3}.$$

The sequence of vectors $\mathbf{a}^{(m)}$ ($m = 1, 2, \dots$) is constructed inductively.

Next we fix $s \geq 1$, and we consider integers m such that $t_m \geq s$.

Main Lemma. *With the notation defined above, we have:*

$$(70) \quad \sum_{0 \leq x_2 \leq 1} \sum_{0 \leq x_3 < k_m} \sum_{0 \leq y_2 < t_m} B(x_2, x_3, y_2, s, \mathbf{a}^{(m)}) = O(m^{2s+3} \log^2 m),$$

$$(71) \quad \sum_{0 \leq x_3 < k_m} \sum_{0 \leq y_4 < t_{m+1}} B_1(x_3, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}) = O(m^{2s+3} \log^2 m),$$

$$(72) \quad \sum_{0 \leq x_5 \leq 1} \sum_{0 \leq x_6 < k_{m+1}} \sum_{y_2 \in (t_m - s, t_m)} B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}) \\ = O(m^{2s+3} \log^2 m),$$

where $m = 1, 2, \dots$, $x_2 = 2\{([x_6/k_m] + x_5 k_{m+1}/k_m)/2\}$, $x_3 = k_m\{x_6/k_m\}$, $y_4 = t_{m+1}\{(t_m(p_1^m - 1) + y_2)/t_{m+1}\}$, and $s \leq t_m$.

Proof. The proof follows from (7), (8), (36), (52), (58), (64), (68) and (69). $\square$

3. Proof of Theorem 1.

In this section, the integer s is fixed. For $m = 1, 2, \dots$, let

$$V_{m,1} = [0, 2k_m p_1^m) \times [t_{m-1} p_2^{m-1}, t_m p_2^m),$$

$$V_{m,2} = [2k_{m-1} p_1^{m-1}, 2k_m p_1^m) \times [1, t_m p_2^m),$$

$$(73) \quad V_m = V_{m,1} \cup V_{m,2}, \quad G_{M,N} = [0, M) \times [1, N),$$

$$(74) \quad G_1^{(m)} = G_{M,N} \cap V_{m,1}, \quad G_2^{(m)} = G_{M,N} \cap (V_{m,2} \setminus V_{m,1}),$$

and let

$$(75) \quad D_E = \#ED^{(s)}\left(\left(\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\}\right)_{(x,y) \in E}\right).$$

It is easy to see that $G_i^{(m)}$ is rectangular domain ($i = 1, 2$) and

$$D_{G_{M,N}} \leq \sum_{m \geq 1} \sum_{i=1,2} D_{G_i^{(m)}}.$$

Thus, to compute $D_{G_{M,N}}$, it is sufficient to find the estimate of D_F , where F is an arbitrary rectangular domain in V_m , $m = 1, 2, \dots$. According to (9), the analytic expression of $\{\alpha_y q^x\}$ depends on the position of (x, y) in V_m . We will consider three possibilities for the position of F in V_m : the middle, the right bourne and the upper bourne. Next we will consider sub-domains $F_{i,j} = \{(x, y) \in F \mid x \equiv i \pmod{2k_m}, y \equiv j \pmod{t_m}\}$ of rectangular domain F , and we will compute the discrepancy on the sub-domain $F_{i,j}$ for all $(i, j) \in [0, 2k_m) \times [0, t_m)$, with $m = 1, 2, \dots$.

First we obtain a simple expression for $\{\alpha_y q^x\}$, where (x, y) belongs to a middle domain of V_m :

Lemma 13. *Let $(x, y) \in V_m$; $x = 2k_m x_1 + k_m x_2 + x_3$, $y = t_m y_1 + y_2$, with $x_2 \in \{0, 1\}$, $x_3 \in \{0, \dots, k_m - 1\}$, $y_1 \in \{0, \dots, p_2^m - 1\}$, $y_2 \in \{0, \dots, t_m - 1\}$, $x_1 + x_2 \leq p_1^m - 1$, and $t_m > s$. Then*

$$(76) \quad \{\alpha_y q^x\}_{2k_m - x_3} = \left\{ \frac{q^{x_3} a_{y_2, x_2}^{(m)} (x_1 p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m - x_3} + \frac{1}{q^{k_m - x_3}} \left\{ \frac{a_{y_2, x_2 + 1}^{(m)} ((x_1 + x_2) p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m}.$$

Proof. From (8) we obtain

$$(77) \quad j \in [t_{r(j)-1} p_1^{r(j)-1}, t_{r(j)} p_1^{r(j)}), \quad j = 1, 2, \dots.$$

Let $(x, y) \in [0, 2k_m p_1^m] \times [t_{m-1} p_2^{m-1}, t_m p_2^m] = V_{m,1}$.

Then equation (8) implies that $r(y) = m$.

According to (9), we find that

$$(78) \quad \alpha_y q^x = q^x \sum_{n=0}^{p_1^m - 1} \sum_{\nu=0}^1 \frac{1}{q^{k_m(2n+\nu)}} \left\{ \frac{a_{y_2, \nu}^{(m)} (p_2^m n + p_1^m y_1)}{p_1^m p_2^m} \right\}_{k_m} + q^x \sum_{i=m+1}^{\infty} \sum_{n=p_1^{i-1} k_{i-1}/k_i}^{p_1^i - 1} \sum_{\nu=0}^1 \frac{1}{q^{k_i(2n+\nu)}} \left\{ \frac{a_{y_2(i), \nu}^{(i)} (p_2^i n + p_1^i y_1(i))}{p_1^i p_2^i} \right\}_{k_i},$$

where $y = t_m y_1 + y_2$; $y_2(i) \in \{0, 1, \dots, t_i - 1\}$, $y_2(i) \equiv y \pmod{t_i}$, and $y_1(i) = (y - y_2(i))/t_i$, $i = m+1, \dots$.

Bearing in mind that $k_i(2n+\nu) - x \geq 2k_{i-1} p_1^i - x \geq 2k_m p_1^m - x$ for $i \geq m+1$ and $n \geq p_1^{i-1} k_{i-1}/k_i$, we obtain that

$$\begin{aligned} & \{\alpha_y q^x\}_{2k_m - x_3} \\ &= \left\{ q^{2k_m x_1 + k_m x_2 + x_3} \sum_{n=0}^{p_1^m - 1} \sum_{\nu=0}^1 \frac{1}{q^{k_m(2n+\nu)}} \left\{ \frac{a_{y_2, \nu}^{(m)} (p_2^m n + p_1^m y_1)}{p_1^m p_2^m} \right\}_{k_m} \right\}_{2k_m - x_3}. \end{aligned}$$

Hence

$$(79) \quad \{\alpha_y q^x\}_{2k_m - x_3} = \left\{ q^{x_3} \left\{ \frac{a_{y_2, x_2}^{(m)} (p_2^m x_1 + p_1^m y_1)}{p_1^m p_2^m} \right\}_{k_m} + q^{x_3} \sum_{\substack{n=x_1 \text{ for } x_2=0 \\ n=x_1+1 \text{ for } x_2=1}} \right. \\ \left. \times \sum_{\substack{\nu=1, \text{ for } x_2=0 \\ \nu=0, \text{ for } x_2=1}} \frac{1}{q^{k_m(2n+\nu-2x_1-x_2)}} \left\{ \frac{a_{y_2, \nu}^{(m)} (p_2^m n + p_1^m y_1)}{p_1^m p_2^m} \right\}_{k_m} \right\}_{2k_m - x_3}$$

$$\begin{aligned}
&= \left\{ \left\{ \frac{q^{x_3} a_{y_2, x_2}^{(m)} (x_1 p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m - x_3} \right. \\
&\quad \left. + \frac{1}{q^{k_m - x_3}} \left\{ \frac{a_{y_2, x_2 + 1}^{(m)} ((x_1 + x_2) p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m} \right\}_{2k_m - x_3} \\
&= \left\{ \frac{q^{x_3} a_{y_2, x_2}^{(m)} (x_1 p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m - x_3} \\
&\quad + \frac{1}{q^{k_m - x_3}} \left\{ \frac{a_{y_2, x_2 + 1}^{(m)} ((x_1 + x_2) p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m}.
\end{aligned}$$

Now let $(x, y) \in [2k_{m-1}p_1^{m-1}, 2k_m p_1^m] \times [1, t_{m-1}p_2^{m-1}] = V_{m,2} \setminus V_{m,1}$. Then equation (77) and the conditions of the lemma show that $r(y) < m$. From (9) we get

$$\begin{aligned}
(80) \quad &\alpha_y q^x = f_{x,y} \\
&+ q^x \sum_{n=p_1^{m-1}k_{m-1}/k_m}^{p_1^m-1} \sum_{\nu=0}^1 \frac{1}{q^{k_m(2n+\nu)}} \left\{ \frac{a_{y_2, \nu}^{(m)} (p_2^m n + p_1^m y_1)}{p_1^m p_2^m} \right\}_{k_m} \\
&+ q^x \sum_{i=m+1}^{\infty} \sum_{n=p_1^{i-1}k_{i-1}/k_i}^{p_1^i-1} \sum_{\nu=0}^1 \frac{1}{q^{k_i(2n+\nu)}} \left\{ \frac{a_{y_2(i), \nu}^{(i)} (p_2^i n + p_1^i y_1(i))}{p_1^i p_2^i} \right\}_{k_i},
\end{aligned}$$

where $f_{x,y} \geq 0$ is an integer. It is easy to see that, here,

$$x \geq 2k_{m-1}p_1^{m-1} = \min_{n \geq p_1^{m-1}k_{m-1}/k_m, \nu=0,1} (k_m(2n+\nu)).$$

Hence we can repeat the calculations (77) - (79). Thus we obtain the desired result. $\square$

Define $G_1 =$

$$\begin{aligned}
(81) \quad G_1(m, x_2, x_3, y_2, K_1, K, L_1, L) = &\left\{ (x, y) \mid x = 2k_m x_1 + k_m x_2 + x_3, \right. \\
&\left. y = t_m y_1 + y_2, x_1 \in [K_1, K_1 + K), y_1 \in [L_1, L_1 + L] \right\},
\end{aligned}$$

with $x_2 \in \{0, 1\}$, $x_3 \in \{0, \dots, k_m - 1\}$ and $y_2 \in \{0, \dots, t_m - 1\}$.

Lemma 14. Let $G_1 \subset V_m$; $K, K_1, L, L_1 \geq 0$ be integers; $K_1 + K + x_2 \leq p_1^m$; $t_m(L_1 + L) + y_2 \leq t_m p_2^m - s$; $t_m > s$. Then

$$\begin{aligned}
(82) \quad \#G_1 D^{(s)} \left((\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\})_{(x,y) \in G_1} \right) \\
\leq 2^s B(x_2, x_3, y_2, s, \mathbf{a}^{(m)}) + s 2^{s+2}.
\end{aligned}$$

Proof. Let $(x, y) \in G_1$,

$$(83) \quad \theta_m(y_2, i) = \begin{cases} 1 & \text{if } y_2 + i \geq t_m \\ 0 & \text{otherwise,} \end{cases}$$

and let

$$a_{i+t_m, v+2}^{(m)} = a_{i, v}^{(m)}, \quad i, v = 0, 1, 2, \dots$$

(We continue periodically the coordinates of vector $\mathbf{a}^{(m)}$.) From Lemma 13 we have that

$$(84) \quad \{\alpha_{y+i} q^x\}_{2k_m-x_3} = \left\{ \frac{q^{x_3} a_{y_2+i, x_2}^{(m)} (x_1 p_2^m + (y_1 + \theta_m(y_2, i)) p_1^m)}{p_1^m p_2^m} \right\}_{k_m-x_3} \\ + \frac{1}{q^{k_m-x_3}} \left\{ \frac{a_{y_2+i, x_2+1}^{(m)} ((x_1 + x_2) p_2^m + (y_1 + \theta_m(y_2, i)) p_1^m)}{p_1^m p_2^m} \right\}_{k_m}, \\ i = 0, 1, \dots, s-1.$$

We denote the left side of (82) by σ . Applying Lemma 5 with $k = k^{(i)} = 2k_m - x_3$ and $q_i = q$ ($i = 1, \dots, s$), we get $\sigma \leq$

$$\#G_1 \left(\frac{s}{q^{2k_m-x_3}} + D^{(s)} \left((\{\alpha_y q^x\}_{2k_m-x_3}, \dots, \{\alpha_{y+s-1} q^x\}_{2k_m-x_3})_{(x,y) \in G_1} \right) \right).$$

Using Lemma 6 with $k = k^{(i)} = k_m - x_3$, $\ell = \ell^{(i)} = k_m$ and $q = q_i$ ($i = 1, \dots, s$), we obtain from (84)

$$(85) \quad \sigma \leq \#G_1 \left(\frac{2s}{q^{2k_m-x_3}} \right. \\ \left. + 2^s D^{(2s)} \left(\left(\left(\left\{ \frac{q^{x_3} a_{y_2+i, x_2}^{(m)} (x_1 p_2^m + (y_1 + \theta_m(y_2, i)) p_1^m)}{p_1^m p_2^m} \right\} \right) \right. \right. \right. \\ \left. \left. \left. \left\{ \frac{a_{y_2+i, x_2+1}^{(m)} ((x_1 + x_2) p_2^m + (y_1 + \theta_m(y_2, i)) p_1^m)}{p_1^m p_2^m} \right\} \right)_{0 \leq i \leq s-1} \right)_{(x,y) \in G_1} \right) \right).$$

Then Corollary 3 (with $M = K, M_1 = K_1, N = L + 1, N_1 = L_1, d = 2s$, and $r = m$) implies that

$$(86) \quad \sigma \leq \#G_1 \left(\frac{2s}{q^{2k_m-x_3}} + \frac{2s2^s}{p_1^m p_2^m} \right) + 2^s \sum_{h_{-1} \in C(p_1^m)} \sum_{h_{-2} \in C(p_2^m)} \\ \times \sum_{\mathbf{h} \in C_{2s}(p_1^m p_2^m)} r^{-1}(h_{-1}, p_1^m) r^{-1}(h_{-2}, p_2^m) r^{-1}(\mathbf{h}, p_1^m p_2^m) |\sigma_1(\mathbf{h})|,$$

where $\#G_1 = K(L+1) \leq p_1^m p_2^m$ and

$$\begin{aligned} \sigma_1(\mathbf{h}) = & \sum_{x_1=0}^{p_1^m-1} \sum_{y_1=0}^{p_2^m-1} e\left(\frac{h_{-1}x_1}{p_1^m} + \frac{h_{-2}y_1}{p_2^m}\right) \\ & + \sum_{i=0}^{s-1} \left(q^{x_3} h_i a_{y_2+i, x_2}^{(m)} (x_1 p_2^m + (y_1 + \theta_m(y_2, i)) p_1^m) \right. \\ & \left. + h_{s+i} a_{y_2+i, x_2+1}^{(m)} ((x_1 + x_2) p_2^m + (y_1 + \theta_m(y_2, i)) p_1^m) \right) / p_1^m p_2^m . \end{aligned}$$

It is easy to see that

$$h_{-1}x_1 p_2^m + h_{-2}y_1 p_1^m \equiv (x_1 p_2^m + y_1 p_1^m)(h_{-1}p_2^m M_2 + h_{-2}p_1^m M_1) \pmod{p_1^m p_2^m} ,$$

where $M_1 p_1^m \equiv 1 \pmod{p_2^m}$ and $M_2 p_2^m \equiv 1 \pmod{p_1^m}$. Taking into account that $x_1 p_2^m + y_1 p_1^m$ passes the complete residue system $\pmod{p_1^m p_2^m}$ and using Lemma 1, we have that

$$\begin{aligned} |\sigma_1(h)| = & p_1^m p_2^m \delta_{p_1^m p_2^m} \left(h_{-1} p_2^m M_2 + h_{-2} p_1^m M_1 + \sum_{i=0}^{s-1} (q^{x_3} h_i a_{y_2+i, x_2}^{(m)} \right. \\ & \left. + h_{s+i} a_{y_2+i, x_2+1}^{(m)}) \right) . \end{aligned}$$

Now from (24), (35) and (86) we find that

$$\sigma \leq (2sq^{x_3-2k_m} + s2^{s+1} p_1^{-m} p_2^{-m}) \#G_1 + 2^s B(x_2, x_3, y_2, s, \mathbf{a}^{(m)}) .$$

Applying (7), (81) and the condition of the lemma, we obtain the desired result. $\square$

We can now use Lemma 14 to compute the discrepancy of the considered double sequence in the rectangular domain $E \subset V_m$.

Lemma 15. *Let $K_1, K_6, L_1, L_6 \geq 0$ be integers, $E = [2k_m K_1, K_6] \times [t_m L_1, L_6] \subset V_m$, $2k_m K_1 < K_6 \leq k_m(2p_1^m - 1)$, $t_m L_1 \leq L_6 \leq t_m p_2^m - s$; $t_m > s$. Then*

$$(87) \quad D_E = O(m^{2s+3} \log^2 m) ,$$

where the constant implied by O only depends on s .

Proof. We consider following rectangular domains:

$$(88) \quad \begin{aligned} E_1 &= [2k_m K_1, K_5] \times [t_m L_1, L_5], & E_2 &= [2k_m K_1, K_5] \times [L_5, L_6], \\ E_3 &= [K_5, K_6] \times [L_5, L_6], & E_4 &= [K_5, K_6] \times [t_m L_1, L_5] , \end{aligned}$$

where $K, K_2, K_3, K_5, L, L_2, L_5 \geq 0$ are integers, and

$$(89) \quad \begin{aligned} K_2 &\in \{0, 1\}, K_3 \in [0, k_m); K_1, K, L, L_1 \geq 0, L_2 \in [0, t_m), \\ K &= [K_6/2k_m] - K_1, K_5 = 2k_m(K_1 + K), \\ K_6 &= K_5 + k_m K_2 + K_3, \\ L &= [L_6/t_m] - L_1, L_5 = t_m L_1 + t_m L, L_6 = L_5 + L_2. \end{aligned}$$

It is evident that

$$E = E_1 \cup E_2 \cup E_3 \cup E_4, \quad \text{and} \quad E_i \cap E_j = \emptyset \quad \text{for} \quad i \neq j.$$

Using (2) and (75), we obtain

$$(90) \quad D_E \leq \sum_{1 \leq i \leq 4} D_{E_i}.$$

From (81) we obtain :

$$(91) \quad \begin{aligned} E_1 &= \bigcup_{\substack{0 \leq x_2 \leq 1 \\ 0 \leq x_3 < k_m \\ 0 \leq y_2 < t_m}} G_1(m, x_2, x_3, y_2, K_1, K, L_1, L), \\ E_2 &= \bigcup_{\substack{0 \leq x_2 \leq 1 \\ 0 \leq x_3 < k_m \\ 0 \leq y_2 < L_2}} G_1(m, x_2, x_3, y_2, K_1, K, L_1 + L, 0), \\ E_4 &= \bigcup_{\substack{0 \leq x_2 k_m + x_3 < K_2 k_m + K_3 \\ 0 \leq y_2 < t_m}} G_1(m, x_2, x_3, y_2, K_1 + K, 1, L_1, L). \end{aligned}$$

We obtain from (89), (88), (7), and (8) that

$$\#E_3 \leq 2k_m t_m = O(m \log m).$$

Applying (75), (70), and Lemma 14, we get from (7) and (91) that

$$\begin{aligned} D_{E_1} &\leq \sum_{0 \leq x_2 \leq 1} \sum_{0 \leq x_3 < k_m} \sum_{0 \leq y_2 < t_m} \#G_1(m, x_2, x_3, y_2, K_1, K, L_1, L) \\ &\quad \times D^{(s)}\left(\left(\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\}\right)_{(x,y) \in G_1(m, x_2, x_3, y_2, K_1, K, L_1, L)}\right) \\ &\leq 2^s \sum_{0 \leq x_2 \leq 1} \sum_{0 \leq x_3 < k_m} \sum_{0 \leq y_2 < t_m} \left(B(x_2, x_3, y_2, s, \mathbf{a}^{(m)}) + 4s\right) \\ &= O(m^{2s+3} \log^2 m). \end{aligned}$$

Similarly, estimates are valid for the cases of sets E_2 and E_4 . Thus

$$\max_{1 \leq i \leq 4} D_{E_i} = O(m^{2s+3} \log^2 m).$$

Now from (90) we obtain the assertion of the lemma. $\square$

Consider the right bourne of V_m . Define

$$(92) \quad G_2 = G_2(m, x_3, y_4, L_2, L_3) = \{(x, y) \mid x = k_m(2p_1^m - 1) + x_3, \\ y = t_{m+1}y_3 + y_4, y_3 \in [L_2, L_2 + L_3] \subset [0, p_2^m]\}.$$

Lemma 16. *Let $G_2 \subset V_m$; $L_2, L_3 \geq 0$ be integers; $t_{m+1}(L_2 + L_3) + y_4 \leq t_m p_2^m - s$; $y_4 \in [0, t_{m+1})$, $x_3 \in [0, k_m)$, $t_m > s$. Then*

$$(93) \quad \#G_2 D^{(s)}\left(\left(\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\}\right)_{(x,y) \in G_2}\right) \\ \leq 2^s B_1(x_3, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}) + s2^{s+3}.$$

Proof. Let $(x, y) \in G_2$. Thus $y < t_m p_2^m$, and from (77) we find that $r(y) \leq m$. Similarly to (78) and (80) we have that

$$\{\alpha_y q^x\} = \left\{ q^x \sum_{n=\delta_m^{(y)} p_1^{m-1} k_{m-1}/k_m}^{p_1^m-1} \sum_{\nu=0}^1 \frac{1}{q^{k_m(2n+\nu)}} \left\{ \frac{a_{y_2, \nu}^{(m)}(p_2^m n + p_1^m y_1)}{p_1^m p_2^m} \right\}_{k_m} \right. \\ \left. + q^x \sum_{i=m+1}^{\infty} \sum_{n=p_1^{i-1} k_{i-1}/k_i}^{p_1^i-1} \sum_{\nu=0}^1 \frac{1}{q^{k_i(2n+\nu)}} \left\{ \frac{a_{y_2(i), \nu}^{(i)}(p_2^i n + p_1^i y_1(i))}{p_1^i p_2^i} \right\}_{k_i} \right\}.$$

Hence

$$\{\alpha_y q^{2k_m(p_1^m-1)+k_m+x_3}\} = \left\{ q^{x_3} \left\{ \frac{a_{y_2, 1}^{(m)}((p_1^m-1)p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m} \right. \\ \left. + \frac{1}{q^{k_m-x_3}} \left\{ \frac{a_{y_4, 0}^{(m+1)}(n_m p_2^{m+1} + y_3 p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\}_{k_{m+1}} + \frac{\varepsilon_{x,y}}{q^{k_{m+1}+k_m-x_3}} \right\},$$

with $y_2 = t_m\{y_4/t_m\}$, $y_1 = y_3 t_{m+1}/t_m + [y_4/t_m]$, $n_m = p_1^m k_m/k_{m+1}$, and $\varepsilon_{x,y} \in [0, 1)$.

Therefore

$$(94) \quad \{\alpha_y q^x\}_{k_{m+1}+k_m-x_3} = \left\{ \frac{q^{x_3} a_{y_2, 1}^{(m)}(x_0 + y_1 p_1^m)}{p_1^m p_2^m} \right\}_{k_m-x_3} \\ + \frac{1}{q^{k_m-x_3}} \left\{ \frac{a_{y_4, 0}^{(m+1)}(n_m p_2^{m+1} + y_3 p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\}_{k_{m+1}},$$

where $x = 2k_m(p_1^m - 1) + k_m + x_3$, and $x_0 = (p_1^m - 1)p_2^m$. We have for $i \in [0, s-1]$ that $(x, y+i) \in V_m$, and that $r(y+i) \leq m$. Thus we can

apply (94) for the pair $(x, y + i)$. Bearing in mind (83), we obtain

$$(95) \quad \{\alpha_{y+i}q^x\}_{k_{m+1}+k_m-x_3} = \left\{ \frac{q^{x_3}a_{y_4+i,1}^{(m)}(x_0 + (y_1 + \theta_m(y_2, i))p_1^m)}{p_1^m p_2^m} \right\}_{k_m-x_3} \\ + \frac{1}{q^{k_m-x_3}} \left\{ \frac{a_{y_4+i,0}^{(m+1)}(n_m p_2^{m+1} + (y_3 + \theta_{m+1}(y_4, i))p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\}_{k_{m+1}},$$

with $y_2 = t_m\{y_4/t_m\}$, $y_1 = y_3 t_{m+1}/t_m + [y_4/t_m]$, $x = 2k_m(p_1^m - 1) + k_m + x_3$, and $x_0 = (p_1^m - 1)p_2^m$. We denote the left side of (93) by σ . Applying Lemma 5 with $k = k^{(i)} = k_{m+1} + k_m - x_3$ and $q_i = q$ ($i = 1, \dots, s$), we obtain from (19)

$$\sigma \leq \#G_2 \left(\frac{s}{q^{k_{m+1}+k_m-x_3}} + D^{(s)} \left(\left(\{\alpha_y q^x\}_{k_{m+1}+k_m-x_3}, \dots, \{\alpha_{y+s-1} q^x\}_{k_{m+1}+k_m-x_3} \right)_{(x,y) \in G_2} \right) \right).$$

Using Lemma 6 with $k = k^{(i)} = k_m - x_3$, $\ell^{(i)} = k_m$ and $q_i = q$ ($i = 1, \dots, s$), we obtain from (95)

$$\sigma \leq \#G_2 \left(\frac{2s}{q^{k_{m+1}+k_m-x_3}} + 2^s D^{(2s)} \left(\left(\left(\left\{ \frac{q^{x_3}a_{y_4+i,1}^{(m)}(x_0 + (y_3 t_{m+1}/t_m + [y_4/t_m] + \theta_m(y_2, i))p_1^m)}{p_1^m p_2^m} \right\}, \right. \right. \right. \\ \left. \left. \left\{ \frac{a_{y_4+i,0}^{(m+1)}(n_m p_2^{m+1} + (y_3 + \theta_{m+1}(y_4, i))p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\} \right)_{0 \leq i \leq s-1} \right)_{L_2 \leq y_3 \leq L_2+L_3} \right) \right).$$

Here we now replace fractions with denominators $p_1^m p_2^m$ and $p_1^{m+1} p_2^{m+1}$ with fractions with denominator p_2^{m+1} , and we again apply Lemma 5 with the parameters $2s$ instead of s , p_2 instead of q_i , and $m+1$ instead of $k^{(i)}$ ($i = 1, \dots, 2s$):

$$(96) \quad \sigma \leq \#G_2 \left(\frac{2s}{q^{k_{m+1}+k_m-x_3}} + \frac{2s \cdot 2^s}{p_2^{m+1}} + 2^s D^{(2s)} \left(\left(\left(\left\{ \frac{p_2 t_{m+1}/t_m y_3 q^{x_3} a_{y_4+i,1}^{(m)} + d_i}{p_2^{m+1}} \right\}, \right. \right. \right. \\ \left. \left. \left\{ \frac{y_3 a_{y_4+i,0}^{(m+1)} + f_i}{p_2^{m+1}} \right\} \right)_{0 \leq i < s} \right)_{L_2 \leq y_3 \leq L_2+L_3} \right) \right),$$

where $d_i = [p_2^{m+1} \{q^{x_3} a_{y_4+i,1}^{(m)} (-p_1^{-m} + ([y_4/t_m])\} + \theta_m(y_2, i)) p_2]$, and $f_i = [p_2^{m+1} \{a_{y_4+i,0}^{(m+1)} n_m p_1^{-m-1}\} + \theta_{m+1}(y_4, i)]$, $i = 0, \dots, s-1$.

Then Corollary 2 (with $d = 2s, T = P = p_2^{m+1}, N = \#G_2 = L_3 + 1$) implies that

$$\begin{aligned} \sigma &\leq \#G_2 \left(\frac{2s}{q^{k_{m+1}+k_m-x_3}} + \frac{4s2^s}{p_2^{m+1}} \right) \\ &+ 2^s \sum_{h_{-2} \in C(p_2^{m+1})} \sum_{\mathbf{h} \in C_{2s}(p_2^{m+1})} r^{-1}(h_{-2}, p_2^{m+1}) r^{-1}(\mathbf{h}, p_2^{m+1}) \left| \sum_{y_3=0}^{p_2^{m+1}-1} e\left(\frac{1}{p_2^{m+1}} \right. \right. \\ &\times \left. \left. \left(h_{-2}y_3 + \sum_{i=0}^{s-1} (h_i(p_2 t_{m+1}/t_m y_3 q^{x_3} a_{y_4+i,1}^{(m)} + d_i) + h_{s+i}(y_3 a_{y_4+i,0}^{(m+1)} + f_i)) \right) \right) \right|. \end{aligned}$$

Taking into account (7), (54), (92), Lemma 1, and the conditions of the lemma we obtain :

$$\begin{aligned} \sigma &\leq s2^{s+3} + 2^s \sum_{h_{-2} \in C(p_2^{m+1})} \sum_{\mathbf{h} \in C_{2s}(p_2^{m+1})} r^{-1}(h_{-2}, p_2^{m+1}) r^{-1}(\mathbf{h}, p_2^{m+1}) \\ &\times p_2^{m+1} \delta_{p_2^{m+1}} \left(h_{-2} + \sum_{i=0}^{s-1} (p_2 t_{m+1}/t_m q^{x_3} h_i a_{y_4+i,1}^{(m)} + h_{s+i} a_{y_4+i,0}^{(m+1)}) \right) \\ &\leq s2^{s+3} + 2^s B_1(x_3, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}) . \end{aligned}$$

Thus we obtain the assertion of the lemma. $\square$

Corollary 7. Let $E = [k_m(2p_1^m - 1), K_6] \times [t_m L_1, L_6] \subset V_m$; $0 \leq t_m L_1 \leq L_6 \leq t_m p_2^m - s$, $k_m(2p_1^m - 1) < K_6 \leq 2k_m p_1^m$, $t_m \geq s$. Then

$$(97) \quad D_E = O(m^{2s+3} \log^2 m) .$$

Proof. Let $K_4 = k_m(2p_1^m - 1)$, $L_2 = [t_m L_1/t_{m+1}] + 1$, $L_3 = [L_6/t_{m+1}] - L_2 - 1$. It is easy to see that

$$(98) \quad t_{m+1}(L_2 + L_3) + 1 < L_6 \leq t_m p_2^m - s .$$

If $L_3 \leq 0$, then $D_E \leq 4k_m t_{m+1} = O(m \log m)$. Let $L_3 > 0$, and let $E_1 = [K_4, K_6] \times [t_m L_1, t_{m+1} L_2]$, $E_2 = [K_4, K_6] \times [t_{m+1} L_2, t_{m+1}(L_2 + L_3)]$, $E_3 = [K_4, K_6] \times (t_{m+1}(L_2 + L_3), L_6]$.

It is evident that

$$E = E_1 \cup E_2 \cup E_3, \quad \text{and} \quad E_i \cap E_j = \emptyset \quad \text{for} \quad i \neq j .$$

Using (2) and (75), we find that

$$(99) \quad D_E \leq D_{E_1} + D_{E_2} + D_{E_3} .$$

From (7) and (8) we obtain :

$$(100) \quad D_{E_i} \leq \#E_i \leq 2k_m t_{m+1} = O(m^2), \quad i = 1, 3 .$$

It follows from (75) that

$$D_{E_2} \leq \sum_{0 \leq x_3 < z} \sum_{0 \leq y_4 < t_{m+1}} \#G_2 D^{(s)} \left((\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\})_{(x,y) \in G_2} \right),$$

where $G_2 = G_2(m, x_3, y_4, L_2, L_3)$, and $z = K_6 - K_4 \leq k_m$. Bearing in mind that (98) is true, we can apply Lemma 16:

$$D_{E_2} \leq \sum_{0 \leq x_3 < k_m} \sum_{0 \leq y_4 < t_{m+1}} (2^s B_1(x_3, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}) + s 2^{s+3})$$

Now from (71), (99), and (100) we obtain the assertion of the corollary. $\square$

We now consider the upper bourne of V_m :

Lemma 17. *Let $K_2, K_3 \geq 0$, $K_2 + K_3 \leq p_1^{m-1} - 2$, $x_5 \in \{0, 1\}$, $x_6 \in \{0, 1, \dots, k_{m+1} - 1\}$, $y_2 \in (t_m - s, t_m)$, $t_m > s$,*

$$(101) \quad G_3 = G_3(m, x_5, x_6, y_2, K_2, K_3) = \{(x, y) \mid x = 2k_{m+1}x_4 + k_{m+1}x_5 + x_6, y = t_m(p_2^m - 1) + y_2, K_2 \leq x_4 < K_2 + K_3\} \subset V_m,$$

for $m = 1, 2, \dots$

Then

$$(102) \quad \#G_3 D^{(s)} \left((\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\})_{(x,y) \in G_3} \right) \leq s 2^{s+3} + 2^s B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}),$$

where $x_2 = 2\{([x_6/k_m] + x_5 k_{m+1}/k_m)/2\}$, $x_3 = k_m \{x_6/k_m\}$, and $y_4 = t_{m+1} \{(t_m(p_2^m - 1) + y_2)/t_{m+1}\}$.

Proof. Let $(x, y) \in G_3$ and let $i \in [0, t_m - y_2)$. The equality (83) and the conditions of the lemma show, that $\theta_m(y_2, i) = 0$. The pair $(x, y + i)$ satisfies the conditions of Lemma 13. The equation (84) implies that

$$(103) \quad \{\alpha_{y+i} q^x\}_{2k_m - x_3} = \left\{ \frac{q^{x_3} a_{y_2+i, x_2}^{(m)} (x_1 p_2^m + (p_2^m - 1) p_1^m)}{p_1^m p_2^m} \right\}_{k_m - x_3} + \frac{1}{q^{k_m - x_3}} \left\{ \frac{a_{y_2+i, x_2+1}^{(m)} ((x_1 + x_2) p_2^m + (p_2^m - 1) p_1^m)}{p_1^m p_2^m} \right\}_{k_m},$$

where

$$(104) \quad \begin{aligned} x &= 2k_{m+1}x_4 + k_{m+1}x_5 + x_6 = 2k_m x_1 + k_m x_2 + x_3, \\ x_3 &= k_m \{x_6/k_m\} \in \{0, 1, \dots, k_m - 1\}, \\ x_1 &= x_4 k_{m+1}/k_m + [(k_{m+1}x_5 + x_6)/k_m], \\ x_2 &\in \{0, 1\}, \quad x_2 \equiv [x_6/k_m] + x_5 k_{m+1}/k_m \pmod{2}. \end{aligned}$$

Now let $i \in [t_m - y_2, s - 1]$. Then $y + i > t_m p_2^m$, and $r(y + i) = m + 1$ (see (8) and (77)). The pair $(x, y + i)$ satisfies the conditions of Lemma 13 (with $m + 1$ instead of m). Hence, we have from (84) and (104) that

$$(105) \quad \{\alpha_{y+i} q^x\}_{2k_{m+1}-x_6} \\ = \left\{ \frac{q^{x_6} a_{y_4+i, x_5}^{(m+1)} (x_4 p_2^{m+1} + (y_3 + \theta_{m+1}(y_4, i)) p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\}_{k_{m+1}-x_6} \\ + \frac{1}{q^{k_{m+1}-x_6}} \left\{ \frac{a_{y_4+i, x_5+1}^{(m+1)} ((x_4 + x_5) p_2^{m+1} + (y_3 + \theta_{m+1}(y_4, i)) p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\}_{k_{m+1}},$$

where

$$y = t_m(p_1^m - 1) + y_2 = t_{m+1}y_3 + y_4, \quad y_4 \in \{0, 1, \dots, t_{m+1} - 1\},$$

$$y_4 \equiv t_m(p_1^m - 1) + y_2 \pmod{t_{m+1}}, \quad y_3 = (t_m(p_1^m - 1) + y_2 - y_4)/t_{m+1}.$$

It is easy to see that if $k_{m+1} = k_m$, then $x_3 = x_6$. Otherwise $k_{m+1} \geq p_1 k_m \geq 2k_m$, and $2k_{m+1} - x_6 \geq k_{m+1} \geq 2k_m \geq 2k_m - x_3$.

We denote the left side of (102) by σ . Applying Lemma 5 with $k = 2k_m - x_3$, $k^{(i)} = 2k_m - x_3$ for $i \in [0, t_m - y_2]$; $k^{(i)} = 2k_{m+1} - x_6$ for $i \in [t_m - y_2, s - 1]$, and $q_i = q$, for $i = 1, \dots, s$, we get

$$\sigma \leq \#G_3 \left(\frac{s}{q^{2k_m - x_3}} + D^{(s)} \left(\left((\{\alpha_{y+i} q^x\}_{k^{(i)}})_{i \in [0, s-1]} \right)_{(x, y) \in G_3} \right) \right).$$

Using Lemma 6 with $k = 2k_m - x_3$, $k^{(i)} = 2k_m - x_3$, $\ell^{(i)} = k_m$ for $i \in [0, t_m - y_2]$, and $k^{(i)} = 2k_{m+1} - x_6$, $\ell^{(i)} = k_{m+1}$ for $i \in [t_m - y_2, s - 1]$, we obtain from (103), and (105)

$$\sigma \leq \#G_3 \left(\frac{2s}{q^{2k_m - x_3}} + 2^s D^{(2s)} \left(\left(\left(\left\{ \frac{q^{x_3} a_{y_2+i, x_2}^{(m)} (x_1 p_2^m + (p_2^m - 1) p_1^m)}{p_1^m p_2^m} \right\}, \right. \right. \right. \right. \\ \left. \left. \left\{ \frac{a_{y_2+i, x_2+1}^{(m)} ((x_1 + x_2) p_2^m + (p_2^m - 1) p_1^m)}{p_1^m p_2^m} \right\} \right)_{i \in [0, t_m - y_2]}, \right. \\ \left. \left(\left\{ \frac{q^{x_6} a_{y_4+i, x_5}^{(m+1)} (x_4 p_2^{m+1} + (y_3 + \theta_{m+1}(y_4, i)) p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\}, \right. \right. \\ \left. \left. \left\{ \frac{a_{y_4+i, x_5+1}^{(m+1)} ((x_4 + x_5) p_2^{m+1} + (y_3 + \theta_{m+1}(y_4, i)) p_1^{m+1})}{p_1^{m+1} p_2^{m+1}} \right\} \right)_{i \in [t_m - y_2, s-1]} \right)_{K_2 \leq x_4 < K_2 + K_3} \right),$$

where $x_1 = x_4 k_{m+1}/k_m + [(k_{m+1} x_5 + x_6)/k_m]$.

Here we now replace fractions with denominators $p_1^m p_2^m$ and $p_1^{m+1} p_2^{m+1}$ with fractions with denominator p_1^{m+1} , and we again apply Lemma 5 with

the parameters $2s$ instead of s , p_1 instead of q_i , and $m+1$ instead of $k^{(i)}$ ($i = 1, \dots, 2s$):

$$\sigma \leq \#G_3 \left(\frac{2s}{q^{2k_m - x_3}} + \frac{2s2^s}{p_1^{m+1}} + 2^s D^{(2s)} \left(\left(\left(\left\{ \frac{p_1 q^{x_3} a_{y_2+i, x_2}^{(m)} x_4 k_{m+1}/k_m + d_i}{p_1^{m+1}} \right\}, \right. \right. \right. \right. \\ \left. \left. \left\{ \frac{p_1 a_{y_2+i, x_2+1}^{(m)} x_4 k_{m+1}/k_m + d'_i}{p_1^{m+1}} \right\} \right)_{i \in [0, t_m - y_2]}, \left(\left\{ \frac{q^{x_6} a_{y_4+i, x_5}^{(m+1)} x_4 + f_i}{p_1^{m+1}} \right\}, \right. \right. \\ \left. \left. \left\{ \frac{a_{y_4+i, x_5+1}^{(m+1)} x_4 + f'_i}{p_1^{m+1}} \right\} \right)_{i \in [t_m - y_2, s-1]} \right)_{K_2 \leq x_4 < K_2 + K_3} \right),$$

where for $i = 0, \dots, t - y_2 - 1$

$$d_i = [q^{x_3} a_{y_2+i, x_2}^{(m)} (p_1 [(k_{m+1} x_5 + x_6)/k_m] - p_1^{m+1} p_2^{-m})],$$

$$d'_i = [a_{y_2+i, x_2+1}^{(m)} (p_1 (x_2 + [(k_{m+1} x_5 + x_6)/k_m]) - p_1^{m+1} p_2^{-m})],$$

and for $i = t - y_2, \dots, s - 1$

$$f_i = [(p_1/p_2)^{m+1} q^{x_6} a_{y_4+i, x_5}^{(m+1)} (y_3 + \theta_{m+1}(y_4, i))],$$

$$f'_i = [a_{y_4+i, x_5+1}^{(m+1)} ((p_1/p_2)^{m+1} (y_3 + \theta_{m+1}(y_4, i)) + x_5)].$$

Then Corollary 2 (with $N = \#G_3 = K_3$, $T = P = p_1^{m+1}$, and $d = 2s$) implies that $\sigma \leq$

$$\#G_3 \left(\frac{2s}{q^{k_m}} + \frac{4s2^s}{p_1^{m+1}} \right) + 2^s \sum_{h_{-1} \in C(p_1^{m+1})} \sum_{\mathbf{h} \in C_{2s}(p_1^{m+1})} r^{-1}(h_{-1}, p_1^{m+1}) r^{-1}(\mathbf{h}, p_1^{m+1}) \\ \times \left| \sum_{x_4=0}^{p_1^{m+1}-1} e \left(\left(h_{-1} x_4 + \sum_{0 \leq i < t_m - y_2} (x_4 p_1 k_{m+1}/k_m q^{x_3} a_{y_2+i, x_2}^{(m)} + d_i) h_i \right. \right. \right. \\ \left. \left. + \sum_{0 \leq i < t_m - y_2} (x_4 p_1 k_{m+1}/k_m a_{y_2+i, x_2+1}^{(m)} + d'_i) h_{s+i} \right. \right. \\ \left. \left. + \sum_{t_m - y_2 \leq i < s} (q^{x_6} a_{y_4+i, x_5}^{(m+1)} + f_i) h_i + \sum_{t_m - y_2 \leq i < s} (x_4 a_{y_4+i, x_5+1}^{(m+1)} + f'_i) h_{s+i} \right) / p_1^{m+1} \right|.$$

Using Lemma 1, (7), (101), and the conditions of the lemma, we find that

$$\sigma \leq 8s2^s + 2^s \sum_{h_{-1} \in C(p_1^{m+1})} \sum_{\mathbf{h} \in C_{2s}(p_1^{m+1})} r^{-1}(h_{-1}, p_1^{m+1}) r^{-1}(\mathbf{h}, p_1^{m+1}) p_1^{m+1} \\ \times \delta_{p_1^{m+1}} \left(h_{-1} + \sum_{0 \leq i < t_m - y_2} (q^{x_3} h_i a_{y_2+i, x_2}^{(m)} + h_{s+i} a_{y_2+i, x_2+1}^{(m)}) p_1 k_{m+1}/k_m \right. \\ \left. + \sum_{t_m - y_2 \leq i < s} (q^{x_6} h_i a_{y_4+i, x_5}^{(m+1)} + h_{s+i} a_{y_4+i, x_5+1}^{(m+1)}) \right).$$

We obtain from (61) that

$$\sigma \leq s2^{s+3} + 2^s B_2(x_2, x_3, x_5, x_6, y_2, y_4, s, \mathbf{a}^{(m)}, \mathbf{a}^{(m+1)}) .$$

Thus we derive the desired result. $\square$

Corollary 8. *Let $E = [2k_m K_1, K_6] \times [t_m p_2^m - s + 1, L_6] \subset V_m$; $2k_m K_1 < K_6 \leq k_m(2p_1^m - 1)$, $t_m p_2^m - s + 1 < L_6 \leq t_m p_2^m$, $t_m > s$. Then*

$$(106) \quad D_E = O(m^{2s+3} \log^2 m) .$$

Proof. Let $L_4 = t_m p_2^m - s + 1$, $K_2 = [k_m K_1 / k_{m+1}] + 1$, $K_3 = [K_6 / k_{m+1}] - K_2 - 1$. It is easy to see that

$$(107) \quad k_{m+1}(K_2 + K_3 + 1) - 1 < K_6 \leq k_m(2p_1^m - 1).$$

If $K_3 \leq 0$, then

$$D_E \leq 2sk_{m+1} = O(m).$$

Now let $K_3 > 0$. Put

$$\begin{aligned} E_1 &= [2k_m K_1, 2k_{m+1} K_2] \times [L_4, L_6], \\ E_2 &= [2k_{m+1} K_2, 2k_{m+1}(K_2 + K_3)] \times [L_4, L_6], \\ E_3 &= [2k_{m+1}(K_2 + K_3), K_6] \times [L_4, L_6]. \end{aligned}$$

It is evident that

$$E = E_1 \cup E_2 \cup E_3, \quad \text{and} \quad E_i \cap E_j = \emptyset \quad \text{for} \quad i \neq j .$$

Using (2) and (75), we find that

$$(108) \quad D_E \leq D_{E_1} + D_{E_2} + D_{E_3} .$$

From (7) and (8) we obtain :

$$(109) \quad D_{E_i} \leq \#E_i \leq 4sk_{m+1} = O(m), \quad i = 1, 3 .$$

It follows from (75) and (101), that

$$\begin{aligned} D_{E_2} &\leq \sum_{0 \leq x_5 \leq 1} \sum_{0 \leq x_6 < k_{m+1}} \sum_{y_2 \in (t_m - s, t_m)} \#G_3 \\ &\quad \times D^{(s)}\left(\left(\{\alpha_y q^x\}, \dots, \{\alpha_{y+s-1} q^x\}\right)_{(x,y) \in G_3}\right), \end{aligned}$$

where $G_3 = G_3(m, x_5, x_6, y_2, K_2, K_3)$.

Because (107), we can apply Lemma 17. Thus from (72), (108), and (109), we obtain (106). $\square$

Now, combining (87), (97), and (106) we obtain :

Lemma 18. *Let $K_1, K_6, L_1, L_6 \geq 0$ be integers and $E = [2k_m K_1, K_6] \times [t_m L_1, L_6] \subset V_m$. Then*

$$D_E = O(m^{2s+3} \log^2 m),$$

where the constant implied by O only depends on s .

Proof. Let

$$\begin{aligned} E_1 &= [2k_m K_1, k_m(2p_1^m - 1)) \times [t_m L_1, t_m p_2^m - s], \\ E_2 &= [k_m(2p_1^m - 1), 2k_m p_1^m) \times [t_m L_1, t_m p_2^m - s], \\ E_3 &= [2k_m K_1, k_m(2p_1^m - 1)) \times [t_m p_2^m - s + 1, t_m p_2^m), \\ E_4 &= [k_m(2p_1^m - 1), 2k_m p_1^m) \times [t_m p_2^m - s + 1, t_m p_2^m), \end{aligned}$$

and let

$$E'_i = E \cap E_i, \quad i = 1, \dots, 4.$$

It is easy to see that

$$(110) \quad E = \cup_{i=1}^4 E'_i, \quad \text{and} \quad E'_i \cap E'_j = \emptyset \quad \text{for} \quad i \neq j.$$

Now equations (7) and (75) imply that

$$D_{E'_4} \leq sk_m = O(m).$$

From Lemma 15, Corollary 7, and Corollary 8, we get

$$D_{E'_i} = O(m^{2s+3} \log^2 m), \quad i = 1, 2, 3.$$

From (110), we obtain the assertion of the lemma. $\square$

End of the proof of Theorem 1. We use notations (73), (74), and (75).

Let

$$(111) \quad G^{(m)} = G_{M,N} \cap V_m \neq \emptyset \quad \text{for} \quad m = 1, \dots, r \quad \text{and} \quad G^{(r+1)} = \emptyset.$$

It is evident that

$$G^{(m)} = G_1^{(m)} \cup G_2^{(m)}, \quad m = 1, 2, \dots \quad G^{(m)} \cap G^{(n)} = \emptyset \quad \text{for} \quad m \neq n,$$

and $G_i^{(m)}$ is the rectangular domain ($i = 1, 2$). Let $(K_\nu^{(m,i)}, L_\mu^{(m,i)})_{\nu, \mu=1,2}$ be coordinates of the vertex of $G_i^{(m)}$:

$$G_i^{(m)} = [K_1^{(m,i)}, K_2^{(m,i)}) \times [L_1^{(m,i)}, L_2^{(m,i)}) \quad i = 1, 2, \quad m = 1, 2, \dots$$

From (7), (8), and (48), we see that

$$k_{m-1} p_1^{m-1} \equiv 0 \pmod{k_m} \quad \text{and} \quad t_{m-1} p_2^{m-1} \equiv 0 \pmod{t_m} \quad m = 2, 3, \dots$$

Hence

$$K_1^{(m,i)} \equiv 0 \pmod{2k_m} \quad \text{and} \quad L_1^{(m,i)} \equiv 0 \pmod{t_m}, \quad m = 2, 3, \dots$$

Applying Lemma 18, we obtain

$$D_{G_i^{(m)}} = O(m^{2s+3} \log^2 m), \quad i = 1, 2, \quad m = 2, 3, \dots$$

Using (2), (75), and (111), we find that

$$\begin{aligned} MND\left(\left(\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\}\right)_{1 \leq n \leq N, 0 \leq m < M}\right) &= D_{G_{M,N}} \\ &\leq \sum_{m=1}^r \sum_{i=1}^2 D_{G_i^{(m)}} = O\left(\sum_{m=1}^r m^{2s+3} \log^2 m\right) = O(r^{2s+4} \log^2 r) \\ &= O((\log MN)^{2s+4} \log^2 \log MN). \end{aligned}$$

Thus we obtain the assertion of Theorem 1. $\square$

Remark. Using (76), (95), and (103) we can find specifically digits $d_{i,j}$ ($i, j = 1, 2, \dots$) of a normal lattice configuration (see (5)). With the notations defined above, if $(x, y) = (2k_m x_1 + k_m x_2 + x_3, t_m y_1 + y_2) \in V_m$, then

$$d_{x,y} = \left[q \left\{ \frac{q^{x_3} a_{y_2, x_2}^{(m)} (x_1 p_2^m + y_1 p_1^m)}{p_1^m p_2^m} \right\} \right].$$

4. Proof of Theorem 2

Let

$$\begin{aligned} (112) \quad A_3(j, m, \nu, c_{1,0}, \dots, c_{s,0}, c_{1,1}, \dots, c_{s,1}) &= p^m \sum_{h_0 \in C(p^m)} \sum_{\mathbf{h} \in C_{2s}(p^m)} \\ &\times r^{-1}(h_0, p^m) r^{-1}(\mathbf{h}, p^m) \delta_{p^m} \left(h_0 + \sum_{i=1}^s (q_i^j h_i c_{i,\nu} + h_{i+s} c_{i,\nu+1}) \right), \end{aligned}$$

$$\begin{aligned} (113) \quad A_4(j, m, \nu, c_{1,0}, \dots, c_{s,0}, c_{1,1}, \dots, c_{s,1}) \\ = p^m \sum_{\mathbf{h} \in C_{2s}(p^m)} r^{-1}(\mathbf{h}, p^m) \delta_{p^m} \left(\sum_{i=1}^s (q_i^j h_i c_{i,\nu} + h_{i+s} c_{i,\nu+1}) \right), \end{aligned}$$

with $c_{i,\nu+2} = c_{i,\nu}$, $\nu = 0, 1$, $i = 1, \dots, s$.

Lemma 19. Let $\nu \in \{0, 1\}$, $j \in [0, k_m)$, $\varphi_0 = \varphi(p)/p$, and

$$K(s) = 4p/\varphi(p) \left(\frac{2}{\pi} \log p + \frac{7}{5} \right)^{2s+1}.$$

Then

$$(114) \quad \frac{1}{\varphi_0^{2s} p^{2ms}} \sum_{(c_{1,0}, \dots, c_{s,1}) \in (\Delta_m^*(p))^{2s}} A_3(j, m, \nu, c_{1,0}, \dots, c_{s,1}) < K(s) m^{2s+1},$$

and

$$(115) \quad \frac{1}{\varphi_0^{2s} p^{2ms}} \sum_{(c_{1,0}, \dots, c_{s,1}) \in (\Delta_m^*(p))^{2s}} A_4(j, m, \nu, c_{1,0}, \dots, c_{s,1}) < K(s) m^{2s}.$$

Proof. We will prove the inequality (114). The proof of (115) repeats that of (114). We denote the left side of (114) by σ . Changing the order of the summation, from (112) we obtain

$$\sigma \leq \sum_{h_0 \in C(p^m)} \sum_{\mathbf{h} \in C_{2s}(p^m)} r^{-1}(h_0, p^m)(\mathbf{h}, p^m) E(h_0, \mathbf{h}),$$

where

$$E(h_0, \mathbf{h}) = \frac{1}{\varphi_0^{2s} p^{2ms}} \sum_{(c_1, \dots, c_{2s}) \in (\Delta_m(p))^{2s}} \times p^m \delta_{p^m} \left(h_0 + \sum_{j=1}^s (q_i^j h_j c_j + h_{j+s} c_{j+s}) \right).$$

Let $(h_1, \dots, h_{2s}, p^m) = p^\alpha$, let $h_i = p^\alpha h'_i$, $i = 1, \dots, 2s$, and $(h'_{\nu_0 s + i_0}, p) = 1$, $\nu_0 = 0, 1$, $i_0 = 1, \dots, s$. If $h_0 \not\equiv 0 \pmod{p^\alpha}$, then $E(h_0, \mathbf{h}) = 0$.

Now let $h_0 = p^\alpha h'_0$. Hence

$$E(h_0, \mathbf{h}) \leq \max_{(c_1, \dots, c_{2s}) \in (\Delta_m(p^m))^{2s}} p^m \frac{p}{\varphi(p)} \sum_{c_{s\nu_0+i_0}=0}^{p^m-1} \times \delta_{p^{m-\alpha}} \left(h'_0 + \sum_{i=1}^s (q_i^j h'_i c_i + h'_{i+s} c_{i+s}) \right).$$

Bearing in mind that $(q_{i_0}^j h'_{s\nu_0+i_0}, p) = 1$, and

$$\sum_{c_{s\nu_0+i_0}=0}^{p^m-1} \delta_{p^{m-\alpha}} (f + q_{i_0}^j h'_{s\nu_0+i_0} c_{s\nu_0+i_0}) = p^\alpha,$$

we find that

$$E(h_0, \mathbf{h}) \leq p^{\alpha+1}/\varphi(p).$$

Applying Lemma 2, we obtain

$$\begin{aligned} \sigma &\leq \sum_{\alpha=0}^m \sum_{\substack{h_0 \in C(p^m) \\ (h_0, p^m) = p^\alpha}} \sum_{\substack{\mathbf{h} \in C_{2s}(p^m) \\ (h_1, \dots, h_{2s}, p^m) = p^\alpha}} r^{-1}(h_0, p^m)(\mathbf{h}, p^m) p^{\alpha+1}/\varphi(p) \\ &\leq p/\varphi(p) \sum_{\alpha=0}^m \left(\frac{2}{\pi} m \log p + \frac{7}{5} \right)^{2s} \left(1 + p^{-\alpha} \left(\frac{2}{\pi} m \log p + \frac{7}{5} \right) \right) \\ &< m^{2s+1} p/\varphi(p) \left(\frac{2}{\pi} \log p + \frac{7}{5} \right)^{2s} \left(\frac{m+1}{m} + \frac{p}{p-1} \left(\frac{2}{\pi} \log p + \frac{7}{5} \right) \right) \\ &\leq K(s) m^{2s+1}. \end{aligned}$$

□

Corollary 9. *There exist integers $a_{1,0}^{(m)}, \dots, a_{s,0}^{(m)}, a_{1,1}^{(m)}, \dots, a_{s,1}^{(m)}$ such that*

$$(116) \quad \sum_{j=0}^{k_m-1} \sum_{\nu=0}^1 A_3(j, m, \nu, a_{1,0}^{(m)}, \dots, a_{s,0}^{(m)}, a_{1,1}^{(m)}, \dots, a_{s,1}^{(m)}) \leq 4k_m K(s) m^{2s+1},$$

and

$$(117) \quad \sum_{j=0}^{k_m-1} \sum_{\nu=0}^1 A_4(j, m, \nu, a_{1,0}^{(m)}, \dots, a_{s,0}^{(m)}, a_{1,1}^{(m)}, \dots, a_{s,1}^{(m)}) \leq 4k_m K(s) m^{2s}, \quad m = 1, 2, \dots$$

We use such integers $a_{1,0}^{(m)}, \dots, a_{s,1}^{(m)}$ ($m = 1, 2, \dots$) to construct the real numbers $\alpha_1, \dots, \alpha_s$ (15).

Lemma 20. *Let $1 \leq M \leq p^m$. Then for $m = 1, 2, \dots$*

$$(118) \quad 2k_m M D\left(\left(\{\alpha_1 q_1^x\}, \dots, \{\alpha_s q_s^x\}\right)_{x \in [n_m, n_m + 2k_m M)}\right) = O(m^{2s+2}),$$

$$(119) \quad 2k_m p^m D\left(\left(\{\alpha_1 q_1^x\}, \dots, \{\alpha_s q_s^x\}\right)_{x \in [n_m, n_{m+1})}\right) = O(m^{2s+1}).$$

Proof. We denote the left side of (118) by σ . Equation (2) implies that

$$(120) \quad \sigma \leq \sum_{x_3=0}^{k_m-1} \sum_{x_2=0}^1 \sigma(x_2, x_3),$$

with

$$\sigma(x_2, x_3) = M D^{(s)}\left(\left(\{\alpha_1 q_1^{n_m + 2k_m x_1 + k_m x_2 + x_3}\}, \dots, \{\alpha_s q_s^{n_m + 2k_m x_1 + k_m x_2 + x_3}\}\right)_{x_1 \in [0, M)}\right).$$

We apply $k^{(i)} = 2k_m - x_3$, $i = 1, \dots, s$ to Lemma 5:

$$\sigma(x_2, x_3) \leq \frac{sM}{q^{2k_m - x_3}} + M D^{(s)}\left(\left(\{\alpha_i q_i^{n_m + 2k_m x_1 + k_m x_2 + x_3}\}_{2k_m - x_3}\right)_{i=1, \dots, s}\right)_{x_1 \in [0, M)},$$

where $q = \min(q_1, \dots, q_s)$. Using (15), we obtain, similarly to (79), that

$$\{\alpha_i q_i^{n_m + 2k_m x_1 + k_m x_2 + x_3}\}_{2k_m - x_3} = \left\{ \frac{q_i^{x_3} a_{i,x_2}^{(m)} x_1}{p^m} \right\}_{k_m - x_3, i} + \frac{1}{q_i^{k_m - x_3}} \left\{ \frac{a_{i,x_2+1}^{(m)} (x_1 + x_2)}{p^m} \right\}_{k_m, i}.$$

Then Lemma 6 (with $k_i = k_m - x_3$ and $l_i = k_m$, $i = 1, \dots, s$) shows that

$$\sigma(x_2, x_3) \leq \frac{2sM}{q^{2k_m - x_3}} + 2^s M D^{(2s)} \left(\left(\left(\left\{ \frac{q_i^{x_3} a_{i,x_2}^{(m)} x_1}{p^m} \right\} \right)_{i=1, \dots, s}, \right. \right. \\ \left. \left. \left(\left\{ \frac{a_{i,x_2+1}^{(m)} (x_1 + x_2)}{p^m} \right\} \right)_{i=1, \dots, s} \right)_{x_1 \in [0, M)} \right).$$

Now Corollary 2 (with $T = P = p^m$) implies that

$$\sigma(x_2, x_3) \leq \frac{2sM}{q^{k_m}} + \frac{2s2^s M}{p^m} + 2^s \sum_{h_0 \in C(p^m)} \sum_{\mathbf{h} \in C_{2s}(p^m)} r^{-1}(h_0, p^m) r^{-1}(\mathbf{h}, p^m) \\ \times \left| \sum_{x_1=0}^{p^m-1} e \left(\left(h_0 x_1 + \sum_{i=1}^s \left(q_i^{x_3} h_i a_{i,x_2}^{(m)} x_1 + h_{s+i} a_{i,x_2+1}^{(m)} (x_1 + x_2) \right) \right) / p^m \right) \right|.$$

Applying Lemma 1, (14), and (112), we obtain

$$\sigma(x_2, x_3) \leq 4s2^{2s} + 2^s A_3(x_3, m, x_2, a_{1,0}^{(m)}, \dots, a_{s,1}^{(m)}).$$

From (14), (116), (120), and Corollary 9, we get (118). Using (117), we similarly obtain (119). $\square$

End of the proof of Theorem 2. Let N be in $[n_r, n_{r+1})$. Define $D(N_1, N_2) = 0$ for $N_2 \leq 0$, and

$$(121) \quad D(N_1, N_2) = N_2 D \left(\left(\{ \alpha_1 q_1^x \}, \dots, \{ \alpha_s q_s^x \} \right)_{x \in [N_1, N_1 + N_2)} \right) \\ \text{for } N_2 > 0.$$

Using (2), (14), and Lemma 20 we have for $M \in [1, 2k_m p^m]$ that

$$(122) \quad D(n_m, M) \leq D(n_m, 2k_m \lfloor M/2k_m \rfloor) + 2k_m = O(m^{2s+2}).$$

Applying (2), (14), (118), (119), (121), and (122), we get the assertion of Theorem 2:

$$D(1, N) \leq D(1, 2k_1 p) + \sum_{m=2}^r D(n_m, 2k_m p^m) + D(n_r, N + 1 - n_r) \\ = O \left(\sum_{m=1}^r m^{2s+1} + r^{2s+2} \right) = O(r^{2s+2}) = O(\log^{2s+2} N). \quad \square$$

5. Appendix

The proof of Proposition 1. We follow [KN p.70]. Let $s_1, s_2 \geq 1$ be integers. Consider a box $v \subset [0, 1)^s$, a block of digits $G_{s_1, s_2} = \{g_{i,j} \in$

$\{0, \dots, q-1\} \mid i = 0, \dots, s_1-1, j = 1, \dots, s_2\}$, a configuration $\omega = (d_{i,j})_{i,j \geq 1}$ with $d_{i,j} \in \{0, \dots, q-1\}$ and real numbers

$$\alpha_m = \sum_{n=1}^{\infty} d_{m,n}/q^n.$$

Now let

$$A_v(M, N) = \#\{(m, n) \in [1, M] \times [1, N] \mid (\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\}) \in v\},$$

$$S(M, N, G_{s_1, s_2}) = \#\{(m, n) \in [1, M] \times [1, N] \mid d_{m+i, n+j} = g_{i,j}, \\ i \in [0, s_1), j \in [1, s_2]\}.$$

The block $(d_{m+i, n+j})_{i=0, j=1}^{s_1-1, s_2}$ is identical with G_{s_1, s_2} if and only if

$$\alpha_{m+i} = [\alpha_{m+i}] + \sum_{k=1}^n \frac{d_{m+i, k}}{q^k} + \frac{g_{i,1}}{q^{n+1}} + \dots + \frac{g_{i, s_2}}{q^{n+s_2}} + \sum_{k=n+s_2+1}^{\infty} \frac{d_{m+i, k}}{q^k},$$

or

$$\{\alpha_{m+i} q^n\} = \frac{g_{i,1}}{q} + \dots + \frac{g_{i, s_2}}{q^{s_2}} + \sum_{k=s_2+1}^{\infty} \frac{d_{m+i, k}}{q^k},$$

or

$$\{\alpha_{m+i} q^n\} \in \left[\frac{g_{i,1} q^{s_2-1} + \dots + g_{i, s_2}}{q^{s_2}}, \frac{g_{i,1} q^{s_2-1} + \dots + g_{i, s_2} + 1}{q^{s_2}} \right) \\ = \Delta_i(G_{s_1, s_2}), \quad i \in [0, s_1).$$

It follows that

$$S(M, N, G_{s_1, s_2}) = A_v(M, N), \quad \text{with } v = \prod_{i=0}^{s_1-1} \Delta_i(G_{s_1, s_2}).$$

Now suppose that the double sequence $(\{\alpha_m q^n\}, \dots, \{\alpha_{m+s-1} q^n\})_{m, n \geq 1}$ is uniformly distributed in $[0, 1)^{s_1}$. Then

$$A_v(M, N) = q^{-s_1 s_2} MN + o(MN), \quad \text{with } \max(M, N) \rightarrow \infty,$$

and so ω is a normal lattice configuration.

Conversely, if ω is a normal configuration, then

$$A_v(M, N) = S(M, N, G_{s_1, s_2}) = MN \text{mes } v + o(MN),$$

where $v = \prod_{i=0}^{s_1-1} \Delta_i(G_{s_1, s_2})$, $\text{mes } v = q^{-s_1 s_2}$ and $\max(M, N) \rightarrow \infty$. This holds for all G_{s_1, s_2} . Therefore

$$(123) \quad A_v(M, N) = MN \text{mes } v + o(MN),$$

for all boxes $v = \prod_{i=1}^{s_1} [0, \gamma_i)$ with $\gamma_i = h_i/q^{s_2}$, $h_i \in \{0, \dots, q^{s_2}-1\}$, $s_2 \geq 1$, $i = 1, \dots, s_1$, and $\max(M, N) \rightarrow \infty$.

Now let v be a box with arbitrary $\gamma_i \in (0, 1]$ ($i = 1, \dots, s_1$), and let $\epsilon \in (0, 1)$ be given. Put $s_2 = 1 + [-\log_q \epsilon]$ and put $h_i = [\gamma_i q^{s_2}]$, $i = 1, \dots, s_1$. Then

$$(124) \quad A_{v_1}(M, N) \leq A_v(M, N) \leq A_{v_2}(M, N),$$

where $v_1 = \prod_{i=1}^{s_1} [0, h_i/q^{s_2}]$ and $v_2 = \prod_{i=1}^{s_1} [0, (h_i + 1)/q^{s_2}]$.

It is easy to see that $\gamma_i \in [h_i/q^{s_2}, (h_i + 1)/q^{s_2})$, with $i = 1, \dots, s_1$, $1/q^{s_2} < \epsilon$. According to [Ni, Lemma 3.9]

$$\max(|\text{mes } v_1 - \text{mes } v|, |\text{mes } v_2 - \text{mes } v|) \leq 1 - (1 - 1/q^{s_2})^{s_1} \leq s_1 q^{-s_2} \leq s_1 \epsilon.$$

From (123) and (124) we deduce that

$$A_v(M, N) = MN(\text{mes } v + \epsilon_1) + o(MN),$$

where $|\epsilon_1| \leq s_1 \epsilon$, and $\max(M, N) \rightarrow \infty$. Hence for all $s_1 \geq 1$ and all boxes $v \subset [0, 1]^{s_1}$, $A_v(M, N) = MN \text{mes } v + o(MN)$, with $\max(M, N) \rightarrow \infty$, and so $(\{\alpha_m q^n\}, \dots, \{\alpha_{m+s_1-1} q^n\})_{m, n \geq 1}$ is a uniformly distributed double sequence in $[0, 1]^{s_1}$. $\square$

Acknowledgment. I am very grateful to the referee for many corrections and suggestions which improved this paper.

References

- [C] D. J. CHAMPERNOWNE, *The construction of decimal normal in the scale ten*. J. London Math. Soc. **8** (1935), 254–260.
- [Ci] J. CIGLER, *Asymptotische Verteilung reeller Zahlen mod 1*. Monatsh. Math. **64** (1960), 201–225.
- [DrTi] M. DRMOTA, R. F. TICHY, *Sequences, Discrepancies and Applications*. Lecture Notes in Mathematics **1651**, Springer-Verlag, Berlin, 1997.
- [Ei] J. EICHENAUER-HERRMANN, *A unified approach to the analysis of compound pseudorandom numbers*. Finite Fields Appl. **1** (1995), 102–114.
- [KT] P. KIRSCHENHOFER, R. F. TICHY, *On uniform distribution of double sequences*, Manuscripta Math. **35** (1981), 195–207.
- [Ko1] N. M. KOROBOV, *Numbers with bounded quotient and their applications to questions of Diophantine approximation*. Izv. Akad. Nauk SSSR Ser. Mat. **19** (1955), 361–380.
- [Ko2] N. M. KOROBOV, *Distribution of fractional parts of exponential function*. Vestnik Moskov. Univ. Ser. 1 Mat. Meh. **21** (1966), no. 4, 42–46.
- [Ko3] N. M. KOROBOV, *Exponential sums and their applications*, Kluwer Academic Publishers, Dordrecht, 1992.
- [KN] L. KUIPERS, H. NIEDREITER, *Uniform distribution of sequences*. John Wiley, New York, 1974.
- [Le1] M. B. LEVIN, *On the uniform distribution of the sequence $\{\alpha \lambda^n\}$* . Math. USSR Sbornik **27** (1975), 183–197.
- [Le2] M. B. LEVIN, *The distribution of fractional parts of the exponential function*. Soviet. Math. (Iz. Vuz.) **21** (1977), no. 11, 41–47.
- [Le3] M. B. LEVIN, *On the discrepancy estimate of normal numbers*. Acta Arith. **88** (1999), 99–111.
- [LS1] M. B. LEVIN, M. SMORODINSKY, *Explicit construction of normal lattice configuration*, preprint.
- [LS2] M. B. LEVIN, M. SMORODINSKY, *A $\mathbb{Z}^d$ generalization of Davenport and Erdős theorem on normal numbers*. Colloq. Math. **84/85** (2000), 431–441.

- [Ni] H. NIEDERREITER, *Random Number Generation and Quasi-Monte Carlo Methods*. SIAM, Philadelphia, 1992.
- [Ro] K. ROTH, *On irregularities of distributions*. *Mathematika* 1 (1954), 73–79.

Mordechay B. LEVIN
Department of Mathematics and Computer Science
Bar-Ilan University
Ramat-Gan, 52900
Israel
E-mail : `mlevin@macs.biu.ac.il`