

L. ANGERMANN

Addendum to the paper “Numerical solution of second-order equations on plane domains”

M2AN - Modélisation mathématique et analyse numérique, tome 27, n° 1 (1993), p. 1-7

http://www.numdam.org/item?id=M2AN_1993__27_1_1_0

© AFCET, 1993, tous droits réservés.

L'accès aux archives de la revue « M2AN - Modélisation mathématique et analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**ADDENDUM TO THE PAPER
 « NUMERICAL SOLUTION OF SECOND-ORDER EQUATIONS
 ON PLANE DOMAINS (*) »**

by L. ANGERMANN ⁽¹⁾

Communicated by R. TEMAM

In the paper [1] we applied a so-called upwind discretization scheme to the problem

$$\begin{cases} -\Delta u + b \cdot \nabla u + cu = f & \text{in } D \\ u = 0 & \text{on } B, \end{cases} \quad (1)$$

where D is a bounded plane domain with polygonal boundary. We could demonstrate some important properties of the resulting scheme. Moreover, we proved a convergence result. Unfortunately, for that proof we had to require the underlying triangulations to be uniformly acute, i.e. in addition to the quite usual assumption on the existence of a uniform *lower* bound for all interior angles of the triangles (Zlámal's angle condition), we needed a uniform *upper* bound being less than $\frac{\pi}{2}$ strongly. In addition, we had to suppose that the family of triangulations satisfies the so-called inverse assumption (quasiuniformity condition).

In this note, we intend to show that the last two assumptions are not related with the discretization principle itself, but rather with the approximation quality of the coefficients which can be achieved.

Namely, if we redefine the quantities N_{ij} , c_i and f_i from [1] as follows :

$$N_{ij} = \frac{1}{m_{ij}} \int_{B_{ij}} (n \cdot b) ds, \quad c_i = \frac{1}{m_i} \int_{D_i} c dx, \quad f_i = \frac{1}{m_i} \int_{D_i} f dx,$$

(*) Manuscript received July 1991.

(¹) Universität Erlangen-Nürnberg, Institut für Angewandte Mathematik, Martensstraße 3, W-8520 Erlangen, Germany.

then instead of the assumption (A2) from [1] it is sufficient to suppose that obtuse triangles do not appear in the triangulations. Such triangulations are called *weakly acute*.

It is not difficult to verify that the statements of Lemmata 1-3 in [1] remain valid. However, the proof of the convergence theorem must be modified essentially. Therefore we completely bring the formulation and the proof of this theorem.

THEOREM 1 : *Let the weak solution u of (1) belong to $H_0^1(D) \cap H^2(D)$. Then the estimate*

$$\|u - u_h; H^1(D)\| \leq Ch$$

holds, where C is a positive constant independent of h . ■

Proof : For arbitrary $w_h \in X_h$, we have

$$C \|u_h - w_h; H^1(D)\|^2 \leq a(u_h - w_h, u_h - w_h). \quad (2)$$

Setting $v_h = u_h - w_h$, the right-hand side of this inequality can be treated as follows :

$$\begin{aligned} a(v_h, v_h) &= a(u - w_h, v_h) + a(u_h - u, v_h) \\ &= a(u - w_h, v_h) + a(u_h, v_h) - a_l(u_h, v_h) + (f, v_h)_l - (f, v_h) \\ &= a(u - w_h, v_h) + \delta a \end{aligned}$$

Using the definition of f_l , the term δa admits the following representation, which is different from [1] :

$$\begin{aligned} \delta a &= \sum_{i \in V} \sum_{j \in V_i} (1 - r_{ij}) N_{ij} (u_{hi} - u_{hj}) v_{hi} m_{ij} + \int_D (b \cdot \nabla u_h) v_h dx \\ &\quad - \sum_{i \in V} \int_{D_i} [c_i u_{hi} v_{hi} - c u_h v_h] dx - \sum_{i \in V} \int_{D_i} [f v_h - f_l v_{hi}] dx \\ &= \sum_{i \in V} \left\{ \sum_{j \in V_i} (1 - r_{ij}) N_{ij} (u_{hi} - u_{hj}) v_{hi} m_{ij} + \int_{D_i} (b \cdot \nabla u_h) v_{hi} dx \right\} \\ &\quad + \sum_{i \in V} \int_{D_i} (b \cdot \nabla u_h) (v_h - v_{hi}) dx - \sum_{i \in V} \int_{D_i} [c_i u_{hi} - c u_h] v_{hi} dx \\ &\quad + \sum_{i \in V} \int_{D_i} c u_h (v_h - v_{hi}) dx - \sum_{i \in V} \int_{D_i} f [v_h - v_{hi}] dx \\ &= \delta_1 + \delta_2 + \delta_3, \end{aligned}$$

where

$$\begin{aligned} \delta_1 &= \sum_{i \in V} \int_{D_i} [b \cdot \nabla u_h + c u_h - f] (v_h - v_{h_i}) dx, \\ \delta_2 &= \sum_{i \in V} v_{h_i} \left\{ \sum_{j \in V_i} u_{h_i} N_{ij} m_{ij} - \int_{D_i} [(\nabla \cdot b - c) u_h + c_i u_{h_i}] dx \right\}, \\ \delta_3 &= \sum_{i \in V} \sum_{j \in V_i} \int_{B_{ij}} \{ (n \cdot b) u_h - [r_{ij} u_{h_i} + (1 - r_{ij}) u_{h_j}] N_{ij} \} v_{h_i} ds. \end{aligned}$$

By Cauchy's inequality, for δ_1 we obtain

$$\delta_1 \leq \|b \cdot \nabla u_h + c u_h - f; L_2(D)\| \left\{ \sum_{i \in V} \int_{D_i} |v_h - v_{h_i}| dx \right\}^{\frac{1}{2}}.$$

The last term can be estimated by means of Lemma 2.1 from [3], that is

$$\delta_1 \leq Ch \|b \cdot \nabla u_h + c u_h - f; L_2(D)\| \|v_h; H^1(D)\|. \tag{3}$$

Now we consider δ_2 . Using the definition of c_i , we have

$$\delta_2 = \sum_{i \in V} v_{h_i} \left\{ \sum_{j \in V_i} u_{h_i} N_{ij} m_{ij} - \int_{D_i} (\nabla \cdot b) u_{h_i} dx - \int_{D_i} (\nabla \cdot b - c)(u_h - u_{h_i}) dx \right\}.$$

Integrating the first integral by parts and using the definition of N_{ij} , we obtain

$$\begin{aligned} \delta_2 &= \sum_{i \in V} \int_{D_i} (c - \nabla \cdot b)(u_h - u_{h_i}) v_{h_i} dx \\ &\leq \|c - \nabla \cdot b; L_\infty(D)\| \sum_{i \in V} \int_{D_i} |u_h - u_{h_i}| |v_{h_i}| dx. \end{aligned}$$

Applying Cauchy's inequality and Lemma 2.1 from [3], it follows

$$\delta_2 \leq Ch \|c - \nabla \cdot b; L_\infty(D)\| \|u_h; H^1(D)\| \|v_h\|_l.$$

In view of the equivalence of the $\|\cdot\|_l$ -norm with the L_2 -norm on X_h (see Lemma 2.2 in [3]), we get

$$\delta_2 \leq Ch \|c - \nabla \cdot b; L_\infty(D)\| \|u_h; H^1(D)\| \|v_h; L_2(D)\|. \tag{4}$$

Thus it remains to consider the third term. By a symmetry argument, we see that

$$\delta_3 = \frac{1}{2} \sum_{i \in V} \sum_{j \in V_i} \int_{B_{ij}} \{ (n \cdot b) u_h - [r_{ij} u_{hi} + (1 - r_{ij}) u_{hj}] N_{ij} \} (v_{hi} - v_{hj}) ds.$$

Obviously, by means of B_{ij} and x_i there can be defined some triangle T_{ij} in such a way that B_{ij} is one edge and x_i the opposite vertex. On the other hand, it is possible to represent this triangle as the union of two triangles $T_{ij}^{(k)}$ ($k = 1, 2$) having in common the vertex x_i and the straight-line segment connecting the node x_i with the point $\frac{1}{2}(x_i + x_j)$.

Now we can write $\delta_3 = \delta_{31} + \delta_{32}$, where

$$\begin{aligned} \delta_{31} &= \frac{1}{2} \sum_{i \in V} \sum_{j \in V_i} \int_{B_{ij}} (n \cdot b - N_{ij}) u_h (v_{hi} - v_{hj}) ds, \\ \delta_{32} &= \frac{1}{2} \sum_{i \in V} \sum_{j \in V_i} \int_{B_{ij}} [u_h - r_{ij} u_{hi} - (1 - r_{ij}) u_{hj}] N_{ij} (v_{hi} - v_{hj}) ds. \end{aligned}$$

We turn to the estimation of the integral

$$J_{ij}(b) = \int_{B_{ij}} (n \cdot b - N_{ij}) u_h ds.$$

To this end we introduce a reference triangle \tilde{T} with vertices $t_1 = (0, 0)$, $t_2 = (1, 0)$, $t_3 = (0, 1)$ and transform T_{ij} onto \tilde{T} in such a way that B_{ij} has the image $\tilde{B} = \overline{t_1 t_2}$. Then we have

$$J_{ij}(b) = m_{ij} J(\tilde{b}),$$

where

$$J(\tilde{b}) = \int_{\tilde{B}} (n \cdot \tilde{b} - \tilde{N}_{ij}) \tilde{u}_h d\tilde{s}.$$

It follows

$$\begin{aligned} |J(\tilde{b})| &\leq \|n \cdot \tilde{b} - \tilde{N}_{ij}; L_2(\tilde{B})\| \|\tilde{u}_h; L_2(\tilde{B})\| \\ &\leq 2 \|n \cdot \tilde{b}; L_2(\tilde{B})\| \|\tilde{u}_h; L_2(\tilde{B})\|. \end{aligned}$$

Using the trace theorem [4], we get

$$|J(\tilde{b})| \leq C \|n \cdot \tilde{b}; H^1(\tilde{T})\| \|\tilde{u}_h; H^1(\tilde{T})\|.$$

Observing that J vanishes for constant \tilde{b} , we conclude from Bramble-Hilbert's lemma [2] the estimate

$$|J(\tilde{b})| \leq C |n \cdot \tilde{b}; H^1(\tilde{T})| \| \tilde{u}_h; H^1(\tilde{T}) \| .$$

The back-transformation (see [2], again) yields

$$\begin{aligned} |J_{ij}(b)| &\leq C m_{ij} h_{T_{ij}} (\text{meas } T_{ij})^{-1} |n \cdot b; H^1(T_{ij})| \| u_h; H^1(T_{ij}) \| \\ &\leq C m_{ij} h_{T_{ij}} (\text{meas } T_{ij})^{-\frac{1}{2}} |n \cdot b; W_\infty^1(T_{ij})| \| u_h; H^1(T_{ij}) \| \\ &\leq C m_{ij} h_{T_{ij}} (\text{meas } T_{ij})^{-\frac{1}{2}} |b; [W_\infty^1(T_{ij})]^2| \| u_h; H^1(T_{ij}) \| . \end{aligned}$$

Thus we have

$$\begin{aligned} \delta_{31} &\leq Ch |b; [W_\infty^1(D)]^2| \sum_{i \in V} \sum_{j \in V_i} m_{ij} (\text{meas } T_{ij})^{-\frac{1}{2}} \times \\ &\qquad \qquad \qquad \times |v_{hi} - v_{hj}| \| u_h; H^1(T_{ij}) \| . \end{aligned}$$

In view of the relations

$$|v_{hi} - v_{hj}| = d_{ij} |n \cdot \nabla v_h| \quad \text{and} \quad d_{ij} m_{ij} = 4 \text{ meas } T_{ij} \tag{5}$$

it follows

$$\delta_{31} \leq Ch |b; [W_\infty^1(D)]^2| \sum_{i \in V} \sum_{j \in V_i} (\text{meas } T_{ij})^{\frac{1}{2}} |n \cdot \nabla v_h| \| u_h; H^1(T_{ij}) \| .$$

Cauchy's inequality implies

$$\delta_{31} \leq Ch |b; [W_\infty^1(D)]^2| |v_h; H^1(D)| \| u_h; H^1(D) \| . \tag{6}$$

Thus it remains to consider the term δ_{32} . Using the notation $B_{ij}^{(k)} = B_{ij} \cap T_{ij}^{(k)}$, we can write

$$\delta_{32} = \frac{1}{2} \sum_{i \in V} \sum_{j \in V_i} \sum_{k=1}^2 \int_{B_{ij}^{(k)}} u_{hij} N_{ij} (v_{hi} - v_{hj}) ds ,$$

where

$$u_{hij} = u_h - r_{ij} u_{hi} - (1 - r_{ij}) u_{hj} .$$

Now, on each triangle $T_{ij}^{(k)}$ we have

$$u_h(x) = \bar{u}_{hi} + \nabla u_h \cdot (x - x_i)$$

with ∇u_h being constant on $T_{ij}^{(k)}$. Thus, u_{hij} can be treated as follows :

$$\begin{aligned} u_{hij} &= (1 - r_{ij})(u_{hi} - u_{hj}) + \nabla u_h \cdot (x - x_i) \\ &= (1 - r_{ij}) d_{ij} (n \cdot \nabla u_h) + \nabla u_h \cdot (x - x_i) \\ &\leq C d_{ij} \|\nabla u_h\|. \end{aligned}$$

Therefore, together with (5) this estimate implies

$$\begin{aligned} \delta_{32} &\leq C \sum_{i \in V} \sum_{j \in V_i} d_{ij}^2 N_{ij} \sum_{k=1}^2 \int_{B_{ij}^{(k)}} \|\nabla u_h\| \|\nabla v_h\| ds \\ &= C \sum_{i \in V} \sum_{j \in V_i} d_{ij}^2 N_{ij} \sum_{k=1}^2 \frac{m_{ij}^{(k)}}{\text{meas } T_{ij}^{(k)}} \int_{T_{ij}^{(k)}} \|\nabla u_h\| \|\nabla v_h\| dx, \end{aligned}$$

where $m_{ij}^{(k)}$ is the length of $B_{ij}^{(k)}$.

Observing that $\text{meas } T_{ij}^{(k)} = \frac{1}{4} m_{ij}^{(k)} d_{ij}$, after the application of various variants of Cauchy's inequality we obtain

$$\begin{aligned} \delta_{32} &\leq C \sum_{i \in V} \sum_{j \in V_i} d_{ij} N_{ij} \sum_{k=1}^2 \int_{T_{ij}^{(k)}} \|\nabla u_h\| \|\nabla v_h\| ds \\ &\leq Ch \|b; [L_\infty(D)]^2\| \sum_{i \in V} \sum_{j \in V_i} |u_h; H^1(T_{ij})| |v_h; H^1(T_{ij})| \\ &\leq Ch \|b; [L_\infty(D)]^2\| \sum_{i \in V} |u_h; H^1(D_i)| |v_h; H^1(D_i)| \\ &\leq Ch \|b; [L_\infty(D)]^2\| |u_h; H^1(D)| |v_h; H^1(D)|. \end{aligned} \quad (7)$$

Summarizing the relations (3), (4), (6), (7) and using Lemma 1 from [1] in order to derive an a priori estimate for $\|u_h; H^1(D)\|$, we get from (2)

$$\|u_h - w_h; H^1(D)\| \leq C [\|u - w_h; H^1(D)\| + h].$$

The triangle inequality yields

$$\begin{aligned} \|u - u_h; H^1(D)\| &\leq \|u - w_h; H^1(D)\| + \|u_h - w_h; H^1(D)\| \\ &\leq C [\|u - w_h; H^1(D)\| + h]. \end{aligned}$$

Choosing for w_h the Lagrangian interpolant in X_h of the exact solution u of (1), the standard interpolation theory gives

$$\|u - w_h; H^1(D)\| \leq Ch |u; H^2(D)|.$$

Now, the desired result follows immediately. ■

REFERENCES

- [1] L. ANGERMANN, Numerical solution of second-order elliptic equations on plane domains, *M²AN Modél. Math. Anal. Numér.*, 25(2) : 169-191, 1991.
- [2] P. CIARLET, *The finite element method for elliptic problems*, North-Holland, Amsterdam-New York-Oxford, 1978.
- [3] T. IKEDA, *Maximum principle in finite element models for convection-diffusion phenomena*, North-Holland, Amsterdam-New York-Oxford, 1983.
- [4] J. WLOKA, *Partielle Differentialgleichungen. Sobolevräume und Randwertaufgaben*, Teubner-Verlag, Stuttgart, 1982.