

A. J. DE JONG

**Erratum : Crystalline Dieudonné module theory via formal and rigid geometry**

*Publications mathématiques de l'I.H.É.S.*, tome 87 (1998), p. 175

[http://www.numdam.org/item?id=PMIHES\\_1998\\_\\_87\\_\\_175\\_0](http://www.numdam.org/item?id=PMIHES_1998__87__175_0)

© Publications mathématiques de l'I.H.É.S., 1998, tous droits réservés.

L'accès aux archives de la revue « Publications mathématiques de l'I.H.É.S. » (<http://www.ihes.fr/IHES/Publications/Publications.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

ERRATUM TO :  
CRYSTALLINE DIEUDONNÉ MODULE THEORY  
VIA FORMAL AND RIGID GEOMETRY

by A. J. de JONG

Lemma 7.1.13.1 of [1] is wrong. This was pointed out to the author by Brian Conrad. The only place where Lemma 7.1.13.1 is used in the article is in the construction of the maps  $\beta_n$  (7.1.13.2). We give a correct construction of  $\beta_n$  (for  $A$  such that  $\pi$  is not a zero divisor).

Let  $c$  be an integer such that

$$\pi A \cap I^n \subset \pi I^{n-c}$$

for all  $n \geq c$ . The existence of  $c$  is the Artin-Rees lemma. Then for any integer  $t \in \mathbf{N}$  we have

$$\pi^t A \cap (\pi^{t-1} I^n + \pi^{t-2} I^{2n} + \dots + I^{tn}) \subset \pi^t I^{n-c}.$$

Next, we come to the definition of  $\beta_n$ . Let  $n \geq c$ . Any  $a \in A[I^n/\pi]$  can be written in the form

$$a = a_0 + a_1/\pi + \dots + a_t/\pi^t, \quad a_i \in I^{in}$$

for some  $t \in \mathbf{N}$ . We simply put  $\beta_n(a) = a_0 \bmod I^{n-c}$ . To show that  $\beta_n$  is well defined, suppose that  $a_0 + a_1/\pi + \dots + a_t/\pi^t$  ( $a_i \in I^{in}$ ) represents zero in  $A[I^n/\pi]$ . This means that  $\pi^t a_0 + \pi^{t-1} a_1 + \dots + a_t = 0$ , as  $A$  has no  $\pi$ -torsion. Thus

$$\pi^t a_0 \in \pi^{t-1} I^n + \pi^{t-2} I^{2n} + \dots + I^{tn}$$

and by the above it follows that  $a_0 \in I^{n-c}$ . Hence

$$\beta_n(a_0 + a_1/\pi + \dots + a_t/\pi^t) = a_0 \bmod I^{n-c} = 0$$

as desired.

[1] A. J. de Jong, Crystalline Dieudonné module theory via formal and rigid geometry, *Publ. math. IHES*, **82**, 5-96 (1995).

Department of Mathematics,  
Fine Hall,  
Washington Rd  
Princeton, NJ 08544-1000  
États-Unis.

*Manuscrit reçu le 9 février 1998.*