

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

J. P. SINGHAL

Operational formulae for certain classical polynomials

Rendiconti del Seminario Matematico della Università di Padova,
tome 38 (1967), p. 33-40

<http://www.numdam.org/item?id=RSMUP_1967__38__33_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1967, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

OPERATIONAL FORMULAE FOR CERTAIN CLASSICAL POLYNOMIALS

J. P. SINGHAL *)

1. Recently Srivastava [7, p. 43] has defined a set of polynomials $A_n^{(\alpha)}(x)$ related to the Laguerre polynomials by means of the relations

$$(1.1) \quad \sum_{r=0}^n A_r^{(\alpha)}(x) L_{n-r}^{(\alpha+r)}(x) = 0, \quad n \geq 1,$$

$$A_0^{(\alpha)}(x) = 1.$$

He also gave the generating function, hypergeometric representation and the Rodrigues' formula for these polynomials [7, pp. 44-45] in the forms :

$$(1.2) \quad (1+t)^{-1-\alpha} e^{xt} = \sum_{r=0}^{\infty} t^r A_r^{(\alpha)}(x),$$

$$(1.3) \quad A_n^{(\alpha)}(x) = \frac{1}{(n)!} \sum_{r=0}^{\infty} \frac{(-n)_r (1+\alpha)_r}{(r)!} x^{n-r},$$

and

$$(1.4) \quad A_n^{(\alpha)}(x) = \frac{x^{n+\alpha+1}}{(n)!} D^n \{x^{-\alpha-1} e^x\}, \quad \left(D = \frac{d}{dx} \right),$$

respectively.

*) Indirizzo dell'A. : Deptt. of Maths. — University of Jodhpur — Jodhpur — India.

In this paper we give some operational formulae for these as well as Laguerre polynomials and employ them to derive many interesting results.

The first operational formula to be proved is

$$(1.5) \quad \prod_{j=1}^n (xD + x - \alpha - j) = (n)! \sum_{r=0}^n \frac{x^r}{(r)!} A_{n-r}^{(\alpha)}(x) D^r.$$

Note that the formula (1.5) corresponds to the one given by Carlitz [3, p. 219] in the case of Laguerre polynomials [8, p. 428].

To prove (1.5) we observe that if

$$\Omega_n = \prod_{j=1}^n (xD + x - \alpha - j), \quad \Omega_0 = 1,$$

it can be proved very easily by the method of induction that

$$\Omega_n(y) = x^{n+\alpha+1} e^{-x} D^n \{e^x x^{-\alpha-1} y\},$$

where y is some differentiable function of x .

Next since

$$\begin{aligned} D^n \{e^x x^{-\alpha-1} \cdot y\} &= \sum_{r=0}^n \binom{n}{r} D^{n-r} \{e^x x^{-\alpha-1}\} D^r y \\ &= \sum_{r=0}^n x^{-n-\alpha-1} e^x (n)! \frac{x^r}{(r)!} A_{n-r}^{(\alpha)}(x) D^r y, \end{aligned}$$

(1.5) follows immediately.

In (1.5) if we take $y = 1$, we obtain

$$(1.6) \quad \prod_{j=1}^n (xD + x - \alpha - j) \cdot 1 = (n)! A_n^{(\alpha)}(x).$$

As an application of (1.5) and (1.6), let us consider

$$\begin{aligned} (m+n)! A_{m+n}^{(\alpha)}(x) &= \prod_{j=1}^m (xD + x - \alpha - n - j) \prod_{j=1}^n (xD + x - \alpha - j) \cdot 1 \\ &= (n)! \prod_{j=1}^m (xD + x - \alpha - n - j) A_n^{(\alpha)}(x) \\ &= (m)! (n)! \sum_{r=0}^n \frac{x^r}{(r)!} A_{m-r}^{(\alpha+n)}(x) D^r A_n^{(\alpha)}(x). \end{aligned}$$

But since

$$D^r A_n^{(\alpha)}(x) = A_{n-r}^{(\alpha)}(x),$$

we readily get

$$(1.7) \quad \binom{m+n}{n} A_{m+n}^{(\alpha)}(x) = \sum_{r=0}^{\min(m,n)} \frac{x^r}{(r)!} A_{m-r}^{(\alpha+n)}(x) A_{n-r}^{(\alpha)}(x).$$

Further, from (1.7) we have

$$\begin{aligned} \sum_{m=0}^{\infty} \binom{m+n}{n} t^m A_{m+n}^{(\alpha)}(x) &= \sum_{r=0}^n \frac{(xt)^r}{(r)!} A_{n-r}^{(\alpha)}(x) \sum_{m=0}^{\infty} t^m A_m^{(\alpha+n)}(x) \\ &= \sum_{r=0}^n \frac{(xt)^r}{(r)!} A_{n-r}^{(\alpha)}(x) \cdot e^{xt} (1+t)^{-(\alpha+n+1)}, \end{aligned}$$

and making use of the relation [7, p. 45]

$$A_n^{(\alpha)}(x+y) = \sum_{r=0}^n \frac{y^r}{(r)!} A_{n-r}^{(\alpha)}(x),$$

we get the known formula [6, p. 7]

$$(1.8) \quad \sum_{m=0}^{\infty} \binom{m+n}{n} t^m A_{m+n}^{(\alpha)}(x) = e^{xt} (1+t)^{-(\alpha+n+1)} A_n^{(\alpha)}\{x(1+t)\}.$$

Another operational formula for the polynomials $A_n^{(\alpha)}(x)$ is

$$(1.9) \quad (1+D)^{-1-\alpha} x^n = (n)! A_n^{(\alpha)}(x), \quad \left(D = \frac{d}{dx} \right).$$

To prove it we note that

$$(1+D)^{-1-\alpha} x^n = \sum_{r=0}^n (-1)^r \frac{(1+\alpha)_r}{(r)!} D^r x^n = \sum_{r=0}^n \frac{(1+\alpha)_r (-n)_r}{(r)!} x^{n-r},$$

which evidently yields the formula (1.9).

From (1.9) we have

$$(n)! A_n^{(\alpha+\beta)}(x) = (1+D)^{-1-\alpha-\beta} x^n,$$

$$(1.10) \quad A_n^{(\alpha+\beta)}(x) = (1+D)^{-\beta} A_n^{(\alpha)}(x);$$

the last formula gives us

$$(1.11) \quad A_n^{(\alpha+\beta)}(x) = \sum_{r=0}^n (-1)^r \frac{(\beta)^r}{(r)!} A_{n-r}^{(\alpha)}(x).$$

Further let us operate on the identity

$$e^{xt} = \sum_{n=0}^{\infty} \frac{t^n}{(n)!} x^n,$$

by $(1+D)^{-1-\alpha}$; the familiar shift rule then gives us

$$(1+D)^{-1-\alpha} e^{xt} = e^{xt} (1+t)^{-1-\alpha} \left\{ 1 + \frac{D}{1+t} \right\}^{-1-\alpha} \cdot 1 = (1+t)^{-1-\alpha} e^{xt}.$$

On the other hand the second member yields

$$\sum_{n=0}^{\infty} t^n A_n^{(\alpha)}(x)$$

with the help of (1.9).

We thus arrive at the familiar generating function (1.2).

Now replace by tD_y , $\left(D_y = \frac{d}{dy}\right)$ in (1.2) and operate on both sides by $(1+D_y)^{-1-\beta}$. The left-hand side gives us

$$(1+D_y)^{-1-\beta} e^{xty} (1+ty) = e^{xty} \{1+xt+D_y\}^{-1-\beta} (1+yt)^{-1-\alpha}$$

$$= e^{xyt} (1+xt)^{-1-\beta} \sum_{r=0}^{\infty} (-1)^r \frac{(1+\beta)_r}{(r)!} \cdot \frac{D_y^r}{(1+xt)^r} (1+yt)^{-1-\alpha}$$

$$\simeq e^{xyt} (1+xt)^{-1-\beta} (1+yt)^{-1-\alpha} {}_2F_0 \left[1+\alpha, 1+\beta; -; \frac{t}{(1+xt)(1+yt)} \right]$$

and the right-hand side yields

$$\sum_{n=0}^{\infty} (n)! t^n A_n^{(\alpha)}(x) A_n^{(\beta)}(y).$$

Combining these two sides we finally get

$$(1.12) \quad \sum_{n=0}^{\infty} (n)! t^n A_n^{(\alpha)}(x) A_n^{(\beta)}(y) \\ \simeq e^{xyt} (1+xt)^{-1-\beta} (1+yt)^{-1-\alpha} {}_2F_0 \left[1+\alpha, 1+\beta; -; \frac{t}{(1+xt)(1+yt)} \right].$$

From the relation (1.12) we have

$$(1+xt)^{-1-\beta} (1+yt)^{-1-\alpha} {}_2F_0 \left[1+\alpha, 1+\beta; -; \frac{t}{(1+xt)(1+yt)} \right] \\ = \sum_{r,s,k=0}^{\infty} \frac{(1+\alpha)_r (1+\beta)_r (1+\alpha+r)_s (1+\beta+r)_k}{(r)! (s)! (k)!} (-1)^{s+k} x^k y^s \cdot t^{r+s+k} \\ = \sum_{n=0}^{\infty} (-t)^n \sum_{k=0}^n \sum_{r=0}^k (-1)^r \frac{(1+\alpha)_{n+r-k} (1+\beta)_k}{(r)! (k-r)! (n-k)!} x^{k-r} y^{n-k} \\ = \sum_{n=0}^{\infty} (-t)^n \sum_{k=0}^n \frac{(1+\beta)_k (1+\alpha)_{n-k}}{(n-k)!} y^{n-k} \sum_{r=0}^k (-1)^r \frac{(1+\alpha+n-k)_r}{(r)! (k-r)!} x^{k-r} \\ = \sum_{n=0}^{\infty} (-t)^n \sum_{k=0}^n \frac{(1+\beta)_k (1+\alpha)_{n-k}}{(n-k)!} y^{n-k} A_k^{(\alpha+n-k)}(x).$$

Thus (1.12) is equivalent to

$$(1.13) \quad \sum_{r=0}^n \frac{(r)!}{(n-r)!} (-1)^r (xy)^{n-r} A_n^{(\alpha)}(x) A_n^{(\beta)}(y) \\ = \sum_{k=0}^n \frac{(1+\beta)_k (1+\alpha)_{n-k}}{(n-k)!} y^{n-k} A_k^{(\alpha+n-k)}(x).$$

2. Making use of the relation [7, p. 45]

$$L_n^{(\alpha+n+1)}(-x) = A_n^{(\alpha)}(x)$$

and our formula (1.8), we get

$$(2.1) \quad (1 - D)^{\alpha+n}(-x)^n = (n)! L_n^{(\alpha)}(x),$$

which yields the following interesting result

$$(2.2) \quad L_n^{(\alpha+\beta)}(x) = (1 - D)^\beta L_n^{(\alpha)}(x).$$

Now consider

$$\begin{aligned} L_n^{(\alpha+\beta+1)}(x) &= (1 - D)^{\beta+1} L_n^{(\alpha)}(x) = \left[1 + \frac{D}{1 - D} \right]^{-\beta-1} L_n^{(\alpha)}(x) \\ &= \sum_{r=0}^n (-1)^r \frac{(\beta+1)_r}{(r)!} (1 - D)^{-r} D^r L_n^{(\alpha)}(x), \end{aligned}$$

and it follows that

$$(2.3) \quad L_n^{(\alpha+\beta+1)}(x) = \sum_{r=0}^n \frac{(\beta+1)_r}{(r)!} L_{n-r}^{(\alpha)}(x).$$

Formula (2.3) was proved earlier in a different way by Al-Salam [1, p. 131], and our proof differs markedly with that of Rainville [5, p. 209].

Next let us consider the expression,

$$(-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)]$$

which by the usual shift rule gives us

$$(-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)] = (1 - D)^n L_n^{(\alpha)}(x).$$

On making use of the relation (2.2) we obtain

$$(-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)] = L_n^{(\alpha+n)}(x),$$

which may be put in the form

$$(2.4) \quad R_n(1 + \alpha, x) = (-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)],$$

where $R_n(a, x)$ is the pseudo Laguerre set defined by Shively [5, p. 298] as

$$R_n(a, x) = \frac{(a)_{2n}}{(n)! (a)_n} {}_1F_1(-n; a + n; x).$$

The formula (2.4) has been proved recently by Khandekar in a different way (see [4], p. 2).

Further, from (2.1) we have

$$\frac{(-x)^n}{(n)!} = (1 - D)^{-\alpha-n} L_n^{(\alpha)}(x) = \sum_{r=0}^n \frac{(\alpha + n)_r}{(r)!} D^r L_n^{(\alpha)}(x)$$

which gives

$$(2.5) \quad \frac{(-x)^n}{(n)!} = \sum_{r=0}^n (-1)^r \frac{(\alpha + n)_r}{(r)!} L_{n-r}^{(\alpha+r)}(x).$$

Next consider the identity

$$e^{-xt} = \sum_{n=0}^{\infty} (-x)^n \frac{t^n}{(n)!},$$

operate on both sides by $(1 - D)^\alpha$, make use of (2.1) and proceed as in the cases of (1.12) and (1.13). We then get the generating function

$$(2.6) \quad (1 + t)^\alpha e^{-xt} = \sum_{n=0}^{\infty} t^n L_n^{(\alpha-n)}(x),$$

due to Erdélyi, and the known formula [2, p. 151]

$$(2.7) \quad \begin{aligned} & \sum_{n=0}^{\infty} (n)! t^n L_n^{(\alpha-n)}(x) L_n^{(\beta-n)}(y) \\ &= \begin{cases} e^{xyt} (1 - yt)^{\alpha-\beta} t^\beta (\beta)! L_\beta^{(\alpha-\beta)} \left(-\frac{(1 - xt)(1 - yt)}{t} \right) \\ e^{xyt} (1 - xt)^{\beta-\alpha} t^\alpha (\alpha)! L_\alpha^{(\beta-\alpha)} \left(-\frac{(1 - xt)(1 - yt)}{t} \right). \end{cases} \end{aligned}$$

ACKNOWLEDGEMENT

I am grateful to Dr. H. M. Srivastava of Jodhpur University for his kind guidance and invaluable help during the preparation of this paper.

REFERENCES

1. W. A. AL-SALAM, *Operational representations for the Laguerre and other polynomials*, Duke Math. J., **31** (1964), 127-142.
2. H. BUCHHOLZ, *Die konfluente hypergeometrische funktionen*, Berlin (1953).
3. L. CARLITZ, *A note on the Laguerre polynomials*, Michigan Math. J., **7** (1960), 219-223.
4. P. R. KHANDEKAR, *Rodrigues' formula for some generalized hypergeometric polynomials*, Math. Stu. of the Indian Math. Soc. (1) **34** (1966), 1-3.
5. E. D. RAINVILLE, *Special Functions*, Macmillan, New York (1960).
6. R. P. SINGH, *Some polynomials of Srivastava related to the Laguerre polynomials*, Math. Japonicae, **9** (1964), 5-9.
7. K. N. SRIVASTAVA, *Some polynomials related to the Laguerre polynomials*, J. Indian Math. Soc. (2), **28** (1964), 43-50.
8. H. M. SRIVASTAVA, *On Bessel, Jacobi and Laguerre polynomials*, Rend. Semin. Mat. Univ. Padova, **35** (1965), 424-432.

Manoscritto pervenuto in redazione il 1^o settembre 1966.